

Cell wall composition affects Cd²⁺ accumulation and intracellular thiol peptides in marine red algae

Virginia García-Ríos^a, Yolanda Freile-Pelegrín^a, Daniel Robledo^a, David Mendoza-Cózatl^b,
Rafael Moreno-Sánchez^b, Gerardo Gold-Bouchot^{a,*}

^a Departamento de Recursos del Mar, CINVESTAV-IPN, Unidad Mérida, Apdo. Postal 73-Cordemex, Mérida, Yucatán 97310, México

^b Departamento de Bioquímica, Instituto Nacional de Cardiología, México D.F. 14080, México

Received 23 September 2006; received in revised form 3 November 2006; accepted 4 November 2006

Abstract

Two red macroalgae species, *Gracilaria cornea* and *Chondrophycus poiteaui*, were evaluated for their intra and extracellular Cd²⁺ accumulation capacity, photosynthetic response and thiol peptide production. Algae were exposed for 3 and 7 days to 0.1 and 1 µg CdCl₂ ml⁻¹ (0.89 and 8.9 µM). Intracellular accumulation of Cd²⁺ by *G. cornea* was relatively low, only comprising 20% of total metal (intracellular + extracellular). In contrast, *C. poiteaui* accumulated intracellularly close to 100% of total Cd²⁺. In both species, metal uptake was dependent on the external Cd²⁺ concentration, metal exposure time and cell wall composition. In response to Cd²⁺ exposure, low amounts of thiol peptides were synthesized and the major difference between *G. cornea* and *C. poiteaui* was in the cell wall composition. The absence of insoluble polysaccharides in the cell wall of *C. poiteaui* suggested that this insoluble fraction might be involved in establishing an efficient barrier for the intracellular accumulation of Cd²⁺. This is the first study in which the cell wall composition, its influence on Cd²⁺ accumulation and intracellular responses in red macroalgae are evaluated. © 2006 Elsevier B.V. All rights reserved.

Keywords: Red macroalgae; Cadmium accumulation; Photosynthesis; Phytochelatins; Polysaccharides; Cell wall composition

1. Introduction

Toxic effects of metals on photosynthetic organisms include growth inhibition, photosynthetic damage and mortality, in addition to binding and blockade of essential groups in biomolecules, genotoxic damage and oxidative stress (Webster and Gadd, 1996; Mendoza-Cózatl et al., 2002; Okamoto et al., 2001). Photosynthetic organisms have developed intra and extra cellular mechanisms for metal detoxification (Clemens, 2001; Hall, 2002) and some of them, such as formation of phytochelatins (PCs) may be used as biomarkers (Ferrat et al., 2003).

The cell wall is the first barrier for metal ion uptake in higher plants and algae (Macfie and Welbourn, 2000). This extra cellular barrier has the capacity to bind metal ions in negatively charged sites. In the cell wall of marine algae, large amounts of sulfated polysaccharides are found, in contrast to freshwater and land plants (Craigie, 1990). Particularly,

red algae polysaccharides have a larger capacity to bind and accumulate metals ions when compared to brown (Phaeophyta) or green (Chlorophyta) algae (Stengel et al., 2004).

One of the most widespread intracellular mechanisms for Cd²⁺ detoxification in photosynthetic organisms involves the chelation of the metal ions by peptides or proteins such as glutathione (GSH), metallothioneins (MTs) and phytochelatins (PCs) (Kotrba et al., 1999; Cobbett, 2000; Mendoza-Cózatl et al., 2005). Resistance to metals in higher plants and some green microalgae has been well documented, being attributed in most cases to the synthesis of PCs (Pawlik-Skowrońska, 2000, 2003; Torricelli et al., 2004; Kobayashi et al., 2006). However, in macroalgae, there are only a few studies on intracellular responses to metals: PC synthesis (Hu and Wu, 1998; Malea et al., 2006), conjugation with GSH (Cairrão et al., 2004) or MT identification (Morris et al., 1999).

In the present study, we analyzed the intracellular and extracellular responses to Cd²⁺ exposure in two red macroalgae with different cell wall compositions, *Gracilaria cornea* and *Chondrophycus poiteaui*. Cell wall composition and thiol peptide synthesis were also analyzed in relation to the Cd²⁺ accumulation mechanisms in both species.

* Corresponding author. Tel.: +52 999 1242161; fax: +52 999 9812334.
E-mail address: ggold@mda.cinvestav.mx (G. Gold-Bouchot).

2. Materials and methods

2.1. Cd^{2+} exposure

Unialgal cultures of *G. cornea* and *C. poiteaui* were established in a culture chamber under laboratory conditions at 23 °C, 80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and 12-h light:12-h dark photoperiod with natural sea water filtered through 1.2 μm Millipore filters and 33‰ salinity. Macroalgae were grown in 1000 ml cylinders for 3–7 days in the presence of 0.1 or 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$ in natural sea water, corresponding to 0.89 and 8.9 μM , respectively. Control cultures with no Cd^{2+} added to the culture media were also carried out.

2.2. Total and intracellular Cd^{2+} quantification

After 3 and 7 days at the indicated added CdCl_2 concentrations, algae were collected and rinsed with distilled water. Samples for the determination of intracellular Cd^{2+} were rinsed twice for 10 min with water containing 5 mM EDTA (pH 8) to remove metals adsorbed by the cell wall. Thereafter, the samples were freeze-dried for determination of total (water-rinsed) and intracellular accumulated Cd^{2+} (EDTA-rinsed) and dry weight. Metal content in the macroalgae was determined with a Perkin-Elmer SIMAA 6100 graphite furnace atomic absorption spectrophotometer. Samples of an *Ulva lactuca* certified reference material (BCR Reference No. 279, Commission of the European Communities) were also analyzed with every batch of samples, with an average 98% recovery with respect to the Cd^{2+} certified concentration.

2.3. Cell wall polysaccharides analysis

Macroalgal tissue was freeze-dried, pulverized, and incubated in 0.1 M sodium acetate buffer (pH 6), 5 mM EDTA, 5 mM cysteine and 0.5 mg papain. Sulfated polysaccharides were precipitated with 10% cetylpyridinium chloride (adapted from Farias et al., 2000). Soluble sulfated polysaccharides were separated from the insoluble fraction after dissolving with distilled water and further filtrating through a sintered-glass plate (4.0–5.5 μm pore size). Fractions obtained were freeze-dried and weighed (Melo et al., 2002). The sulfate content of polysaccharides was determined by a turbidimetric method (Jackson and McCandless, 1978).

2.4. Photosynthesis light-response curves

Photosynthetic responses in macroalgae exposed to Cd^{2+} were measured. Photosynthesis measurements were performed in a DW/2 chamber at 21 °C using a Clark-type oxygen electrode and the Oxygraph program (Hansatech, King's Lynn, UK). After equilibration in darkness for 3 min, respiration rate was measured as O_2 consumption. Oxygen evolution was measured during periods of 2 min at light intensities, varying from 50 to 1400 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Maximal net photosynthetic activity (P_{max}) was calculated for each individual sample from their photosynthetic light-response curves (Henley, 1993).

2.5. HPLC analysis of intracellular thiols

Algal samples were homogenized with a mortar and pestle in liquid nitrogen and 1.5 ml of 50 mM Tris-HCl buffer (pH 8), containing 1 mM EGTA. The homogenate was centrifuged for 20 min at 4 °C and $50,000 \times g$. The supernatant was kept under reduced conditions by adding DTT (2 mM final concentration), mixed and centrifuged again for 20 min at 4 °C and $50,000 \times g$. Proteins were precipitated with perchloric acid (3% final concentration), vortexed and centrifuged as described before. The clear supernatant was filtered (using 0.22 μm pore diameter cut-off filters, Millex Millipore) and injected into a Spherisorb column (C18, reverse phase, 5 μm ODS 2, 4.6 mm \times 150 mm). The column was equilibrated with trifluoroacetic acid (0.1%) and peptides were eluted in a 0–20% acetonitrile linear gradient at a flow rate of 1 ml min⁻¹. Post-column derivatization with Ellman's reagent was used to detect thiol-containing compounds at 412 nm, as described elsewhere (Rausser, 1991). A synthesized PC_2 standard was used to identify PC-related compounds.

2.6. Data analysis

Statistical analysis of data was carried out with STATISTICA software (Version 6, StatSoft, Inc., 2003). To determine significant differences in treatments, a non-parametric Kruskal–Wallis analysis of variance was used. The significant level was established at $\alpha=0.05$. Data is presented as the median of four independent replicates \pm one interquartile range.

3. Results

3.1. Total and intracellular Cd^{2+} quantification

Cd^{2+} uptake in *G. cornea* and *C. poiteaui* was proportional to the CdCl_2 concentration added to the culture medium and to the time of exposure (Table 1). Both *G. cornea* and *C. poiteaui* exhibited a high uptake when exposed for 7 days to 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$. *C. poiteaui* accumulated four-fold more Cd^{2+} than *G. cornea* (Table 1). However, most of the metal found in *G. cornea* was allocated extracellularly (80%). Interestingly, *C. poiteaui* accumulated most of the Cd^{2+} inside the cell (near 100% of total Cd^{2+}). Both *G. cornea* and *C. poiteaui* exhibited a higher uptake when exposed to 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$ and after 7 days exposure.

3.2. Cell wall polysaccharides analysis

Content of soluble and insoluble polysaccharide in *G. cornea* and *C. poiteaui* is shown in Fig. 1. For *G. cornea*, polysaccharides constituted $35 \pm 7\%$ of total dry wt, 50% of which were soluble and 50% insoluble. In this species, no significant changes in the soluble/insoluble ratio were found after 3 or 7 days of exposure. In *C. poiteaui*, a lower polysaccharide content was obtained ($23 \pm 3\%$ of total dry wt) when compared to *G. cornea*, but a marked difference between these species was the negligible content of the insoluble polysaccharide fraction in *C. poiteaui* (Fig. 1).

Table 1

Total and intracellular Cd²⁺ accumulation ($\mu\text{g g}^{-1}$ dry wt) and total thiol peptide concentrations (nmol SH mg⁻¹ dry wt) in *G. cornea* and *C. poiteaui*

Exposure conditions	Total Cd ²⁺	Intracellular Cd ²⁺	Total thiol	Thiol peptides		
				PC ₂	X ₁	
<i>G. cornea</i>						
Control						
3 days	0.4 ± 0.4	0.5 ± 0.5	0.7 ± 0.2			
7 days	0.4 ± 0.1	0.14 ± 0.05	0.8 ± 0.5			
0.1 µg Cd ml ⁻¹						
3 days	3.4 ± 2	1.4 ± 0.7	0.7 ± 0.4			
7 days	6 ± 1	1.1 ± 0.7	0.81 ± 0.17			
1 µg Cd ml ⁻¹						
3 days	14.9 ± 2.7 ^a	3 ± 1.6 ^a	0.8 ± 0.3	0.02 ± 0.03		
7 days	22.9 ± 2.7 ^a	5 ± 1.8	1 ± 0.7	0.09 ± 0.10	0.01 ± 0.01	
Kruskal–Wallis test						
3 days	<i>H</i> = 9.846 <i>p</i> = 0.007	<i>H</i> = 9.269 <i>p</i> = 0.020	<i>H</i> = 0.356 <i>p</i> = 0.837			
7 days	<i>H</i> = 9.846 <i>p</i> = 0.007	<i>H</i> = 4.718 <i>p</i> = 0.094	<i>H</i> = 0.089 <i>p</i> = 0.956			
Exposure conditions	Total Cd ²⁺	Intracellular Cd ²⁺	Total thiol	Thiol peptides		
				PC ₂	X ₁	X ₂
<i>C. poiteaui</i>						
Control						
3 days	0.19 ± 0.03	0.24 ± 0.10	1.0 ± 0.9			
7 days	0.18 ± 0.03	0.33 ± 0.04	0.47 ± 0.5			
0.1 µg Cd ml ⁻¹						
3 days	14 ± 4.6	14 ± 3.1	1.0 ± 0.5	0.01 ± 0.00	0.01 ± 0.01	0.02 ± 0.01
7 days	21 ± 2	23.5 ± 1.4	0.64 ± 1.44	0.01 ± 0.01	0.01 ± 0.02	0.02 ± 0.01
1 µg Cd ml ⁻¹						
3 days	78.6 ± 23 ^a	72 ± 25 ^a	1.43 ± 0.61	0.03 ± 0.01 ^a	0.02 ± 0.01	0.02 ± 0.01 ^a
7 days	93.4 ± 11 ^a	64 ± 12 ^a	1.07 ± 0.23	0.03 ± 0.03	0.03 ± 0.02 ^a	0.02 ± 0.01
Kruskal–Wallis test						
3 days	<i>H</i> = 7.854 <i>p</i> = 0.020	<i>H</i> = 8.910 <i>p</i> = 0.012	<i>H</i> = 0.694 <i>p</i> = 0.707	<i>H</i> = 6.562 <i>p</i> = 0.038	<i>H</i> = 5.833 <i>p</i> = 0.054	<i>H</i> = 6.562 <i>p</i> = 0.038
7 days	<i>H</i> = 9.846 <i>p</i> = 0.007	<i>H</i> = 9.846 <i>p</i> = 0.007	<i>H</i> = 2.472 <i>p</i> = 0.290	<i>H</i> = 5.426 <i>p</i> = 0.066	<i>H</i> = 6.325 <i>p</i> = 0.042	<i>H</i> = 4.076 <i>p</i> = 0.130

Median ± quartile range; $n=4$.^a Significant difference with respect to control.

In the control group (no Cd²⁺ exposure), polysaccharide sulfate content was 2.6 times higher in *C. poiteaui* ($20 \pm 5 \mu\text{g sulfate mg}^{-1}$ polysaccharide) than in *G. cornea* ($7.7 \pm 2 \mu\text{g sulfate mg}^{-1}$ polysaccharide; Fig. 2). Soluble polysaccharides in *G. cornea* had a significant increase in sulfate content after 3 days exposure to $0.1 \mu\text{g CdCl}_2 \text{ ml}^{-1}$; after 7 days exposure, sulfate content returned back to basal levels (Fig. 2). *C. poiteaui* did not show any significant changes in polysaccharides sulfate content after 3 or 7 days of Cd²⁺ exposure.

3.3. Photosynthetic parameters

Maximal photosynthesis (P_{max}) values in *G. cornea* and *C. poiteaui* are shown in Fig. 3. No significant differences in P_{max} were observed in *G. cornea* between treatments after 3 and 7 days of CdCl₂ exposure (Fig. 3). On the other hand, in *C.*

poiteaui, exposure to $1 \mu\text{g CdCl}_2 \text{ ml}^{-1}$ caused a considerable decrease of P_{max} after 3 days (Fig. 3). After 7 days, P_{max} values returned to control levels without significant differences with the control group.

3.4. Intracellular thiols

After 3 days exposure of *G. cornea* to $1 \mu\text{g CdCl}_2 \text{ ml}^{-1}$, small production of thiol-peptides with similar HPLC retention time to phytochelatin-2 (PC₂) was detected; after 7 days exposure, a second thiol-peptide with a longer retention time was also detected (X₁), suggesting the occurrence of larger PCs in *G. cornea* (Fig. 4). However, no significant difference in total thiol content in *G. cornea* was observed (Table 1).

In *C. poiteaui*, after 3 days exposure to 0.1 or $1 \mu\text{g CdCl}_2 \text{ ml}^{-1}$, three thiol peptides were found (Fig. 5). The




Fig. 1. Total, soluble and insoluble polysaccharides content in *G. cornea* and absence of insoluble polysaccharides in *C. poiteaui* exposed to 0, 0.1 and 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$ in the seawater. Median \pm one interquartile range ($n=4$). *Significant differences with respect to control in *C. poiteaui* ($H=7.848$; $p=0.020$).

first compound exhibited a retention time similar to PC_2 ; two other compounds with higher retention times, possibly larger PCs, were also detected and quantified (X_1 and X_2). There was a significant increase in the content of these Cd^{2+} induced

thiol-peptides after 3 and 7 days of Cd^{2+} exposure (Table 1). Although *C. poiteaui* exhibited a higher number of different thiol-compounds than *G. cornea*, the content of thiol peptides did not show significant changes between treatments.




Fig. 2. Increase and recovery of sulfate content in polysaccharides of *G. cornea* and *C. poiteaui* exposed to 0, 0.1 and 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$ in the seawater. Median \pm one interquartile range ($n=4$). *Significant differences with respect to control in *G. cornea* ($H=6.727$; $p=0.035$).




Fig. 3. Maximal photosynthesis (P_{\max}) of: (A) *G. cornea* and (B) *C. poiteaui* after 3 days (grey bars) and 7 days (stripped bars) exposure to 0, 0.1 and 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$. Median \pm one interquartile range ($n=4$). * Significant differences with respect to control of *C. poiteaui* ($H=7.385$; $p=0.025$).

4. Discussion

4.1. Cd^{2+} uptake

The different capacity to accumulate Cd^{2+} by *G. cornea* and *C. poiteaui*, extra and intracellularly indicated that these species have different protection mechanisms developed against metal exposure. Apparently, the complex polysaccharides of the *G. cornea* cell wall were an efficient barrier to avoid metal toxicity, thus leading to a low intracellular metal accumulation. *G. cornea* showed a limited capacity to accumulate Cd^{2+} intracellularly, while extracellular binding correlated with the CdCl_2 concentration in the medium. In contrast, *C. poiteaui* predominantly

accumulated Cd^{2+} intracellularly, and this process correlated with the Cd^{2+} concentration and time of exposure, although it is worth noting that the metal accumulation– CdCl_2 concentration relationship was not linear. Moreover, *C. poiteaui* showed a marked cellular response to the metal exposure regarding photosynthesis and production of thiol peptides. On this regard, Wang and Dei (1999) indicated that metal uptake in *Gracilaria blodgettii* was higher at lower metal concentration in the medium.

Metal uptake is determined by the physical and chemical characteristics of the metal ion, and by the biological characteristics of the exposed organisms (Ariza et al., 1999). Maximal total Cd^{2+} accumulation in *C. poiteaui* ($93.5 \mu\text{g g}^{-1}$ dry wt) and *G. cornea* ($22.9 \mu\text{g g}^{-1}$ dry wt) was far below data obtained in other




Fig. 4. Representative chromatograms of thiol peptides of *G. cornea* exposed to 0, 0.1 and 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$ in the seawater.




Fig. 5. Representative chromatograms of thiol peptides of *C. poiteaui* exposed to 0, 0.1 and 1 $\mu\text{g CdCl}_2 \text{ ml}^{-1}$ in the seawater.

studies with microalgae and brown macroalgae. For example, in *Chlamydomonas reinhardtii*, Cd^{2+} accumulation (Kobayashi et al., 2006) was 500 and 1800-fold higher than that obtained by *C. poiteaui* and *G. cornea*, respectively, at similar added Cd^{2+} concentrations in the medium. In *Fucus serratus*, copper accumulation reached $1000 \mu\text{g g}^{-1}$ dry wt when exposed to $0.8 \mu\text{M}$ copper (Nielsen et al., 2003).

4.2. Cell wall polysaccharide and sulfur content

No evident changes in soluble or insoluble polysaccharides content were found after Cd^{2+} exposure in *G. cornea*. In contrast, soluble polysaccharides in *C. poiteaui* decreased after 7 days of exposure to both Cd^{2+} concentrations used. This response may be explained by intracellular metal accumulation and the associated perturbation of cellular functions. In *G. cornea*, its high content of cell wall insoluble polysaccharides might have facilitated the extracellular binding of Cd^{2+} and hence intracellular synthesis of the cell wall and other intracellular processes remained protected. On the other hand, absence of cell wall insoluble polysaccharides in *C. poiteaui* correlated with a negligible external Cd^{2+} binding and significant intracellular accumulation of Cd^{2+} , thus inducing alteration in cell wall synthesis and other intracellular functions.

It has not been clearly established whether cell wall composition determines metal accumulation, or it plays a protective role in microalgae. On this regard, increased polysaccharide production and low resistance, or moderate production of polysaccharides and very efficient exclusion mechanisms have been observed in phytoplankton by Pistocchi et al. (2000). In two *Chlamydomonas reinhardtii* strains, variation in the composition of the cell wall did not correlate with the different tolerance

to metal toxicity (Macfie and Welbourn, 2000). A comparison of Cd^{2+} bio-absorption capacity among different macroalgae showed that alginic acid in the cell wall of brown macroalgae was more efficient to bind Cd^{2+} than polysaccharides found in red or green macroalgae (Hashim and Chu, 2004). In contrast, Stengel et al. (2004) showed that red macroalgae were more efficient to accumulate zinc than the other algal groups. The lack of insoluble polysaccharides in *C. poiteaui* (see Fig. 1) may be related to the observed high intracellular accumulation of Cd^{2+} in this species; however, more studies are necessary to determine the role of cell wall insoluble polysaccharides of red algae in Cd^{2+} accumulation.

Although sulfur containing compounds are responsible for Cd^{2+} chelation in brown algae (Raize et al., 2004), sulfate content was not related to the extracellular Cd^{2+} binding capacity between the two species used in the present study. *C. poiteaui* exhibited higher polysaccharide sulfate content; however, it did not retain Cd^{2+} in the cell wall. On the contrary, sulfate content in the soluble fraction of *G. cornea* polysaccharides increased after 3 days of Cd^{2+} exposure, returned to basal concentrations after 7 days. Seasonal sulfate content changes occurred in *G. cornea* (Freile-Pelegrin and Robledo, 1997). Increased sulfate content could be related to a general cell response to stress of multiple origins and not to a specific protective response to Cd^{2+} exposure.

4.3. Cellular response to Cd^{2+}

Accumulation of Cd^{2+} in chloroplasts and its effects on photosynthesis have been documented in microalgae (Okamoto et al., 2001; Mendoza-Cózatl et al., 2002). In *Gonyaulax polyedra*, acute exposure to Cd^{2+} generated oxidative stress in the chloro-

plasts, whereas under chronic exposure the antioxidant system was able to protect (Okamoto et al., 2001). The decrease of P_{\max} in *C. poiteaui* after 3 days of exposure and its recover after 7 days may reflect the activation of the antioxidant system. Analysis of the degree of oxidative stress and antioxidant enzymes activity are necessary to determine the effect of metal exposure on photosynthetic responses. On this issue, P_{\max} recovery in *C. poiteaui* after 7 days of Cd^{2+} exposure may be related to metal detoxification mechanisms. This suggestion is consistent with the synthesis of thiol-peptides beginning after 3 days and increasing after 7 days of Cd^{2+} exposure. As PC synthesis was induced at low, almost negligible levels, it may be proposed that PC synthesis is not a significant part of the Cd^{2+} detoxification mechanisms in this species. Higher induction of PC synthesis in microalgae exposed to Cd^{2+} concentrations similar to those used in this study has been previously reported (Mendoza-Cózatl et al., 2005; Kobayashi et al., 2006), as well as after exposure to other metals such as zinc and lead (Pawlik-Skowrońska, 2000, 2003; Tsuji et al., 2003).

GSH is the major reservoir (up to 90%) of non-proteic sulfur in the cell, and is the substrate for PC synthesis. Consumption of GSH by PC synthesis induces an increase in the rate of GSH synthesis to restore basal levels. For that reason, depending on the GSH demand rate, GSH levels may increase or decrease after Cd^{2+} exposure (reviewed by Mendoza-Cózatl et al., 2005). In our results, total thiol content did not change with Cd^{2+} exposure; in other words, there was no significant PC synthesis. In the two marine macroalgae analyzed in the present study, Cd^{2+} exposure caused a low PC synthesis without GSH depletion.

No significant synthesis of PCs in *G. cornea* and *C. poiteaui* was achieved at the low intracellular Cd^{2+} concentration accumulated. In higher plants and some yeast, the number of Cd^{2+} ions bound by the thiol group is variable (<1–4), and depends on the thiol-compound chemical nature (i.e. GSH, PCs, etc.) (Zenk, 1996). In the present study, the thiol group content/intracellular Cd^{2+} ratio in *C. poiteaui* was 2:1, which suffices to chelate all intracellular Cd^{2+} without PC synthesis, although Cd^{2+} should not be fully inactivated as a thiol/ Cd^{2+} ratio of 4 is required. This response suggests that red macroalgae have detoxification mechanisms that involve active plasma membrane metal uptake, and an efficient intracellular metal binding by compounds other than PCs, perhaps smaller thiol-compounds. A low PC induction has also been observed in the green macroalgae *Enteromorpha* spp. (Malea et al., 2006). Other possible intracellular mechanisms different to PCs may also be responsible for metal detoxification in some marine macroalgae. Non-protein reduced sulfur like cysteine and glutathione might be the major thiol compounds involved in the detoxification of all intracellular Cd^{2+} in this macroalgae whereas PCs may play only a minor role.

5. Conclusions

Soluble and insoluble polysaccharides present in the cell wall of *G. cornea* bound most of the Cd^{2+} extracellularly. The absence of insoluble polysaccharides in the cell wall of *C. poiteaui* was the major difference when compared to the *G. cornea* cell wall. In *C. poiteaui*, most of the Cd^{2+} was accumulated intracellularly,

which however did not induce a significant PC synthesis and therefore other detoxification mechanisms may be involved. The recovery of the photosynthetic performance of *C. poiteaui* after 7 days of Cd^{2+} exposure suggested the activation of intracellular detoxification mechanisms. However, more studies on oxidative stress and antioxidant enzymes activity are required to determine the effect of metal exposure on these physiological responses.

Acknowledgements

The present work was partially supported by grants 43811-Q from CONACYT-México and 2002-C01-1057 from CONACYT-SAGARPA. We appreciate C. Chavez, M. Zaldivar and V. Ceja for their contribution in the polysaccharides, photosynthesis and metal analysis. We thank deeply Dr. Peter Chapman for his valuable comments.

References

- Ariza, M.E., Bijur, G.N., Williams, M.V., 1999. Environmental Metal Pollutants, Reactive Oxygen Intermediaries and Genotoxicity. Kluwer Academic Publishers, Boston, USA.
- Cairrão, E., Couderchet, M., Soares, A.M.V.M., Guilhermino, L., 2004. Glutathione-S-transferase activity of *Fucus* spp. as a biomarker of environmental contamination. Aquat. Toxicol. 70, 277–286.
- Clemens, S., 2001. Molecular mechanisms of plant metal tolerance and homeostasis. Planta 212, 475–486.
- Cobbett, C.S., 2000. Phytochelatin and their roles in heavy metal detoxification. Plant Physiol. 123, 825–832.
- Craigie, J.S., 1990. Cell walls. In: Kathleen, M.C., Robert, G.S. (Eds.), Biology of the Red Algae. Cambridge University Press, pp. 221–258.
- Farias, W.R.L., Valente, A.P., Pereira, M.S., Mourao, P.A.S., 2000. Structure and anticoagulant activity of sulfated galactans. J. Biol. Chem. 275, 29299–29307.
- Ferrat, L., Pergent-Martini, C., Roméo, M., 2003. Assessment of the use of biomarkers in aquatic plants for the evaluation of environmental quality: application to seagrasses. Aquat. Toxicol. 65, 187–204.
- Freile-Pelegrin, Y., Robledo, D., 1997. Effects of season on the agar content and chemical characteristics of *Gracilaria cornea* from Yucatan, Mexico. Botanica Marina 40, 285–290.
- Hall, J.L., 2002. Cellular mechanisms for heavy metal detoxification and tolerance. J. Exp. Bot. 53, 1–11.
- Hashim, M.A., Chu, K.H., 2004. Biosorption of cadmium by brown, green, and red seaweeds. Chem. Eng. J. 97, 249–255.
- Henley, W.J., 1993. Measurement and interpretation of photosynthetic light response curves in algae in the context of photoinhibition and diel changes. J. Phycol. 29, 729–739.
- Hu, S., Wu, M., 1998. Cadmium sequestration in the marine macroalga *Kappaphycus alvarezii*. Mol. Mar. Biol. Biotechnol. 7, 97–104.
- Jackson, S.G., McCandless, E.L., 1978. Simple, rapid, turbidometric determination of inorganic sulfate and/or protein. Anal. Biochem. 90, 802–808.
- Kobayashi, I., Fujiwara, S., Saegusa, H., Inoue, M., Matsumoto, H., Tsuzuki, M., 2006. Relief of arsenate toxicity by Cd-stimulated phytochelatin synthesis in the green alga *Chlamydomonas reinhardtii*. Mar. Biotechnol. 8, 94–101.
- Kotrba, P., Macek, T., Ruml, T., 1999. Heavy metal-binding peptides and proteins in plants: a review. Collect. Czech. Chem. Commun. 64, 1057–1086.
- Macfie, S.M., Welbourn, P.M., 2000. The cell wall as a barrier to uptake of metal ions in the unicellular green alga *Chlamydomonas reinhardtii* (Chlorophyceae). Arch. Environ. Contam. Toxicol. 39, 413–419.
- Malea, P., Rijstenbil, J.W., Haritonidis, S., 2006. Effects of cadmium, zinc and nitrogen status on non-protein thiols in the macroalgae *Enteromorpha* spp. from the Scheldt Estuary (SW Netherlands, Belgium) and Thermaikos Gulf (N Aegean Sea, Greece). Mar. Environ. Res. 62, 45–60.

- Melo, M.R.S., Feitosa, J.P.A., Freitas, A.L.P., De Paula, R.C.M., 2002. Isolation and characterization of soluble sulfated polysaccharide from the red seaweed *Gracilaria cornea*. Carbohydr. Polym. 49, 491–498.
- Mendoza-Cózatl, D., Devars, S., Loza-Tavera, H., Moreno-Sánchez, R., 2002. Cadmium accumulation in the chloroplast of *Euglena gracilis*. Physiol. Plant. 115, 276–283.
- Mendoza-Cózatl, D., Loza-Tavera, H., Hernández-Navarro, A., Moreno-Sánchez, R., 2005. Sulfur assimilation and glutathione metabolism under cadmium stress in yeast, protists and plants. FEMS Microbiol. Rev. 29, 653–671.
- Morris, C.A., Nicolaus, B., Sampson, V., Harwood, J.L., Kille, P., 1999. Identification and characterization of a recombinant metallothionein protein from a marine alga, *Fucus vesiculosus*. Biochem. J. 338, 553–560.
- Nielsen, H.D., Brownlee, C., Coelho, S.M., Brown, M.T., 2003. Inter-population differences in inherited copper tolerance involve photosynthetic adaptation and exclusion mechanisms in *Fucus serratus*. New Phytol. 160, 157–165.
- Okamoto, O.K., Pinto, E., Latorre, L.R., Bechara, E.J.H., Colepicolo, P., 2001. Antioxidant modulation in response to metal-induced oxidative stress in algal chloroplasts. Arch. Environ. Contam. Toxicol. 40, 18–24.
- Pawlik-Skowrońska, B., 2000. Relationships between acid-soluble thiol peptides and accumulated Pb in the green alga *Stichococcus bacillaris*. Aquat. Toxicol. 50, 221–230.
- Pawlik-Skowrońska, B., 2003. When adapted to high zinc concentrations the periphytic green alga *Stigeoclonium tenue* produces high amounts of novel phytochelatin-related peptides. Aquat. Toxicol. 62, 155–163.
- Pistocchi, R., Mormile, A.M., Guerrini, F., Isani, G., Boni, L., 2000. Increased production of extra- and intracellular metal-ligands in phytoplankton exposed to copper and cadmium. J. Appl. Phycol. 12, 469–477.
- Raize, O., Argaman, Y., Yannai, S., 2004. Mechanisms of biosorption of different heavy metals by brown marine macroalgae. Biotechnol. Bioeng. 87, 451–458.
- Rausser, W.E., 1991. Cadmium-binding peptides from plants. Methods Enzymol. 205, 319–333.
- Stengel, D.B., Macken, A., Morrison, L., Morley, N., 2004. Zinc concentrations in marine macroalgae and lichen from western Ireland in relation to phylogenetic grouping, habitat and morphology. Mar. Poll. Bull. 48, 902–909.
- Torricelli, E., Gorbi, G., Pawlik-Skowrońska, B., di Toppi, L.S., Corradi, M.G., 2004. Cadmium tolerance, cysteine and thiol peptide levels in wild type and chromium-tolerant strains of *Scenedesmus acutus* (Chlorophyceae). Aquat. Toxicol. 68, 315–323.
- Tsuji, N., Hirayanagi, N., Iwabe, O., Namba, T., Tagawa, M., Miyamoto, S., Miyasaka, H., Takagi, M., Hirata, K., Miyamoto, K., 2003. Regulation of phytochelatin synthesis by zinc and cadmium in marine green alga, *Dunaliella tertiolecta*. Phytochemistry 62, 453–459.
- Wang, W.X., Dei, R.C.H., 1999. Kinetic measurements of metal accumulation in two marine macroalgae. Mar. Biol. 135, 11–23.
- Webster, E.A., Gadd, G.M., 1996. Perturbation of monovalent cation composition in *Ulva lactuca* by cadmium, copper and zinc. BioMetals 9, 51–56.
- Zenk, M.H., 1996. Heavy metal detoxification in higher plants—a review. Gene 179, 21–30.