

# Ocean Color Reveals Increased Blooms in Various Parts of the World

The magnitude of phytoplankton blooms has increased significantly in many areas of the world during the past 11 years, as shown in data from ocean color sensors on board satellites. These areas with increased blooms are likely to be environmentally stressed and undergoing undesirable environmental changes such as a higher frequency of harmful algal blooms and oxygen depletion in bottom layers of oceans, estuaries, and lakes. These changes can disrupt traditional fisheries and recreational use in many coastal areas.

An algal bloom is a rapid increase in the concentration of phytoplankton algae that occurs when conditions turn favorable for algal growth. A typical example of a bloom is the phytoplankton spring bloom. Algal blooms are natural phenomena that remove dissolved carbon and nutrients. In addition, they produce new biomass that supports higher trophic levels including fish and fisheries. However, the blooms also lead to excessive turbidity, oxygen depletion in the bottom layers, and the possible death of fish, benthic animals, and bottom vegetation.

Satellite measurement of spectral reflectance (ocean color) is a cost-effective method to monitor phytoplankton by its proxy, chlorophyll *a* concentration (the green pigment that is present in all algae, Chl-*a*). Understanding the effects of the increasing atmospheric carbon dioxide concentrations and higher surface temperatures on ocean biota is a major theme of NASA's Ocean Biology and Biogeochemistry Program [McClain *et al.*, 2006]. In September 2007, the NASA Sea-viewing Wide Field-of-view Sensor (SeaWiFS) completed 10 years

in orbit (and efforts to revise the sensor are still under way following an incident on 1 January 2008 that has prevented data downlink). SeaWiFS, together with its predecessor, the Japan Space Exploration Agency's Ocean Color and Temperature Scanner (OCTS), which operated for only 8 months prior to a power failure, provide approximately 11 years of high-quality global ocean color observations.

While measurements of Chl-*a* of individual blooms are variable (blooms are very patchy in both space and time) and satellite sensors miss some blooms due to persistent cloud cover, the monthly mean composite Chl-*a* is representative of the mean phytoplankton concentration. It is well known that satellite estimation of Chl-*a* in nearshore waters is complicated and that other optically active substances, such as detrital material, dissolved organic substances, and suspended sediments, can interfere in this process. In spite of these potential errors, the monthly composite Chl-*a* is a robust index of water quality and corresponds to the combined effects of Chl-*a* and other optically active substances.

We used global Chl-*a* data from OCTS and SeaWiFS processed to 9-kilometer pixel resolution, and we found the highest monthly Chl-*a* value for each year and each pixel. We then compiled time series of these annual maxima from 1997 to 2007. We interpret the time series of the annual maxima as a change in bloom magnitude. We used the Sen slope nonparametric estimate of the slope to detect trends and their significance in bloom magnitude. Estimates of the slope of bloom magnitude were also obtained with the linear least squares regression, but the Sen slope estimator is preferable due to its insensitivity to outliers. In Figure 1, areas of increased bloom magnitude are shown in red and those with decreased bloom magnitude are shown in blue. The global map of increased bloom magnitude is shown at <http://spg.ucsd.edu/blooms.png>, and a Google™ Earth version is shown at <http://spg.ucsd.edu/blooms.kmz>.

## Phytoplankton Blooms

During the past 11 years, bloom magnitudes have increased in extensive areas of eastern boundary upwelling currents along the Washington-Oregon-California coast of North America, the northern Peru coast off South America, sections of the coast of Namibia off Africa, and off the southwestern tip of India. Eastern boundary currents are characterized by upwelling of nutrient-rich waters that makes these areas very productive and important for fisheries. Increased blooms also are evident in enclosed and semienlosed basins such as the Baltic Sea, Lake Maracaibo in Venezuela, the Sea of Azov, and the northeastern area of the Caspian Sea. In the Baltic Sea, the Chl-*a* algorithm is known to have large errors and the bloom magnitude primarily reflects the abundance of surface-floating cyanobacteria [Kahru *et al.*, 2007].

Increased blooms are also observed in the northern Arabian Sea and the Gulf of




Fig. 1. Areas of increased (red) and decreased (blue) phytoplankton bloom magnitude. The trend and its significance are estimated using the Sen slope test at 95% confidence level. Individual time series of spatially averaged Chl-*a* (milligrams per meter cubed) are shown as insets for the 50-kilometer nearshore band (a) off the Washington-Oregon-Northern California coast, (b) off Northern Peru, (c) in the Sea of Azov, and (d) off southwest India.

Oman, as well as in outflow areas of big rivers such as Brazil's Amazon and Tocantins and Africa's Congo. In the Mississippi River outflow area, blooms have increased in the western part of the outflow and decreased in the eastern part. In East Asia, blooms have increased off Shanghai and along the west coast of Korea. In the Yellow Sea and the adjoining Bohai Sea, blooms have actually decreased, due possibly to increased turbidity. Blooms have also decreased in the outflow area of the Po River in the Adriatic Sea, likely as a result of efforts to control pollution. Significant changes (i.e., increases or decreases in bloom magnitude) over large areas are evident at high latitudes (the Barents and Bering seas, the Sea of Okhotsk, and so forth), but the reliability of these estimates is less certain due to persistent cloud cover and infrequent satellite measurements of these areas.

In summary, ocean color data from 1997 to 2007 show increased phytoplankton bloom magnitude in eastern boundary upwelling systems and in a number of areas with known eutrophication. Increased blooms in eastern boundary currents may be caused by increased upwelling, but we are unaware of any direct evidence supporting that. The increased blooms off Oregon are linked to the increase in the "dead zones" of oxygen-depleted water [Service, 2004, 2007]. It has been suggested [Goes *et al.*, 2005] that the intensification of the southwest monsoon due to global warming has increased the productivity in the Arabian Sea, but this has been disputed by Prakash and Ramesh [2007]. Figure 1 shows that the increase in bloom magnitude is observed in the relatively narrow region of the Gulf of Oman and in nearshore areas of the northern Arabian Sea. Some of the observed trend in bloom magnitude is attrib-

utable to the strong El Niño of 1997–1998 in the start of the time series. However, bloom magnitudes have increased in many areas even after 1997–1998. We hope that experts familiar with conditions in specific areas can interpret these patterns using local data

## Acknowledgments

SeaWiFS and OCTS data were made available by the NASA Ocean Color Processing Group. Support by the NASA Ocean Biogeochemistry and ECOHAB (Ecology and Oceanography of Harmful Algal Blooms) programs is acknowledged.

## References

- Goes, J., P. G. Thoppil, H. do R. Gomes, and J. T. Fasullo (2005), Warming of the Eurasian landmass is making the Arabian Sea more productive, *Science*, 308(5721), 545–547.
- Kahru, M., O. P. Savchuk, and R. Elmgren (2007), Satellite measurements of cyanobacterial bloom frequency in the Baltic Sea: Interannual and spatial variability, *Mar. Ecol. Prog. Ser.*, 343, 15–23, doi:10.3354/meps06943.
- McClain, C., S. Hooker, G. Feldman, and P. Bontempo (2006), Satellite data for ocean biology, biogeochemistry, and climate research, *Eos Trans. AGU*, 87(34), 337, 343.
- Prakash, S., and R. Ramesh (2007) Is the Arabian Sea getting more productive?, *Curr. Sci.*, 92(5), 667–671.
- Service, R. F. (2004), New dead zone off Oregon coast hints at sea change in currents, *Science*, 305(5687), 1099.
- Service, F. S. (2007), "Dead zone" reappears off Oregon coast, *ScienceNOW*, 3, 31 July. (Available at <http://sciencenow.sciencemag.org/cgi/content/full/2007/731/3>)

—MATI KAHRU and B. GREG MITCHELL, Scripps Institution of Oceanography, University of California, San Diego, La Jolla; E-mail: mkahru@ucsd.edu

# EOS

TRANSACTIONS  
AMERICAN GEOPHYSICAL UNION  
The Newspaper of the Earth and Space Sciences

## Editors

John W. Geissman, Wendy S. Gordon,  
Manuel Grande, Hassan Virji

## Editor in Chief

A. F. Spilhaus, Jr.

©2008 American Geophysical Union. Material in this issue may be photocopied by individual scientists for research or classroom use. Permission is also granted to use short quotes, figures, and tables for publication in scientific books and journals. For permission for any other uses, contact the AGU Publications Office.

*Eos*, Transactions, American Geophysical Union (ISSN 0096-3941) is published weekly by the American Geophysical Union, 2000 Florida Ave., NW, Washington, DC 20009 USA. Periodical Class postage paid at Washington, D.C., and at additional mailing offices. POSTMASTER: Send address changes to Member Service Center, 2000 Florida Ave., NW, Washington, DC 20009 USA.

Views expressed in this publication do not necessarily reflect official positions of the American Geophysical Union unless expressly stated.

<http://www.agu.org/pubs/eos>