

POLISH POLAR RESEARCH	19	1–2	37–60	1998
-----------------------	----	-----	-------	------

Stanisław RAKUSA-SUSZCZEWSKI¹, Krzysztof JAŻDŻEWSKI², Andrzej MYRCHA³
and Maria OLECH⁴

¹ Department of Antarctic Biology
Polish Academy of Sciences
Ustrzycka 10
02-141 Warszawa, POLAND

³ Institute of Biology,
University of Białystok
Świerkowa 20B
25-950 Białystok, POLAND

² Department of Polar Biology and Oceanobiology
University of Łódź
Banacha 12/16
90-237 Łódź, POLAND

⁴ Institute of Botany
Jagiellonian University
Lubicz 46,
31-512 Kraków, POLAND

Biological and ecological studies carried out at the Polish Antarctic Station *Henryk Arctowski*, 1977–1997

ABSTRACT: The authors describe the scope of Polish studies in the field of biology and ecology carried on during 20 years of activity of Polish Antarctic Station. Principal results are briefly summarized and ample literature is presented.

Key words: Antarctica, *Arctowski* Station, biological studies.

Introduction

The scientific successs of the consecutive Polish Antarctic Biological Expeditions (1968/70, 1971/72, 1973/74) as well as of the Ist Polish Antarctic Marine Research Expedition (1975/76) led to the decision of the Polish Governement to built a Polish Station in Western Antarctica¹.

It was built on King George Island (South Shetland Islands) at Point Thomas, Admiralty Bay, during two months of the austral summer 1976/77. The Station was named *Henryk Arctowski* after the scientific leader of the famous *Belgica*

¹ One of principal goals of this decision was Polish membership in the Antarctic Treaty Consultative Parties. In 1958, Poland received from Soviet Union their base *Oasis* situated at Bunger Hills, Eastern Antarctica, opened it in 1959 and renamed after *Antoni Bolesław Dobrowolski*. This station has not been occupied permanently and thus was not fulfilling the prerequisite conditions, even though Poland signed the Antarctic Treaty in 1961.

expedition (1897–99) commanded by Adrien De Gerlache. The Station was opened on 26 February 1977. It is located on a rather vast coastal plain with several freshwater ponds and mossy tundra carpets, bordered from the sea by storm ridges and some cliff rocks (Figs 1 and 2).

The Station consists of one large dwelling house used as a hotel by wintering groups (kitchen, bathroom, WC, dining-rest room, reading room and single-person bed-rooms for 13 people), several laboratory buildings (meteo, biology, geophysics), 2 large storage halls, a power-plant building and some “summer” houses for seasonal groups. A large tank for diesel oil is situated some 800 m to the north of the Station.

The *Henryk Arctowski* Station owns a small cutter *Stoń Morski* (length 9 m), equipped with an echosounder and winch enabling trawling, dredging and using bottom samplers down to the greatest depths of Admiralty Bay (over 500 m), as well as for hauling planktonic nets and using a variety of hydrological equipment (Fig. 3). A former boat used in oceanobiological research at the Station was named *Dziunia* (length 7 m); the boat is now exhibited at the marine museum in Hel (Fig. 4).

Scope of scientific research

From the very beginning, the studies carried on at the *Arctowski* Station were multidisciplinary. They included geophysics, geology, pedology, palaeontology, meteorology, geodesy, hydrology, biology and ecology. The latter three fields of studies, in accordance with the scientific interests of the supervising institutions (formerly the Institute of Ecology of Polish Academy of Sciences, now Department of Antarctic Biology Pol. Acad. Sci.), were hitherto represented by the highest number of researchers. The total number of scientific papers published based on the material collected at *Arctowski* Station in these fields is estimated at some 400.

Biological and ecological research: background studies

The main scientific efforts of Polish biologists and ecologists working at the *Arctowski* base has been directed toward recognition and description of the structure and functioning of the marine ecosystem of Admiralty Bay and land biota of King George Island, mainly of the nearest vicinity of the Station, *i.e.* along the south-western shores of Admiralty Bay.

Admiralty Bay is a trifurcated fjord with maximum depth of over 500 m and a total surface of ca 120 km². Its shoreline is some 83 km long; ice-free shores

constitute over 53% of the whole shoreline (Rakusa-Suszczewski 1980, Rakusa-Suszczewski *et al.* 1993a).

The ultimate goal of all biological studies, in accordance with an ecological approach of the leading institutions managing the *Arctowski* Station, is to gain an understanding of the functioning of the Antarctic near-shore ecosystem by the quantitative evaluation of all possible links of the trophic chain. Research has been carried out mostly in unknown regions, where the background for ecological studies needed to be studied first. These basic studies included investigations in the fields of geodesy, geomorphology, climatology, meteorology and hydrology (Rakusa-Suszczewski 1993c–e, Rakusa-Suszczewski *et al.* 1993b, c). Detailed maps were prepared (Furmańczyk and Marsz 1980; Battke 1981, 1990), and morphometry of the shores and basic ecological characteristics of Admiralty Bay region were presented (Marsz 1984, Marsz and Rakusa-Suszczewski 1987, Rakusa-Suszczewski 1995a). Climatic conditions were studied both in macro-scale (Styszyńska 1990) as well as in microscale (Moczydłowski 1986, Schroeter *et al.* 1995).

Recognition of physical and chemical properties of the environment of Admiralty Bay, sometimes also of the adjoining Bransfield Strait, constituted a basis for all further ecological studies (Pęcherzewski 1980, Pruszek 1980, Samp 1980, Szafranski and Lipski 1982, Bojanowski 1984, Catewicz and Kowalik 1984, Woźniak *et al.* 1984, Lipski 1987, Tokarczyk 1987, Sarukhanyan and Tokarczyk 1988, Madejski and Rakusa-Suszczewski 1990, Lipski and Rakusa-Suszczewski 1990, Rakusa-Suszczewski 1996). These studies have proved the low stability of the Admiralty Bay waters, the lack of a distinct thermocline, halocline and pycnocline. It was found that the wide contact of Admiralty Bay with Bransfield Strait waters and the system of currents allow Admiralty Bay water to exchange within 2 weeks.

Plankton of Admiralty Bay

Microbiological studies of Admiralty Bay were carried out mainly by Zdanowski (1981, 1988, 1995), Zdanowski and Donachie (1993) and Donachie (1995). A considerable variability in the amount of saprophytic bacteria over longer than seasonal scales was observed. It was also suggested that the reason for this variability could be the varying abundance of krill in the Bay. Special attention was paid to the microorganisms occurring in krill stomachs (Rakusa-Suszczewski and Zdanowski 1989) and to the role of bacteria in the process of krill decomposition and autolysis (Zdanowski 1981, 1988, 1995; Donachie 1995). A large culture collection of ca 500 bacterial strains has been assembled, deriving from various marine habitats, in order to compile nutritional profiles for these strains.

It appears that 93% of primary production takes place in the 25-m thick euphotic layer of Admiralty Bay. Primary production and chlorophyll contents have been studied by Hapter *et al.* (1984), Tokarczyk (1986), Lipski (1987) and Domanov and Lipski (1990). Daily primary production fluctuates in a wide range, depending on the season and the region of Admiralty Bay, being the lowest in inner inlets where water transparency was reduced. In general, the average values of primary production approach those of the open waters around South Shetland Islands. Total annual production in surface waters of the central basin of Admiralty Bay is estimated at 6.6 g C m^{-3} .

The phytoplankton of Admiralty Bay was investigated by Kopczyńska (1980, 1981) and Ligowski (1986, 1993a). The nanoplankton is composed of flagellates (12–15 μm), Prasinophyceae, Cryptophyceae and “monads” (4–6 μm). Maximum numbers were recorded during the austral summer in February; algal concentrations ranged from 35×10^3 to $1600 \times 10^3 \text{ cells dm}^{-3}$. In net phytoplankton diatoms dominated, with such taxa as *Thalassiosira antarctica*, *Nitzschia cylindrus*, *Corethron criophilum* and *Thalassiothrix* spp. as main components. Altogether 168 diatom taxa in the phytoplankton were recognized. As was expected, the highest phytoplankton densities occurred in the austral spring and summer, the lowest – during winter (April – October).

Sea-ice algae in Admiralty Bay were studied by Ligowski (1987, 1993a,c). Three different diatom communities were discovered in ice-floes with different characteristic species. In all, as many as 138 diatom taxa were identified in the sea-ice from Admiralty Bay. Earlier assumptions that the physical concentration of algae takes place during sea-ice formation have been confirmed. The diatom cells' density in ice was found to be even 2500 times higher than in the surrounding water.

Ligowski (1985), during his studies of the diet of krill in Admiralty Bay, was able to demonstrate that *Euphausia superba* consumed considerable amounts both of sea-ice algae and of epibenthic/epiphytic diatoms. It was concluded that in the near-shore, neritic conditions krill supplements its primarily planktonic diet with sea-ice- and benthic micro-algae.

In zooplankton some groups were more thoroughly studied. Protozoans of the Tintinnina suborder were investigated by Wasik and Mikołajczyk (1993, 1994), who found 17 tintinnid species in the Admiralty Bay and in adjacent Bransfield Strait waters, with *Cymatocyclis convallaria* and *Codonellopsis balechi* being dominants. Tintinnids occurred only in the uppermost 50 m layer, with maximum density up to 4000 ind. m^{-3} . The ultrastructure of *C. convallaria* was also studied, discovering new interesting morphological data.

The Copepoda of Admiralty Bay have been studied by several Polish authors (Chojnacki and Węgleńska 1984, Żmijewska 1993). Thirty nine species have been recorded, with clear dominance of *Metridia gerlachei*; next in terms of abundance were *Euchaeta antarctica*, *Calanoides acutus* and *C. propinquus*.

The yearly cycle of zooplankton in Admiralty Bay waters was investigated by Menshenina and Rakusa-Suszczewski (1992).

Special attention in zooplankton studies was paid to Euphausiacea. Five species of this group were recorded in Admiralty Bay: *Euphausia superba*, *E. frigida*, *E. triacantha*, *E. crystallorophias* and *Thysanoessa macrura*. Studies were carried out on the morphology of postlarval stages (Kittel and Presler 1980), and on the diet of some euphausiid species (Kittel and Ligowski 1980, Ligowski 1993b). Krill – *Euphausia superba* – was obviously the most thoroughly studied species. Its population structure was investigated by Jackowska (1980), Kittel (1980), Rakusa-Suszczewski and Stępnik (1982) and Stępnik (1982); the growth and mortality of *E. superba* by McClatchie *et al.* (1991); whereas its commensals and parasites – by Stawiszyńska-Janasz and Kittel (1982), Rakusa-Suszczewski and Filcek (1988), Rakusa-Suszczewski and Nemoto (1989) and Stankovič and Rakusa-Suszczewski (1996).

Benthos of Admiralty Bay

In the sediments and on benthic macroalgae of Admiralty Bay, 120 diatom taxa were found (Ligowski 1993b,d). In different depths various diatoms dominated. Attempts to estimate the total microphytobenthos production in the basin studied clearly showed that the role of this primary production is very significant.

Zieliński (1981, 1990) studied the macrophytobenthos of Admiralty Bay. His studies have brought a list of 36 macroalgal taxa, among them 20 species of Rhodophyta, 10 species of Thallophyta, 5 species of Chlorophyta and 1 species of Chrysophyta. The most common taxa were *Monostroma harriotti*, *Iridaea cordata*, *Adenocystis utricularis*, *Ascoseira mirabilis*, *Desmarestia* spp. and *Himantothallus grandifolius*.

Macroalgae occur in Admiralty Bay down to the depths of 90–100 m. The extension of their meadows was estimated by air photographs analysis (Furmańczyk and Zieliński 1982) at some 30% of Admiralty Bay surface bottom.

Polish studies of zoobenthos of Admiralty Bay are ranked among the most comprehensive investigations of the Antarctic bottom fauna (Knox 1994). Along with qualitative faunistic studies (Jażdżewski 1981, 1993; Arnaud *et al.* 1986; Siciński 1986, 1993; Jażdżewski *et al.* 1991, 1992, 1995; Siciński and Janowska 1992; Presler 1993b; Błażewicz and Jażdżewski 1996, 1997; Presler and Figielska 1998) in which over 400 invertebrate taxa have been recorded, our investigations have brought the quantitative estimation of the abundance and biomass of zoobenthos of Admiralty Bay (Jażdżewski *et al.* 1986, 1991). Polychaeta and Amphipoda appear to be the most diverse animal groups – about 120 species of each group. In terms of abundance Bivalvia and Polychaeta predominate, where-

as in terms of biomass, in shallow waters, bivalve molluscs also predominate, yielding in the deeper sublittoral to Ascidiacea, Echinodermata, and Bryozoa. Maximum densities of zoobenthos observed ranged up to over 36000 ind. m⁻² and over 2500 g m⁻², but estimates drawn from some underwater pictures indicate the presence of higher biomass values per m². The average biomass of the benthos increases from ca 250 g m⁻² in the shallow sublittoral (10–15 m) to over 1000 g m⁻² at the depths 100–200 m, and then decreases in the deepest parts of the Bay (400–500 m) to about 300 g m⁻² (Jażdżewski and Siciński 1993).

A significant role in the Antarctic benthic communities is played by necrophagous animals. This ecological group was studied by Presler (1986, 1993a) who recorded 23 scavenging species (among others 5 lysianassoid amphipods and 11 asteroids) in Admiralty Bay and has presented their depth preferences and seasonal changes in the distribution of particular necrophagous taxa.

Autecological study of a littoral gastropod, *Nacella concinna*, was carried out by Filcek (1993).

Fishes of Admiralty Bay

The studies of fishes inhabiting Admiralty Bay commenced with the first wintering (1977/1978) at the *Arctowski* Station. They included faunistic studies (Żukowski 1980) as well as observations on the growth, respiration, and food habits of nototheniid fishes (Linkowski and Żukowski 1980, Linkowski *et al.* 1983, Kulesz 1994). Further ichthyological studies were carried out by Skóra and Neyelov (1992), Skóra (1993, 1995), Kulesz and Zadróźny (1996), and Zadróźny (1996). Thirty-nine fish species belonging to 11 families have been hitherto recorded in Admiralty Bay; one third of them belonged to the family Nototheniidae. The most common species are *Notothenia neglecta*, *N. rossi* and *Gobionotothen gibberifrons*. One fish species is new for science (*Acanthodraco dewitti*). Depth-related changes in ichthyofauna of Admiralty Bay have been described. It was observed that the abundance and biomass of fishes in Admiralty Bay were distinctly lower than those of adjacent Bransfield Strait shelf of King George Island.

Birds and mammals of Admiralty Bay

Birds and mammals breeding at the shores of King George Island, and mainly in the *Arctowski* Station area, have been monitored from the very beginning of the station's activity (1977/1978). Altogether twelve nesting bird species and 16 further species visiting King George Island have been recorded by Polish and American ornithologists working in this area (Jabłoński 1980, Presler 1980,

Trivelpiece *et al.* 1987). The census of 3 species of pygoscelid penguins (*Pygoscelis adeliae*, *P. antarctica* and *P. papua*) and of other nesting birds was carried out by Jabłoński (1984a, b, 1986, 1987), Sierakowski (1991) and Lesiński (1993). Significant fluctuations of the number of breeding pairs of pygoscelid penguins were recorded over the period 1978–1990. The most serious decline in the number of breeding pairs, nearly 3-fold, was observed in *Pygoscelis adeliae*. Moczydłowski (1989) has investigated the nesting strategy of pygoscelid penguins. Detailed studies on the breeding biology of the Wilson's storm petrel (*Oceanites oceanicus*) were carried out by Wasilewski (1986) and on the Antarctic tern (*Sterna vittata*) by Jabłoński (1995).

Short summary of ornithological observations in the Admiralty Bay area was presented by Myrcha (1993).

Five species of pinnipedian mammals occur in the region of Admiralty Bay; one species of fur seal – *Arctocephalus gazella*, and 4 species of seals – *Mirounga leonina*, *Leptonychotes weddelli*, *Lobodon carcinophagus* and *Hydrurga leptonyx*, the latter two species being closely associated with pack ice. Monitoring of seals has been consequently carried out for years (Myrcha and Teliga 1980, Presler 1980, Wojciechowski 1980, Krzemiński 1981, Jabłoński *et al.* 1987, Sierakowski 1991, Lesiński 1993, Rakusa-Suszczewski and Sierakowski 1993). The results of this census indicate also the significant fluctuations of populations of all the monitored species.

Parasites of vertebrates

Parasites of vertebrate animals of King George Island and Admiralty Bay were thoroughly studied by a number of Polish scientists (Zdzitowiecki 1978, 1979, 1985, 1986a–c, 1987, 1991; Zdzitowiecki and Drózd 1980; Cielecka and Zdzitowiecki 1981, 1989; Jarecka 1984; Jarecka and Ostas 1984; Zdzitowiecki and Rokosz 1986; Zdzitowiecki *et al.* 1989; Wojciechowska 1993a–c; Cielecka *et al.* 1995; Zdzitowiecki and Cielecka 1996). Many species new to science have been described and some full life cycles recognized. The contribution of our parasitologists and especially that of K. Zdzitowiecki, to the knowledge of parasites of Antarctic vertebrates, must be recognized as one of major achievements of the Polish scientific activity in the Southern Ocean (see Zdzitowiecki 1991b, 1997).

Soil formation studies

Land biota were also investigated by Polish biologists and ecologists working at the *Arctowski* Station. A pioneer research was carried out on the fate of faeces

of numerous penguins in the rookeries of King George Island. Ecological studies by Tatur and Myrcha (1983, 1984, 1989, 1993), Tatur and Barczuk (1984, 1985), Myrcha *et al.* (1985), Tatur (1987, 1989), Myrcha and Tatur (1991) and Tatur *et al.* (1997) have demonstrated that penguins in the rookery regions deposit as much as 10 kg dry weight of guano per square m during one breeding season. It was calculated that about 5% of the ice-free area of Admiralty Bay is affected by the deposition of guano.

This strong fertilization with nutrients leads to the formation of specific ornithogenic soils. Minerals forming these soils were studied in detail; it appears that the composition of phosphates is strikingly diverse. The monomineral accumulations of phosphates and the sequence of changes in mineral composition of ornithogenic soils is unique on a global scale.

The presence of penguin rookeries substantially modifies the fertility of soils in the maritime Antarctic. In the rookery itself, high nutrient contents prevent the development of vegetation that develops, however, at the peripheries of rookeries and then invades the abandoned rookeries. The succession begins with the development of the coprophagous alga *Prasiola crispa* leading to the compact plant communities with diverse composition of species.

Soil microorganisms were studied by Zabawski and Piasecki (1981), Pietr (1986, 1993), Czekanowska and Zabawski (1988), and Zabawski (1995). It was demonstrated that as a result of bacterial activity fresh guano deposited by penguins undergoes mineralization over a period of 3 weeks. Proteolytic bacteria and ammonifiers dominate, while chitinolytic bacteria are also prominent. Eleven species of penicillian fungi were isolated from soil samples. Fungal contribution to the mineralization of penguin guano appears to be very low. In general, in comparison with regions having different climatic regimes, microbial populations of the soils in the Admiralty Bay region displayed greater qualitative and quantitative differences over small areas.

Botanical studies on land

Botanical research of King George Island has been extensively carried out during nearly all Polish expeditions to the *Arctowski* Station. Freshwater and aerophytic algae and Cyanophyceae were studied by Kawecka and Olech (1993, 1996), Luścińska and Kyć (1993) Massalski *et al.* (1994, 1995) and Kawecka *et al.* (1996). 132 species of Bacillariophyceae, 30 species of other algae (Chlorophyceae, Chrysophyceae, Euglenophyceae), and 26 species of Cyanophyceae were recorded (Kawecka and Olech 1996, Mrozińska *et al.* 1996). Macromycetes were investigated by Gumińska *et al.* (1994); 6 species were found and their distribution was described.

Lichenicolous fungi were studied by Olech and Alstrup (1989, 1995), and Alstrup and Olech (*in press*). 65 taxa were discovered, among them nearly half were new to science. A key for determination of this group of organisms in the Antarctic was prepared.

Diverse and thorough research on lichens (lichenized fungi) of King George Island and adjacent regions of maritime Antarctic was carried out by Olech (1989, 1990, 1991a, b), Olech and Alstrup (1989), and Olech and Sochting (1993). In all, over 200 species of lichens were recorded, among them 10 taxa new to science, 51 new to Antarctica, and 30 new to the Southern Hemisphere. A monograph of the genus *Caloplaca* in the Antarctic was prepared (Sochting and Olech 1995).

The distribution of 10 species of liverworts (Hepaticae) in the region of Admiralty Bay was described in details by Ochyra and Vana (1989).

Ochyra and Bell (1984), Ochyra (1985, 1987, 1990), Ochi and Ochyra (1985), Ochyra and Afonina (1986), Ochyra and Ochi (1986), and Ochyra and Lewis-Smith (1996) have studied mosses (Bryophyta) in the wide surroundings of the *Arctowski* Station. They have found over 50 species of mosses there, among them some taxa new to science.

Polish phytosociological research on King George Island was initiated by Furmańczyk and Ochyra (1982), then followed by Olech (1990), who has described 4 ornithocoprophagous communities in the neighbourhood of penguin rookeries. Olech (1994) investigated also the distribution of lichens in the region of Lions Rump (SSSI No. 34), using a 250 × 250 m grid, preparing a map of vegetation.

Vascular plants and their distribution and growth under different experimental conditions, their flowering and fruiting, were studied by Zarzycki (1993), whereas the peat forming vegetation, peat mounds, and the origin and development of mires in the region of *Arctowski* Station were described by Fabiszewski and Wojtuń (1993).

Land and freshwater invertebrates

Land and freshwater invertebrates were also investigated. Niedbała (1986) found 6 taxa of oribatid mites in the material collected at the *Arctowski* Station, among them one species new to science. Jurasz *et al.* (1983), and Janiec (1991) have studied the occurrence, distribution and life cycle of freshwater anostracan crustacean, *Branchinecta gaini*. Janiec (1993, 1996a) has described in detail freshwater invertebrate communities composed of Nematoda, Tardigrada, Rotatoria and Crustacea occurring in ponds and peat bogs of King George Island. Worth mentioning is the fact that this paper belongs to the class of those rare publications where the same author studied similar biotopes in the Arctic and in

the Antarctic, making the comparisons especially valuable. In another paper by Janiec (1996b) attention was focussed on the process of invasion of different taxa into new deglaciated areas and to the succession of the communities.

Biochemical and physiological studies

Physiological and biochemical studies carried out at the *Arctowski* Station have been diversified. Heavy metal contents in Antarctic lichens was investigated by Olech (1991c); lichens were also used as indicators of air pollution in the area of *Arctowski* Station (Szymczyk *et al.* 1994).

The contamination of various Antarctic organisms by the chlorinated hydrocarbons residues (CHs) was studied by Łukowski (1983a, b), Łukowski *et al.* (1987), Karolewski *et al.* (1987) and Łukowski and Ligowski (1987). CHs accumulate in organisms and are concentrated passing from one trophic level to the next. DDT concentration in krill were, on an average, 0.06 ppm, in pygoscelid penguins on average 0.198 ppm, and in leopard seal 0.250 ppm. Due to the studies of planktonic diatoms coming from various areas it was also found that the ice sheet and glaciers in the Antarctic constitute an additional source of contamination by CHs which have accumulated there in the past. It appeared also that the accumulation of CHs is greatest in the diatoms developing in sea-ice.

CHs contamination of flying birds appeared to be dependent on the type of food consumed and on the habits throughout the year. Contamination was higher in species migrating to the northern hemisphere. The diminishing of the contamination of Adelie penguin (*Pygoscelis adeliae*) by DDT residues in last decades was also observed, reflecting worldwide decrease in the use of this insecticide. Studies of pinnipedian contamination by these agents revealed that the highest levels occur in seals feeding on penguins (leopard seals) and on fish (Weddell seal).

Physiological properties of blood in several birds inhabiting King George Island were investigated by Myrcha and Kostelecka-Myrcha (1980a, b) and Kostelecka-Myrcha and Myrcha (1980, 1989). These authors demonstrated significant differences between particular species. Body calorific values of pygoscelid penguins during their development were studied by Myrcha and Kamiński (1982).

Ample biochemical studies of krill (*E. superba*) from Admiralty Bay were also been carried out at the *Arctowski* Station. Many papers were devoted to the lipid content, its seasonal changes in krill, as well as to the lipids oxidability (Kořakowska 1986, 1991a, b). Proteins of krill, their amino-acid composition and their autoproteolysis were studied by Kořakowski (1986a, b, 1987, 1989, 1993), whereas the technical possibility of using autoproteolysis to obtain pro-

tein precipitate was investigated by Kołakowski *et al.* (1980a,b), and Kołakowski and Lachowicz (1982).

Metabolism of krill was studied under different thermic conditions (Rakusa-Suszczewski 1990). High seasonal variability in respiration rate and biomass was observed; in winter a 40% decrease of metabolism was recorded. Respiration of some amphipods and gastropods inhabiting Admiralty Bay has also been investigated (Rakusa-Suszczewski and Lach 1991, Rakusa-Suszczewski 1992).

Ecosystem functioning

Early attempts to present a general pattern of functioning of the ecosystem of Admiralty Bay and its shores were made by Rakusa-Suszczewski (1980a, b, 1987). 15 years after the foundation of the *Arctowski* Station our knowledge has substantially increased due to all the afore-mentioned studies and also many others; it was therefore possible to produce a monograph entitled *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay* (Rakusa-Suszczewski 1993a), summarizing most of the hitherto obtained Polish biological and ecological studies carried out at the *Arctowski* Station. In the introductory and closing articles short quantitative evaluations of Admiralty Bay's functioning as a semi-closed hydrological unit and a very good example of the Antarctic near-shore ecosystem were presented (Rakusa-Suszczewski 1993b, f). This same author has estimated the flow of matter in the Admiralty Bay area (Rakusa-Suszczewski 1995b).

Life in small areas of ice-free Antarctic shores is closely connected with the productivity of the ocean and with the transport of organic matter and nutrients from sea to land. Primary production in Admiralty Bay in austral summer is estimated at over 2300 tons of phytoplankton net weight. The standing crop of benthic microalgae has been calculated at some 13 tons of chlorophyll, whereas that of macroalgae – at about 300 tons. Secondary production of zooplankton, according to the well-known principles, can be estimated at about 230 tons. The biomass of all three pygoscelid penguins in the breeding period was calculated to be about 250 tons. Their daily consumption of krill was estimated at 37 tons. It is therefore evident that Admiralty Bay does not suffice as a foraging ground and penguins have to penetrate much more distant areas.

The competition for food between birds and mammals is probably not very high, since their abundance peaks fall in different months. Main consumers of krill and fishes are sea-elephants and fur seals; the amount of food taken by these two pinnipeds in the region of Admiralty Bay was calculated as 1400 tons. Crab-eater seals alone consume some 630 tons of krill.

Birds and mammals bring organic matter in form of faeces, egg shells, feathers, mollusc shells and hairs on to the land. It was estimated that in the region of rookeries 1.8 kg of organic matter per square m is transported. Ornithogenic soils of abandoned rookeries are then occupied by land vegetation. Freshwater bodies enriched by nutrients are also richer in micro- and meio-organisms.

Impact of human activity on Antarctic ecosystem

One should remember that the land ecosystems in the Antarctic are rather fragile and susceptible to anthropogenic impact. Since the founding of the *Arctowski* Station some 15000 tons of materials, fuel and food were brought on land, whereas only 1500 tons were taken back. Fuel consumption at the *Arctowski* Station fluctuated around 200 t yearly, whereas daily food consumption was about 5 kg, and water consumption about 120 l per person. The functioning of the station and its logistic needs can be better imagined keeping in mind that from the very beginnings till now some 250 persons have wintered at the *Arctowski* Station, and some 600 persons stayed there during austral summer (usually 2–3 months). Several thousand tourists have visited the station and their number is increasing far too quickly (Donachie 1993). Major problems of the anthropogenic impact, mainly upon land biota of King George Island, were signalized by Krzyszowska (1990), Olech (1991c, 1994, 1996a, b), Rakusa-Suszczewski and Krzyszowska (1991) and Rakusa-Suszczewski (1993a, b).

Protection and monitoring of Antarctic environment

The Polish initiative has led to the creation and acceptance by SCAR of the Site of Special Scientific Interest (SSSI) No. 8. This is an area in the vicinity of *Arctowski* Station, on the western shores of Admiralty Bay. In 1990 a successive, scientifically well-substantiated initiative was presented to SCAR, namely the foundation of another SSSI on King George Island (No. 34), situated on western shores of King George Bay; an area remote from human activity and therefore very useful in comparative studies.

In this region of the South Shetland Islands commercial fishing of krill and fish are carried out. The presence of fishery fleets and exploitation of the living resources of the region create serious dangers for the shelf and near shore ecosystem. Polish monitoring of penguins and pinnipedian mammals are in response to SCAR and CCAMLR recommendations. Most of the biological studies carried out at the *Arctowski* Station are complementary to the research conducted during several multinational marine expeditions in the BIOMASS

programme, a fact which has gained high recognition in the international scientific community. Land and inland freshwater studies conducted on King George Island by Polish biologists fit well to the international BIOTAS programme (Biological Investigations of Terrestrial Antarctic Systems).

Final remarks

At present Polish biological studies in the Antarctic are restricted to the near-shore zone where Poland participates in the international scientific program CZ-ASIZ (Coastal Zone – Ecology of Antarctic Sea Ice Zone). In this program the *Arctowski* Station is recognized as a core station.

The wide scope of Polish biological and ecological studies carried out so far over twenty years at the *Arctowski* Station has aided in the recognition and estimation of the natural variability of the maritime Antarctic geo-ecosystem, the understanding of causes of its variability, and the influence of physical parameters and biological responses at the levels of species, populations, or communities.

References

- ALSTRUP V. and OLECH M. [*in press*]. The lichenicolous fungi of the South Shetland Islands with a key to the species known from Antarctica and Subantarctica. — *Cryptogamie, Bryol. Lichenol.*
- ARNAUD P.M., JAŻDŻEWSKI K., PRESLER P. and SICIŃSKI J. 1986. Preliminary survey of benthic invertebrates collected by Polish Antarctic Expeditions in the Admiralty Bay (King George Island, South Shetlands, Antarctica). — *Pol. Polar Res.*, 7: 7–24.
- BATTKE Z. 1981. Admiralty Bay. Scale 1:50.000. H. Arctowski Antarctic Station. — Printed by WZKart., Warsaw.
- BATTKE Z. 1990. Admiralty Bay. King George Island. Scale 1:50.000. — Printed by E. Romer State Cartographical Publishing House, Warsaw.
- BŁAŻEWICZ M. and JAŻDŻEWSKI K. 1996 (1995). Cumacea (Crustacea, Malacostraca) of Admiralty Bay (King George Island, South Shetland Islands), preliminary note. — *Pol. Polar Res.* 16: 71–85.
- BŁAŻEWICZ M. and JAŻDŻEWSKI K. 1997 (1996). A contribution to the knowledge of Tanaidacea (Crustacea, Malacostraca) of Admiralty Bay, King George Island, Antarctic. — *Pol. Polar Res.* 17: 213–220.
- BOJANOWSKI R. 1984. Hydrochemical observations at an Anchored Station in Ezcurra Inlet. — *Oceanologia*, 15: 21–64.
- CATEWICZ Z. and KOWALIK Z. 1984. Harmonic analysis of tides in Admiralty Bay. — *Oceanologia*, 15: 97–109.
- CHOJNACKI J. and WĘGLEŃSKA T. 1984. Periodicity of composition, abundance and vertical distribution of summer zooplankton (1977/1978) in Ezcurra Inlet, Admiralty Bay (King George Island, South Shetlands). — *J. Plankt. Res.*, 6: 997–1017.

- CIELECKA D. and ZDZITOWIECKI K. 1981. The tapeworm *Microsomacanthus shetlandicus* sp.n. (Hymenolepididae) from the Dominican Gull of King George Island (South Shetlands, Antarctic). — Bull. Acad. Pol. Sci., Ser. Sci. Biol., Cl.II, 29: 173–180.
- CIELECKA D., WOJCIECHOWSKA A. and ZDZITOWIECKI K. 1992. Cestodes from penguins on King George Island (South Shetlands, Antarctic). — Acta Parasitologica, 37: 65–72.
- CZEKANOWSKA A. and ZABAWSKI J. 1988. Soil actinomycetes of the Admiralty Bay area at King George Island, Antarctic. In: A. Jahn, J. Pereyma and A. Szczepankiewicz-Szmyrka (eds.), XVIth Polar Symposium: *Present state and selected problems of Polish polar studies*. — Wrocław Univ. Publ. House, Wrocław; 279–288.
- DOMANOV M. M. and LIPSKI M. 1990. Annual cycle of chlorophyll *a* and primary production of phytoplankton in Admiralty Bay (Antarctica). — Pol. Arch. Hydrobiol., 37: 471–478.
- DONACHE S. P. 1995. Ecophysiological description of marine bacteria from Admiralty Bay (Antarctica), and the digestive tracts of selected Euphausiidae. In: S. Rakusa-Suszczewski and S. P. Donachie (eds.), *Microbiology of Antarctic marine environments and krill intestine, its decomposition and digestive enzymes*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 101–196.
- FABISZEWSKI J. and WOJTUŃ B. 1993. 21. Peat-forming vegetation. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 189–195.
- FILCEK K. 1993. 12. Zoobenthos. 12.6. Patellidae. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 120–122.
- FURMAŃCZYK K. and MARSZ A. 1980. Szetlandy Południowe (South Shetland Islands), Wyspa Króla Jerzego (King George Is. – Isla 25 de Mayo – Ostrov Waterloo), Zatoka Admirałicji (Admiralty Bay – Bahia Lasserre). Scale 1:25.000. — IGUG, Szczecin, Printed by OPGK.
- FURMAŃCZYK K. and OCHYRA R. 1982. Plant communities of the Admiralty Bay region (King George Island, South Shetland Islands, Antarctic). — Pol. Polar Res., 3: 25–39.
- FURMAŃCZYK K. and ZIELIŃSKI K. 1982. Distribution of macroalgae groupings in shallow waters of Admiralty Bay (King George Island, South Shetland Islands, Antarctic), plotted with the help of air photographs analysis. — Pol. Polar Res., 3: 41–47.
- GUMIŃSKA B., HEINRICH Z. and OLECH M. 1993. Macromycetes of the south Shetland Islands (Antarctica). — Pol. Polar Res., 15: 103–110.
- HAPTER R., WOŹNIAK B. and DOBROWOLSKI K. 1984. Primary production in Ezcurra Inlet during austral summer of 1977/78. — Oceanologia, 15: 175–183.
- JABŁOŃSKI B. 1980. Distribution and numbers of birds and pinnipeds of Penguin Island (South Shetland Islands) in January 1979. — Pol. Polar Res. 1: 109–116.
- JABŁOŃSKI B. 1984a. Distribution and breeding preferences of penguins in the region of the Admiralty Bay (King George Island, South Shetland Islands) in the season 1979/80. — Pol. Polar Res., 5: 5–16.
- JABŁOŃSKI B. 1984b. Distribution and numbers of penguins in the region of King George Island (South Shetland Islands) in the breeding season 1980/81. — Pol. Polar Res., 5: 17–30.
- JABŁOŃSKI B. 1986. Distribution, abundance and biomass of a summer community of birds in the region of the Admiralty Bay (King George Island, South Shetlands, Antarctica) in 1978/79. — Pol. Polar Res., 7: 217–260.
- JABŁOŃSKI B. 1987. Diurnal pattern of changes in the number of penguins on land and the estimation of their abundance (Admiralty Bay, King George I., South Shetland Islands). — Acta zool. cracov., 30: 119–136.
- JABŁOŃSKI B. 1995. Distribution, abundance and biology of the Antarctic Tern, *Sterna vittata* Gmelin, 1789 on King George Island (South Shetland Islands). — Acta zool. cracov., 38: 399–460.

- JABŁOŃSKI B., KRZEMIŃSKI W. and ZDZITOWIECKI K. 1987. Distribution and number of fur seals, *Arctocephalus gazella* (Peters, 1875) of King George Island (South Shetlands). — *Acta zool. cracov.*, 30,9: 119–136.
- JACKOWSKA H. 1980. Krill monitoring in Admiralty Bay (King George Island, South Shetland Islands) in summer 1979/1980. — *Pol. Polar Res.* 1: 117–125.
- JANIEC K. 1991. A further note on *Branchinecta gaini* Daday, 1910 population from the region of the *Arctowski* Station (King George Island, South Shetland Islands). — *Pol. Polar Res.* 12: 123–127.
- JANIEC K. 1993. The freshwater micro- and meiofauna of Admiralty Bay, King George Island, South Shetland Islands. — *Proc. NIPR Symp. Polar Biol.*, 6: 133–138.
- JANIEC K. 1996a. The comparison of freshwater invertebrates of Spitsbergen (Arctic) and King George Island (Antarctic). — *Pol. Polar Res.* 17: 173–202.
- JANIEC K. 1996b. Short distance wind transport of macrofauna in maritime Antarctic (King George Island, South Shetland Islands). — *Pol. Polar Res.* 17: 203–211.
- JARECKA L. 1984. Development of *Hymenolepis arctowskii* Jarecka et Ostas, 1984 (Cestoda, Hymenolepididae) in the intermediate host *Branchinecta gaini* Daday (Branchiopoda) of the Antarctic. — *Acta Parasitol. Pol.*, 29: 337–342.
- JARECKA L. and OSTAS J. 1984. *Hymenolepis arctowskii* n.sp. (Cestoda, Hymenolepididae) from *Larus dominicanus* Licht. of the Antarctic. — *Acta Parasitol. Pol.*, 29: 189–196.
- JAŻDŻEWSKI K. 1981. Amphipod crustaceans in the diet of pygoscelid penguins of the King George Island, South Shetland Islands, Antarctica. — *Pol. Polar Res.*, 2: 133–144.
- JAŻDŻEWSKI K. 1993. 12. Zoobenthos. 12.4. Amphipoda. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 108–116.
- JAŻDŻEWSKI K. and SICIŃSKI J. 1993. 12. Zoobenthos. 12.1. General remarks. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 83–95.
- JAŻDŻEWSKI K., WĘSLAWSKI J. M. and DE BROYER C. 1995. A comparison of the amphipod faunal diversity in two polar fjords : Admiralty Bay, King George Island (Antarctic) and Hornsund, Spitsbergen (Arctic). — *Pol. Arch. Hydrobiol.*, 42: 367–384.
- JAŻDŻEWSKI K., DE BROYER C., TEODORCZYK W. and KONOPACKA A. 1992. Survey and distributional patterns of the amphipod fauna of Admiralty Bay, King George Island, South Shetland Islands. — *Pol. Polar Res.* 12: 461–472.
- JAŻDŻEWSKI K., TEODORCZYK W., SICIŃSKI J. and KONTEK B. 1991. Amphipod crustaceans as an important component of zoobenthos of the shallow Antarctic sublittoral. — *Hydrobiologia*, 223: 105–117.
- JAŻDŻEWSKI K., JURASZ W., KITTEL W., PRESLER E., PRESLER P. and SICIŃSKI J. 1986. Abundance and Biomass Estimates of the Benthic Fauna in Admiralty Bay, King George Island, South Shetland Islands. — *Polar Biol.*, 6: 5–16.
- JURASZ W., KITTEL W. and PRESLER P. 1983. Life cycle of *Branchinecta gaini*, Daday, 1910 (Branchiopoda, Anostraca) from King George Island, South Shetland Islands. — *Pol. Polar Res.*, 4: 143–154.
- KAROLEWSKI M. A., ŁUKOWSKI A. B. and HALBA R. 1987. Residues of chlorinated hydrocarbons in the adipose tissue of the Antarctic pinnipeds. — *Pol. Polar Res.*, 8: 189–196.
- KAWECKA B. and OLECH M. 1993. Diatom communities in the Vanishing and Ornithologis Creeks (King George Island, South Shetlands, Antarctica). — *Hydrobiologia*, 269/270: 327–333.
- KAWECKA B. and OLECH M. 1996. Diatom communities in the inland waters of Admiralty Bay (King George Island, South Shetlands, Antarctica). In: M. OLECH (ed.), *Antarctic Research on Taxonomy and Ecology of Algae*. — Proc. Workshop, Cracow; 9.
- KAWECKA B., OLECH M. and NOWOGRODZKA-ZAGÓRSKA M. 1996. Morphological variability of diatom *Luticola muticopsis* in the inland waters of King George Island (South Shetlands, Antarctica). — *Pol. Polar Res.* 17: 23–30.

- KITTEL W. 1980. Populational studies on *Euphausia superba* Dana, 1852 (Euphausiacea, Crustacea) in waters of the Admiralty Bay during Antarctic summer of 1978. — Pol. Arch. Hydrobiol., 27: 267–272.
- KITTEL W. and LIGOWSKI R. 1980. Algae found in the food of *Euphausia crystallorophias* (Crustacea). — Pol. Polar Res. 1: 129–137.
- KITTEL W. and PRESLER P. 1980. Morphology of the postlarval developmental stages of *Euphausia crystallorophias* Holt et Tattersall, 1960 (Euphausiacea, Crustacea). — Pol. Arch. Hydrobiol., 27: 259–265.
- KNOX G. A. 1994. The Biology of Southern Ocean. Ocean Studies in Polar Research. — University Cambridge Press, Cambridge, 444 pp.
- KOŁAKOWSKA A. 1986. Lipid composition of fresh and frozen-stored krill. — Z. Lebensm. Unters. Forsch. 182: 475–478.
- KOŁAKOWSKA A. 1991a. The influence of sex and maturity stage of krill (*Euphausia superba* Dana) upon the content and composition of its lipids. — Pol. Polar Res. 12: 73–78.
- KOŁAKOWSKA A. 1991b. The lipid composition of some Antarctic Crustacea. — Pol. Polar Res. 12: 79–89.
- KOŁAKOWSKI E. 1986a. Changes of non-protein nitrogen fractions in Antarctic krill (*Euphausia superba* Dana) during storage at 3°C and 20°C. — Z. Lebensm. Unters. Forsch., 183: 421–425.
- KOŁAKOWSKI E. 1986b. Seasonal variation of autoprolytic activity in the Antarctic krill, *Euphausia superba* Dana. — Pol. Polar Res., 7: 275–282.
- KOŁAKOWSKI E. 1987. Seasonal changes in nitrogen fractions in the Antarctic krill (*Euphausia superba* Dana). Part 1. Basic nitrogen fractions. — Pol. Polar Res., 8: 159–165.
- KOŁAKOWSKI E. 1989. Proteolytic activity of Antarctic krill in relation to its feeding intensity in spring and summer. — Pol. Polar Res. 10: 141–150.
- KOŁAKOWSKI E. 1993. Composition and changes of free amino acids in Antarctic krill during storage at 3°C. — Z. Lebensm. Unters. Forsch..
- KOŁAKOWSKI E. and LACHOWICZ K. 1982. Application of partial autoprolysis to extraction of protein from Antarctic krill (*Euphausia superba* Dana). Part 3. — Nahrung, 16: 933–939.
- KOŁAKOWSKI E., GAJOWIECKI L., SZYBOWICZ Z. and CHODORSKA T. 1980a. Application of partial autoprolysis to extraction of protein from Antarctic krill (*Euphausia superba* Dana). Part 1. — Nahrung, 24: 499–506.
- KOŁAKOWSKI E., GAJOWIECKI L., SZYBOWICZ Z. and CHODORSKA T. 1980b. Application of partial autoprolysis to extraction of protein from Antarctic krill (*Euphausia superba* Dana). Part 2. — Nahrung, 24: 507–512.
- KOPCZYŃSKA E. E. 1980. Small-scale vertical distribution of phytoplankton in Ezcurra Inlet, Admiralty Bay, South Shetland Islands. — Pol. Polar Res. 1: 77–96.
- KOPCZYŃSKA E. E. 1981. Periodicity and composition of summer phytoplankton in Ezcurra Inlet, Admiralty Bay, South Shetland Islands. — Pol. Polar Res., 2: 55–70.
- KOSTELECKA-MYRCHA A. and MYRCHA A. 1980. Hematological studies on Antarctic birds. II. Changes of the hematological indices during the development of the pygoscelid penguins. — Pol. Polar Res. 1: 175–182.
- KOSTELECKA-MYRCHA A. and MYRCHA A. 1989. Change of the red blood picture during nesting development of Wilson's storm petrel (*Oceanites oceanicus* Kühl). — Pol. Polar Res. 10: 151–162.
- KRZEMIŃSKI W. 1981. Southern elephant seal (*Mirounga leonina* L.) of Admiralty Bay (King George Island, South Shetland Islands). Its numbers and activity during the moulting season in summer 1978/1979. — Pol. Polar Res., 2: 143–152.
- KRZYSZOWSKA A. 1990. The content of fuel oil in soil and effect of sewage on water nearby the H.Arctowski Polish Antarctic Station (Admiralty Bay, King George Island). — Pol. Arch. Hydrobiol., 37: 313–326.

- KULESZ J. 1994. Seasonal biology of *Notothenia gibberifrons*, *N. rossi* and *Trematomus newnesi*, as well as respiration of young fish from Admiralty Bay (King George, South Shetland Islands). — Pol. Arch. Hydrobiol., 41: 79–102.
- KULESZ J. and ZADRÓŻNY T. 1996. Biology of *Trematomus newnesi* Boulenger, 1902 (King George Island, Antarctica). — Pol. Arch. Hydrobiol., 43: 355–364.
- LESIŃSKI G. 1993. Monitoring of birds and pinnipeds on King George Island (South Shetland Islands) in 1989/1990. — Pol. Polar Res. 14: 75–90.
- LIGOWSKI R. 1985. Discrement of plant food of krill from the Admiralty Bay (King George Island). — XIIth Polar Symposium, Szczecin: 101–105.
- LIGOWSKI R. 1986. Net phytoplankton of the Admiralty Bay (King George Island, South Shetland Islands) in 1983. — Pol. Polar Res., 7: 127–154.
- LIGOWSKI R. 1987. Sea ice microalgae community of the floating ice in the Admiralty Bay (South Shetland Islands). — Pol. Polar Res., 8: 367–380.
- LIGOWSKI R. 1993a. Morskie okrzemki (Bacillariophyceae) w ekosystemie Antarktyki i ich znaczenie jako wskaźnika źródła pokarmu kryla (*Euphausia superba* Dana). — Wyd. Uniwersytetu Łódzkiego, Łódź: 242 pp.
- LIGOWSKI R. 1993b. Distribution of Antarctic diatoms in different habitats and its importance as food for krill. In: A. Leventer (ed.), Proceedings of the Fourth Polar Diatom Colloquium, Department of Quaternary Research, Stockholm, Sweden, August 24–28, 1992. — BPRC Misc. Series M-323, Byrd Polar Research Center, Columbus; 8–10.
- LIGOWSKI R. 1993c. 8. Sea ice algae. In: S. Rakusa-Suszczewski (ed.), *The Maritime Coastal Antarctic Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 49–51.
- LIGOWSKI R. 1993d. 9. Microphytobenthos. In: S. Rakusa-Suszczewski (ed.), *The Maritime Coastal Antarctic Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 53–56.
- LINKOWSKI T. B. and ŻUKOWSKI C. 1980. Observation on the growth of *Notothenia coriiceps neglecta* Nybelin and *Notothenia rossi marmorata* Fisher in Admiralty Bay (King George Island, South Shetland Islands). — Pol. Polar Res. 1: 155–162.
- LINKOWSKI T. B., PRESLER P. and ŻUKOWSKI C. 1983. Food habits of nototheniid fishes (Nototheniidae) in Admiralty Bay (King George Island, South Shetland Islands). — Pol. Polar Res., 4: 79–95.
- LIPSKI M. 1987. Variations of physical conditions and chlorophyll *a* contents in Admiralty Bay (King George Island, South Shetland Islands, 1979). — Pol. Polar Res., 8: 307–332.
- LIPSKI M. and RAKUSA-SUSZCZEWSKI S. 1990. Early summer pattern of vertical distribution of chlorophyll *a* (Bransfield Strait, Antarctica, November 1986). — Pol. Arch. Hydrobiol., 37: 287–293.
- LUŚCIŃSKA M. and KYĆ A. 1993. Algae inhabiting creeks of the region of “H. Arctowski” Polish Antarctic Station, King George Island, South Shetlands. — Pol. Polar Res. 14: 393–428.
- ŁUKOWSKI A. B. 1983a. DDT residues in the tissues and eggs of three species of penguins from breeding colonies at Admiralty Bay (King George Island, South Shetland Islands). — Pol. Polar Res., 4: 127–134.
- ŁUKOWSKI A. B. 1983b. DDT and its metabolites in the tissues and eggs of migrating Antarctic seabirds from the regions of the South Shetland Islands. — Pol. Polar Res., 4: 135–141.
- ŁUKOWSKI A. B. and LIGOWSKI R. 1987. Cumulation of chloroorganic insecticides by Antarctic marine diatoms. — Pol. Polar Res., 8: 167–177.
- ŁUKOWSKI A. B. and LIGOWSKI R. 1988. Contamination of Antarctic marine phytoplankton by chlorinated hydrocarbons (BIOMASS III). — Pol. Polar Res., 9: 399–408.
- ŁUKOWSKI A. B., KAROLEWSKI M. A. and GÓRSKI T. 1987. Polychlorinated biphenyls in the tissues of Antarctic marine migratory birds and penguins from the breeding colony on King George Island (South Shetland Islands). — Pol. Polar Res., 8: 179–187.

- MADEJSKI P. and RAKUSA-SUSZCZEWSKI S. 1990. Icebergs as tracers of water movement in Bransfield Strait. — *Antarctic Science*, 2: 259–263.
- MARSZ A. 1984. From surveys of the geomorphology of the shores and bottom of the Ezcurra Inlet. — *Oceanologia*, 15: 209–220.
- MARSZ A. and RAKUSA-SUSZCZEWSKI S. 1987. Ecological characteristics of the Admiralty Bay region (King George, South Shetland Islands). — *Kosmos*, 36: 103–127.
- MASSALSKI A., MROZIŃSKA T. and OLECH M. 1994. Ultrastructure of *Lobosphaera reniformis* (Watanabe) Komarek et Tott (Chlorellales) from King George Island, South Shetland Islands, Antarctica. — *Acta Soc. Bot. Pol.*, 63: 205–210.
- MASSALSKI A., MROZIŃSKA T. and OLECH M. 1995. *Lobococcus irregularis* (Boye-Pet.) Reisigl var. *antarcticus* var. nov., from King George Island, South Shetland Islands, Antarctica, and its ultrastructure. — *Nova Hedwigia*, 61: 199–206.
- MCCLATCHIE S., RAKUSA-SUSZCZEWSKI S. and FILCEK K. 1991. Seasonal growth and mortality of *Euphausia superba* in Admiralty Bay, South Shetland Islands, Antarctica. — *ICES J. mar. Sci.*, 48: 335–342.
- MENSHENINA L. and RAKUSA-SUSZCZEWSKI S. 1992. Zooplankton changes during the year in Admiralty Bay (February 1990–January 1991). — *Pol. Arch. Hydrobiol.*, 39: 49–58.
- MOCZYDŁOWSKI E. 1986. Microclimate of the nest-sites of pygoscelid penguins (Admiralty Bay, South Shetland Islands). — *Pol. Polar Res.*, 7: 377–394.
- MOCZYDŁOWSKI E. 1989. Protection of eggs and chicks against flooding as a part of nesting strategy of pygoscelid penguins at King George Island, South Shetlands. — *Pol. Polar Res.* 10: 163–182.
- MROZIŃSKA T., OLECH M. and MASSALSKI A. 1986. Algae of ponds and stream on moraines of the Ecology Glacier (King George Island, South Shetlands, Antarctica. In: M. Olech (ed.), *Antarctic Research on Taxonomy and Ecology of Algae*. — Proc. Workshop, Cracow; 14.
- MYRCHA A. 1993. 14. Birds. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 161–165.
- MYRCHA A. and KAMIŃSKI P. 1982. Changes in body calorific values during nestling development of penguins of the genus *Pygoscelis*. — *Pol. Polar Res.*, 3: 81–88.
- MYRCHA A. and KOSTELECKA-MYRCHA A. 1980a. Hematological studies on Antarctic birds. 1. Hematological indices in some species of the birds studied during austral summer. — *Pol. Polar Res.* 1: 169–174.
- MYRCHA A. and KOSTELECKA-MYRCHA A. 1980b. Blood picture in some species of Antarctic birds. — *Bull. Pol. Acad. Sci.*, CI II, 27: 911–915.
- MYRCHA A. and TATUR A. 1991. Ecological role of current and abandoned penguin rookeries in the land environment of the maritime Antarctic. — *Pol. Polar Res.* 12: 3–24.
- MYRCHA A. and TELIGA K. 1980. Observation of pinnipedian mammals in the vicinity of Arctowski Station (King George Island) in 1978. *Pol. Polar Res.* 1: 117–126.
- MYRCHA A., PIETR S. J. and TATUR A. 1985. The role of Pygoscelid penguin rookeries in nutrient cycles at Admiralty Bay, King George Island. In: W.R. Siegfried, P.R. Condry and R.M. Laws (eds.), *Antarctic Nutrient Cycle and Food Webs*. — Springer Verlag Berlin–Heidelberg 1985: 156–162.
- NIEDBAŁA W. 1986. Several species of moss-mites (Acari, Oribatida) of the Antarctic coastal zone. — *Pol. Polar Res.*, 7: 119–126.
- OCHI H. and OCHYRA R. 1985. *Bryum dichotomum* Hedw., a species new to the Antarctic. — *Lindbergia*, 11: 157–160.
- OCHYRA R. 1985. On the Antarctic species of the family Orthotrichaceae. — *Lindbergia*, 11: 141–146.
- OCHYRA R. 1987. *Schistidium steerei* (Grimmiaceae) a Remarkable New Species from the Antarctic, with a Note on *S. obtusifolium*. — *Mem. New York Bot. Gard.*, 45: 607–614.

- OCHYRA R. 1990. The discovery of the South Georgian endemic species *Schistidium urnulaceum* (C.Muell.) B.G.Bell. (Musci: Grimmiaceae) in the Antarctic. — Pol. Polar Res., 11: 133–146.
- OCHYRA R. and AFONINA O. M. 1986. The taxonomic position and geographical distribution of *Grimmia andraeopsis* C. Muell. (Grimmiaceae, Musci). — Pol. Polar Res., 7: 319–331.
- OCHYRA R. and BELL B. G. 1984. A record of *Schistidium falcatum* (Bryophyta; Musci) from the Antarctic. — Brit. Antarct. Surv. Bull., 64: 77–79.
- OCHYRA R. and LEWIS-SMITH R. I. 1996. Antipodal mosses: VI. *Stegonia latifolia* (Musci, Pottiaceae) in the Antarctic. — Fragm. Flor. Geobot., 41: 985–994.
- OCHYRA R. and OCHI H. 1986. New or otherwise interesting species of the genus *Bryum* (Musci, Bryaceae) in the Antarctic. — Acta Bot. Hung., 32: 209–219.
- OCHYRA R. and VANA J. 1989. The hepatics of King George Island, South Shetland Islands, Antarctica, with particular reference to the Admiralty Bay region. — Pol. Polar Res. 10: 183–210.
- OLECH M. 1989. Lichens from the Admiralty Bay region, King George Island (South Shetland Islands, Antarctica). — Acta Soc. Bot. Pol., 58: 493–512.
- OLECH M. 1990. Preliminary studies on ornithocoprophilous lichens of the Arctic and Antarctic regions. — Proc. NIPR Symp. Polar Biol. Tokyo, 3: 218–23.
- OLECH M. 1991a. *Japewia tornoensis* (Lichenes), a species new to the Antarctic. — Pol. Bot. Stud., 2: 217–218.
- OLECH M. 1991b. Differentiation and ecological character of vegetation of Western Antarctica (King George Island). — Antarctic Science – Global Concerns. Abstr. p.108.
- OLECH M. 1991c. Preliminary observations on the content of heavy metals in thalli of *Usnea antarctica* Du Rietz. Lichenes in the vicinity of the H. Arctowski Polish Antarctic Station. — Pol. Polar Res. 12: 129–131.
- OLECH M. 1994. Apofity we florze Sztetlandów Południowych (Antarktyka). In: J. Kiszka and J. Piórecki (eds.), *Porosty apofityczne jako wynik antropopresji*. — Mat. Symp.: 31–37.
- OLECH M. 1996a. Human impact on terrestrial ecosystems in West Antarctica. — Proc. NIPR Symp., Polar Biol., 9: 299–306.
- OLECH M. 1996b. Anthropogenic causes of colonisation of terrestrial biotope in West Antarctica. — 23 Polar Symposium, Sosnowiec; 173–174.
- OLECH M. and ALSTRUP V. 1989. *Thelocarpon cyaneum* sp.nov. — Nord. J. Bot., Copenhagen, 9: 575–576.
- OLECH M. and ALSTRUP V. 1995. Grzyby pasożytujące na plechach porostów w rejonach polarnych. — 22 Symp. Polar., Zamek Książ; 119–122.
- OLECH M., DUTKIEWICZ E. M., KWIATEK W. M., KAJFOSZ J. and SZYMCHYK S. 1994. Local Lead Pollution in the Antarctic, Annual Report, Institute of Nuclear Physics, Cracow; 37–38.
- OLECH M. and SOCHTING U. 1993. Four new species of *Caloplaca* from Antarctica. — Lichenologist, 25: 261–269.
- OLECH M., SZYMCHYK S. and KAJFOSZ J. 1993. Lokalne zanieczyszczenia ołowiem środowiska w rejonie Antarktyki. — Prace Mineralogiczne, 83: 51–54.
- PECHERZEWSKI K. 1980. Distribution and quantity of suspended matter in Admiralty Bay (King George Island, South Shetland Islands). — Pol. Polar Res. 1: 75–82.
- PIETR S. J. 1986. The physiological groups of microorganisms in different soils at Admiralty Bay region (King George Island, South Shetland Islands, Antarctica). — Pol. Polar Res., 7: 395–406.
- PIETR S. J. 1993. Soil microorganisms. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 166–172.
- PRESLER P. 1980. Phenological and physiological observations carried out during the first wintering at the Arctowski Station in 1977. — Pol. Arch. Hydrobiol., 27: 245–252.
- PRESLER P. 1986. Necrophagous invertebrates of the Admiralty Bay of King George Island (South Shetland Islands, Antarctica). — Pol. Polar Res., 7: 25–61.

- PRESLER P. 1993a. 12. Zoobenthos. 12.2. Necrophagous invertebrates. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 96–100.
- PRESLER P. 1993b. 12. Zoobenthos. 12.5. Ophiuroidea. In: S. Rakusa-Suszczewski S. (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 117–119.
- PRESLER P. and FIGIELSKA E. 1998 (1997). New data on the Asteroidea of Admiralty Bay, King George Island, South Shetland Islands. — *Pol. Polar Res.* 18: 107–117.
- PRUSZAK Z. 1980. Currents circulation in the waters of Admiralty Bay (region of Arctowski Station on King George Island). — *Pol. Polar Res.* 1: 55–74.
- RAKUSA-SUSZCZEWSKI S. 1980a. Environmental conditions and the functioning of Admiralty Bay (South Shetland Islands) as part of the near shore Antarctic ecosystem. — *Pol. Polar Res.* 1: 11–27.
- RAKUSA-SUSZCZEWSKI S. 1980b. The role of near-shore research in gaining an understanding of the functioning of Antarctic ecosystem. — *Pol. Arch. Hydrobiol.*, 27: 229–233.
- RAKUSA-SUSZCZEWSKI S. 1987. The matter transport in the near-shore ecosystem of the Admiralty Bay (King George, South Shetland Islands). — *Colloque sur l'écologie marine des les subantarctiques et antarctiques*, Paris, 25 juin 1985, CNFRA: 7–15.
- RAKUSA-SUSZCZEWSKI S. 1990. Seasonal changes in respiration and biomass of *Euphausia superba* Dana from Admiralty Bay (South Shetland Islands). — *Pol. Arch. Hydrobiol.*, 37: 305–311.
- RAKUSA-SUSZCZEWSKI S. 1992. Respiration of *Nacella concinna* Strebel, 1908 from the intertidal zone of Admiralty Bay (South Shetland Islands, Antarctica). — *Pol. Arch. Hydrobiol.*, 39: 205–209.
- RAKUSA-SUSZCZEWSKI S. (ed.), 1993a. *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 216 pp.
- RAKUSA-SUSZCZEWSKI S. 1993b. 1. Introduction. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 7–14.
- RAKUSA-SUSZCZEWSKI S. 1993c. 5. Marine environment. 5.1. Water exchange. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 31–32.
- RAKUSA-SUSZCZEWSKI S. 1993d. 5. Marine environment. 5.2. Hydrography and hydro-chemistry. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 32–34.
- RAKUSA-SUSZCZEWSKI S. 1993e. 5. Marine environment. 5.3. Mineral suspension. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 34–35.
- RAKUSA-SUSZCZEWSKI S. 1993f. 23. Matter transport and budgets in the Admiralty Bay area. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 199–212.
- RAKUSA-SUSZCZEWSKI S. 1995a. The hydrography of Admiralty Bay and its inlets, coves and lagoons (King George Island, Antarctica), *Pol. Polar Res.* 16: 61–70.
- RAKUSA-SUSZCZEWSKI S. 1995b. Flow of matter in the Admiralty Bay area, King George Island, Maritime Antarctic. — *Proc. NIPR Symp. Polar Biol.*, 8: 101–113.
- RAKUSA-SUSZCZEWSKI S. 1996. Spatial and seasonal variability of temperature and salinity in Bransfield Strait and Admiralty Bay, Antarctica. — *Pol. Polar Res.* 17: 29–42.
- RAKUSA-SUSZCZEWSKI S. and FILCEK K. 1988. Protozoa on the body of *Euphausia superba* Dana from Admiralty Bay (the South Shetland Islands). — *Acta Protozool.*, 27: 23–29.
- RAKUSA-SUSZCZEWSKI S. and KRZYSZOWSKA A. 1991. Assessment of the environmental impact of the "H. Arctowski" Polish Antarctic Station (Admiralty Bay, King George Island, South Shetland Islands). — *Pol. Polar Res.* 12: 105–121.

- RAKUSA-SUSZCZEWSKI S. and LACH A. 1991. Respiration of *Orchomene plebs* (Hurley, 1965) and *Waldeckia obesa* (Chevreux, 1905) from Admiralty Bay (South Shetland Islands, Antarctic). — *Hydrobiologia*, 223: 177–180.
- RAKUSA-SUSZCZEWSKI S. and NEMOTO T. 1989. Ciliates associations on the body of krill *Euphausia superba* Dana. — *Acta Protozool.*, 28: 77–86.
- RAKUSA-SUSZCZEWSKI S. and SIERAKOWSKI K. 1993. Pinnipeds in Admiralty Bay, King George Island, South Shetlands (1988–1992). — *Pol. Polar Res.* 14: 439–453.
- RAKUSA-SUSZCZEWSKI S. and STEPNIK R. 1982. Three species of krill from Admiralty Bay (King George Island, South Shetlands) in summer 1978/79. — *Pol. Arch. Hydrobiol.*, 27: 273–284.
- RAKUSA-SUSZCZEWSKI S. and ZDANOWSKI M. 1989. Bacteria in krill (*Euphausia superba* Dana) stomach. — *Acta Protozool.*, 28: 87–90.
- RAKUSA-SUSZCZEWSKI S., BATCKE Z. and CISAK J. 1993a. 4. Morphometry of the Admiralty Bay shores and basin. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 27–30.
- RAKUSA-SUSZCZEWSKI S., MIETUS M. and PIASECKI J. 1993b. 3. Weather and climate. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 19–25.
- SAMP R. 1980. Selected environmental factors in the waters of Admiralty Bay (King George Island, South Shetland Islands). — *Pol. Polar Res.* 1: 53–66.
- SARUKHANYAN E. I. and TOKARCZYK R. 1988. Coarse scale hydrological condition in Admiralty Bay in summer 1982 (King George Island, South Shetlands). — *Pol. Polar Res.*, 9: 121–132.
- SCHROETER B., OLECH M., KAPPEN L. and HEITLAND W. 1995. Ecophysiological investigations of *Usnea antarctica* in the maritime Antarctic. I Annual microclimatic conditions and potential primary production. — *Antarctic Science*, 7: 251–260.
- SICIŃSKI J. 1986. Benthic assemblages of Polychaeta in chosen regions of the Admiralty Bay (King George Island, South Shetland Islands). — *Pol. Polar Res.*, 7: 63–78.
- SICIŃSKI J. 1993. 12. Zoobenthos. 12.3. Polychaeta. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 101–107.
- SICIŃSKI J. and JANOWSKA E. 1993. Polychaetes of the shallow sublittoral of Admiralty Bay, King George Island, South Shetlands. — *Antarctic Science*, 5: 161–167.
- SIERAKOWSKI K. 1991. Birds and mammals in SSSI No.8 in the season 1988/1989 (South Shetland Islands, King George Island, Admiralty Bay). — *Pol. Polar Res.* 12: 25–54.
- SKÓRA K. E. 1993. 13. Fish. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 123–128.
- SKÓRA K. E. 1995. *Acanthodraco dewitti* gen. et sp. n. (Pisces, Bathydraconidae) from Admiralty Bay (King George Island, South Shetland Islands, Antarctica). — *Arch. Fish. Mar. Res.*, 3: 283–289.
- SKÓRA K. E. and NEYLOV A. V. 1992. Fish of Admiralty Bay (King George Island, South Shetland Islands, Antarctica). — *Polar Biol.* 12: 469–476.
- SOCHTING U. and OLECH M. 1995. The lichen genus *Caloplaca* in polar regions. — *Lichenologist*, 27: 463–471.
- STANKOVIĆ A. M. and RAKUSA-SUSZCZEWSKI S. 1996. Parasitic Protozoa on appendages and inside the body of *Euphausia superba* Dana. — *Pol. Polar Res.* 17: 169–172.
- STAWISZYŃSKA-JANAS M. and KITTEL W. 1982. Epizoiczne sysydłaczki (Protozoa, Suctorida) na *Euphausia superba* Dana i *Euphausia crystallorophias* Holt et Tattersall. — *Polskie Badania Polarne 1970–1982*, Toruń 1983: 239–248.
- STEPNIK R. 1982. All-year populational studies of Euphausiacea (Crustacea) in the Admiralty Bay (King George Island, South Shetland Islands, Antarctic). — *Pol. Polar Res.*, 3: 49–68.
- STYSZYŃSKA A. 1990. The effect of wind direction and orography on air temperature at the “Arctowski” Station. — *Pol. Polar Res.* 11: 69–93.

- SZAFRAŃSKI J. and LIPSKI M. 1982. Characteristics of water temperature and salinity at Admiralty Bay (King George Island) during austral summer 1978–1979. — *Pol. Polar Res.*, 3: 7–24.
- SZYMCZYK S., OLECH M. and KAJFOSZ J. 1991. Air pollution studies by PIXE analysis of lichens. — Annual Report, Institute of Nuclear Physics, Cracow; 145–146.
- TATUR A. 1987. Fluorine in ornithogenic soils and minerals on King George Island, West Antarctica. — *Pol. Polar Res.*, 8: 65–74.
- TATUR A. 1989. Ornithogenic soils of the maritime Antarctic. — *Pol. Polar Res.* 10: 481–532.
- TATUR A. and BARCZUK A. 1984. Phosphates of ornithogenic soils on the volcanic King George Island (Maritime Antarctic). — *Pol. Polar Res.*, 5: 61–97.
- TATUR A. and BARCZUK A. 1985. Ornithogenic phosphates on King George Island in the maritime Antarctic Zone. In: W.R. Siefried, P.R. Condy & R.M. Laws (eds.), *Antarctic Nutrient Cycle and Food Webs*. — Springer Verlag, Berlin — Heidelberg 1985: 163–168.
- TATUR A. and MYRCHA A. 1983. Changes in chemical composition of waters running off from the penguin rookeries in Admiralty Bay region (King George Island, South Shetland Islands, Antarctica). — *Pol. Polar Res.*, 4: 113–125.
- TATUR A. and MYRCHA A. 1984. Ornithogenic soils on King George Island, South Shetland Islands (Maritime Antarctic Zone). — *Pol. Polar Res.*, 5: 31–60.
- TATUR A. and MYRCHA A. 1989. Soils and vegetation in abandoned penguin rookeries. — *Proc. NIPR Symp. Polar Biol.*, 2: 181–189.
- TATUR A. and MYRCHA A. 1993. 17. Ornithogenic soils. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem in Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 161–165.
- TATUR A., MYRCHA A. and NIEGODZISZ J. 1997. Formation of abandoned penguin rookery ecosystems in the maritime Antarctic. — *Polar Biol.* 17: 405–417.
- TOKARCZYK R. 1986. Annual cycle of chlorophyll *a* in Admiralty Bay, 1981–1982 (King George, South Shetlands). — *Pol. Arch. Hydrobiol.*, 33: 177–188.
- TOKARCZYK R. 1987. Classification of water masses in the Bransfield Strait and southern part of the Drake Passage using a method of statistical multidimensional analysis. — *Pol. Polar Res.*, 8: 333–366.
- TRIVELPIECE S. G., GEUPEL G.R., KJELMYR J., MYRCHA A., SICIŃSKI J., TRIVELPIECE W.Z. and VOLKMAN N.J. 1987. Rare bird sightings from Admiralty Bay, King George Island, South Shetlands, Antarctica, 1976–1987. — *Cormorant*, 15: 59–66.
- WASIK A. and MIKOŁAJCZYK E. 1993. 11. Zooplankton. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 61–64.
- WASIK A. and MIKOŁAJCZYK E. 1994. Annual cycle of tintinnids in Admiralty Bay with an emphasis on seasonal variability in *Cymatocyclis affinis* / *Convallaria lorica* morphology. — *J. Plankton Res.* 16: 1–8.
- WASILEWSKI A. 1986. Ecological aspects of the breeding cycle in the Wilson's storm petrel, *Oceanites oceanicus* (Kuhl) at King George Island (South Shetland Islands, Antarctic). — *Pol. Polar Res.*, 7: 173–216.
- WOJCIECHOWSKA A. 1993a. The tetraphyllidean and tetrabothriid cercoids from Antarctic bony fishes. I. Morphology. Identification with adult forms. — *Acta Parasitologica*, 38: 15–22.
- WOJCIECHOWSKA A. 1993b. The tetraphyllidean and tetrabothriid cercoids from Antarctic bony fishes. II. Occurrence of cercoids in various fish species. — *Acta Parasitologica*, 38: 113–118.
- WOJCIECHOWSKA A. 1993c. The tetraphyllidean and tetrabothriid cercoids from Antarctic bony fishes. III. Infection of *Notothenia neglecta* and *N. rossii* in Admiralty Bay. — *Acta Parasitologica*, 38: 166–169.
- WOJCIECHOWSKA A. and ZDZITOWIECKI K. 1995. Cestodes of Antarctic seals. — *Acta Parasitologica*, 40: 125–131.

- WOYCIECHOWSKI M. 1980. Numbers and activity of Weddell seal (*Leptonychotes weddelli* Lesson) in the Admiralty Bay (King George Island, South Shetland Islands) in summer 1978/1979. — Pol. Polar Res. 1: 207–216.
- WOŹNIAK B., HAPTER R. and MAJ B. 1984. The inflow of solar energy and the irradiance of the euphotic zone in the region of Ezcurra Inlet during the Antarctic summer of 1977/78. — Oceanologia, 15: 141–170.
- ZABAWSKI J. 1995. Grzyby z rodzaju *Penicillium* w niektórych glebach rejonu Wyspy Króla Jerzego (Szetlandy Półn., Antarktyka). — XXII Sympozjum Polarne, Wrocław-Książ, 27–28.XI.1995; 143–149.
- ZABAWSKI J. and PIASECKI J. 1981. Studies of soil microflora of King George Island. In: E. Maliszewska (ed.), Microbiological transformations of nitrogen compounds in soil under various ecological conditions. — Institute of Cultivation, Fertilization and Pedology, Puławy; 131–138.
- ZADRÓŻNY T. 1996. Fishes of Admiralty Bay caught in 1994 and 1995 (King George Island, South Shetland Islands, Antarctica). — Pol. Arch. Hydrobiol., 43: 347–354.
- ZARZYCKI K. 1993. 20. Vascular Plants and terrestrial biotopes. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 180–187.
- ZDANOWSKI M. K. 1981. Growth of bacteria in the course of decomposition of *Euphausia superba* Dana. — Bull. Acad. Pol. Biol., 29: 155–161.
- ZDANOWSKI M. K. 1988. Matter conversion in the course of krill (*Euphausia superba* Dana) decomposition in the Antarctic ecosystem. Part 1. — Pol. Arch. Hydrobiol., 35: 65–96.
- ZDANOWSKI M. K. 1995. Characteristics of bacteria in selected Antarctic marine habitats. In: S. Rakusa-Suszczewski and S. P. Donachie (eds.), *Microbiology of Antarctic marine environments and krill intestine, its decomposition and digestive enzymes*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 7–100.
- ZDANOWSKI M. K. and DONACHIE S. P. 1993. 6. Bacterioplankton. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 39–44.
- ZDZITOWIECKI K. 1978. Preliminary studies of fish parasites in the areas of South Shetland Islands and Southern Georgia (Antarctica). — Kosmos, Ser. A, 27: 651–659.
- ZDZITOWIECKI K. 1979. Digenetic trematodes in alimentary tracts of fishes of South Georgia and South Shetlands (Antarctica). — Acta Ichthyol. Piscat., 9: 15–31.
- ZDZITOWIECKI K. 1985. Acanthocephalans of birds from South Shetlands (Antarctic). — Acta Parasitol. Pol., 30: 11–24.
- ZDZITOWIECKI K. 1986. Acanthocephala of the Antarctic. — Pol. Polar Res., 7: 79–117.
- ZDZITOWIECKI K. 1986b. Prevalence of acanthocephalans in fishes of South Shetlands (Antarctic). I. Juvenile *Corynosoma* spp. — Acta Parasitol. Pol., 30: 143–160.
- ZDZITOWIECKI K. 1986c. Prevalence of acanthocephalans in fishes of South Shetlands (Antarctic). III. *Metacanthocephalus johnstoni* Zdzitowiecki, 1983; *M. dalmori* Zdzitowiecki, 1983 and notes on other species; general conclusions. — Acta Parasitol. Pol., 31: 125–141.
- ZDZITOWIECKI K. 1987. Digenetic trematodes from the alimentary tracts of fishes off South Shetlands (Antarctic). — Acta Parasitol. Pol., 33: 155–167.
- ZDZITOWIECKI K. 1991. Antarctic Acanthocephala. In: J.W. Wägele and J. Siegg (eds.), *Synopses of the Antarctic Benthos*, vol. 3. — Koeltz Scientific Books, Koenigstein, 116 pp.
- ZDZITOWIECKI K. 1997. Antarctic Digenea, Parasites of fishes. — In: J.W. Wägele and J. Siegg (eds.), *Synopses of the Antarctic Benthos*, vol. 8. Koeltz Scientific Books, Koenigstein. 156 pp.
- ZDZITOWIECKI K. and CIELECKA D. 1996. Morphology and occurrence of *Dichelyne* (*Cucullanelus*) *fraseri* (Baylis, 1929), a parasitic nematode of Antarctic and sub-Antarctic fishes. — Acta Parasitologica, 41: 30–37.

- ZDZITOWIECKI K. and DRÓZDŹ J. 1980. Redescription of *Stegophorus macronektes* (Johnston et Mawson, 1942) and description of *Stegophorus arctowskii* sp.n. (Nematoda, Spirurida) from birds of South Shetlands (the Antarctic). — *Acta Parasitol. Pol.*, 27: 205–212.
- ZDZITOWIECKI K. and ROKOSZ B. 1986. Prevalence of acanthocephalans in fishes of South Shetlands (Antarctic). II. *Aspensertis austrinus* Van Cleave, 1929 and remarks on the validity of *Heteracanthocephalus hureaui* Dollfus, 1965. — *Acta Parasitol. Pol.*, 30: 161–171.
- ZDZITOWIECKI K., NIEWIADOMSKA K. and DRÓZDŹ J. 1989. Trematodes of birds and mammals in the environs of H.Arctowski Station (South Shetlands, Antarctic) — *Acta Parasitol. Pol.*, 34: 247–257.
- ZIELIŃSKI K. 1981. Benthic macroalgae of Admiralty Bay (King George Island, South Shetland Islands) and circulation of algae matter between the water and the shore. — *Pol. Polar Res.*, 2: 71–94.
- ZIELIŃSKI K. 1990. Bottom macroalgae of Admiralty Bay (King George Island, South Shetland Islands, Antarctica). — *Pol. Polar Res.* 11: 95–131.
- ŻMIJEWSKA M. I. 1993. 11. Zooplankton. 11.2. Copepoda. In: S. Rakusa-Suszczewski (ed.), *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. — Department of Antarctic Biology, Pol. Acad. Sci., Warsaw; 64–66.
- ŻUKOWSKI C. 1980. Catches of fishes of the genus *Notothenia* and *Trematomus* at Admiralty Bay (King George Island, South Shetland Islands) in the winter-spring season 1977. — *Pol. Polar Res.* 1: 163–167.