

# Differential shuffling of native genetic diversity across introduced regions in a brown alga: Aquaculture vs. maritime traffic effects

Marie Voisin, Carolyn R. Engel, and Frédérique Viard\*

Equipe Evolution et Génétique des Populations Marines, Unité Mixte de Recherche, Centre National de la Recherche Scientifique/Université Pierre et Marie Curie 7144, Station Biologique de Roscoff, Place Georges-Teissier, BP 74, 29682 Roscoff Cedex, France

Communicated by Juan Carlos Castilla, Pontificia Universidad Católica de Chile, Santiago, Chile, March 3, 2005 (received for review October 1, 2004)

**Worldwide marine invaders, such as the brown alga *Undaria pinnatifida*, offer challenging models for unraveling the apparent paradox of sustainable settlement of exotic species over a large spectrum of environments. Two intergenic noncoding mitochondrial loci were found to be highly informative at the within-species level. Twenty-five haplotypes were found over the whole dataset (333 base pairs, 524 individuals, and 24 populations). The native range showed striking population genetic structure stemming from low diversity within and high differentiation among populations, a pattern not observed in the introduced range of this seaweed. Contrary to classical expectations of founding effects associated with accidental introduction of exotic species, most of the introduced populations showed high genetic diversity. At the regional scale, genetic diversity and sequence divergence showed contrasting patterns in the two main areas of introduction (Europe and Australasia), suggesting different processes of introduction in the two regions. Gene genealogy analyses point to aquaculture as a major vector of introduction and spread in Europe but implicate maritime traffic in promoting recurrent migration events from the native range to Australasia. The multiplicity of processes and genetic signatures associated with the successful invasion confirms that multiple facets of global change, e.g., aquaculture practices, alteration of habitats, and increased traffic, act in synergy at the worldwide level, facilitating successful pandemic introductions.**

coastal habitats | gene genealogy | introduced species | kelp | mitochondrial DNA

**B**iological invasions are closely linked to global change (1–3). The spread and success of exotic species may be facilitated by (i) climate change (4), (ii) increase in disturbed and/or anthropogenic habitats (5, 6), and (iii) synergistic effects resulting from increased biotic exchanges (7, 8). In turn, biological invasions may feed back into global change (9) by modifying interactions among species and even whole ecosystems (10, 11), inducing extinction in the native fauna/flora (12, 13) and homogenizing global diversity (14, 15). Whether consequence or cause of global change, the range expansion of exotic species depends on defying the barriers of natural dispersal (16, 17) and on adaptation to a novel local abiotic and biotic environment in which the organism did not evolve (6).

In the marine environment, where the rate of introduction of alien species has dramatically increased over the past century (e.g., refs. 18 and 19), ship-ballasting procedures, increased volume of transoceanic shipping and intracoastal boating, and modern aquaculture practices all contribute to the establishment of veritable invasion corridors (20, 21). Likewise, the propagule pressure hypothesis (*sensu* Williamson, ref. 17, p. 45; and ref. 6) postulates that, even if preceded by many failed attempts, the recurrence of introductions amplifies the very low initial probability of successful establishment. Furthermore, successive migration events may also increase overall genetic diversity, thereby improving the likelihood of encountering adaptive genetic variants and the rate of adaptive evolution (6, 22–24).

Successful settlement of many marine invaders may indeed be explained by recurrent introductions (16). Alternatively, successful introductions of exotic species may be linked to changes in the recipient region (2, 25). For example, in the marine environment, many successful invaders have settled in anthropogenic habitats, such as marinas, commercial shipping ports, shellfish farms, or heavily disturbed/polluted natural areas that impose novel ecological conditions even for the native species community (6). In this context, the widely introduced brown alga *Undaria pinnatifida* provides an excellent opportunity for exploring the mechanisms and processes behind the apparent paradox (6) of rapid adaptation of exotic species to a wide spectrum of environments, competing with native species that have naturally evolved there.


*U. pinnatifida* (Laminariales, Alariaceae), otherwise known as the edible seaweed *wakame*, is an Asian species native to Japanese, Korean, and Chinese waters, where it is extensively cultivated. The first observation of this species outside its native Asian range was recent (1971, in the Mediterranean Sea). Since then, this kelp has been introduced to both Northern and Southern hemispheres, spreading to the Atlantic, Mediterranean, Adriatic, and North Sea coasts of Europe (see ref. 26); New Zealand (27); Australia (28); Argentina (29); and, most recently, to the American Pacific coast (California, ref. 26; Mexico, ref. 30). The patterns and intensity of invasion of *U. pinnatifida* vary around the world, ranging from recent rapid long-distance spread in Californian harbors (31); slow long-distance spread in European marinas, concomitant with the presence of *wakame* farms in Brittany, France (32); and heavy proliferation and naturalization in both harbors and natural subtidal zones in Australasia (28, 33, 34) and Argentina (35). The rapid spread of the species within regions of introduction is thought to be linked to its biological characteristics (i.e., annual life history, high growth rates, and high fecundity), typical of pioneer species (36). Because this kelp can be both accidentally transported by various vectors such as hull fouling, ballast water, and shellfish-farming transplants and also deliberately introduced for aquaculture purposes, the source, number, and diversity of introductions may contribute to the various degrees of successful settlement around the world (11, 25).

Despite increased public awareness and, in some cases, legislative measures, until now the few empirical studies of *U. pinnatifida* have concentrated on spread and local ecological dynamics (26, 31, 33). However, phylogeographic analyses lend insight into the mechanisms of introduction, colonization, and spread of invaders (23). Although the mitochondrial genome of a kelp has been entirely sequenced (37), no gene genealogy studies have been attempted in brown algae at the population

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. AY821889–AY821906).

\*To whom correspondence should be addressed. E-mail: viard@sb-roscoff.fr.

© 2005 by The National Academy of Sciences of the USA



**Fig. 1.** Distribution of the 25 mtDNA haplotypes detected in the 24 wild and cultivated *U. pinnatifida* populations sampled around the world. Color-motif pie charts display relative frequencies of each haplotype in each population. Foreground colors of the pie charts refer to the clusters defined as in Fig. 2: red, cluster A; red and blue, cluster B1; and green, cluster B2. The international code of the country of origin of each population is indicated in parentheses.

level. Here, we use two polymorphic mitochondrial markers (i.e., intergenic spacer regions) to compare the diversity, sequence divergence (gene genealogies), and patterns of invasion among regions of introduction of native, cultivated, and spontaneous introduced populations of *U. pinnatifida* from around the world.

### Materials and Methods


Samples representing the present-day distribution of *U. pinnatifida* were collected from five natural populations located in the native range (Japan, Korea, and China), 16 natural populations in all documented (before 2004) regions of introduction (Europe, mainland Australia, Tasmania, New Zealand, Argentina, and California; Fig. 1; details in Table 2, which is published as supporting information on the PNAS web site), and cultivated populations from Japan and Korea (commercial specimens) and in France (sampled directly from the farm). In each population, 23–24 individuals were collected, but commercial specimens comprised fewer individuals. Excised thallus tissue ( $\approx 5$  cm<sup>2</sup>) was dried in silica gel before DNA analysis. Total DNA was extracted from  $\approx 5$  mg of dried tissue by using the Nucleospin Multi96 Plant extraction kit (Macherey & Nagel) according to the manufacturer's instructions. Using primers designed from the complete sequence of the mitochondrial genome of *Laminaria digitata* (Laminariales, Laminariaceae) (37), two mitochondrial intergenic spacer regions were PCR-amplified: (i) between the *atp8* and *trnS* genes and (ii) between the *trnW* and *trnI* genes, hereafter called *atp8-S* and *W-I*, respectively. Primers were designed in highly conserved coding regions flanking each spacer. For *atp8-S*, 239 base pairs were amplified by using *atp8-trnS-F* (3' end of *atp8*, 5'-TGTACGTTTCATATTACCTTCTTTAGC-3') and *atp8-trnS-R* (5' end of *trnS*, 5'-TAGCAAACCAAGGCTTTCAAC-3') primers and for *W-I*, 292 base pairs were amplified by using *trnW-trnI-F* (3' end of *trnW*, 5'-GGGGTTCAAATCCCTCTCTT-3') and *trnW-trnI-R* (5' end of *trnI*, 5'-CCTACATTGTTAGCTTCATGAGAA-3') primers.

For both loci, PCRs were carried out in 20  $\mu$ l containing 5  $\mu$ l of sample DNA diluted 1:100/0.1 mg/ml BSA/2.5 mM MgCl<sub>2</sub>/75

mM Tris-HCl/20 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>/200  $\mu$ M dNTPs/250 nM each primer/0.5 unit Thermoprime Plus DNA polymerase (Abgene, Epsom, U.K.) and run on a MJ Research (Cambridge, MA) PTC200 thermocycler. After an initial denaturation step (95°C, 5 min), touchdown PCR was carried out for five cycles of 30 s at 95°C, 30 s at 60°C reduced by 1°C each subsequent cycle, and 45 s at 72°C, followed by 30 cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 45 s.

To detect single-site substitutions at the *atp8-S* locus, all PCR products were sequenced in both directions by using the amplification primers on an Applied Biosystems PRISM 3100 Automated DNA Sequencer (PE Applied Biosystems). For *W-I*, a more efficient screening strategy was developed by using single-stranded DNA conformation polymorphism (SSCP) analysis (38). PCRs were performed by using a modified forward primer (*trnW-trnI-F2*, 5'-ATGAGGGTAAGTTTTTCACAGT-3'), designed based on preliminary sequencing of seven individuals. To check the accuracy of SSCP typing, an average of nine individuals possessing each distinct *W-I* SSCP profile were sequenced. Each SSCP profile was found to have the same unique sequence, and some differed by as little as 1 base pair.

Sequences were aligned by using CLUSTALW (39) in BIOEDIT (40). No ambiguous base compositions were encountered, suggesting that heteroplasmy is absent in *Undaria* (see ref. 41). Due to the nonindependence of the two loci ( $D' = 0.77$ ,  $P < 0.001$ ), statistical analyses were carried out on concatenated intergenic sequences with a total length of 333 bp (see Table 3, which is published as supporting information on the PNAS web site). Estimates of haplotypic ( $H_c$ ) and nucleotide ( $\pi$ ) diversity were calculated for each population, within three main regional groups [native region (Far-East Asia, i.e., Japanese, Korean, and Chinese populations); Europe, Australasia (mainland Australia, Tasmania, and New Zealand)]; and for the entire dataset by using ARLEQUIN v. 2.0 (<http://anthropologie.unige.ch/arlequin>). This software was also used to carry out an analysis of molecular variance by computing  $\Phi$  statistics, analogous to Wright's fixation indices,  $F$  statistics, and exact tests of differentiation to investigate population structure (42) (i) among populations within each region, (ii) among and within



**Fig. 2.** Haplotypic network and regional distribution of the 25 *U. pinnatifida* mtDNA haplotypes. Circle sizes are proportional to the number of individuals observed for each haplotype, and color pie charts indicate the distribution of the haplotype among wild populations in the five world-wide regions and crops. Connecting lines show mutational pathways among haplotypes; triangles represent single-site substitutions; perpendicular bars, indels; and boxed bars, blocks of 16 and 2 tandem indels. Unlabeled red nodes indicate inferred steps not found in the sampled populations. Clusters group closely related haplotypes.

native and introduced regions, and (iii) among and within the two introduced regions. Gene genealogies were reconstructed with a reduced-median network (43) by using NETWORK 4.1 ([www.fluxus-engineering.com/sharenet.htm](http://www.fluxus-engineering.com/sharenet.htm)). According to the author's recommendations, superfluous links were eliminated by using the maximum parsimony calculation after down-weighting by half rapidly mutating sites (two single-site substitutions). Analyses were carried out treating the two multiple-site insertion deletions (indels; see *Results*) either as tandem independent sites (total sequence length, 333 bp) or as two single blocks (i.e., one block coding a 2-bp indel and one, a 16-bp indel, decreasing the total sequence length to 316 positions) by using several block-weighting schemes. Because we obtained the same results with either coding method, only values from the more conservative block-coding method are explicitly given and discussed.

## Results

**Genetic Diversity Across Populations and Regional Groups.** Over the whole dataset (524 individuals), 10 polymorphic sites, nine substitutions, and one indel were identified at the *atp8-S* locus and 24 polymorphic sites, five substitutions, and three indels of 16, 1, and 2 bp at the *W-I* locus. At each locus, nine different sequences were identified. The combined dataset (concatenated

sequence) was made up of 25 haplotypes, resulting in high overall haplotype ( $H_e = 0.86$ ) and nucleotide ( $\pi = 0.0139$ ) diversity. More haplotypes were detected than would be expected in the complete absence of homoplasy (17; i.e., the sum of the number of haplotypes at both loci minus one). In addition, although highly significant, the linkage disequilibrium value between the two loci was not close to unity, probably reflecting the occurrence of several evolutionary pathways among haplotypes rather than recombination. Statistical parsimony associated with network reconstruction is particularly recommended for managing homoplasy (43). In any case, the limited homoplasy detected (see loops in Fig. 2) was not uniquely due to the concatenation of the two loci (data not shown).

The distribution of the 25 haplotypes is detailed in Fig. 1. At the regional scale, the native region (Asia) showed high genetic diversity with eight haplotypes and  $H_e = 0.84$  ( $\pi = 0.0121$ ; Table 1). Introduced regions displayed contrasting levels of genetic diversity. Australasia, with 14 different haplotypes, showed the highest genetic diversity ( $H_e = 0.81$ ;  $\pi = 0.0166$ ; Table 1). In Europe, where only nine different haplotypes were observed, haplotype diversity was only moderate ( $H_e = 0.55$ ), and nucleotide diversity was reduced ( $\pi = 0.0021$ ; Table 1). This European pattern was due to the high frequency of one haplotype (Up01) in all but two populations (St. Malo and Cudillero; Fig. 1).

At the population level,  $H_e$  ranged from 0.0 to 0.88 (Table 2). Monomorphic populations were found in both native and introduced groups. Nevertheless, highly contrasting patterns were observed in Asia compared with Europe and Australasia. Native populations showed the lowest within-population genetic diversity [mean  $H_e$ ,  $0.17 \pm 0.20$ , number of populations ( $N_{pop}$ ) = 5], whereas the two main introduced regions showed considerably higher values (Europe, mean  $H_e$ ,  $0.36 \pm 0.23$ ,  $N_{pop} = 9$ ; Australasia, mean  $H_e$ ,  $0.46 \pm 0.33$ ,  $N_{pop} = 5$ ), with the most genetically diverse population found in Australasia (i.e., Dunedin; Table 2). Over all cultivated populations, the genetic diversity was very low ( $H_e = 0.24$ , Table 1) due to both the monomorphism of the French crop (Fig. 1) and the consistent presence of the Up01 haplotype whatever the sample.

**Among-Region Effects in the Population Genetic Structure.** Genetic differentiation was significant among populations within each region, but differentiation was at least twice as great among populations in either Far-East Asia or Australasia as compared with Europe (native, among-populations fixation index  $\Phi_{ST} = 0.960$ ; Australasia,  $\Phi_{ST} = 0.800$ ; Europe,  $\Phi_{ST} = 0.386$ ; all  $P < 0.001$ ). When comparing Australasia with native populations, significant genetic structure was found at the within-region level (among-populations within-regions fixation index,  $\Phi_{SC} = 0.868$ ,  $P < 10^{-5}$ ) but not at the among-regions level (among-regions fixation index,  $\Phi_{CT} = -0.019$ ,  $P = 0.533$ ) due to very high variation among-populations within each region as well as shared haplotypes (e.g., Up14) among regions. Conversely, European populations were highly genetically differentiated from the

**Table 1. Molecular diversity indices for each geographic region and crops**

Region	$N_{ind}$	$N_{pop}$	$P_{pop}$	$N_{hap}$	$H_e$	$S$	$\pi \times 10^2$ (SD)
Asia	118	5	3	8	0.84	11	1.21 (0.68)
Europe	208	9	8	9	0.55	4	0.21 (0.18)
Australasia	120	5	4	14	0.81	14	1.66 (0.90)
South America	24	1	1	2	0.29	1	0.09 (0.11)
North America	23	1	0	1	0.00	0	0.00 (0.00)
Crops	31	3	2	4	0.24	6	0.29 (0.23)
Total	524	24	18	25	0.86	17	1.39 (0.76)

$N_{ind}$ , number of individuals;  $N_{pop}$ , number of populations;  $P_{pop}$ , number of polymorphic populations;  $N_{hap}$ , number of haplotypes;  $H_e$ , haplotypic diversity;  $S$ , number of polymorphic sites;  $\pi$ , nucleotide diversity.



is also cultivated (59), Cudillero shows dominance of an alternate A haplotype. Cultivation may be at least partly responsible for the successful establishment of *U. pinnatifida* in Europe. Breeding programs reduce genetic variability and select for strains adapted for vigorous growth on ropes or nets, characteristics that may promote growth on artificial floating structures, typical of anthropogenic environments (ports, marinas, boat hulls, etc.), where spontaneous European populations of *U. pinnatifida* are usually found. Taken together, these pieces of evidence suggest that aquaculture was the source of both primary and secondary introductions of *U. pinnatifida* and its successful adaptation in Europe.

**Multiple Ships and Divergent Sources in Australasia.** All three haplotype clusters were well represented in Australasia, resulting in the highest values of sequence divergence observed at both the regional and population levels in this study, particularly in New Zealand. Based on morphological data and precise records of ship activities, it was suggested that two to three introductions to New Zealand from Asia by way of maritime traffic occurred independently in two distant ports (Wellington, in 1987 and Timaru, before 1988), with subsequent spread to intermediate points (27, 34). Our genetic data, however, suggest a greater number of recurrent (local) introductions all over New Zealand: of 12 haplotypes detected in New Zealand samples, only two were shared between Otago and Diamond Harbors. The heavy proliferation on natural substrates in New Zealand may be strongly linked to the redistribution of genetic variation from among (native) populations to within (introduced) populations, thereby stimulating novel, locally adaptive, genetic combinations.

Nevertheless, in contrast to New Zealand populations, those in Tasmania and mainland Australia showed relatively low within-population diversity, suggesting few locally successful, yet highly divergent, introduction events. Furthermore, although close geographically and connected via shipping traffic, Tasmania (first record of *U. pinnatifida* 1988, ref. 60) and Melbourne (1996, ref. 28) presented haplotypes that belonged to two different subclusters (B2 and B1, respectively; Figs. 1 and 2), suggesting independent primary introduction events (see ref. 34). Melbourne possesses haplotypes that are found in New Zealand, possibly representing a case of secondary relay from previously introduced populations, although a primary (perhaps cultivated) Asian origin cannot be excluded. Interestingly, haplotype composition in Argentina (first record of *U. pinnatifida* 1992, ref. 29) was highly similar to Melbourne, suggesting a single common origin for both introductions. Tasmania, on the other hand, appears to have arisen from an isolated and independent primary introduction. It was the only introduced area to show B2 haplotypes, separated by at least eight mutational steps from any other haplotype found in introduced populations. Furthermore, contrary to all other introduced areas, the initial populations in Tasmania were not found on floating structures in sheltered sites but on natural substrates in exposed sites (27, 60). In addition, the two Tasmanian populations have none of the four haplotypes found in the crop samples. The unusual Tasmanian haplotypes and habit could be interpreted as a cryptic (sub)species introduction. Given the highly localized distribution of B2 in Asia, the close phylogenetic relationship among the three *Undaria* species (61) and the great morphological plasticity noted within and

among species in this genus (62, 63), cluster B2 may in fact represent an alternate *Undaria* subspecies. Likewise, specific ecological adaptations appear to have promoted other cryptic invasions, such as in the polychaete *Marenzelleria* sp. (64) or the suggested derived incipient species status of invasive *Caulerpa taxifolia* (65). Testing the hypothesis of reproductive isolation between mitochondrial haplotype clusters, assuming they correspond to divergent genetic strains, in *U. pinnatifida* calls for the use of nuclear markers with analysis of cytonuclear disequilibrium (66).

## Conclusion

**Genetic Characteristics and Invasibility.** Genetic characteristics associated with invasion success are typically classified into two categories according to species: either (pre)adapted but relatively reduced genetic diversity resulting from population bottlenecks (e.g., ant, ref. 67; reed, ref. 68) or rapid evolution associated with enhanced levels of diversity due to recurrent introduction events (e.g., lizard, ref. 24; mollusks, refs. 69 and 70). In contrast, we uncovered a multiplicity of genetic characteristics in introduced *U. pinnatifida* populations. In particular, contrary to the well studied worldwide introduced alga *C. taxifolia* (65, 71), the pervasiveness of *U. pinnatifida* does not stem from the (rapid) evolution of a particular invasive type(s) with secondary relay around the world. We have effectively shown that the successful establishment of *U. pinnatifida* varies from (i) introduction of cultivated strains preadapted to anthropogenic habitats (i.e., floating structures and artificial substrata in Europe); (ii) divergent multiple-source introductions resulting in high genetic diversity promoting rapid local adaptation (New Zealand); (iii) putative secondary relay from these locally adapted invasive types (Melbourne and Argentina); and, finally, (iv) single-source, but cryptic, introductions (Tasmania). These different types of introduction processes have occurred independently around the world and promoted the settlement of the species over a large spectrum of environments.

We have demonstrated that a variety of microevolutionary processes can account for the success of a single worldwide invasive species. It is thus clearly difficult, based on localized genetic architecture only, to predict the invasive potential of a given species, and studies over the global range of introduction are essential. The multiplicity of processes and genetic signatures associated with the successful invasion of a given species confirms that multiple facets of global change, e.g., aquaculture practices, alteration of habitats, and increased traffic, act in synergy at the worldwide level, facilitating successful pandemic introductions.

This study would not have been possible without generous help from our colleagues from around the world (listed in Table 2) who sampled (or helped us sample) *Undaria* populations, for which we are extremely grateful. We thank C. Daguin, D. Jollivet, C. Lemaire, C. Maggs, and M. Valero for helpful comments on the manuscript. We are indebted to M. Perennou for help in using the facilities provided by the Génomère Ouest/Génomère sequencing platform. This study was part of the European Network of Excellence "Marine Genomics" (Contract 505403). This work was also supported by a Ph.D. fellowship ("Renouvellement des Compétences" Program) from the Région Bretagne (to M.V.) and the Centre National de la Recherche Scientifique through an Action Thématique et Incitative sur Programme (ATIP) grant (to F.V.) and a postdoctoral fellowship (to C.R.E.).

- Chapin, F. S. I., Zavaleta, E. S., Eviner, V. T., Naylor, R. L., Vitousek, P. M., Reynolds, H. L., Hopper, D. U., Lavorel, S., Sala, O. E., Hobbie, S. E., et al. (2000) *Nature* **405**, 234–242.
- Carlton, J. T. (2000) in *Invasive Species in a Changing World*, eds. Mooney, H. A. & Hobbs, R. (Island Press, Washington, DC), pp. 31–53.
- Occhipinti-Ambrogi, A. & Savini, D. (2003) *Marine Pollut. Bull.* **46**, 542–551.
- Stachowicz, J. J., Terwin, J. R., Whitlatch, R. B. & Osman, R. W. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 15497–15500.
- Suarez, A. V., Bolger, D. T. & Case, T. J. (1998) *Ecology* **79**, 2041–2056.
- Sax, D. F. & Brown, J. H. (2000) *Global Ecol. Biogeogr.* **9**, 363–371.
- Sala, O. E., Chapin, F. S., III, Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Hueneke, L. F., Jackson, R. B., Kinzig, A., et al. (2000) *Science* **287**, 1770–1774.
- Simberloff, D. & Von Holle, B. (1999) *Biol. Invasions* **1**, 21–32.
- Dukes, J. S. & Mooney, H. A. (1999) *Trends Ecol. Evol.* **14**, 135–139.
- Levine, J. M. & D'Antonio, C. M. (1999) *Oikos* **87**, 15–26.

11. Bax, N., Williamson, A., Aguero, M., Gonzalez, E. & Geeves, W. (2003) *Marine Pollut.* **27**, 313–323.
12. Sax, D. F. & Gaines, S. D. (2003) *Trends Ecol. Evol.* **18**, 561–566.
13. Roberts, C. M. & Hawkins, J. P. (1999) *Trends Ecol. Evol.* **14**, 241–246.
14. McKinney, M. L. & Lockwood, J. L. (1999) *Trends Ecol. Evol.* **14**, 450–453.
15. Myers, N. (1997) *Science* **278**, 597–598.
16. Kolar, C. S. & Lodge, D. (2001) *Trends Ecol. Evol.* **16**, 199–204.
17. Williamson, M. (1996) *Biological Invasions* (Chapman & Hall, London).
18. Ruiz, G. M., Fofonoff, P., Carlton, J. T., Wonham, M. J. & Hines, A., H. (2000) *Annu. Rev. Ecol. Syst.* **31**, 481–531.
19. Ricciardi, A. & Maclsaac, H. J. (2000) *Trends Ecol. Evol.* **15**, 62–65.
20. Drake, J. M. & Lodge, D. M. (2004) *Proc. R. Soc. London* **271**, 575–580.
21. Lee, C. E. (2002) *Trends Ecol. Evol.* **17**, 386–391.
22. Sakai, A. K., Allendorf, F. W., Holt, J. S., Lodge, M., Molofsky, J., With, K. A., Baughman, S., Cabin, R. J., Cohen, J. E., Ellstrand, N. C., McCauley, D. E., *et al.* (2001) *Annu. Rev. Ecol. Syst.* **32**, 305–332.
23. Kolbe, J. J., Glor, R. E., Schettino, L. S., Lara, A. C., Larson, A. & Losos, J. B. (2004) *Nature* **431**, 177–181.
24. Carlton, J. T. (1996) *Biol. Conserv.* **78**, 97–106.
25. Silva, P. C., Woodfield, A. R., Cohen, A. N., Harris, L. H. & Goddard, J. H. R. (2002) *Biol. Invasions* **4**, 333–338.
26. Hay, C. H. (1990) *Br. Phycol. J.* **25**, 301–313.
27. Campbell, S. J., Bité, J. S. & Burridge, T. R. (1999) *Bot. Mar.* **42**, 231–241.
28. Casas, G. N. & Piriz, M. L. (1996) *Hydrobiologia* **326/327**, 213–215.
29. Aguilar-Rosas, R., Aguilar-Rosas, L. E., Avila-Serrano, G. & Marcos-Ramirez, R. (2004) *Bot. Mar.* **47**, 255–258.
30. Thornber, C. S., Kinlan, B. P., Graham, M. H. & Stachowicz, J. J. (2004) *Mar. Ecol. Prog. Ser.* **268**, 69–80.
31. Floch, J.-Y., Pajot, R. & Mouret, V. (1996) *Hydrobiologia* **326/327**, 217–222.
32. Valentine, J. P. & Johnson, C. R. (2003) *J. Exp. Mar. Biol. Ecol.* **295**, 63–90.
33. Hay, C. H. & Sanderson, J. C. (1999) in *Islands in the Pacific Century: Pacific Science Inter-Congress* (University of the South Pacific, Suva, Fiji).
34. Orensanz, J. M. L., Schwindt, E., Pastorino, G., Bortolus, A., Casas, G., Darrigran, G., Elias, R., Lopez Gappa, J. J., Obenat, S., Pascual, M., *et al.* (2002) *Biol. Invasions* **4**, 115–143.
35. Grime, J. P. (1977) *Am. Nat.* **111**, 1169–1194.
36. Oudot-Le Secq, M. P., Kloareg, B. & Loiseaux de Goër, S. (2002) *J. Phycol.* **37**, 163–172.
37. Sunnucks, P., Wilson, A. C. C., Beheregaray, L. B. & Zenger, K. (2000) *Mol. Ecol.* **9**, 1699–1710.
38. Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994) *Nucleic Acids Res.* **22**, 4673–4680.
39. Hall, T. A. (1999) *Nuclear Acids Symp. Ser.* **41**, 95–98.
40. Coyer, J. A., Hoarau, G., Stam, W. T. & Olsen, J. L. (2004) *Mol. Ecol.* **13**, 1323–1326.
41. Excoffier, L., Smouse, P. E. & Quattro, J. M. (1992) *Mol. Mar. Biol. Biotechnol.* **5**, 295–298.
42. Posada, D. & Crandall, K. A. (2001) *Trends Ecol. Evol.* **16**, 37–45.
43. Coyer, J. A., Peters, A. F., Hoarau, G., Stam, W. T. & Olsen, J. L. (2002) *Eur. J. Phycol.* **37**, 173–178.
44. Coyer, J. A., Peters, A. F., Hoarau, G., Stam, W. T. & Olsen, J. L. (2002) *Proc. R. Soc. London* **269**, 1829–1834.
45. Ehara, M., Kitayama, T., Watanabe, K. I., Inagaki, Y., Hayashi-Ishimaru, Y. & Ohama, T. (1999) *Phycol. Res.* **47**, 225–231.
46. Peters, A. F., Scornet, D., Müller, D. G., Kloareg, B. & Cock, J. M. (2004) *Eur. J. Phycol.* **39**, 235–242.
47. Yoon, H. S. & Boo, S. M. (1999) *Hydrobiologia* **398/399**, 47–55.
48. Allendorf, F. W. & Lundquist, L. L. (2003) *Conserv. Biol.* **17**, 24–30.
49. Davies, N., Villanblanca, F. X. & Roderick, G. K. (1999) *Genetics* **153**, 351–360.
50. Perez, R., Lee, J. Y. & Juge, C. (1981) *Science Pêche* **315**, 1–12.
51. Wolff, W. J. & Reise, K. (2002) in *Invasive Aquatic Species of Europe*, eds. Leppäkoski, E., Gollasch, S. & Olenin, S. (Kluwer, Dordrecht, The Netherlands), pp. 193–205.
52. Boudouresque, C. F., Gerbal, M. & Knoepffler-Peguy, M. (1985) *Phycologia* **24**, 364–366.
53. Ohno, M. & Largo, D. B. (1998) in *Seaweed Resources of the World*, eds. Critchley, A. T. & Ohno, M. (Japanese International Cooperation Agency, Yokosura, Japan), pp. 1–14.
54. Perez, R., Kaas, R. & Barbaroux, O. (1984) *Science Pêche* **343**, 3–15.
55. Fletcher, R. L. & Farrel, P. (1999) *Helgolander Meeresuntersuchungen* **52**, 259–275.
56. Critchley, A. T. & Ohno, M. (1998) in *Seaweed Resources of the World* (Japanese International Cooperation Agency, Yokosura, Japan), p. 431.
57. International Council for the Exploration of the Sea (2001) *ICES Coop. Res. Rep.* **248**, 58–59.
58. Juanes, J. A. & Sosa, P. A. (1998) in *Seaweed Resources of the World*, eds. Critchley, A. T. & Ohno, M. (Japan International Cooperation Agency, Yokosura, Japan), pp. 164–175.
59. Sanderson, J. C. (1990) *Bot. Mar.* **33**, 153–157.
60. Yoon, H. S., Lee, J. Y., Boo, S. M. & Bhattacharya, D. (2001) *Mol. Phylogenet. Evol.* **21**, 231–243.
61. Saito, Y. (1972) in *Contribution to the Systematics of the Benthic Marine Algae of the North Pacific*, eds. Abbott, A. & Kurogi, M. (Japanese Society of Phycology, Kobe, Japan), pp. 117–132.
62. Stuart, M. D., Hurd, C. L. & Brown, M. T. (1999) *Hydrobiologia* **398/399**, 191–199.
63. Bastrop, R., Jürss, K. & Sturmbauer, C. (1998) *Mol. Biol. Evol.* **15**, 97–103.
64. Meusnier, I., Valero, M., Olsen, J. L. & Stam, W. T. (2004) *Eur. J. Phycol.* **39**, 83–92.
65. Basten, C. J. & Asmussen, M. A. (1997) *Genetics* **146**, 1165–1171.
66. Tsutsui, N. D., Suarez, A. V., Holway, D. A. & Case, T. J. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 5948–5953.
67. Saltonstall, K. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 2445–2449.
68. Bachelet, G., Simon-Bouhet, B., Desclaux, C., Garcia-Meunier, P., Mairesse, G., de Montaudouin, X., Raigné, H., Randriambao, K., Sauriau, P.-G. & Viard, F. (2004) *Mar. Ecol. Prog. Ser.* **276**, 147–159.
69. Dupont, L., Jollivet, D. & Viard, F. (2003) *Mar. Ecol. Prog. Ser.* **253**, 183–195.
70. Jousson, O., Pawlowski, J., Zaninetti, L., Zechman, F., Dini, F., Di Giuseppe, G., Woodfield, G., Millar, R. & Meinesz, A. (2000) *Nature* **408**, 157.