

A NEW EARLY OLIGOCENE CRAB (DECAPODA, BRACHYURA, DROMIACEA) FROM NORTHWEST BELGIUM, WITH COMMENTS ON ITS PALAEOBIOLOGY

BARRY W. M. VAN BAKEL^{1,*}, PEDRO ARTAL², RENÉ H. B. FRAAIJE¹ & JOHN W. M. JAGT³

(3 figures, 1 table)

1. Oertijdmuseum De Groene Poort, Bosscheweg 80, NL-5283 WB Boxtel, the Netherlands;

E-mail: barryvanbakel@gmail.com [* corresponding author], info@oertijdmuseum.nl

2. Museo Geológico del Seminario de Barcelona, Diputación 231, E-08007 Barcelona, Spain;

E-mail: partal@optimus.es

3. Natuurhistorisch Museum Maastricht (SCZ), de Bosquetplein 6-7, NL-6211 KJ Maastricht, the Netherlands;

E-mail: john.jagt@maastricht.nl

ABSTRACT. A new genus and species of dromiacean crab, *Lucanthonisia praemium*, is recorded from the basal Belsele-Waas Clay Member (Boom Clay Formation; Rupelian, lower Oligocene) as formerly exposed at the Scheerders van Kerchove (SVK) clay pit near Sint-Niklaas, province of Oost-Vlaanderen (northwest Belgium). Extant dromioid crabs either carry foreign objects for camouflage (Dromiidae, Homolodromiidae) or are associated with corals (Dynomenidae). Although having been collected from the basal portion of the Belsele-Waas Clay Member which is rich in phosphatic nodules, preservation of the new form is such that long-term transport is unlikely. Therefore, the species is assumed to have lived hidden amongst and below such nodules, but it cannot be ruled out that it also carried sponges for defence. To our knowledge, representatives of such biota have not yet been recorded from the Boom Clay Formation. *Dromilites eotvoesi* from middle Miocene ('upper Badenian', Serravallian) strata in the Budapest area (central Hungary) is transferred to the new genus, *Lucanthonisia*. *Lucanthonisia eotvoesi* n. comb. differs from *L. praemium*, the type species, in having shorter, more anteriorly directed lateral spines, more closely spaced and less divergent rostral horns, better-developed epigastric nodes and a less convex carapace longitudinally. Preservation of a partial thoracic sternum in *L. praemium* gen. nov., sp. nov. allows comparison with dromioid families and tentative placement of the new genus, as well as the allied *Basinotopus*, in the Dynomenidae.

KEYWORDS: Crustacea, Decapoda, Dromioidea, Dynomenidae, Oligocene, Belgium, new taxa, palaeobiology.

1. Introduction

Dromiaceans often hide in crevices or carry foreign objects or organisms such as sponges, shells, ascidians, and rarely sea anemones to prevent being detected by potential predators (Guinot et al., 1995; Müller, 2004). In the Dromiidae diverse objects or organisms are grasped by modified posterior legs, placed on the carapace and held there. Some sponge species and most sea anemones taken by dromioids are toxic and thus provide additional protection against predators. In the fossil record, dromioid diversity (i.e., members of the families Dromiidae de Haan, 1833 and Dynomenidae Ortmann, 1892) increased following Cretaceous-Paleogene (K/Pg) boundary perturbations (Table 1). However, in comparison to other brachyurans, dromiacean crabs constitute only a minor percentage of fossil decapod crustacean faunas (Bell, 1858; Bachmayer & Tollmann, 1953; Quayle & Collins, 1981; Busulini et al., 1983; Solé & Via, 1989; Beschin et al., 1994, 2001, 2002, 2005, 2007a, b; De Angeli, 1995; Blow & Manning, 1996; Kato & Karasawa, 1998; De Angeli & Beschin, 2001; Karasawa, 2001; Fraaije, 2003; Müller, 2004; Schweitzer & Feldmann, 2005), which is comparable to extant faunas (Ng et al., 2008).

In Europe, decapod crustaceans are fairly common in strata of Eocene and Miocene age, but are much rarer in Oligocene deposits. Not only are such strata less well exposed, they also represent mainly siliciclastic, slightly cold-water platform settings, in which sandy and/or clay-rich, volcanoclastic sediments were laid down (Vandenbergh, 1978; Zimmerle, 1994). Notable exceptions are the rich faunas recorded from the lower Oligocene of northern Germany (Förster & Mundlos, 1982) and Piemonte, northern Italy (Allasinaz, 1987). Carbonate nearshore or reefal settings are uncommon during this time interval (compare Müller, 2004), which may explain the rarity of dromioid crabs.

Predominantly clayey Rupelian strata are well known and widely distributed in northwest Belgium, in particular in the Boom area (Fig. 1). Formerly, these clays were excavated on a large scale for local brickworks. In these strata, van Bakel et al. (2006) recognised one species each of macrurans and anomurans, plus four brachyuran taxa, inclusive of the new genus and species described below, to which they referred as 'dromiid n. gen., n. sp.'. The sole crab species of Rupelian age recorded so far from Belgium was *Coeloma* (*Paracoeloma*) *rupeliense*




Figure 1. Map of Belgium, indicating the location of the former Scheerders van Kerchove (SVK) clay pit near Sint-Niklaas (province of Oost-Vlaanderen).

Stainier, 1887, of which hundreds of nodule-preserved specimens are housed in various museum collections (Van Straelen, 1921; Feldmann & Dhondt, 1991). Verheyden (2002) described collections held at the Institut royal des Sciences naturelles de Belgique (IRScNB, Brussels) in detail, correctly interpreted *Portunus nodosus* Van Beneden, 1883 to be a *nomen nudum* and showed *Coeloma rupeliense* to be the valid name of this taxon. In addition, he illustrated large cheliped remains of a giant macruran, *Homarus percyi* Van Beneden, 1872. Both *Coeloma*

(*Paracoeloma*) *rupeliense* and *Homarus percyi* have subsequently also been recorded from nodules reworked into the base of the Pliocene sequence at Kallo, near Antwerp (van Bakel et al., 2006).

In recent years, Rupelian strata at the Scheerders van Kerchove (SVK) clay pit (Fig. 2) have yielded a rich crustacean fauna, the composition of which differs from assemblages known from adjacent outcrops. A diogenid hermit crab with stridulatory apparatus, *Ciliopagurus obesus*, was described by van Bakel et al. (2003). Of *Coeloma* (*Coeloma*) sp. only fragmentary specimens are known, and a cirolanid isopod and stomatopod remains are currently being studied by us. In the present paper, we describe a new genus and species of dromiacean, which could have been derived from the genus *Basinotopus* M'Coy, 1849 (type species: *Inachus lamarckii* Desmarest, 1822). Both genera may be related to the extant genus *Paradynomene* Sakai, 1963 (type species: *P. tuberculata* Sakai, 1963).

2. Systematic palaeontology

Classification and terminology follow Guinot & Bouchard (1998), Martin & Davis (2001), Guinot & Tavares (2003), Ahyong et al. (2007) and Ng et al. (2008). Commonly used morphological terms are the following: episternite – lateral extension of the sternite, in the form of a more or less pointed projection which bears the gynglyme receiving the condyle of the corresponding appendage




Figure 2. Stratigraphy of the section formerly exposed at the Scheerders van Kerchove (SVK) clay pit (modified from Mollen, 2007), with arrow marking provenance of the type specimen of *Lucanthonisia praeium* gen. nov., sp. nov.

(see Guinot & Tavares, 2003: 47); sternocoxal depressions – deeply excavated depressions at the lateral sides of the thoracic sternum in which the coxae of the pereopods can move, as defined by Guinot et al. (1995) and Guinot & Bouchard (1998). Sternites and episternites are indicated by number 1 to 8, as are the pereopods 1 to 5 (e.g., P1 for chelipeds, P2-P5 for walking pereopods). We follow Ng et al. (2008) in assigning authorship of the taxon *Brachyura* to Linnaeus (1758) rather than to Latreille (1802).

To denote the repositories of specimens referred to in the text, the following abbreviations are used: MAB – Oertijdmuseum De Groene Poort, Boxtel, the Netherlands; MGUH – Geological Museum, Copenhagen University, Copenhagen, Denmark; MNM – Magyar Nemzeti Múzeum, Óslénytár, Hungary.

Infraorder Brachyura Linnaeus, 1758
 Section Dromiacea de Haan, 1833
 Superfamily Dromioidea de Haan, 1833
 Family ?Dynomenidae Ortmann, 1892
 Genus *Lucanthonisia* gen. nov.

Derivation of name. In honour of Mr Luc Anthonis (Bouwel, Belgium), who collected and donated the type and sole specimen known to date.

Type species. *Lucanthonisia praemium* sp. nov.

Diagnosis. Carapace subovate with a wide posterior margin; length slightly exceeding width, convex in longitudinal and transverse cross sections; rostrum produced, tridentate with thorn-shaped spines; lateral margins with four equal-sized, slender spines; carapace surface with numerous pits, areolation weak; cardiac region broad, cervical groove does not reach the lateral carapace margin, but the branchial groove does. Orbitofrontal margin more than half the carapace width (excluding lateral spines); orbits deep, ovate, with a small notch below the outer orbital angle, suborbital area with two tubercles. Sternal plate flat, with fused elements 4-7, which are constant in width, except for the episternites which are purely laterally directed; sternite 4 broad.

Discussion. As a complement to the extensive revision by McLay (1993), Guinot & Tavares (2003: 44) proposed a




Figure 3. *Lucanthonisia praemium* gen. nov., sp. nov., MAB k.2448 (holotype), basal portion of the Belsele-Waas Clay Member, Boom Clay Formation (Rupelian, lower Oligocene), SVK clay pit, Sint-Niklaas (province of Oost-Vlaanderen, Belgium), in orbital (A), right lateral (B), dorsal (C) and ventral (D) views. Specimen coated with ammonium chloride prior to photography. Scale bar equals 3 mm.

new subfamilial arrangement of the Dromiidae and recognised thirty-eight genera, primarily based on, 'the morphology of the thoracic sternum, spermathecae at the extremity of sternal sutures 7/8, uropods, vestigial male pleopods on abdominal somites 3-5, coxa of the fifth pereopod and the penis'. The same characters were primarily used in a key to dromiacean families by Guinot & Tavares (2003: 120). When dealing with fossil material, where mostly only dorsal carapaces are available, proxy characters can be used for classification (Schweitzer, 2003).

***Lucanthonisia praemium* sp. nov.**

Derivation of name. From Latin, *praemium*, reward or recompense, alluding to the enormous amount of sieving residue processed to find the specimen.

Type. Holotype, of indeterminate sex, is MAB k.2448 (leg. L. Anthonis). Carapace length (including rostrum) is 6,3 mm, carapace width (including lateral spines) is 5,7 mm; it is interpreted as a juvenile.

Diagnosis. Carapace subovate in outline, slightly longer than wide; rostrum produced, wide; tridentate through robust inner orbital spines; lateral margins spinose, posterior margin broad; only central carapace regions well defined.

Locality and stratigraphy. Collected from sieve residues of samples taken in 2005 from the basal portion of the Belsele-Waas Clay Member (Boom Clay Formation; see Fig. 2), dated early to middle Rupelian (Oligocene) on the basis of calcareous nannofossils (Steurbaut, 1986, 1992; Laga et al., 2002), typical of calcareous nannoplankton zone NP 23 (sensu Martini, 1971), as exposed at the Scheeders van Kerchove's Verenigde Fabrieken (SVK) clay pit, southwest of Sint-Niklaas, province of Oost-Vlaanderen (Belgium), now no longer worked. For more detailed data on this locality and the section formerly exposed, reference is made to Vandenberghe (1978), Vandenberghe et al. (2002) and Mollen (2007).

Description. Carapace subovate, slightly longer than wide, greatest carapace width a third from posterior margin, convex in longitudinal and transverse cross sections; orbitofrontal margin 64 % of maximum carapace width, marginal spines included. Front produced, broad, occupying more than half of orbitofrontal width; inner orbital spines stout and divergent, at slightly lower (vertical) level than sharp, thorn-shaped rostrum; rostrum downturned, but with its tip directed anteriorly; orbits deep, broadly ovate, with weakly raised and entire orbital margin leading to a distinct outer orbital corner. A small notch is situated just below the outer orbital corner; the suborbital margin with two tubercles, the outer slightly larger than the inner. Anterolateral margin long, bearing four upwardly directed, equal-sized conical spines; first three spines slightly anteriorly directed and evenly spaced, separated by broadly U-shaped notches; posterior lateral

spine more widely separated from anterior ones, directed perpendicularly to carapace axis. Posterolateral margin short, strongly curved towards posterior margin. Posterior margin equals in width the orbitofrontal margin and is bounded by a distinct rim and a rather deep groove.

Epigastric elevations hardly noticeable, bounding the weakly defined, narrowly triangular, anterior mesogastric process. Posterior mesogastric process more strongly elevated, broadly triangular and subdivided by obscure median sulcus. The large and gently raised protogastric regions barely separated from the narrow hepatic strip; urogastric lobe low, anterior margin concave, posterior margin straight and more clearly defined; lateral margins strongly divergent. Cardiac region large, shield shaped, posteriorly indistinctly bounded. Meso- and metabranchial regions obliquely directed, raised axially, paired by cervical and branchiocardiac grooves, and only basally separated by anterior branch of branchiocardiac groove; only a rim of pterygostomian region is preserved. Cervical groove faint laterally, interrupted axially between clearly visible gastric pits. The cervical groove does not notch the lateral margin, but is replaced by a Y-shaped groove which extends downwards on the subhepatic area, starting at the two anteriormost lateral spines. The branchial groove runs through uninterrupted and makes a notch laterally on margin; it is prolonged ventrally.

Proepistome and epistome preserved : proepistome triangular with apex directed forwards, the lateral sides being concave; epistome wide, with typical wing-shaped lateral projections; mandibular sternite attached to epistome, anteriorly defined by a triangular fossa.

Thoracic sternum well preserved, interpreted to be formed at least by sternites 4-6, broken just posterior to sternite 6. Sternal plate undivided and relatively wide, its width being constant along sternites 4-6 (except for extended episternites). Sternites 1-3 not preserved, the small triangle visible in Fig. 3D is formed by sediment only. Sutures not visible. Sternites 4 and 6 easily recognised on account of the landmarks represented by well-developed gynglymes; sternite 4 quadrate with a concave anterior margin and with a markedly projected episternite. Sternite 6 short; sternites 5 and 6 with a very weak median depression. Posterior sternites not preserved. Episternites 4 and 5 (corresponding to P1 and P2, respectively) elongated, triangular, directed outwardly, their anterior side being hollowed in deep sternocoxal depressions.

Carapace surface with numerous small, evenly distributed pits.

Discussion. *Lucanthonisia* gen. nov. is included in the Dromiacea and may be placed either in the Dromiidae (see McLay, 1991, 1993) or Dynomenidae (see McLay, 1999). Since we assume the new genus to be related to *Basinotopus*, the existence of dorsal uropods in *B. lamareckii* (see Bell, 1858: pl. 5, figs. 1, 7-9) eliminates the family Homolodromiidae Alcock, 1900 (see Guinot & Richer de Forges, 1995) in which the uropods always show as ventral lobes and in which the abdominal somites

have lateral pleurae.

Lucanthonisia might be assigned to the Dromiidae. With respect to the thoracic sternum, the following can be stated. If the holotype of *Lucanthonisia praemium* gen. nov., sp. nov. represents a female specimen, it cannot be referred to the Dromiinae de Haan, 1833, a subfamily in which the female sternal sutures 7 and 8 move forwards, and thus are very long, the apertures of the spermathecae opening far beyond the level of P3 coxa (see Guinot & Tavares, 2003: 49, 121; Guinot & Quenette, 2005: 272, fig. 1). Only the atypical dromiine *Stebbingdromia* Guinot & Tavares, 2003 has short sutures 7 and 8 in females, but the carapace is more rounded, the rostrum is bifid and the anterolateral margin has only two flat teeth. If the type of *L. praemium* gen. nov., sp. nov. represents a male (i.e., in males, the sternal plate is not modified by suture 7/8; see Guinot & Quenette, 2005: 272), it may be a dromiine (compare e.g. the sternum in *Platydrodromia* Brocchi, 1877 in Guinot & Tavares, 2003: fig. 16, and that of *Exodromidia* Stebbing, 1905 in Guinot & Bouchard, 1998: fig. 5). However, many dromiines have a narrower sternite 4 (compare e.g. *Austrodromidia australis* (Rathbun, 1923) as illustrated by Guinot & Tavares, 2003: fig. 1A) and generally have the anterior termination of sternite 4 as a single apex (as in *Stebbingdromia* Guinot & Tavares, 2003; see Guinot & Tavares, 2003: fig. 17A, C). In extant *Dromidiopsis* Borradaile, 1900, the carapace is more circular, the median rostral tooth does not project beyond the adjacent pair, and the posterior corners are pronounced rather than rounded (see Cleve et al., 2007: fig. 9C). Sternite 4 is very narrow in *Dromidiopsis*, and the episternites not much extended (see Guinot & Tavares, 2003: fig. 6A), which differs significantly from *Lucanthonisia* gen. nov.

If *Lucanthonisia* gen. nov. is not a dromiine, it might belong to the subfamily Sphaerodromiinae Guinot & Tavares, 2003, a group in which the female sternal suture 7/8 is very short, only posterior and located laterally, the apertures of spermathecae being posterior to the level of the P3 coxal gonopore. The carapace outline in sphaerodromiines is ovoid as in *Lucanthonisia* gen. nov., but with a narrower posterior margin.

The atypical dromiacean genus, *Frodromia* McLay, 1993 (see McLay, 1993: figs 6, 17D; considered to be of uncertain status by Guinot & Tavares, 2003; now in the tribe Frodromiini Števčić, 2005, within the Sphaerodromiinae) also has short sutures 7/8 in females. In *Lucanthonisia* gen. nov., sternite 4 is wider than that in *Frodromia* (compare Guinot & Tavares 2003: fig. 25b).

In the genus *Basinotopus*, the broadest portion of the cardiac region typically equals a third of total carapace width, measured from the bases of the lateral spines. Collins (2003) and Collins & Jakobsen (2004) distinguished *Dromilites* H. Milne Edwards, 1837 (type species: *D. bucklandii* H. Milne Edwards, 1837) from *Basinotopus* by noting that *Dromilites* has a bifid rostrum (in dorsal view), while *Basinotopus* differs markedly in showing a produced rostrum between inner orbital horns,

i.e. has a tridentate front. Adopting the definition of *Basinotopus* by Collins (2003) and Collins & Jakobsen (2004), *Dromilites eotvoesi* Müller, 1975 (p. 508, pl. 1, fig. 5; pl. 2, figs. 1, 4; holotype is MNM MF 11; see for a better illustration Müller, 1976: 150, pl. 2, figs. 1-3), which was transferred to the genus *Dromia* Weber, 1795 (Müller, 1984: 63, pl. 29, figs. 8, 9; pl. 30, figs. 1-7), could be assigned to *Basinotopus*. However, *Dromilites eotvoesi* cannot be accommodated in *Basinotopus* because, although carapace outline, frontal structure and division of carapace regions are roughly similar, *Dromilites eotvoesi* has a smooth carapace with numerous setal pits. This morphology indicates a hairy appearance in life. In addition, the lateral spines are smaller and more slender. The morphology of *Dromilites eotvoesi* suggests it hid and lived in crevices. The new species from the lower Oligocene of Belgium is closely similar, and both it and *Dromilites eotvoesi* are placed in a new genus, *Lucanthonisia* gen. nov.

Lucanthonisia eotvoesi n. comb. was originally described from middle Miocene ('lower upper Badenian', equivalent of lower Serravallian; see Harzhauser et al., 2002) of the Budapest area in central Hungary. It has shorter, more anteriorly directed lateral spines than *L. praemium* sp. nov., its rostral horns are more closely spaced and less divergent, it has better-developed epigastric nodes, while the branchiocardiac groove in *L. praemium* sp. nov. divides the carapace surface more distinctly in the centre. The posterior margin is more clearly rimmed and lined by a deeper groove in *L. praemium*, and the carapace is more convex in longitudinal cross section. Müller (1984: 63) noted that *D. eotvoesi* was known only from 'lower upper Badenian' levels, generally co-occurring with the calappid *Calappa heberti* Brocchi, 1883, and absent from reefal structures.

Of *Basinotopus tricornis* Collins & Jakobsen, 2004 (holotype is MGUH 26778) a number of well-preserved specimens are known from the Lillebælt Clay Formation of Jylland (northern Denmark) of early Lutetian (middle Eocene) age (see Janssen et al., 2007). These specimens show details of abdomen, pereopods, mouthparts and thoracic sternum (Collins & Jakobsen, 2004: text-fig. 3; pl. 2, figs. 1-7). Whereas only a 'possible male abdomen' is mentioned in their description, one specimen (MGUH 26776; see their pl. 2, fig. 1b) is in fact a female with an obvious gonopore on the (right) P3 coxa. The spermathecae are diminutive, at about the same vertical level as the gonopore, or slightly lower; therefore, sternal suture 7/8 is relatively short. In *B. tricornis* the anterior axial portions of sternite 7 are raised. Sternite 4 is narrower than that in *Lucanthonisia* gen. nov., the episternites are hollowed by sternocoxal depressions as in the new genus. The inclined sulcus between sternites 4 and 5 is absent in *Lucanthonisia praemium* gen. nov., sp. nov. *Basinotopus tricornis* appears closely related to the type species, *B. lamarckii*, from the lower Eocene (Ypresian) of England and the middle Eocene of northern Italy (see Quayle & Collins, 1981; Busulini et al., 1983; Collins, 2003; Beschin et al., 2005), as well as to *B. alpina*

(Glaessner, 1929) from the middle Eocene (Lutetian) of St Pankraz near Salzburg (Austria) and the lower Eocene of Spain (pers. obs.). *Dromilites lothi* Förster & Mundlos, 1982, from the upper Eocene-lower Oligocene of Handorf (north-east Germany), has a tridentate front, and the carapace areolation is utterly different. The delicate nature of the front could have been misinterpreted on account of the poor preservation in the type specimen; in any case, the inner orbital horns are weakly developed. The carapace outline, with its wide posterior margin, and broad sternite 4, makes placement of *Lucanthonisia* gen. nov. and *Basinotopus* in the subfamily Sphaerodromiinae unlikely.

The new genus could belong to the Dynomenidae, a family with a good fossil record (see e.g., Wright & Collins, 1972: fig. 9). In this group, the female sternal suture 7/8 is short, as in males. The thoracic sternum of *Lucanthonisia* shows about the same organisation as that of a dynomenid (compare Guinot & Bouchard, 1998: fig. 7A, C, D): sternites 4-7 fused in an undivided plate of constant width, thus the anterior side of sternite 4 broad, a flat medial area, and the episternites directed purely laterally. The carapace shape in *Lucanthonisia* gen. nov. is more elongate than in most extant dynomenids, but it should be remarked that at least in one dynomenid genus, the most primitive one, *Acanthodromia* A. Milne-Edwards, 1879, the carapace is long as well. *Paradynomene* Sakai, 1963 resembles *Basinotopus* (e.g., *B. lamarckii* and *B. tricornis*) in several respects (see McLay, 1999; McLay & Ng, 2004), although in the former the rostrum is shorter and stouter, the grooves and the branchial ones in particular, are steeper, and the ornament is different.

The combination of carapace ornament, the slender nature of the lateral spines, oval carapace outline and rostral structure, distinguishes the new form from other dromioid genera. Due to its size, which is smaller than average for a dromiacean, the holotype might be considered to represent a juvenile. Carapace size in the family Dynomenidae, to which *Lucanthonisia* gen. nov. and *Basinotopus* could possibly belong (see discussion below), is generally small (see McLay, 1999) but rarely as small as the specimen described here. A detailed comparison with numerous specimens of the related *Basinotopus alpina* from the Ilerdian (= lower Ypresian) of northeast Spain and of *B. lamarckii* from the Ypresian of southeast England, has revealed no indications of changes in carapace ornament or morphology during ontogeny, which is why we exclude the possibility that the type of *L. praemium* gen. nov., sp. nov. is an immature individual of *Basinotopus*.

Noetlingia Beurlen, 1928 has an elongate carapace, with only diminutive lateral nodes. Its type species, *Dromia claudiopolitana* Bittner, 1893, has a length/width ratio of 1.35 vs 1.13 in *L. praemium* gen. nov., sp. nov. The length measured includes the rostrum, the width is inclusive of the lateral spines. In addition, in *Noetlingia* the branchial and cardiac grooves are not clearly distinguished.

In conclusion, we tentatively assign *Lucanthonisia* gen. nov. to the Dynomenidae, as based on combined thoracic and carapace features shared with primitive dynomenid genera such as *Acanthodromia* and *Paradynomene*. However, this conclusion remains tentative, because we have not been able to examine male and female uropods, abdominal holding structures, spermathecae, or the condition of pereopods 4 and 5; diagnostic of familial and subfamilial assignments within Dromiacea.

3. Palaeobiology

Carapace ornament is highly functional in dromiacean crabs, setal pits being indicative of a hairy surface, and whenever this occurs, it extends to the pereopods, which all together give the crab a woolly appearance. A comparable morphology is seen in e.g. porcellanids and majids. In the homoloid *Poupinia hirsuta* Guinot, 1991, the last pair of legs are extremely mobile and covered with long setae, and can be bent over the carapace to camouflage it (Guinot et al., 1995). A different way for dromiids to camouflage their bodies is to hold an object with their modified posterior pereopods (P4 and P5) (see e.g., Lewinsohn, 1984; McLay, 1991, 1993, 2001; McLay & Crosnier, 1991; Guinot & Bouchard, 1998).

The Eocene *Basinotopus* is characterised by a scabrous posterior carapace surface, distinctly differentiated regions with crests, and strong lateral spines of considerable length. Representatives of this genus were likely covered by epizooobionts (*sensu* Taylor & Wilson, 2002) such as ascidians, and even after being noted by a predator would not be considered a palatable choice. Such epizooobiont cover could be referred to as a symbiosis, it being an ecological relationship between organisms of two different species that live together in direct contact (Zann, 1980; Campbell et al., 1999). In extant Dromiidae and Homolodromiidae, permanently with a camouflaging host carried on by P4 and P5, it is an obligatory relationship, from which in particular the crab benefits; thus, it is better referred to as commensalism.

Long and stout lateral spines, a thick shell, and sharp spines on the dorsal surface would complicate consumption and thus protect the crab's soft parts; however, it also means severe restrictions for the crab to enter small crevices and cavities. A carapace surface roughened by tubercles and with well-differentiated regions would facilitate anchorage of epizooobionts such as sponges, sea anemones and seagrasses. Subdorsal, mobile and subchelate posterior pereopods could place these foreign objects onto the carapace, or even carry them. Elevation of carapace regions is a constant and distinctive character throughout dromioid evolution. McLay (1993) was of the opinion that dromiids were evolving towards a loss of carrying behaviour, coupled with a tendency to develop a strongly ornamented carapace and with a loss of modified P4 and P5.

Having been collected from the basal portion of the Belsele-Waas Clay Member, which is rich in phosphatic nodules, the holotype of *Lucanthonisia praemium* gen.

nov., sp. nov. is remarkably well preserved. This makes long-term transport highly unlikely, and suggests the species lived hidden amongst or under such nodules. However, it cannot be ruled out that it also carried foreign objects such as sponges, but these have not been documented so far from the Boom Clay Formation. Strata correlative with the Boom Clay Formation (Stover & Hardenbol, 1994) elsewhere in Europe (see e.g., von Daniels et al., 1994; Gaemers, 1994) may be expected to yield similar decapod crustacean faunas, possibly inclusive of dromioids.

4. Acknowledgements

We thank L. Anthonis (Bouwel, Belgium) for donation of the specimen, F. H. Mollen (Berlaar, Belgium) and E. R. H. Wille (Wuustwezel-Gooresind, Belgium) for bringing material to our attention and for supplying items of literature, P. Verhaert and W. Van Branden (both SVK Verenigde Fabrieken, Sint-Niklaas, Belgium) for allowing access to their former clay pit, J. Herman (Belgian Geological Survey, Brussels) for co-ordination of collecting bed by bed at Sint-Niklaas, D. Guinot (Muséum national d'Histoire naturelle, Paris) for invaluable assistance in the interpretation of the thoracic sternum of the new genus, and for commenting in detail on an earlier typescript.

5. References

- AHYONG, S.T., LAI, J.C.Y., SHARKEY, D., COLGAN, D.J. & NG, P.K.L., 2007. Phylogenetics of the brachyuran crabs (Crustacea: Decapoda): The status of Podotremata based on small subunit nuclear ribosomal RNA. *Molecular Phylogenetics and Evolution*, 45: 576-586.
- ALCOCK, A.W., 1900. Materials for a carcinological fauna of India, No. 5. The Brachyura primigenia, or Dromiacea. *Journal of the Asiatic Society of Bengal, Natural Sciences*, 68: 123-169.
- ALLASINAZ, A., 1987. Brachyura Decapoda oligocenici (Rupeliano) del Bacino Ligure Piemontese. *Bollettino del Museo regionale di Scienze naturali, Torino*, 5: 509-566.
- ARTAL, P. & GILLES, A., 2007. New Miocene crabs from Pignat (southeast France). In Garassino, A., Feldmann, R.M. & Teruzzi, G. (eds.), 3rd Symposium on Mesozoic and Cenozoic decapod crustaceans, Museo di Storia Naturale di Milano, May 23-25, 2007. *Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 35: 8-11.
- BACHMAYER, F. & TOLLMANN, A., 1953. Die Crustaceen-Fauna aus dem tortonischen Leithakalk (Steinbrüche der Firma Fenk) bei Groß-Höflein im Burgenland. In *Kober-Festschrift 1953. Skizzen zum Antlitz der Erde. Geologische Arbeiten, herausgegeben aus Anlaß des 70. Geburtstages von Prof. Dr. L. Kober, Universität Wien*, 308-314, pl. 1. Gesellschaftsbuchdruckerei Brüder Hollinek, Wien.
- BAKEL, B.W.M. van, JAGT, J.W.M. & FRAAIJE, R.H.B., 2003. *Ciliopagurus obesus*, a new species of Oligocene hermit crab from northwest Belgium. *Cainozoic Research*, 2 (for 2002): 109-116.
- BAKEL, B.W.M. van, FRAAIJE, R.H.B. & JAGT, J.W.M., 2006. Synopsis of Cenozoic decapod crustaceans from Belgium. *Revista Mexicana de Ciencias Geológicas*, 23: 370-374.
- BARNOLAS, A., 1973. *Dromilites vicensis* n. sp., nuevo braquiuro del Eocene marino de Cataluña. *Instituto de Investigaciones Geológicas de la Universidad de Barcelona*, 28: 5-14.
- BELL, T., 1858. A monograph of the fossil malacostracous Crustacea of Great Britain. Part I. Crustacea of the London Clay. *Monograph of the Palaeontographical Society, London*, 10 (for 1856): viii + 44 pp., pls 1-11.
- BESCHIN, C., BUSULINI, A., DE ANGELI, A. & TESSIER, G., 1994. I crostacei eocenici della Cava «Boschetto» di Nogarole Vicentino (Vicenza-Italia settentrionale). *Società veneziana di Scienze naturali, Lavori*, 19: 159-215.
- BESCHIN, C., BUSULINI, A., DE ANGELI, A. & TESSIER, G., 2002. Aggiornamento ai crostacei eocenici di Cava «Main» di Arzignano (Vicenza-Italia settentrionale) (Crustacea, Decapoda). *Studi e Ricerche, Associazione Amici del Museo, Museo civico 'G. Zannato'*, 2002: 7-28.
- BESCHIN, C., BUSULINI, A. & TESSIER, G., 2007a. First report of a new Eocene crustacean fauna from the Veronese Lessini (N Italy). In Garassino, A., Feldmann, R.M. & Teruzzi, G. (eds.), 3rd Symposium on Mesozoic and Cenozoic decapod crustaceans, Museo di Storia Naturale di Milano, May 23-25, 2007. *Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 35: 12-14.
- BESCHIN, C., BUSULINI, A., DE ANGELI, A. & TESSIER, G., 2007b. *I decapodi dell'Eocene inferiore di Contrada Gecchelina (Vicenza, Italia settentrionale) (Anomura e Brachyura)*, 76 pp. Montecchio Maggiore (Vicenza), Museo di Archeologia e Scienze Naturali 'G. Zannato'.
- BESCHIN, C., DE ANGELI, A. & CHECCHI, A., 2001. Crostacei decapodi associati a coralli della "Formazione di Castelmomberto" (Oligocene) (Vicenza-Italia settentrionale). *Studi e Ricerche, Associazione Amici del Museo, Museo civico 'G. Zannato'*, 2001: 13-30.
- BESCHIN, C., DE ANGELI, A., CHECCHI, A. & ZARANTONELLO, G., 2005. Crostacei eocenici di Grola presso Spagnago (Vicenza, Italia settentrionale). *Studi e Ricerche, Associazione Amici del Museo, Museo civico 'G. Zannato'*, 12: 5-35.
- BEURLEN, K., 1928. Die Decapoden des Schwäbischen Jura, mit Ausnahme der aus den oberjurassischen Plattenkalken stammenden. *Palaeontographica*, A70: 115-278, pls 6-8.

- BITTNER, A., 1883. Neue Beiträge zur Kenntniss der Brachyuren-Fauna des Alttertiärs von Vicenza und Verona. *Denkschriften der kaiserlichen Akademie der Wissenschaften zu Wien*, 46: 299-316.
- BITTNER, A., 1886. Neue Brachyuren des Eocaens von Verona. *Sitzungsberichte der kaiserlichen Akademie der Wissenschaften zu Wien*, 94: 44-55.
- BITTNER, A., 1893. Decapoden des pannonischen Tertiärs. *Sitzungsberichte der kaiserlichen Akademie der Wissenschaften zu Wien*, 102: 10-37.
- BLOW, W.C. & MANNING, R.B., 1996. Preliminary descriptions of 25 new decapod crustaceans from the Middle Eocene of the Carolinas, U.S.A. *Tulane Studies in Geology and Paleontology*, 29: 1-26.
- BORRADAILE, L., 1900. On some Crustaceans from the South Pacific. Part IV. The crabs. *Proceedings of the Zoological Society of London*, 4: 568-596.
- BROCCHI, P., 1877. Sur un Dromien nouveau, genre *Platydomia*. *Bulletin de la Société philomatique de Paris*, (6)12 (for 1875): 53-54.
- BROCCHI, P., 1883. Note sur les crustacés fossiles des terrains tertiaires de la Hongrie. *Annales des Sciences géologiques*, 13: 1-8, pls 4, 5.
- BUSULINI, A., TESSIER, G., VICENTIN, M., BESCHIN, C., DE ANGELI, A. & ROSSI, A., 1983. Nuovo contributo alla conoscenza dei brachiuri eocenici di Cava Main (Arzignano) – Lessini orientali (Vicenza) (Crustacea, Decapoda). *Lavori della Società veneziana di Scienze naturali*, 8: 55-73.
- CAMPBELL, N.A., REECE, J.B. & MITCHELL, L.G., 1999. *Biology*. xxvii + 1175 pp. Benjamin/Cummings, Addison Wesley Longman, Inc., Menlo Park, California.
- CHECCHIA-RISPOLI, G., 1905. I crostacei dell'Eocene dei dintorni di Monreale in provincia di Palermo. *Giornale di Scienze Naturali ed Economiche di Palermo*, 25: 309-325.
- CLEVA, R., GUINOT, D. & ALBENGA, V., 2007. Annotated catalogue of brachyuran type specimens (Crustacea, Decapoda, Brachyura) deposited in the Muséum national d'Histoire naturelle, Paris. Part I. Podotremata. *Zoosystema*, 29: 1-51.
- COLLINS, J.S.H., 2003. A taxonomic review of British decapod Crustacea. *Bulletin of the Mizunami Fossil Museum*, 29 (for 2002): 81-92.
- COLLINS, J.S.H. & DONOVAN, S.K., 2007. New decapod crustaceans from the Palaeogene of Jamaica. *Bulletin of the Mizunami Fossil Museum*, 33 (for 2006): 59-65.
- COLLINS, J.S.H. & JAKOBSEN, S.L., 1995. A synopsis of the biostratigraphic distribution of the crab genera (Crustacea, Decapoda) of the Danian (Palaeocene) of Denmark and Sweden. *Bulletin of the Mizunami Fossil Museum*, 21 (for 1994): 35-46.
- COLLINS, J.S.H. & JAKOBSEN, S.L., 2004. New crabs (Crustacea, Decapoda) from the Eocene (Ypresian/Lutetian) Lillebælt Clay Formation of Jutland, Denmark. *Bulletin of the Mizunami Fossil Museum*, 30 (for 2003): 63-96.
- COLLINS, J.S.H. & WIENBERG RASMUSSEN, H., 1992. Upper Cretaceous-Lower Tertiary decapod crustaceans from West Greenland. *Grønlands Geologiske Undersøgelse, Bulletin*, 162: 1-46.
- DANIELS, C.H. von, GRAMANN, F. & KÖTHE, A., 1994. The Oligocene "Septarienton" of Lower Saxony. Biostratigraphy of an equivalent of the Boom Formation of Belgium, with special considerations to its upper and lower boundaries. *Bulletin de la Société belge de Géologie*, 102 (for 1993): 79-89.
- DAVIDSON, E., 1966. A new Paleocene crab from Texas. *Journal of Paleontology*, 40: 211-213.
- DE ANGELI, A., 1995. Crostacei dell'Eocene superiore di "Fontanella" di Grancona (Vicenza-Italia settentrionale). *Studi e Ricerche, Associazione Amici del Museo, Museo civico 'G. Zannato'*, 1995: 7-24.
- DE ANGELI, A. & BESCHIN, C., 2001. I crostacei fossili del territorio Vicentino. *Natura Vicentina*, 5: 5-54.
- DE ANGELI, A. & GARASSINO, A., 2006. Catalog and bibliography of the fossil Stomatopoda and Decapoda from Italy. *Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 35: 1-95.
- DESMAREST, A.-G., 1822. Les crustacés proprement dits. In Brongniart, A. & Desmarest, A.-G. (eds.), *Histoire naturelle des crustacés fossiles, sous les rapports zoologiques et géologiques*, 67-142, pls 5-11. L. T. Cellot, Paris.
- FELDMANN, R.M. & DHONDT, A.V., 1991. The Van Straelen decapod collection, Institut royal des Sciences naturelles de Belgique, Bruxelles. *Journal of Paleontology*, 65: 514-515.
- FISCHER-BENZON, R. von, 1866. *Ueber das relative Alter des Faxekalkes und über die in demselben vorkommenden Anomuren und Brachyuren*, 1-30, pls 1, 1a, 2-5. Schwer'sche Buchhandlung, Kiel.
- FÖRSTER, R. & MUNDLOS, R., 1982. Krebse aus dem Alttertiär von Helmstedt und Handorf (Niedersachsen). *Palaeontographica*, A179: 148-184.
- FRAAIJE, R.H.B., 2003. Evolution of reef-associated decapod crustaceans through time, with particular reference to the Maastrichtian type area. *Contributions to Zoology*, 72: 119-130.
- FRIEBE, J.G., 1987. Eine Krabben-Fauna aus dem Leithakalk (Badenien) von Wurzing bei Wildon, Steiermark. *Mitteilungen des naturwissenschaftlichen Vereins in Steiermark*, 117: 57-65.
- GAEMERS, P.A.M., 1994. Refined correlations by means of lithostratigraphy and gadid otolith zonation of the Rupelian of the North Sea Basin: a progress report. *Bulletin de la Société belge de Géologie*, 102 (for 1993): 147-157.
- GLAESSNER, M.F., 1929. Dekapodenstudien. IV. Die Dekapodenfauna des Alttertiärs der Nordalpen. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, B63: 158-167.

- GUÉRIN-MÉNEVILLE, F.É. 1829-1844. *Iconographie du règne animal de G. Cuvier, Crustacés*, 1-48, 36 pls. J.-B. Baillière, Paris.
- GUINOT, D., 1991. Établissement de la famille des Poupiniidae pour *Poupinia hirsuta* n. gen., n. sp. de Polynésie (Crustacea, Decapoda, Brachyura, Homoloidea). *Bulletin du Muséum national d'Histoire naturelle, Paris*, (4)12 (for 1990): 577-605.
- GUINOT, D. & BOUCHARD, J.-M., 1998. Evolution of the abdominal holding system of brachyuran crabs (Crustacea, Decapoda, Brachyura). *Zoosystema*, 20: 613-694.
- GUINOT, D., DOUMENC, D. & CHINTIROGLOU, C.C., 1995. A review of the carrying behaviour in brachyuran crabs, with additional information on the symbioses with sea anemones. *The Raffles Bulletin of Zoology*, 43: 377-416.
- GUINOT, D. & QUENETTE, G., 2005. The spermatheca in podotreme crabs (Crustacea, Decapoda, Brachyura, Podotremata) and its phylogenetic implications. *Zoosystema*, 27: 267-342.
- GUINOT, D. & RICHER DE FORGES, B., 1995. Crustacea Decapoda Brachyura: Révision de la famille des Homolodromiidae Alcock, 1900. In Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, volume 13. *Mémoires du Muséum national d'Histoire naturelle, Paris*, 163: 155-282.
- GUINOT, D. & TAVARES, M., 2003. A new subfamilial arrangement of the Dromiidae de Haan, 1833, with diagnoses and descriptions of new genera and species (Crustacea, Decapoda, Brachyura). *Zoosystema*, 25: 43-129.
- HAAN, W. de, 1833-1850. Crustacea. In Siebold, P.F. von (ed.), *Fauna japonica sive descriptio animalium, quae in itinere per Japoniam, jussu et auspiciis superiorum, qui summum in India Batava Imperium tenent, suscepto, annis 1823-1830 collegit, notis, observationibus e adumbrationibus illustravit*, xvii + xxxi + 243 pp. A. Arnz, Lugduni Batavorum.
- HARZHAUSER, M., BÖHME, M., MANDIC, O. & HOFMANN, C.C., 2002. The Karpatian (Late Burdigalian) of the Korneuburg Basin. In Sovis, W. & Schmid, B. (eds.), *Das Karpat des Korneuburger Beckens, Teil 2. Beiträge zur Paläontologie*, 27: 441-456.
- HASWELL, W.A., 1882. Description of some new species of Australian Decapoda. *Proceedings of the Linnean Society of New South Wales*, 6: 750-763.
- IHLE, J.E.W., 1913. Die Decapoda Brachyura der Siboga-Expedition, I. Dromiacea. *Siboga Expeditie, Monografie*, 39b: 1-96.
- JAGT, J.W.M., BAKEL, B.W.M. van, FRAAIJE, R.H.B. & NEUMANN, C., 2006. *In situ* fossil hermit crabs (Paguroidea) from northwest Europe and Russia. Preliminary data on new records. *Revista Mexicana de Ciencias Geológicas*, 23: 364-369.
- JAKOBSEN, S.L. & COLLINS, J.S.H., 1997. New Middle Danian species of anomuran and brachyuran crabs from Fakse, Denmark. *Bulletin of the Geological Society of Denmark*, 44: 89-100.
- JANSSEN, A.W., SCHNETLER, K.I. & HEILMANN-CLAUSEN, C., 2007. Notes on the systematics, morphology and biostratigraphy of fossil holoplanktonic Mollusca, 19. Pteropods (Gastropoda, Euthecosomata) from the Eocene Lillebaelt [sic] Clay Formation (Denmark, Jylland). *Basteria*, 71: 157-168.
- KARASAWA, H., 2001. Coral-associated decapod Crustacea from the Pliocene Daito Limestone Formation and Pleistocene Ryukyu Group, Ryukyu Islands, Japan. *Bulletin of the Mizunami Fossil Museum*, 27 (for 2000): 167-189.
- KATO, H. & KARASAWA, H., 1998. Pleistocene fossil decapod Crustacea from the Boso Peninsula, Japan. *Natural History Research, Special Issue*, 5: 1-31.
- LAGA, P., LOUWY, S. & GEETS, S., 2002. Paleogene and Neogene lithostratigraphic units (Belgium). In Bultynck, P. & Dejonghe, L. (eds.), *Guide to a revised lithostratigraphic scale of Belgium. Geologica Belgica*, 4 (for 2001): 135-152.
- LAMARCK, J. B. P. A. D. de, 1819. *Histoire naturelle des animaux sans vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédée d'une introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie*, 5: 1-612. Déterville & Verdière, Paris.
- LATREILLE, P.A., 1802-1803. *Histoire naturelle, générale et particulière, des crustacés et des insectes. Ouvrage faisant suite aux œuvres de Leclerc de Buffon, et partie du cours complet d'histoire naturelle rédigé par C. S. Sonnini, membre de plusieurs sociétés savantes*, 3: xii + 467 pp. F. Dufart, Paris.
- LATREILLE, P.A., 1812. Crustacés et insectes. in Milbert, J., *Voyage pittoresque à l'Île-de-France, au Cap de Bonne-Espérance et à l'Île de Ténériffe*. Vol. 2 : 270-280. A. Nepveu, Paris.
- LEWINSOHN, C., 1984. Dromiidae (Crustacea, Decapoda, Brachyura) from Madagascar and the Seychelles. *Bulletin du Muséum national d'Histoire naturelle, Paris*, (4)6: 89-129.
- LINNAEUS, C., 1758. *Systema Naturae, per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata*, 1(6). Laurentii Salvii, Holmiae, iii + 824 pp.
- MARTIN, J.W. & DAVIS, G.E., 2001. An updated classification of the Recent Crustacea. *Natural History Museum of Los Angeles, Science Series*, 39: 1-124.
- MARTINI, E., 1971. Standard Tertiary and Quaternary calcareous nannoplankton zonation. In Farinacci, A. (ed.), *Proceedings of the Second Planktonic Conference* (Roma), 2: 739-785. Edizioni Tecnoscienza, Roma.
- M'COY, F., 1849. On the classification of some British fossil Crustacea, with notices of new forms in the university collection at Cambridge. *Annals and Magazine of Natural History*, (2)4: 161-179, 330-335, 412-414.

- McLAY, C.L., 1991. A small collection of deep water sponge crabs (Brachyura, Dromiidae) from French Polynesia, including a new species of *Sphaerodromia* Alcock, 1899. *Bulletin du Muséum national d'Histoire naturelle, Paris*, (4)13: 457-481.
- McLAY, C.L., 1993. Crustacea Decapoda: The sponge crabs (Dromiidae) of New Caledonia and the Philippines with a review of the genera. In Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, volume 10. *Mémoires du Muséum national d'Histoire naturelle, Paris*, 156: 111-251.
- McLAY, C.L., 1999. Crustacea Decapoda: Revision of the family Dynomenidae. In Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, volume 20. *Mémoires du Muséum national d'Histoire naturelle, Paris*, 180: 427-569.
- McLAY, C.L., 2001. The Dromiidae of French Polynesia and a new collection of crabs (Crustacea, Decapoda, Brachyura) from the Marquesas Islands. *Zoosystema*, 23: 77-100.
- McLAY, C.L. & CROSNIER, A., 1991. Description of a new and unusual species of *Sphaerodromia* (Brachyura, Dromiidae) from the Seychelles Islands. *Bulletin du Muséum national d'Histoire naturelle, Paris*, (4)13: 181-188.
- McLAY, C.L. & NG, P., 2004. A taxonomic revision of the genus *Paradynomene* Sakai, 1963 (Crustacea: Decapoda: Brachyura: Dynomenidae). *Zootaxa*, 657: 1-24.
- MILNE-EDWARDS, A., 1879. Mémoire sur les crustacés décapodes du genre *Dynomène*. *Annales des Sciences naturelles, Zoologie*, (6)8: 1-11.
- MILNE EDWARDS, H., 1837. *Histoire naturelle des crustacés, comprenant l'anatomie, la physiologie et la classification de ces animaux*, 2: 1-532, 42 pls. Roret, Paris.
- MOLLEN, F., 2007. A new species of *Abdounia* (Elasmobranchii, Carcharhinidae) from the base of the Boom Clay Formation (Oligocene) in northwest Belgium. *Geologica Belgica*, 10: 69-77.
- MÜLLER, P., 1975. Decapoda (Crustacea) fauna a budapesti miocénböl (3). *Földtani Közlöny*, 105: 506-515.
- MÜLLER, P., 1976. Decapoda (Crustacea) fauna a budapesti miocénböl (4). *Földtani Közlöny*, 106: 149-160.
- MÜLLER, P., 1979a. Decapoda (Crustacea) fauna a budapesti miocénböl (5). *Földtani Közlöny*, 108: 272-312, pls 1-23.
- MÜLLER, P., 1979b. Décapodes du Badénien et Sarmatien de Bulgarie. *Palaeontology, Stratigraphy, Lithology (Sofiya)*, 10: 3-7.
- MÜLLER, P., 1984. Decapod Crustacea of the Badenian. *Geologica Hungarica, Series Palaeontologica*, 42: 1-317.
- MÜLLER, P., 2004. History of reef-dwelling decapod crustaceans from the Palaeocene to the Miocene with comments about Mesozoic occurrences. *Földtani Közlöny*, 134: 237-255.
- MÜLLER, P. & COLLINS, J.S.H., 1991. Late Eocene coral-associated decapods (Crustacea) from Hungary. *Contributions to Tertiary and Quaternary Geology*, 28: 47-92.
- NG, P.K.L., GUINOT, D. & DAVIE, P.J.F., 2008. Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *The Raffles Bulletin of Zoology*, 17: 1-286.
- NOETLING, F., 1885. Die Fauna des samländischen Tertiärs, I. Theil, II. Crustacea und Vermes. *Abhandlungen zur geologischen Specialkarte von Preussen und den Thüringischen Staaten*, 6(3): 381-442 [111-172]; atlas, 9 pls.
- ORTMANN, A.E., 1892. Die Dekapoden-Krebse des Strassburger Museums, mit besonderer Berücksichtigung der von Herrn Dr Döderlein bei Japan und bei den Liu-Kiu-Inseln gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen, V. Theil: Die Abteilungen Hippiden, Dromiiden und Oxystomata. *Zoologisches Jahrbuch (Systematik)*, 6: 532-588, pl. 26.
- PORTELL, R.W. & COLLINS, J.S.H., 2004. Decapod crustaceans of the Lower Miocene Montpelier Formation, White Limestone Group of Jamaica. In Donovan, S.K. (ed.), The Mid-Cainozoic White Limestone Group of Jamaica. *Cainozoic Research*, 3 (for 2003): 109-126.
- QUAYLE, W.J. & COLLINS, J.S.H., 1981. New Eocene crabs from the Hampshire Basin. *Palaeontology*, 24: 733-758, pls 104-105.
- RATHBUN, M.J., 1923. Report on the crabs obtained by the F.I.S. 'Endeavour' on the coasts of Queensland, New South Wales, Victoria, South Australia and Tasmania. Report on the Brachyryncha, Oxystomata and Dromiacea. *Biological Results of the Fishing Experiments carried on the F.I.S. 'Endeavour'*, 1909-14, 5(3): 95-156, pls 16-42.
- REUSS, A.E., 1859. Zur Kenntniss fossiler Krabben. *Denkschriften der kaiserlichen Akademie der Wissenschaften zu Wien, mathematisch-naturwissenschaftliche Klasse*, 17: 1-90, 24 pls.
- ROBERTS, H.B., 1956. Early Tertiary decapod crustaceans from the Vincentown Formation in New Jersey. *Bulletin of the Wagner Free Institute of Science*, 31: 5-12, pl. 2.
- SAKAI, T., 1963. Description of two new genera and fourteen new species from the collection of his Majesty the Emperor of Japan. *Crustaceana*, 5: 213-233.
- SCHLOTTHHEIM, E.F. von, 1820. *Die Petrefaktenkunde auf ihrem jetzigen Standpunkte durch die Beschreibung seiner Sammlung versteinerter und fossiler Überreste des Thier- und Pflanzenreiches der Vorwelt erläutert*. Becker, Gotha, lxii + 437 pp.
- SCHWEITZER, C.E., 2003. Utility of proxy characters for classification of fossils: an example from the fossil Xanthoidea (Crustacea: Decapoda: Brachyura). *Journal of Paleontology*, 77: 1107-1128.

- SCHWEITZER, C.E. & FELDMANN, R.M., 2005. Decapod crustaceans, the K/P event, and Palaeocene recovery. In Koenemann, S. & Jenner, R.A. (eds.), *Crustacea and arthropod relationships. Festschrift for Fredrick R. Schram*, 17-53. CRC Press/Taylor and Francis, Boca Raton.
- SOLÉ, J. & VIA, L., 1989. Crustacis decàpodes fòssils del Paísos Catalans (Recopilació i actualització de dades des de 1855 a 1988). *Batalleria*, 2: 23-42.
- STAINIER, X., 1887. *Caeloma rupeliense*. Brachyoure nouveau de l'argile rupélienne. *Annales de la Société géologique de la Belgique*, 14: 86-96, pl. 5.
- STEBBING, T.R.R., 1905. South African Crustacea (Part III). In: *Marine investigations in South Africa*, 4: 21-123, pls 17-26. Cape Times Ltd., Cape Town.
- STEURBAUT, E., 1986. Late Middle Eocene to Middle Oligocene calcareous nannoplankton from the Kallo well, some boreholes and exposures in Belgium and a description of the Ruisbroek Sand Member. *Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie*, 23: 49-83.
- STEURBAUT, E., 1992. Integrated stratigraphic analysis of lower Rupelian deposits (Oligocene) in the Belgian basin. *Annales de la Société géologique de la Belgique*, 115: 287-306.
- ŠTEVČIĆ, Z., 2005. The reclassification of brachyuran crabs (Crustacea: Decapoda: Brachyura). *Natura Croatica*, 14 (Supplement 1): 1-159.
- STOVER, L.E. & HARDENBOL, J., 1994. Dinoflagellates and depositional sequences in the Lower Oligocene (Rupelian) Boom Clay Formation, Belgium. *Bulletin de la Société belge de Géologie*, 102 (for 1993): 5-77.
- TAYLOR, P.D. & WILSON, M.A., 2002. A new terminology for marine organisms inhabiting hard substrates. *Palaios*, 17: 522-525.
- VAN BENEDEN, P., 1872. Sur la découverte d'un homard fossile dans l'argile du Rupelmonde. *Bulletin de l'Académie royale de la Belgique*, 33: 316-321.
- VAN BENEDEN, P., 1883. Sur quelques formes nouvelles des terrains du pays. *Bulletin de l'Académie royale de la Belgique*, 6: 132-134.
- VANDENBERGHE, N., 1978. Sedimentology of the Boom Clay (Rupelian) in Belgium. *Verhandelingen van de Koninklijke Academie voor Wetenschappen, Letteren en Schone Kunsten van België*, Klasse der Wetenschappen, 40(147): 1-137.
- VANDENBERGHE, N., HERMAN, J. & STEURBAUT, E., 2002. Detailed analysis of the Rupelian Ru-1 transgressive surface in the type area (Belgium). In Gürs, K. (ed.), *Northern European Cenozoic Stratigraphy*. Proceedings of the 8th Biannual Meeting of the RCNNS/RCPNS, 67-81. Landesamt für Natur und Umwelt des Landes Schleswig-Holstein, Flintbek.
- VAN STRAELEN, V., 1921. Catalogue des crustacés décapodes des terrains tertiaires de la Belgique. *Annales de la Société royale de Zoologie et de Malacologie de Belgique*, 51: 111-131.
- VERHEYDEN, T., 2002. Decapods from the Boom Clay (Rupelian, Oligocene) in Belgium. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, 72: 171-191.
- VIA BOADA, L., 1959. Decápodos fósiles del Eoceno español. *Boletín del Instituto geológico y minero de España*, 70: 331-402, 7 pls.
- VOGELTANZ, R., 1968. Beitrag zur Kenntnis der fossilen Crustacea Decapoda aus dem Eozän des Südhelvetikums von Salzburg. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 130: 78-105.
- WANK, M., 1986. Neue Fossilien aus den Steinbrüchen der Wietersdorfer Zementwerke. *Carinthia II*, 176: 61-67.
- WEBER, F., 1795. *Nomenclator entomologicus secundum entomologiam systematicum ill. Fabricii adjectis speciebus recens detectis et varietatibus*, viii + 171 pp. Carolum Ernestum Bohn, Chilonii/Hamburgi.
- WRIGHT, C.W. & COLLINS, J.S.H., 1972. British Cretaceous crabs. *Monograph of the Palaeontographical Society, London*, 126(533): 1-114, 22 pls.
- ZANN, L.P., 1980. *Living together in the sea*, 416 pp. T.F.H. Publications, Neptune NJ.
- ZIMMERLE, W., 1994. On the lithology and provenance of the Rupelian Boom Clay in northern Belgium, a volcanoclastic deposit. *Bulletin de la Société belge de Géologie*, 102 (for 1993): 91-103.

Table 1: Overview of Paleogene and Neogene dromiid and dynomenid (dromioid) crabs described to date, arranged alphabetically, indicating provenance, stratigraphic level and pertinent references.

Ameridromia hyneorum Blow & Manning, 1996
 South Carolina, USA; middle Eocene (Blow & Manning, 1996)

Basinotopus alpina (Glaessner, 1929)
 Spain; lower Eocene. Austria; middle Eocene (Glaessner, 1929; Vogeltanz, 1968; this paper)

Basinotopus lamarckii (Desmarest, 1822)
 England; lower Eocene. Veneto and Sicily, Italy; middle Eocene (De Angeli & Beschin, 2001; Collins, 2003; Beschin et al., 2005; De Angeli & Garassino, 2006)

Basinotopus tricornis Collins & Jakobsen, 2004
 Denmark; middle Eocene (Collins & Jakobsen, 2004; Janssen et al., 2007)

Cyamocarcinus angustifrons Bittner, 1883
 Veneto and Sicily, Italy; Hungary, Austria; lower-upper Eocene (Wank, 1968; De Angeli & Beschin, 2001; De Angeli & Garassino, 2006; Beschin et al., 2007b)

Cryptodromia aff. *fallax* (Lamarck, 1819)
 Japan; Plio-Pleistocene (Karasawa, 2001)

Dromia neogenica Müller, 1979a
 Hungary; middle Miocene (Müller, 1979a, 1984)

Dromia personata (Linnaeus, 1758)
 Sicily, Italy; upper Pleistocene (De Angeli & Garassino, 2006, as *D. vulgaris* H. Milne Edwards, 1837)

Dromidia bedetteae Blow & Manning, 1996
 South Carolina, USA; middle Eocene (Blow & Manning, 1996)

Dromiidae gen. et sp. indet. 1
 Japan; Pleistocene (Kato & Karasawa, 1998)

Dromiidae gen. et sp. indet. 2
 Japan; Pleistocene (Kato & Karasawa, 1998)

Dromiidae sp.
 France; Miocene (Artal & Gilles, 2007)

Dromilites bucklandii H. Milne Edwards, 1837
 England; lower Eocene (Collins, 2003)

Dromilites corvini (Bittner, 1893)
 Hungary; upper Eocene. Vicenza, Italy; upper Eocene-lower Oligocene (Beschin et al., 2001; De Angeli & Beschin, 2001; De Angeli & Garassino, 2006)

Dromilites fossata Müller & Collins, 1991
 Hungary; upper Eocene (Müller & Collins, 1991)

Dromilites hilarionis (Bittner, 1883)
 Vicenza, Italy; Spain; middle-upper Eocene (Busulini et al., 1983; De Angeli, 1995; De Angeli & Beschin, 2001; Beschin et al., 2005; De Angeli & Garassino, 2006)

Dromilites humerosus Quayle & Collins, 1981
 England; middle-upper Eocene (Quayle & Collins, 1981; Collins, 2003)

Dromilites lothi Förster & Mundlos, 1982
 Germany; upper Eocene-lower Oligocene (Förster & Mundlos, 1982)

Dromilites pastoris Via Boada, 1959
 Vicenza, Italy; Spain; middle Eocene (Beschin et al., 1994; De Angeli & Beschin, 2001; Beschin et al., 2005; De Angeli & Garassino, 2006)

Dromilites simplex Quayle & Collins, 1981
 England; middle-upper Eocene (Quayle & Collins, 1981)

Dromilites subglobosa Müller & Collins, 1991
 Hungary; upper Eocene (Müller & Collins, 1991)

Dromilites vicensis Barnolas, 1973
 Spain; Eocene (Solé & Via, 1989)

Dromiopsis americana Roberts, 1956
 New Jersey, USA; Paleocene (Roberts, 1956; Collins & Wienberg Rasmussen, 1992)

Dromiopsis elegans Reuss, 1859
 Denmark; lower Paleocene (Collins & Jakobsen, 1995)

Dromiopsis granulata Collins & Wienberg Rasmussen, 1992
 West Greenland; Maastrichtian (Collins & Wienberg Rasmussen, 1992)

Dromiopsis laevior Reuss, 1859
 Denmark; lower Paleocene (Collins & Jakobsen, 1995)

Dromiopsis minor von Fischer-Benzon, 1866
 Denmark; lower Paleocene (Collins & Jakobsen, 1995)

Dromiopsis paucigranosa Beschin, Busulini, De Angeli & Tessier, 2007
 Vicenza, Italy; lower Eocene (Beschin et al., 2007b)

Dromiopsis rugosa (von Schlottheim, 1820)
 Denmark, southern Sweden; lower Paleocene (Jakobsen & Collins, 1997; Collins & Jakobsen, 2005)

***Dynomene emiliae* Müller, 1979b**

Austria, Bulgaria, Poland, Hungary; middle Miocene (Müller, 1979b, 1984; Friebe, 1987)

***Dynomene hispida* (Latreille in Milbert, 1812)**

Japan; Plio-Pleistocene (Karasawa, 2001, as *D. hispida* Guérin-Méneville, 1832) (see for synonymy: Ng et al., 2008: 38)

***Dynomene lessinea* Beschin, De Angeli & Checchi, 2001**

Vicenza, Italy; lower Oligocene (Beschín et al., 2001; De Angeli & Beschín, 2001; De Angeli & Garassino, 2006)

***Dynomene* aff. *praedator* A. Milne-Edwards, 1879**

Japan; Plio-Pleistocene (Karasawa, 2001)

***Dynomene variabilis* Portell & Collins, 2004**

Jamaica; lower Miocene (Portell & Collins, 2004)

***Eotrachynotocarcinus airaghii* Beschín, Busulini, De Angeli & Tessier, 2007**

Vicenza, Italy; lower Eocene (Beschín et al., 2007b)

***Epigodromia areolata* (Ihle, 1913)**

Japan; Plio-Pleistocene (Karasawa, 2001)

***Gemmellarocarcinus loerentheyi* Checchia-Rispoli, 1905**

Sicily, Italy; middle Eocene. Hungary; upper Eocene (De Angeli & Garassino, 2006)

***Kierionopsis nodosa* Davidson, 1966**

Texas, USA; Paleocene (Davidson, 1966)

***Kromtitis koberi* (Bachmayer & Tollmann, 1953)**

Austria; upper Miocene (Bachmayer & Tollmann, 1953; Müller, 1984)

***Kromtitis koberiformis* Beschín, Busulini, De Angeli & Tessier, 2007**

Vicenza, Italy; lower Eocene (Beschín et al., 2007b)

***Kromtitis levigatus* Beschín, Busulini, De Angeli & Tessier, 2007**

Vicenza, Italy; lower Eocene (Beschín et al., 2007b)

***Kromtitis subovatus* Beschín, Busulini, De Angeli & Tessier, 2007**

Vicenza, Italy; lower Eocene (Beschín et al., 2007b)

***Kromtitis pentagonalis* Müller & Collins, 1991**

Hungary; upper Eocene (Müller & Collins, 1991)

***Kromtitis spinulata* Portell & Collins, 2004**

Jamaica; lower Miocene (Portell & Collins, 2004)

***Kromtitis tetratuberculatus* Beschín, Busulini, De Angeli & Tessier, 2002**

Vicenza, Italy; middle Eocene (Beschín et al., 2002; De Angeli & Garassino, 2006)

***Kromtitis* sp.**

Vicenza, Italy; middle-upper Eocene (Beschín et al., 2007a)

***Lucanthonisia eotvoesi* (Müller, 1975)**

Hungary; middle Miocene (Müller, 1975, 1984; this paper)

***Lucanthonisia praemium* sp. nov.**

Belgium; lower Oligocene (this paper)

***Noetlingia claudiopolitana* (Bittner, 1893)**

Hungary; upper Eocene. Veneto, Italy; middle Eocene (Beschín et al., 1994; De Angeli & Beschín, 2001; De Angeli & Garassino, 2006)

***Noetlingia succini* (Noetling, 1885)**

Samland Peninsula, Kaliningrad, Russia; middle Eocene (Noetling, 1885; Jagt et al., 2006)

***Noetlingia veronensis* (Bittner, 1886)**

Veneto, Italy; middle/upper Eocene (De Angeli & Garassino, 2006)

***Trechmannius circularis* Collins & Donovan, 2007**

Jamaica; lower Paleocene (Collins & Donovan, 2007)