

All rights reserved

Internal reports are not to be reprinted or cited, it is only allowed by consent of the Netherlands Institute For Sea Research.

printed by  texel

PRELIMINARY RESEARCH ON THE CLAY MINERAL COMPOSITION
OF SOME SEDIMENT SAMPLES IN THE ZAIRE ESTUARY AND FAN

door

G.J. DE LANGE*

INHOUD

Summary & Résumé	1
1. Introduction	2
2. Results and Discussion	8
3. References	13

SUMMARY

Samples from the top-layer of some selected trip-cores from the Zaire estuary and fan were analysed for their clay mineral composition. The main feature to be distinguished is a pronounced seaward decrease in the relative percentage of kaolinite and a simultaneous increase in smectite.

RÉSUMÉ

Des échantillons de la couche supérieure de quelques carottes courtes de l'estuaire et "fan" du Zaire ont été analysés pour leur composition en minéraux argileux. La caractéristique dominante, lorsqu'on s'éloigne de la côte, est le décroissement prononcé du pourcentage relatif en kaolinite et, simultanément, l'acroissement en smectite.

* Present address: Instituut voor Aardwetenschappen, afdeling Geochemie, Budapestlaan 4, 3584 CD Utrecht.

1. INTRODUCTION

In order to examine the clay mineral composition of recent sediments from the Zaire estuary and fan (Figs 1 and 2) samples were taken from the top 4 cm of some selected trip-cores (Table I) collected during the May 1978 cruise (JANSEN *et al.*, 1983).




Fig. 1. Location of samples in the Zaire estuary and fan.

Analyses of the clay mineral composition were performed on the $< 2 \mu\text{m}$ fraction with a Philips X-ray diffractometer, using $\text{Co K}_{\alpha 1,2}$ radiation wavelength $1.7902 \text{ \AA}$.

After separation of the $< 2 \mu\text{m}$ fraction by centrifugation and subsequent coagulation as a Ca-clay no other pre-treatments, such as decalcification (DE LANGE, 1976) or iron extraction (MEHRA & JACKSON, 1960; HOLMGREN, 1967; DE LANGE, 1976), were applied to these samples. Removal of the organic material using the bromine method (VAN LANGEVELD, 1977; VAN LANGEVELD *et al.*, 1978) did not improve the diffraction pattern and this




Fig. 2. South Atlantic Ocean; currents after GIRESE (1978); AABW = Antarctic Bottom Water, B = Bergwinds (after BORNHOLD, 1973).

method was therefore only used in the test series on different grain sizes from one sample (Table II).

Powdered, oriented and non-oriented samples were subjected

Table I
Percentage of particles smaller than 2 μm , and the approximate water-depth, of the samples.

Sample number	% < 2 μm	Waterdepth (m)
38	85	1455
40	92	5450
41	83	4770
45	83	4790
27	56	4060
49	78	4330
18	73	-
K	26	*

* Bank of the river Zaire at Kinshasa

to X-ray analysis, under 50% relative humidity or low pressure conditions (Figs 3 and 4) and were X-rayed again after glycolation. The main feature that could be distinguished was similar for all the different techniques and treatments. Therefore the presented results will only contain the numbers obtained from the X-ray diffraction patterns using the rotating specimen ($\emptyset$ 2 cm, $d = 0.5$ mm) with powdered, non-oriented samples (Fig. 5) and from X-ray analysis under 50% relative humidity with oriented samples (Fig. 6).

Determination of the relative percentages was achieved by comparing the peak-heights of smectite (17° as 2θ), illite (10°

Table II
The relative percentage of clay minerals in some fractions of sample 40. Sample 1, 3 and 5 are treated with a bromium-solution (LANGEVELD, VAN DER GAAST & EISMA, 1978), samples 2, 4 and 6 are untreated. (M=smectite; I=illite; K=kaolinite).

	(50% RH, oriented samples)			
Sample number	Relative percentages of			Particle fraction
	M	I	K	
40-1	45	12	43	< 2 μm
40-2	42	14	44	< 2 μm
40-3	33	8	59	< 1 μm
40-4	39	11	50	< 1 μm
40-5	46	7	54	< 0.5 μm
40-6	40	7	52	< 0.5 μm




Fig. 3. X-ray diffraction patterns of oriented samples, measured in a 50% relative humidity atmosphere.




Fig. 4. X-ray diffraction patterns of oriented samples at lower pressure (ca 5 mm Hg).




Fig. 5. The relation between the clay mineral composition and the distance from the coast. The relative clay mineral percentages were calculated from the X-ray diffraction patterns with the rotating specimen and powdered, non-oriented samples, and an atmosphere of 50% relative humidity.

as 20) and kaolinite (14° as 20). The sum of the three clay minerals was taken as 100%. No corrections were made for the individual clay minerals, as done by others; the resulting relative percentages are nevertheless comparable for this region (GOLDBERG & GRIFFIN, 1964; BISCAYE, 1965; GRIFFIN *et al.*, 1968; RATEEV *et al.*, 1969; ROBERT, 1980). Chlorite could not be detected at 29.6° (20) at $1/4^{\circ}$ min, with oriented untreated samples.

Acknowledgements.—The author thanks F. Jansen, T. van Weering and D. Eisma for making this research possible. He is especially grateful to S. van der Gaast for the many interesting and stimulating discussions and exchange of views.




Fig. 6. The relation between the clay mineral composition and the distance from the coast. The relative clay mineral percentages were calculated from the X-ray diffraction patterns of oriented samples, with an atmosphere of 50% relative humidity.

2. RESULTS AND DISCUSSION

In the relative clay mineral composition there is a pronounced decrease in the percentage of kaolinite with increasing distance from the coast (Figs 5 and 6) and a simultaneous in-

crease in smectite. This phenomenon has already been observed by some investigators in other regions (POSTMA *et al.*, 1954; WHITEHOUSE *et al.*, 1960; BISCAYE, 1965; PORRENGA, 1967; CHAMLEY, 1979; GIBBS, 1967). This has been attributed to processes such as:

- 1) differential settling tendencies (POSTMA *et al.*, 1954; WHITEHOUSE *et al.*, 1960),
- 2) decreasing terrestrial influence (BISCAYE, 1965),
- 3) increasing "other source" material (CHAMLEY, 1967, 1971; HOFFERT, 1980).

It seems evident that the river Zaire, or more generally the African continent is the source of kaolinite (BORNHOLD, 1973; EISMA *et al.*, 1978) although authigenese or hydrothermal formation of kaolinite has been reported for some regions (ZEN, 1959; MULLER, 1961; SHAW, 1980; SHIEH & SUTER, 1979).

The occurrence of high relative percentages of kaolinite seems to be restricted mainly to the continental margin, due to the large settling velocity of this mineral (WHITEHOUSE *et al.*, 1960; PORRENGA, 1967). The relationship to the continents is clearly shown by a comparison of ocean sediments (GOLDBERG & GRIFFIN, 1964; BISCAYE, 1965; GRIFFIN *et al.*, 1968; RATEEV *et al.*, 1969) and by the resemblance between the zonation of soils and climates on the continents and the ocean sediments (PÉDRO, 1968).

An indication for the authigenic formation of smectite in the sediments of the South Atlantic Ocean is the relative percentage of smectite in aeolian dust and surface ocean water. In the North Atlantic Ocean the concentrations of smectite in the aeolian dust/surface ocean water and the sediments show a good correlation. As the relative percentage of smectite in the sediments of the South Atlantic Ocean has doubled in com-

parison to the concentration in the aeolian dust/surface water, BEHAIRY *et al.* (1975) conclude that half of the smectite in the sediments of the South Atlantic Ocean must be authigenic.

Besides a terrestrial origin smectite can have various other sources, as mentioned by many authors:

Volcanic/hydrothermal origin.—High percentages of smectite in the sediment have been reported near areas with hydrothermal activity (HÉKINIAN & FEVRIER, 1979; KASTNER, 1981 and refs herein). Formation by alteration of volcanic material has been widely reported from many areas (KENNETH, 1981 and refs herein): Pacific Ocean (PETERSON *et al.*, 1964; KASTNER, 1976, 1981; KURNOSOV *et al.*, 1980); Mediterranean (GRIM & VERNET, 1961; CHAMLEY, 1967, 1971); Atlantic Ocean (ROBERT *et al.*, 1979; ROBERT, 1980).

Possible volcanic areas that could contribute to our samples are the Cameroon Ridge, Mid Atlantic Ridge and Walvis Ridge. The Scotia Ridge being a source of smectite for the Argentine Basin (ZIMMERMAN, 1977) seems too far away to make any significant contribution to the Zaire region. Favourable currents do exist for the other ridges, like the Equatorial Counter Current and the Benguela Current; the influence of these currents on the Zaire region are dependant on amongst other things the presence of "glacials" and "non-glacials" (GIRESSE, 1978), the Equatorial Counter Current descending more to the south in interglacials, and the Benguela Current going far up north in glacials. As the warmer periods coincide with relatively larger concentrations of smectite in the sediment (GIRESSE, 1978; VAN DER GAAST (pers. comm.) one is

tempted to look for the possible volcanic origin of smectite at the Cameroon Ridge. A present day southward flowing bottom current has also been mentioned by SHANNON (1970) in the eastern Angola Basin and postulated by SIESSER (1973, 1980) for the Miocene period.

Authigenese of smectite on silica.—It has been reported that smectite can form not only on volcanic material but also on diatom-frustules (CHAMLEY & MILLOT, 1972; VAN BENNEKOM & VAN DER GAAST, 1976; HOFFERT *et al.*, 1980). This way of formation of smectite could mean that there is a significant source of smectite in the oceans, since diatom-frustules are known to be present on a large scale in oceanic sediments. Unfortunately the frequency of occurrence and the velocity of formation are not yet known. Therefore the importance of this way of formation of smectite cannot be estimated quantitatively. However, since no striking differences have been observed between the clay mineral composition of silicious oozes and calcareous oozes, this formation method does not seem to be of major importance, unless other parameters are necessary as well. Newly formed smectite is generally highly ferrous and aluminous (PORRENGA, 1967; GIRESE, 1975). The river Zaire is rather rich in "dissolved" (partly colloidal) iron (FIGUÈRES *et al.*, 1978) a fact that might promote the new formation of smectite in this region. In some of the cases where there is evidence for authigenese of smectite, there is also volcanic material at hand (CHAMLEY & MILLOT, 1972). CHAMLEY suggests that volcanic material is needed for the formation of smectite from the dissolution of diatoms.

Aeolian transport.—It is known that material from the Sahara

is transported by wind to the ocean over many of the countries along the west coast of Africa, from Sierra Leone to Nigeria (BISCAYE, 1965; McMASTER *et al.*, 1977). Dust collected in the northern part of the South Atlantic Ocean contains a considerable amount of smectite (CHESTER *et al.*, 1972, 1974). Aeolian input from the southern part of Africa seems to be of no importance since no terrestrial material was found on the Walvis Ridge (ZIMMERMAN, 1977). Others claim it to be of importance (BORNHOLD, 1973; CHESTER *et al.*, 1972) especially for smectite and illite. Since no detailed dust collection has been made, especially during the Bergwinds, no approximation can be made of the importance of this possible source. In view of the relative concentrations of smectite and illite in the sediments in the eastern part of the South Atlantic Ocean, it seems reasonable to assume only there is only a very small influx into the Zaire region.

Bottom currents.—An input of material into the Angola Basin could perhaps occur via bottom currents. The northward flowing Antarctic Bottom water contains only minor amounts of clay minerals; contribution to the sediments of the Argentine Basin was considered minor or not present (ZIMMERMAN, 1977). Therefore no input into the Zaire region is to be expected from this current. The possible existence of southward flowing bottom currents as well as the northward flowing Benguela Current has already been mentioned.

Differential settling.—From one sample (40) three different particle sizes were examined for their clay mineral composition (Table II). No differential settling could be deduced, in accordance with the findings of VAN DER GAAST *et al.* (1983),

who found no pronounced differences between the relative percentages of smectite and kaolinite, even in the low size ranges of some samples from the Zaire region.

In summary it can be concluded that all possible explanations for the seaward increase of smectite compared to kaolinite can be ruled out, or they are only of minor or unknown importance. The question of the origin of smectite in the Zaire region must therefore remain partly unanswered. If future research is done on the clay mineral composition of suspended matter in the South Atlantic Ocean, including depth differences (*cf.* McMASTER *et al.*, 1977), and this is combined with a detailed study of the stable isotope composition of smectite (SAVIN & EPSTEIN, 1970) and/or rare earth metals, like cerium (HOFFERT, 1980; DESPRAIRIES *et al.*, 1980; CHAMLEY *et al.*, 1981; PIPER, 1974; STEINBERG & COURTOIS, 1976) we might be more certain about the major paths and origin(s) of smectite.

3. REFERENCES

- ANDEL, T. van & H. POSTMA, 1954. Recent sediments in Gulf of Paria; North Holland Publishing Company, Amsterdam pp. 143.
- BEHAIRY, A.K., R. CHESTER, A.J. GRIFFITHS, L.R. JOHNSON & J.H. STONER, 1975. The clay mineralogy of particulate material from some surface seawaters of the eastern Atlantic Ocean.—*Mar. Geol.*, 18: M45-M56.
- BENNEKOM, A.J. van & S.J. van der GAAST, 1976. Possible clay structures in frustules of living diatoms.—*Geochim. Cosmochim. Acta*, 40: 1149-1152.
- BISCAYE, P.E., 1965. Mineralogy and sedimentation of recent deep-sea clay in the Atlantic Ocean and adjacent seas and oceans.—*Geol. Soc. Am. Bull.*, 76: 803-832.
- BORNHOLD, B.D., 1973. Late Quaternary sedimentation in the eastern Angola Basin.—*Thesis*, Woods Hole (Mass.).
- CHAMLEY, H., 1967. Quelques modalités de la sédimentation argileuse marine aux environs de l'île de Crète.—*Bull. Mus. Anthropol. préh. Monaco*, 14: 25-48.
- , 1971. Recherches sur la sédimentation argileuse en Méditerranée.—*Sci. Geol. Mém.*, Strasbourg, 35: pp. 225.

- , 1979. North Atlantic clay sedimentation and paleo-environment since the late jurassic. In: Deep drilling results in the Atlantic Ocean continental margins and paleo environment (TALWANE M. et al., eds): 342-360.
- CHAMLEY, H. & G. MILLOT, 1972. Néof ormation de montmorillonite à partir de diatomées et de cendres dans les sédiments marins de Santorin. (Méditerranée orientale). C.R. Acad. Sci., 274D: 1132-1134.
- CHAMLEY, H. & C. BONNOT-COURTOIS, 1981. Argiles authigènes et terrigènes de l'Atlantique et du Pacifique NW (Legs 11 et 58 DSDP), rapport des terres rares. Oceanol. Acta, 4: 229-238.
- CHESTER, R., J.J. GRIFFIN & R.C. PADGHAN, 1972. Eolian dust along the eastern margins of the Atlantic Ocean.—Mar. Geol., 13: 78-91.
- CHESTER, R., H. ELDERFIELD, J.J. GRIFFIN, L.R. JOHNSON & R.C. PADGHAN, 1972. Eolian dust along the eastern margins of the Atlantic Ocean.—Mar. Geol. 13:91-105.
- CHESTER, R., J.H. STONER & L.R. JOHNSON, 1974. Montmorillonite in surface detritus. Nature, 249: 335-336.
- DESPRAIRIES, A. & C. BONNOT-COURTOIS, 1980. Relation entre la composition des smectites d'altération sous-marine et leur cortège de terres rares.—Earth Planet. Sci. Lett., 48: 124-130.
- EISMA, D., J. KALF & S.J. van der GAAST, 1978. Suspended matter in the Zaire estuary and the adjacent Atlantic Ocean.—Neth. J. Sea Res., 12: 382-406.
- FIGUÈRES, G., J.M. MARTIN & M. MEYBECK, 1978. Iron behaviour in the Zaire estuary.—Neth J. Sea Res., 12: 329-337.
- GIBBS, R.J., 1967. The geochemistry of the Amazon river system, part I. The factors that control the salinity and the composition and concentration of the suspended solids.—Bull. geol. Soc. Am., 73: 1203-1232.
- GIRESE P., 1975. Essai de chronomètre de la glauconitisation dans la Golfe de Guinée : exemple de vitesse digénétique au Quaternaire supérieur.—C.R. somm. Soc. Géol. Fr., (1975): 163-164.
- , 1978. Le controle climatique de la sédimentation marine et continentale en Afrique centrale atlantique a la fin du Quaternaire - problèmes de corrélation.—Palaeogéogr. Palaeodim. Palaeoecol., 23: 57-77.
- GOLDBERG, E.D. & J.J. GRIFFIN, 1964. Sedimentation rates and mineralogy in the South Atlantic.—J. Geophys. Res., 69: 4293-4309.
- GRIFFIN, J.J., H. WINDOM & E.D. GOLDBERG, 1968. The distribution of clay minerals in the World Ocean.—Deep Sea Res., 15: 433-459.
- GRIM, R.E. & J.P. VERNET, 1961. Etude par diffraction des minéraux argileux de vases méditerranéennes.—Schweiz. Min. Petrol. Mitt., 41: 65-70.
- HÉKINIAN, R. & M. FEVRIER, 1979. Comparison between deep-sea hydrothermal deposits recovered from recent spreading ridges.—Coll. Int. CNRS, 289: 167-178.
- HOFFERT, M., 1980. Les "argiles rouges des grands fonds" dans le Pacifique centre-est. Authigénèse, transport, digénèse. Thèse, Strasbourg.
- HOFFERT, M., A.M. KARPOFF, A. SCHAAF & G. PAUTOT, 1979. The sedimentary deposits of the Tiki Basin passage from the carbonate oozes to "metalliferous sediments".—Coll. Int. CNRS, 289: 101-112.

- HOLMGREN, G.G.S., 1967. A rapid citrate-dithionite extractable iron procedure.—*Proc. Soil Soc. Am.*, 31: 210-211.
- JANSEN, J.H.F., T.C.E. van WEERING, R. GIELES & J. van IPEREN, 1983. Middle and Late Quaternary paleoceanography and climatology of the Zaire-Congo fan and the adjacent eastern Angola Basin.—*Neth. J. Sea Res.* 17 (2).
- KASTNER, M., 1976. Diagenesis of basal sediments and basalts of sites 322, 323, leg 35, Bellinghausen Abyssal Plain. In: *Initial Reports of the DSDP*, 35: 513-519.
- , 1981. Authigenic silicates in Deep Sea sediments. Formation and diagenesis. In: *The Oceanic Lithosphere; The Sea*, vol. 7 (Emiliani C., ed): 915-881.
- KENNETH, J.R., 1981. Marine Tephrochronology. In: *The Oceanic Lithosphere; The Sea*, vol. 7 (Emiliani C., ed.): 1373-1437.
- KURNOSOV, V., N. TSEITLIN & G. NARNOV, 1980. Clay minerals; paleogeographic and diagenetic aspects. In: *Initial Reports of the DSDP*, 56-57: 979-1004.
- LANGE, G.J. de., 1976. Röntgen-diffractie aan kleimineralen van enkele Noordzee-monsters. Intern Verslag NIOZ, 1976-6.
- LANGEVELD, A.D. van., 1977. Het verwijderen van organisch materiaal uit kleimineralen. Intern Verslag NIOZ, 1977-13.
- LANGEVELD, A.D. van., S.J. van der GAAST & D. EISMA, 1978. A comparison of the effectiveness of eight methods for the removal of organic matter from clay.—*Clays Clay Minerals*, 26: 361-364.
- McMASTER, R.L., P.R. BETZER, K.L. CARDER, L. MILLER & D.W. EGGIMANN, 1977. Suspended particle mineralogy and transport in water masses of the West African Shelf adjacent to Sierra Leone and Liberia.—*Deep Sea Res.* 24: 651-665.
- MEHRA, O.P. & M.L. JACKSON, 1958. Iron oxide removal from soils and clays by a dithionite-citrate system buffered with sodium carbonate. *Clays Clay Minerals*, 5: 317-327.
- MULJER, G., 1961. Die rezenten Sedimente im Golf von Neapel. 2. Mineral-Neu- und Umbildungen in den rezenten Sedimenten des Golfes von Neapel. Ein Beitrag zur Umwandlung vulkanischer Glaser durch Halmyrolyse.—*Beitr. Miner. Petr.*, 8: 1-20.
- PÉDRO, G., 1968. Distribution des principaux types d'altération chimique à la surface du globe. Présentation d'une esquisse géographique.—*Rev. Geogr. phys. Géol. dyn.*, Paris, 10: 457-470.
- PETERSON, M.N.A. & J.J. GRIFFIN, 1964. Volcanism and clay minerals in the southeastern Pacific.—*J. Marine Res.*, 22: 13-21.
- PIPER, D.Z., 1974. Rare earth elements in the sedimentary cycle = a summary.—*Chem. Geol.*, 14: 285-304.
- PORRENGA, D.H., 1967. Clay mineralogy and geochemistry of recent sediments in tropical areas. Thesis, Amsterdam, pp 145.
- , 1967. Glauconite and chamosite as depth indicators in the marine environment.—*Mar. Geol.*, 5: 495-501.
- RATEEV, M.A., Z.N. GORBUNOVA, A.P. LISITZYN & G.L. NASOV, 1969. The distribution of clay minerals in the oceans.—*Sedimentology*, 13: 21-43.
- ROBERT C., 1980. Climats et courants cénozoïques dans l'Atlantique Sud d'après l'étude des minéraux argileux (legs 3,39 et 40 DSDP).—*Oceanol. Acta* 3: 369-376.

- ROBERT C., J.P. HERBIN, G. DEROO, G. GIROUT d'ARGOUD & H. CHAMLEY, 1979. L'Atlantique Sud au Cretacee d'après l'étude des minéraux argileux et de la matière organique (legs 39 et 40 DSDP).—*Oceanol. Acta*, 2: 209-218.
- SAVIN, S.M. & S. EPSTEIN, 1970. The oxygen and hydrogen- isotope geochemistry of clay minerals.—*Geochim. Cosmochim. Acta*, 34: 25-42.
- , 1970. The oxygen and hydrogen isotope geochemistry of ocean sediments and shales.—*Geochim. Cosmochim. Acta*, 34: 43-63.
- SHANNON, L.V., 1970. Oceanic circulation off South Africa.—*Fish. Bull. S. Africa*, 6: 27-33.
- SHAW, H.F., 1980. Clay minerals in sediments and sedimentary rock. In: *Developments in petroleum geology-2* (ed. HODSON G.D., ASP London).
- SHIEH, Y.N. & T.G. SUTER, 1979. Formation conditions of authigenic Kaolinite and calcite in coals by stable isotope determinations.—*Clays Clay Minerals*, 27: 154-156.
- SIESSER, W.G., 1973. Native copper in DSDP leg 40 sediments. In: *Initial reports of the DSDP*, 40: 761-765.
- , 1980. Late Miocene origin of the Benguela upwelling system off Northern Namibia. *Science*, 208: 283-285.
- STEINBERG, M. & C. COURTOIS, 1976. Le comportement des terres rares au cours de l'altération et ses conséquences.—*Bull. Soc. géol. Fr.*, t. 18: 13-20.
- WHITEHOUSE, U.G., L.M. JEFFREY & J.D. DEBBRECHT, 1961. Differential settling Clays *Clay Minerals*, *proc. 7th Natl. Conf.*: 1-79.
- ZEN E.-an, 1959. Mineralogy and petrography of marine bottom sediment samples off the coast of Peru and Chile.—*J. Sedim. Petrol.*, 29: 513-539.
- ZIMMERMAN, H.B., 1977. Clay mineral stratigraphy and distribution in the South Atlantic Ocean. In: *Initial Reports of the DSDP*, 39: 395-400.