

THE
BALTIC MARINE BIOLOGISTS
PUBLICATION No. 9



A Compilation of Biometric Conversion Factors for Benthic Invertebrates of the Baltic Sea

by

Heye Rumohr, Thomas Brey and Sven Ankar

1987

ISSN 0282-8839

THE
BALTIC MARINE BIOLOGISTS
Publication No. 9

A Compilation of Biometric Conversion Factors
for Benthic Invertebrates of the Baltic Sea

by

Heye Rumohr¹, Thomas Brey¹ and Sven Ankar²

1987

- 1) Institut für Meereskunde, Düsternbrooker Weg 20,
D-2300 Kiel, F.R.G.
- 2) Askö Laboratory, Inst. of Marine Ecology, Univ. of Stockholm,
S-10691 Stockholm, Sweden

Contents	page
Introduction	1
Methods	2
Results	3
Discussion	3
Conclusions	5
References	7
Appendix I: Data	8
1. Average conversion factors for main taxonomic groups	8
1.1. Average conversion factors: weight -> weight	9
1.2. Average conversion factors: weight -> energy content	10
2. Conversion factors for individual Baltic species	11
2.1. Mollusca	11
2.1.1. Bivalvia	11
Abra spp. (A. alba, longicallus, nitida)	11
Arctica islandica	12
Astarte spp. (A. borealis, elliptica, montagui, sulcata)	12
Cardium spp. (C. edule, fasciatum, glaucum, hauniense)	13
Corbula (Aloidis) gibba	14
Leda pernula	14
Macoma spp. (M. balthica, calcarea)	14
Modiola modiolus	15
Mya spp. (M. arenaria, truncata)	15
Mysella (Montacuta) bidentata	16
Mytilus edulis	16
2.1.2. Gastropoda	18
Aporrhais pespelicani	18
Buccinum undatum	18
Hydrobia spp.	18
Littorina littorea	18
Lymnaea spp. (L. spp., peregra)	19
Paludetrina (Potamopyrgus) jenkinsi	19
Theodoxus fluviatilis	19
2.2. Crustacea	20
2.2.1. Amphipoda	20
Bathyporeia pilosa	20
Corophium spp. (C. volutator, insidiosum)	20

	page
Gammarus spp. (G. spp., duebeni, oceanicus, salinus, zaddachii)	20
Haploops tubicola	21
Phoxocephalus holboelli	21
Pontoporeia spp. (P. affinis, femorata)	21
2.2.2. Isopoda	22
Asellus aquaticus	22
Idothea spp. (I. spp., aleutica, baltica, chelipes, ochotensis)	22
Jaera albifrons	23
Mesidotea entomon	23
2.2.3. Mysidacea	23
Mysidae	23
Gastrosaccus spinifer	23
Mysis spp. (M. mixta, relicta)	24
Neomysis integer	24
2.2.4. Other Crustacea	24
Balanus spp. (B. balanoides, balanus, crenatus, improvisus)	24
Crangon spp. (C. crangon, septemspinosa)	25
Diastylis rathkei	25
Eupagurus bernhardi	25
Hyas coarctatus	26
Ostracoda	26
2.3. Polychaeta	27
2.3.1. Polychaeta errantia	27
Aphrodite aculeata	27
Eteone longa	27
Harmothoe sarsi	27
Nephtys spp. (N. spp., ciliata, incisa)	27
Nereis spp. (N. diversicolor, pelagica)	28
Scoloplos armiger	29
2.3.2. Polychaeta sedentaria	29
Arenicola marina	29
Aricidea spp. (A. jeffreysii, suecia)	29
Pectinaria spp. (P. hypoboreia, koreni)	29
Pygospio elegans	29

	Page
Polydora spp.	30
Sosane gracilis	30
Terebellides stroemi	30
2.4. Other Taxa	31
2.4.1. Turbellaria	31
2.4.2. Nematoda	31
2.4.3. Nemertini	31
2.4.4. Priapulida	31
Halicryptus spinulosus	31
Priapulid caudatus	31
2.4.5. Oligochaeta	32
2.4.6. Insecta	32
Chironomida larvae	32
Trichoptera larvae	32
2.4.7. Echinodermata	32
Asterozoa (A. spp., rubens, vulgaris)	32
Ophiurozoa (Ophiura albida; Amphiura chiajei, filiformis)	33
Echinozoa (Echinocardium; Echinocyamus; Strongylocentrotus)	33
Holothurozoa (Cucumaria)	34
3. Length - weight - relationships	35
3.1. Mollusca	36
3.1.1. Bivalvia	36
Abra spp. (A. alba, nitida, longicallus)	36
Arctica islandica	36
Astarte spp. (A. borealis, elliptica, montagui)	36
Cardium spp. (C. edule, glaucum, hauniense)	37
Macoma balthica	38
Mya arenaria	39
Mytilus edulis	39
Pisidium sp.	41
3.1.2. Gastropoda	42
Hydrobia spp. (H. spp., ulvae)	42
Littorina littorea	42
Lymnaea peregra	43
Paludetrina (Potamopyrgus) jenkinsi	43

	page
Theodoxus fluviatilis	43
Valvata piscinalis	43
3.2. Crustacea	44
3.2.1. Amphipoda	44
Bathyporeia spp. (B. pilosa, sarsi)	44
Corophium spp. (C. volutator, insidiosum)	44
Gammarus spp. (G. spp., salinus)	45
Microdeutopus gryllotalpa	45
Pontoporeia spp. (P. affinis, femorata)	45
3.2.2. Other Crustacea	47
Crangon crangon	47
Diastylis rathkei	47
Idothea baltica	47
Jaera spp.	47
Mesidotea (Saduria) entomon	47
Mysis mixta	47
Neomysis integer	47
3.3. Polychaeta	48
3.3.1. Polychaeta errantia	48
Harmothoe sarsi	48
Nereis diversicolor	48
3.3.2. Polychaeta sedentaria	48
Aricidea suecia	48
Terebellides stroemi	48
3.4. Other Taxa	49
3.4.1. Priapulida	49
Halicryptus spinulosus	49
Priapulus caudatus	49
3.4.2. Insecta	49
Chironomus larvae	49
3.4.3. Echinodermata	49
Asterias rubens	49
Appendix II: References to the data in Appendix I	50
Table of applied methods	55

Introduction

There have been a number of attempts to relate basic physical measurements of aquatic invertebrates i.e. length and weight to each other and to other measures that are easy to obtain (i.e. Slobodkin & Richmond 1961, Brawn et al. 1968, Lie 1968, Prus 1970, Kreutzberg & v.Oertzen 1973, Tyler 1973, Lappalainen & Kangas 1975, Bast & v.Oertzen 1976, Petersen 1977, Chambers & Milne 1979, Norrbin & Bamstedt 1984).

An important reason for this is to reduce laboratory work required for preparing data for benthic production studies.

This paper presents a compilation of published and unpublished data on conversion factors with special reference to Baltic invertebrates, supplemented by data from other marine areas. General conversion factors for main macrobenthic taxa have been computed and their validity assessed.

Beside the general importance of such a compilation of conversion factors they may also serve Baltic benthos investigators to convert "old" wet weight biomass data into more appropriate units of biomass: dry weight, ash free dry weight or energy content (Joule).

This study was promoted by the Baltic Marine Biologists Working Group 11 ("Secondary Production"), to facilitate comparative benthic production studies.

During the preparation of this compilation many members of the working group and other colleagues have opened their drawers and hidden files to supply us with their unpublished results (H. Cederwall, S. Hansson, P. Kangas, N. Kautsky, A. Lappalainen, J. Ostrowski, E. Persson, B. Saebø, M. Weigelt; see also list of contributors in Appendix II).

The collected conversion factors are presented at three levels of precision:

- (1) Average conversion factors 'weight -> weight' and 'weight -> energy content' for main taxonomic groups.
- (2) Conversion factors 'weight -> weight' and 'weight -> energy content' for individual Baltic species.
- (3) Length - weight - relationships for individual Baltic species.

The applicability of these factors is discussed in the light of seasonal variations and geographic gradients which include factors like exposition, spawning cycles and other biotic and abiotic influences. Following the results of this study tentative recommendations will be proposed that will also go into the revised edition of the BMB-recommendations.

Methods

A variety of methods have been applied by various authors to obtain many kinds of conversion factors. The methods used for the data in this compilation are listed in Appendix II.

A critical factor in comparing different studies concern whether fresh or formalin preserved material was used since all preservation media have a distinct influence on the weight (e.g. Mills et al. 1982, Brey 1986).

Further, it is necessary to know the time of exposition and the temperature used in drying and combustion since the weight can be altered quite significantly when certain limits are passed (e.g. Winberg 1971).

The most important question concerns the date at which the sample was taken, since the spawning season and other seasonal influences have distinct effects on the weight to length ratio of many species.

Geographical information is also needed, especially when the limits of distribution of a certain species are reached, and when hydrographical features differ considerably between the sampling areas.

Additionally laboratory methods have to be considered carefully since even the measuring technique may be a source of error in a study.

Results

The results are presented in Appendix I in groups according to their precision.

- (1) Average conversion factors 'weight -> weight' and 'weight -> energy content' for main taxonomic groups.

Data on Gastropoda, Bivalvia, Crustacea and Polychaeta have been analyzed. Bivalves with extraordinary thick shells like *Arctica islandica* and *Astarte* spp., and crustaceans with heavy calcareous skeletons as well as very large species and juveniles have been neglected.

Appendix I: 1.1. shows the average conversion factors for wet weight -> shell free dry weight (WW -> SFDW), dry weight -> ash free dry weight (DW -> AFDW) in molluscs, and the factors for WW -> DW, DW -> AFDW and WW -> AFDW for crustaceans and polychaetes. The WW -> AFDW factors have been calculated from the others.

Appendix I: 1.2. shows the average conversion factors for DW -> Joule/mg, SFDW -> Joule/mg and AFDW -> Joule/mg for the different taxonomic groups.

- (2) Conversion factors 'weight -> weight' and 'weight -> energy content' for individual Baltic species.

Summary data for 92 species are presented in Appendix I: 2., 29 mollusc species, 33 crustacean species, 14 polychaete species and 17 species of other taxa. All data available, including seasonal and geographical variations, are presented.

(3) Length - weight - relationships for individual Baltic species. These data (Appendix I: 3.) are for 44 species (21 mollusc species, 15 crustacean species, 4 polychaete species and 4 species of other taxa).

Discussion

The average conversion factors for main taxa have small confidence limits (see Chapter 1.1. and 1.2. in Appendix I), although very different methods have been used (see Appendix II) to determine weights and energy content.

The factors for converting WW to AFDW, which are calculated here from the other factors, are close to the factors published by Lie (1968). Lie found the average AFDW in %WW factors for bivalves, crustaceans and polychaetes to be 5.5, 15.0 and 13.3. The factors computed in this paper are 6.9, 14.5 and 13.2, respectively. Only the conversion factors for bivalves differ markedly. We assume this to be related to geographical effects, because Lie's data refer to animals from the Pacific coast of the USA, whereas the greater part of the data compiled here is from the Baltic Sea with lower salinity and unsaturated with CaCO_3 . The shells of bivalves are usually thinner in areas with lower salinity (e.g. Arntz & Weber 1970) and this leads to a higher AFDW/WW ratio.

The conversion factors for the species (Appendix I, 2.) show that even in single species there is a wide range of values due to geographical, seasonal and individual differences, and possibly also to interannual variations in the condition (see e.g., Lappalainen & Kangas 1975 [Fig.2] or Davis & Wilson 1983 [Fig.1]).

This is especially true for the conversion of weight into energy, where seasonal differences in condition, different life stages or sex result in large differences in energy content. Throughout the course of a year, maximum to minimum flesh weight of a given size may differ seasonally by a factor of 2 or 3 (e.g. Beukema & deBruin 1977, Chambers & Milne 1979).

The length - weight relationships (Appendix I, 3.) show the largest variation among the data compiled here, although the relationships given by individual workers indicate always a strong correlation between length and weight.

Length - weight relationships show the same effects as mentioned above, but additional factors influencing the equations should be noted.

Length measurements are certainly not standardized, e.g. in the literature one can find that either shell length or shell height of bivalves have been correlated with weight. Some investigators stretch bent animals before measuring, others do not, etc. There is a lot of variation in body length determinations, all affecting the calculation of the length - weight equations.

The relationships obtained are also influenced by the size range measured as well as by the weighing procedure, e.g. sometimes the

shell halves of bivalves are opened and the water in the mantle cavity is removed before weighing. In other cases, especially in small bivalves, this is impossible without damaging the animal. Furthermore, the equations are only valid for the body length intervals which they are based on, and should not be used outside this size range.

The equations should be used with caution, if data on sampling date, geographic sampling area, life stages, sex, size range and laboratory methods used are missing.

Beside all these variations, which are induced either by nature or by the methods used, benthic investigators should consider some basic statistical problems of length - weight relationships:

Length - weight relationships are usually established by fitting a linear regression to the logarithmic data:

$$\log(w_i) = a + b * \log(l_i)$$

The estimates of the slope b and the y -axis intercept a by a "prediction model" could be seriously biased. They should be substituted by an estimate through a "functional model" according to Ricker (1973) or corrected by a factor according to Baskerville (1972) and Sprugel (1983).

Conclusions

It is obvious that much information is still required. Data are particularly required for DW $\rightarrow$ AFDW and AFDW $\rightarrow$ energy relations in polychaetes. Also AFDW $\rightarrow$ energy relations of the following species are scarce or even lacking: Arctica, Astarte, Crangon, Nephtys, Terebellides, Spionids (no data!), Priapulids.

These gaps should be filled in the near future, to make model computations more easy and reliable.

How to use this compilation:

- (1) Average conversion factors 'weight $\rightarrow$ weight' and 'weight $\rightarrow$ energy content' for main taxonomic groups.

These factors seem to be hard facts. They will be useful in all cases where the aim of the investigations carried out is above the species level, e.g. community biomass or biomass of

trophic groups.

- (2) Conversion factors 'weight -> weight' and 'weight -> energy content' for individual Baltic species.

The general validity of factors which depend on any term including calcareous parts (e.g. WW and DW of Molluscs) is poor because shell weight is related to salinity.

Geographical, seasonal and individual differences should always be taken into account, if any conversion factor for single species is used.

- (3) Length - weight - relationships for individual Baltic species. Individual weight is usually only required for a high level of precision in benthic research work. Therefore, the equations compiled here must be used with utmost care. In many cases, it may be preferable for workers who need such relationships, to produce their own.

References (see also Appendix II)

- Arntz, W.E. and W. Weber (1970): *Cyprina islandica* L. (Mollusca, Bivalvia) als Nahrung von Dorsch und Kliesche in der Kieler Bucht. Ber. Deutsch. Wiss. Komm. 21: 193-209.
- Baskerville, G.L. (1972): Use of logarithmic regression in the estimation of plant biomass. Can. J. Forest Res. 2: 49-53.
- Beukema, J.J. and W. deBruin (1977): Seasonal changes in dry weight and chemical composition of the soft parts of the tellinid bivalve *Macoma balthica* in the Dutch Wadden Sea. Netherl. J. Sea Res. 11: 42-55.
- Brey, T. (1986): Formalin and Formaldehyde-depot chemicals: Effects on dry weight and ash free dry weight of two marine bivalve species. Meeresforsch./Rep. on Mar. Res. 31: 52-57.
- Davis, J.P. and J.G. Wilson (1983): Seasonal changes in tissue weight and biochemical composition of the bivalve *Nucula turgida* in Dublin Bay with reference to gametogenesis. Netherl. J. Sea Res. 17: 84-95.
- Mills, E.L., K. Pittmann and B. Munroe (1982): Effects of preservation on the weight of marine benthic invertebrates. Can. J. Fish. Aquat. Sci. 39: 221-224.
- Ricker, W.E. (1973): Linear regressions in fishery research. J. Fish. Res. Bd. Canada 30: 409-434.
- Winberg, G.G. (1971): Methods for the estimation of the production of aquatic animals. Academic Press, London, N.Y., 175p.
- Sprugel, D.G. (1983): Correcting for bias in log-transformed allometric equations. Ecology 64: 209-210.

Appendix I: Data

(In the following the sign '#' indicates a reference number, e.g. #10 is reference No 10 in Appendix II.)

1. Average conversion factors for main taxonomic groups

The average conversion factors for main taxonomic groups (which are Gastropoda, Bivalvia, Crustacea, Polychaeta) are compiled in two tables in chapter 1.1. (conversion factors weight -> weight), and chapter 1.2. (conversion factors weight -> energy content). In the case of the weight -> energy content factors, the median is used instead of the arithmetic mean, because some of the data sets are not normal distributed.

We did not include DW -> energy content factors for Bivalvia and Gastropoda, because these data are scarce and show strong dispersion. However, an additional conversion factor AFDW -> Joule/mg for all taxa was computed.

We did not calculate any general factor for Echinodermata, because there are large differences in body composition (e.g. CaCO_3 -content) in this group.

The 'number of species' given for each conversion factor refers to the minimum number of species included. Due to the fact that many authors published factors related to higher taxonomic groups (e.g. family or genus), the real number of species included here may be distinctly higher.

1.1. Average conversion factors: weight -> weight.

Arithmetic mean and 95% confidence interval.

WW = wet weight, AFDW = ash free dry weight

DW = dry weight, SFDW = shell free dry weight

1) AFDW%WW is calculated from the other factors.

2) except *Arctica islandica* and *Astarte* spp.

3) except *Balanomorpha*, large Decapoda and juvenile stages.

Taxon	type	factor %	95%-conf. interval	number values	number species	references
Bivalvia ²⁾	SFDW%WW	8.1	±1.1	53	24	< a >
	AFDW%DW	15.6	±1.5	31	18	< b >
	AFDW%SFDW	81.3	±2.1	53	22	< c >
	AFDW%WW	6.9 ¹⁾	-	-	-	-
Gastropoda	SFDW%WW	10.2	±1.1	46	27	< d >
	AFDW%DW	23.5	±7.3	8	6	< e >
	AFDW%SFDW	83.7	±1.9	49	27	< f >
	AFDW%WW	8.5 ¹⁾	-	-	-	-
Crustacea ³⁾	DW%WW	20.2	±1.0	65	39	< g >
	AFDW%DW	71.7	±3.4	53	35	< h >
	AFDW%WW	14.5 ¹⁾	-	-	-	-
Polychaeta	DW%WW	17.6	±1.2	80	59	< i >
	AFDW%DW	75.2	±3.4	69	48	< k >
	AFDW%WW	13.2 ¹⁾	-	-	-	-

< a >: #5, #7, #10, #13, #49.

< b >: #14, #32, #33, #41, #51, #54.

< c >: #10, #17, #49.

< d >: #10, #13, #17.

< e >: #14, #32.

< f >: #10, #17, #36.

< g >: #5, #7, #10, #13, #32, #33, #39, #41, #49, #50, #51, #54.

< h >: #10, #12, #14, #15, #32, #33, #37, #39, #49, #51.

< i >: #5, #7, #10, #13, #15, #32, #33, #41, #44, #49, #51.

< k >: #10, #14, #15, #16, #32, #33, #37, #49, #51.

1.2. Average conversion factors: weight -> energy content.

Median and 95% confidence limits of energy content (Joule/mg).

DW: dry weight, SFDW: shell free DW, AFDW: ash free DW.

1 Joule = 0.239 cal, 1 cal = 4.187 Joule.

1) except Balanomorpha, large Decapoda and juvenile stages.

2) except large Decapoda and juvenile stages.

Taxon	weight	median	95%-conf.-limits		number	number	ref.
	type	[Joule/mg]	lower	upper	values	species	
Bivalvia	SFDW	18.85	18.63	19.33	50	38	< a >
	AFDW	22.79	22.07	23.45	37	32	< b >
Gastropoda	SFDW	17.16	16.49	18.29	44	43	< c >
	AFDW	22.63	21.77	23.58	42	42	< d >
Crustacea	DW 1)	15.25	14.49	16.89	60	45	< e >
	AFDW2)	22.45	21.51	23.44	45	40	< f >
Polychaeta	DW	16.46	14.19	17.50	36	28	< g >
	AFDW	23.33	22.22	24.89	19	17	< h >
All taxa	AFDW	22.78	21.95	23.33	143	131	< i >

< a >: #13, #15, #22, #30, #37, #42, #49, #50.

< b >: #10, #15, #17, #18, #24, #25, #29, #30, #37, #41, #42, #49.

< c >: #22, #37, #42, #49.

< d >: #10, #18, #37, #42, #49.

< e >: #12, #13, #15, #18, #20, #22, #29, #30, #37, #41, #42, #49, #50.

< f >: #10, #12, #15, #18, #29, #30, #37, #41, #49, #50.

< g >: #12, #13, #15, #18, #22, #30, #37, #41, #49, #50.

< h >: #10, #12, #15, #17, #18, #37, #42, #49.

< i >: #10, #12, #15, #17, #18, #20, #24, #25, #29, #30, #37, #41, #42, #43, #49, #50.

2. Conversion factors for individual Baltic species

Abbreviations:

WW = wet weight, DW = dry weight
AFDW = ash free dry weight, SF = shell free
N = number of values included,
[additional number in brackets = number of individuals]
1 Joule = 0.239 cal, 1 cal = 4.187 Joule

If two values are connected by a hyphen, these two values are the lower and upper limit of the whole range of values given.
If two values are separated by a slash, these values represent two measurements.

2.1. Mollusca

2.1.1. Bivalvia

Abra spp. (*A. alba*, *longicallus*, *nitida*)

- *A. (Syndosmva) alba*

DW = 31.3 % WW (N=104 [1345 ind], #51)
SFDW = 12.4 - 16.5 % SFWW (seasonal var., #15)
AFDW = 83.6 - 88.0 % SFDW (seasonal var., #15)
= 85.9 (mean, #15)
AFDW = 18.0 % DW (N= 217 [2246 ind], #51)
Joule/mg SFDW: 17.02 - 20.26 (seasonal var., #15)
18.80 (mean, #15)
Joule/mg AFDW: 20.23 - 23.61 (seasonal var., #15)
21.89 (mean, #15)

- *A. longicallus*

AFDW = 16.8 % DW (#54)

- *A. nitida*

AFDW = 18.5 % DW (#54)

Arctica islandica

SFWW	=	19.3 - 22.2 % WW	(#9)
	=	12.8	(#40)
DW	=	53.2 % WW	(N= 45 [153 ind], #51)
SFDW	=	3.3 % WW	(#8)
SFDW	=	13.9 - 19.5 % SFWW	(Jun.-Nov., #44)
AFDW	=	2.9 % WW	(#8)
	=	3.4	(#44)
AFDW	=	11.9 % SFWW	(#44)
AFDW	=	9.1 % DW	(N= 11[42 ind], #14)
	=	10.0	(N=583, #51)
AFDW	=	88.4 % SFDW	(#8)
Joule/mg WW:		0.63	(#8)
Joule/mg SFDW:		18.76	(#8)
		16.88	(#50)
Joule/mg AFDW:		21.22	(#8)

Astarte spp. (*A. borealis*, *elliptica*, *montagui*, *sulcata*)

- *A. borealis*

DW	=	69.9 % WW	(N= 56, #41)
	=	72.9	(N= 19, #51)
SFDW	=	5.0 % WW	(Aug., #10)
AFDW	=	7.2 % DW	(N=127, #51)
Joule/mg AFDW:		21.27 / 22.83	(N=3/3, Aug., #10)

- *A. elliptica*

DW	=	76.6 % WW	(N= 20, #51)
SFDW	=	2.1 % WW	(N=105, #46)
SFDW	=	3.1 % DW	(N= 98, #46)
AFDW	=	7.7 % DW	(N=106, #51)

- *A. montagui*

DW	=	70.6 % WW	(N= 12, #51)
SFDW	=	3.0 % WW	(Aug., #10)
AFDW	=	7.1 % DW	(N= 85, #51)
Joule/mg AFDW:		21.96	(N= 2, Aug., #10)

- A. sulcata

AFDW = 77.0 % SFDW (N= 7, #37)
 Joule/mg SFDW: 18.35 (N= 7, #37)
 Joule/mg AFDW: 23.84 (N= 7, #37)

Cardium spp. (C. edule, fasciatum, glaucum, hauniense)

- C. edule

AFDW = 90.4 - 94.3 % SFDW (seasonal var., #17)
 = 92.2 (mean, #17)
 Joule/mg DW: 2.26 - 5.57 (#22)
 Joule/mg SFDW: 19.13 (#22)
 18.62 - 19.48 (seasonal var., #17)
 19.08 (mean, #17)
 Joule/mg AFDW: 20.02 - 21.53 (seasonal var., #17)
 20.59 (mean, #17)

- C. fasciatum

DW = 31.7 % WW (N= 26, #51)
 AFDW = 14.9 % DW (N= 41, #41)
 = 22.9 (N= 79, #51)
 Joule/mg DW: 2.72 - 5.07 (#22)
 Joule/mg SFDW: 18.63 (#22)

- C. glaucum

DW = 52.7 % WW (N= 49, #32)
 = 54.1 (N= 95, #32)
 = 56.5 (N= 34, Oct., #41)
 SFDW = 12.1 / 19.7 % SFWW (Jul./Dec., #44)
 AFDW = 4.4 % DW (Dec., #44)
 AFDW = 16.8 % SFWW (Dec., #44)
 AFDW = 11.5 % DW (N= 23, #32)
 = 9.7 (N= 38, #32)
 AFDW = 85.3 % SFDW (#44)
 Joule/mg DW: 15.5 (#30)
 14.49 (N= 12, Oct., #41)

- C. hauniense

SFWW = 36.0 % WW (N= 30, Oct., #7)
SFDW = 7.2 % WW (N= 30, Oct., #7)

Corbula (Aloidis) gibba

DW = 56.7 % WW (N= 41 [222 ind], #51)
AFDW = 6.0 % WW (N= 8 [406 ind], #14)
= 6.5 (#51)
AFDW = 12.2 % DW (N= 41 [1879 ind], #14)
= 11.5 (N=329 [12142 ind], #51)

Leda pernula

SFDW = 20.5 / 18.3 % SFWW (Mar./Jul., #44)
AFDW = 3.9 % WW (Mar., #44)
AFDW = 14.1 % SFWW (Mar., #44)

Macoma spp. (M. balthica, calcarea)

- Macoma balthica

SFWW = 40.8 % WW (N=137, #9)
DW = 48.2 % WW (N=295, #32)
= 52.8 (N= 15, Apr., #41)
= 53.5 (M. balthica & M. calcarea, #55)
SFDW = 6.1 % WW (N=137, #9)
= 6.5 (0.556-1.620mm length, N= 9, #7)
SFDW = 13.3 % SFWW (N= 19, #39)
SFDW = 17.0 % DW (#6)
AFDW = 12.0 % SFWW (N= 19, #39)
AFDW = 11.4 % DW (N= 58 [8447 ind], #14)
= 16.7 - 22.8 (seasonal var., #32)
= 21.0 (mean, #32)
AFDW = 18.5 % DW (N= 84 [873 ind], #51)
AFDW = 76.9 - 97.2 % SFDW (seasonal var., #32)
= 91.1 (mean, #32)
= 90.4 (N= 19, #39)
Joule/mg WW: 1.64 / 1.72 (#49)
Joule/mg DW: 4.90 - 12.06 (#22)
15.44 (N= 9, Oct., #41)

(Macoma continued)

Joule/mg SFDW: 16.12 - 19.05 (seasonal var., #17)
 17.79 (mean, #17)
 20.60 (#22)
 16.73 / 19.08 (#49)
 Joule/mg AFDW: 18.93 - 22.23 (seasonal var., #17)
 19.97 (mean, #17)
 16.95 (#21)
 18.39 / 24.18 (#49)

- M. calcareo

SFDW = 1.8 % DW (N= 7, #33)
 AFDW = 14.4 % DW (N= 7, #33)
 = 19.2 (N= 16 [90 ind], #51)
 = 90.4 (N= 18, #39)

Modiola modiolus

SFDW = 15.4 % SFWW (Apr., #44)
 AFDW = 3.2 % WW (Apr., #44)
 AFDW = 12.4 % SFWW (Apr., #44)

Mya spp. (M. arenaria, truncata [1 value])

SFWW = 51.3 % WW (N= 26, Apr., #5)
 DW = 53.8 % WW (N= 52, #32)
 = 45.2 (N=129, Aug., #41)
 = 34.2 (M. trunc., N= 11 [18 ind], #51)
 SFDW = 7.6 % WW (N= 26, Apr., #5)
 SFDW = 14.8 % SFWW (N= 26, Apr., #5)
 SFDW = 14.7 % DW (N= 26, Apr., #5)
 AFDW = 6.3 % WW (N= 11, #33)
 AFDW = 15.7 % DW (N= 44 [3053 ind], #14)
 = 15.4 (N= 29, #32)
 = 16.1 (N= 11, #33)
 = 19.9 (N=135, #51)
 Joule/mg DW: 4.35 - 6.82 (#22)
 14.01 (N= 8, Oct., #41)
 Joule/mg SFDW: 19.30 (#22)
 18.87 (#30)

Myrsella (Montacuta) bidentata

DW	=	47.9	% WW	(N= 44 [609 ind], #51)
AFDW	=	5.2	% WW	(N= 5 [795 ind], #14)
	=	7.2		(#51)
AFDW	=	11.3	% DW	(N= 35 [3650 ind], #14)
	=	15.1	% DW	(N=327 [5151 ind], #51)

Mytilus edulis

SFWW	=	30.5	% WW	(#7)
DW	=	43.6 / 40.2	% WW	(N=389/123, #32)
	=	33.9		(N=301, Oct., #41)
	=	38.2		(N= 18 [58 ind], #51)
SFDW	=	5.0 / 5.0	% WW	(Aug., #10)
	=	4.6		(#26)
SFDW	=	14.0 - 17.7	% SFWW	(N= 29, seasonal var., #24)
		15.9		(mean, #24)
SFDW	=	15.0	% DW	(#19)
AFDW	=	3.9	% WW	(Dec., #44)
AFDW	=	17.7 - 21.2	% DW	(seasonal var., #32)
	=	20.0 / 22.5		(N=114/65, #32)
	=	23.8		(N= 63 [230 ind], #51)
AFDW	=	88.0 / 86.0	% SFDW	(#10)
	=	83.2		(Dec., #44)
	=	90.9 - 93.9		(seasonal var., #17)
		92.1		(mean, #17)
Joule/mg DW:		3.56 - 8.46		(#22)
		14.88		(N= 15, Oct., #41)
Joule/mg SFDW:		20.78		(#8)
		19.99 - 20.89		(seasonal var., #17)
		20.45		(mean, #17)
		21.73		(#22)
		20.73 - 23.41		(N= 28, seasonal var., #24)
		21.50		(mean, #24)
		20.98 - 21.35		(#30)

(Mytilus continued)

Joule/mg AFDW: 21.74 ~ 22.53 (seasonal var., #17)
22.18 (mean, #17)
21.61 ~ 24.63 (N= 28, seasonal var., #24)
22.46 (mean, #24)
23.59 / 23.80 (N=3/3, Aug., #10)

2.1.2. Gastropoda

Apporhais pespelicani

SFDW = 24.0 / 25.1 % SFWW (Aug., #44)
 AFDW = 4.7 % WW (Aug., #44)
 AFDW = 20.4 % SFWW (Aug., #44)
 AFDW = 85.0 % SFDW (Aug., #44)

Buccinum undatum

SFDW = 22.6 / 24.3 % SFWW (Jul./Dec., #44)
 AFDW = 11.1 % WW (Dec., #44)
 AFDW = 22.0 % SFWW (Dec., #44)
 AFDW = 90.5 % SFDW (Dec., #44)

Hydrobia spp.

SFWW = 43.7 % WW (N= 30, Apr., #5)
 DW = 67.0 % WW (N= 30, Apr., #5)
 = 65.6 / 65.9 (N=114/215, #32)
 SFDW = 10.7 % WW (N= 30, Apr., #5)
 SFDW = 24.5 % SFWW (N= 30, Apr., #5)
 SFDW = 15.6 % DW (N= 30, Apr., #5)
 AFDW = 20.5 % DW (N= 35 [3650 ind], #14)
 = 15.1 - 17.9 (#22)
 = 17.9 / 15.1 (N= 28/ 86, #32)
 = 20.5 (N= 45 [209 ind], #51)
 Joule/mg AFDW: 24.61 (#20)

Littorina littora

AFDW = 87.0 - 92.0 % SFDW (seasonal var., #17)
 90.7 (mean, #17)
 Joule/mg SFDW: 18.70 - 21.41 (seasonal var., #17)
 19.76 (mean, #17)
 Joule/mg AFDW: 21.03 - 22.99 (seasonal var., #17)
 21.77 (mean, #17)

Lymnaea spp. (L. spp., peregra)

SFWW	=	37.2	% WW	(L. spp., N= 74, #32)
SFDW	=	6.9		(L. peregra, #20)
SFDW	=	28.5	% SFWW	(L. peregra, #20)
AFDW	=	38.3	% DW	(L. spp., N= 25, #32)
	=	40.5		(L. peregra, #20)

Paludetrina (Potamopyrgus) ienkinsi

DW	=	51.8 / 52.7	% WW	(N= 23 / 66, #32)
AFDW	=	22.3	% DW	(N= 14, #32)
	=	24.9		(#20)

Theodoxus fluviatilis

DW	=	62.0	% WW	(#7)
	=	67.6 / 70.5	% WW	(N=239/62, #32)
SFDW	=	6.2	% WW	(#7)
	=	4.6		(#26)
SFDW	=	9.3	% DW	(#26)
AFDW	=	12.9 / 15.3	% DW	(N= 24/77, #32)
	=	15.4		(Jul.-Oct., #20)

2.2. Crustacea

2.2.1. Amphipoda

Bathyporeia pilosa

DW = 24.0 % WW (N=188, Aug., #41)

Joule/mg DW: 13.47 (N= 2, Oct., #41)

Corophium spp. (C. volutator, insidiosum)

- C. volutator

DW = 21.9 % WW (N= 70, #32)

= 13.7 (N= 39, Oct., #41)

= 21.2 (N= 35, #5)

= 71.1 - 81.6 (seasonal var., #17)

= 77.3 (mean, #17)

= 74.5 (N= 29, #32)

Joule/mg DW: 13.27 - 16.85 (seasonal var., #17)

15.41 (mean, #17)

11.79 (N= 6, Oct., #41)

Joule/mg AFDW: 18.68 - 20-90 (seasonal var., #17)

19.90 (mean, #17)

- C. insidiosum

DW = 12.9 % WW (N= 9 [30 ind], #51)

AFDW = 79.7 % DW (N= 16 [454 ind], #14)

= 83.8 (N= 22 [72 ind], #51)

Gammarus spp. (G. spp, duebeni, oceanicus, salinus, zaddachi)

DW = 15.4 - 26.4 % WW (G. spp., N= 34, #24)

= 19.2 (G. spp., N= 99, #32)

= 18.2 (G. spp., N=391, #32)

= 19.1 (G. oceanicus, N=205, #32)

= 21.1 (G. salinus, N= 74, #32)

= 21.7 (G. zaddachi, N= 38, #32)

AFDW = 71.9 % DW (G. spp., N= 76, #32)

= 71.8 (G. oceanicus, N= 46, #32)

= 71.2 (G. zaddachi, #12)

(Gammarus continued)

Joule/mg DW	Joule/mg AFDW	
10.23	-	(G. sp., N= 34, Oct., #41)
16.11	-	(G. spp., #7)
13.75-16.68	18.69-21.74	(G. spp., #24)
19.66	27.57	(G. spp., juv. 4mm, #26A)
18.64	24.84	(G. spp., juv. 2-5 mm, #12)
18.47	21.50	(G. duebe., females + eggs, #30)
15.90	19.49	(G. ocean., males 6-14mm, #12)
16.12	20.16	(G. salin., males 5-11mm, #12)
15.72	22.07	(G. zadda., juv. 2-5 mm, #12)
16.37	20.51	(G. zadda., females 5-10mm, #12)
15.57	22.18	(G. zadda., males 7-18mm, #12)
12.97	20.92	(G. zadda., males 12-19mm, #12)

Haploops tubicola

DW	=	24.2	% WW	(Feb., #44)
AFDW	=	17.4	% WW	(Feb., #44)
AFDW	=	71.9	% DW	(Feb., #44)

Phoxocephalus holboelli

DW	=	13.6	% WW	(N= 4 [319 ind], #14)
AFDW	=	11.7	% WW	(N= 4 [319 ind], #14)
AFDW	=	84.9	% DW	(N= 33 [2431 ind], #14)

Pontoporeia spp. (P. affinis, femorata)

- P. affinis

DW	=	19.4	% WW	(N=517, May/Aug., #41)
	=	23.6		(N= 37, Aug., #7)
Joule/mg DW	Joule/mg AFDW			
14.6	-			(N= 8, Oct., #41)
19.01/27.22	23.95/32.57			(adult/juv, #27)
22.3	-			(N= 1, #7)

(Pontoporeia continued)

- P. femorata

DW	=	19.2	% WW	(N=292, May/Aug., #41)
	=	15.5		(N= 50, #39)
	=	18.0		(#52)
	=	20.9		(N= 22, Aug., #7)
AFDW	=	13.8	% WW	(N= 50, #39)
AFDW	=	88.6	% DW	(N= 50, #39)
	=	84.5		(N=104 [350 ind], #51)

2.2.2. Isopoda

Asellus aquaticus

DW	=	20.7	% WW	(N= 37, #32)
AFDW	=	68.2	% DW	(N= 15, #32)

Idothea spp. (I. spp., aleutica, baltica, chelipes, ochotensis)

DW	≈ 22.6 / 23.0	% WW	(I. spp., N=159/36, #32)
	≈ 20.6		(I. baltica, N= 55, Apr., #41)
	= 21.0 - 24.0		(I. chelipes, fem., #47)
	= 22.3		(mean, #47)
AFDW	= 90.0	% DW	(I. spp., N= 35, #32)
	= 75.9		(I. balt., N= 5 [42 ind], #51)
Joule/mg	Joule/mg		
DW	AFDW		
24.87	25.92		(I. aleutica, eggs, #12)
23.45	24.70		(I. aleut., embryos, #12)
21.98	24.41		(I. aleut., 0 days old, #12)
25.33	26.46		(I. balt. basteri, eggs, #12)
23.24	25.12		(I. balt. bast., embryos, #12)
20.47	24.95		(I. balt. bast., 0 days old, #12)
8.63/10.22	-		(I. balt., juv. 0/20 days, #12)
6.70/11.26	8.71/17.92		(I. balt., juv. 1.95/4.0mm, #12)
13.98	-		(I. balt., #22)
14.19	20.94		(I. balt. bast., all sizes, #12)
13.13/17.54	18.01/19.68		(I. chel., juv. 2-5/4mm, #12)
16.60	21.66		(I. chel., fem. & eggs, #12)

(Idothea continued)

Joule/mg		Joule/mg	
DW		AFDW	
14.66		19.25	(I. chel., males 5-15mm, #12)
24.16		25.54	(I. ochotensis, eggs, #12)
20.68		-	(I. ochot., embryones, #12)
19.00		23.57	(I. ochot., 0 days old, #12)

Jaera albifrons

DW	=	13.3	% WW	(N= 24, May/Aug./Oct., #41)
	=	23.3		(N= 63, #32)
AFDW	=	70.8	% DW	(N= 10, #32)

Mesidotea entomon

DW	=	19.9	% WW	(N= 42, May/Aug., #41)
	=	22.2		(N= 36, #32)
AFDW	=	65.2	% DW	(Jul.-Oct., #20)
Joule/mg		Joule/mg		
DW		AFDW		
-		24.36		(Aug., #10)
27.55/23.07		28.89/26.46		(eggs/juv. 0 days, #12)

2.2.3. Mysidacea

Mysidae

DW	=	19.9	% WW	(N= 90, #32)
AFDW	=	90.0	% DW	(N= 10, #32)

Gastrosaccus spinifer

DW	=	18.0	% WW	(N= 11 [30 ind], #51)
AFDW	=	12.2	% WW	(N= 5 [10 ind], #14)
AFDW	=	83.6	% DW	(N= 37 [324 ind], #14)
	=	86.9		(N= 84 [182 ind], #51)

Mysis spp. (*M. mixta*, *relicta*)

- *M. mixta*

DW = 13.3 % WW (N= 16, May/Aug./Oct., #41)

Joule/mg DW: 15.12 (N= 5, Oct., #41)

- *M. relicta*

AFDW = 90.0 - 92.5 % DW (seasonal var., embryos, #23)

= 89.9 - 92.5 (" ", juveniles, #23)

= 90.5 - 92.5 (" ", females + eggs, #23)

= 91.9 - 93.4 (" ", females, #23)

= 88.4 - 94.0 (" ", males, #23)

Joule/mg AFDW: 30.47 (embryos, #23)

22.73 (juveniles, #23)

23.50 (females + eggs, #23)

Neomysis integer

DW = 12.8 % WW (N= 16, May/Aug./Oct., #41)

Joule/mg DW: 14.97 / 13.54 (N= 4/ 24, Oct., #41)

2.2.4. Other Crustacea

Balanus spp. (*B. balanoides*, *balanus*, *crenatus*, *improvisus*)

- *B. balanoides*

Joule/mg SFDW: 19.86 (N= 11, #49)

Joule/mg AFDW: 22.20 (N= 11, #49)

- *B. balanus*

Joule/mg SFDW: 15.46 (N= 6, #10)

Joule/mg AFDW: 22.57 (N= 6, #10)

- *B. crenatus*

DW = 47.4 % WW (Feb., #44)

AFDW = 3.9 % WW (Feb., #44)

AFDW = 8.2 % DW (Feb., #44)

- B. improvisus

DW	=	50.7	% WW	(N= 44, #32)
SFDW	=	3.2	% WW	(#26)
SFDW	=	8.9	% DW	(#26)
AFDW	=	3.9	% WW	(#26)
AFDW	=	10.8	% DW	(N= 10, #32)

Crangon spp. (*C. crangon*, *septemspinosus*)

DW	=	19.1	% WW	(<i>C. crangon</i> , N= 48, Dec., #41)
	=	26.0 / 24.0		(<i>C. septemspinosus</i> , Spring / November, #50)
AFDW	=	90.7	% DW	(<i>C. crang.</i> , N= 9 [11 ind], #51)
Joule/mg DW:		17.41		(<i>C. crang.</i> , #12)
		17.25		(<i>C. crang.</i> , #22)
		11.42		(<i>C. crang.</i> , N= 26, Oct., #41)
		17.90 / 17.12		(<i>C. septe.</i> , Spring / Nov., #50)
Joule/mg AFDW:		20.39		(<i>C. crang.</i> , #12)

Diastylis rathkei

DW	=	20.8	% WW	(#12)
	=	19.9		(N=101, Aug., #41)
	=	17.2		(#55)
	=	11.8		(N= 61 [8293 ind], #51)
AFDW	=	14.0	% WW	(#30)
AFDW	=	56.7	% DW	(N= 20 [916 ind], #14)
	=	59.6 - 69.0		(seasonal var., #15)
	=	65.7		(mean, #15)
	=	66.5		(N=438 [31530 ind], #51)
Joule/mg DW:		14.58 - 17.69		(seasonal var., #15)
		16.55		(mean, #15)
		15.12		(#22)
		12.02		(N= 19, Oct., #41)

(Diastylis continued)

Joule/mg AFDW:	24.47	-	25.64	(seasonal var., #15)
	25.19			(mean, #15)
	25.1			(eggs, #43)
	16.9			(juv., #43)
	18.7			(males & females, #43)
	16.4			(females, #43)
	19.0			(females & eggs, #43)

Eupagurus bernhardus

DW	=	27.1	% WW	(Nov., #44)
AFDW	=	16.7	% WW	(Nov., #44)
AFDW	=	61.6	% DW	(Nov., #44)

Hyas coarctatus

DW	=	38.1	% WW	(Feb., #44)
AFDW	=	22.8	% WW	(Feb., #44)
AFDW	=	59.8	% DW	(Feb., #44)

Ostracoda

DW	=	44.0	% WW	(Cyprideis litoralis, #7)
	=	63.0		(Heterocyprideis sorbyana, #7)
	=	61.0		(Paracyprideis fennica, #7)
AFDW	=	8.0	% WW	(Candona spp., #7)
AFDW	=	20.0	% DW	(Cyprideis litoralis, #7)
	=	46 / 50		(N=130/70, Aug., #53)
Joule/mg AFDW:		25.10		(Cyprideis litoralis, #7)
		24.62		(Candona spp., #7)

2.3. Polychaeta

2.3.1. Polychaeta errantia

Aphrodite aculeata

DW	=	22.4	% WW	(Jul., #44)
AFDW	=	15.6	% WW	(Jul., #44)
AFDW	=	69.6	% DW	(Jul., #44)

Eteone longa

AFDW	=	13.6	% WW	(N= 6 [40 ind], #14)
	=	25.1		(N= 32 [30 ind], #51)
AFDW	=	83.3	% DW	(N= 38 [623 ind], #14)
	=	83.1		(N= 27 [50 ind], #51)

Harmothoe sarsi

DW	=	15.3	% WW	(N= 21, #7)
	=	13.5		(N= 48, Apr.-Aug., #41)
	=	14.7		(N= 37 [110 ind], #51)
AFDW	=	15.7	% WW	(N= 9 [108 ind], #14)
AFDW	=	79.1	% DW	(N= 24 [291 ind], #14)
	=	90.6		(N=356 [1303ind], #51)
Joule/mg DW:		15.29		(N= 3, Oct., #41)

Nephtys spp. (N. spp., ciliata, incisa)

DW	=	14.4 - 17.8	% WW	(seasonal var., #15)
	=	16.8		(mean, #15)
	=	21.8		(N. incisa, Apr., #44)
	=	17.1		(N= 3, #11)
	=	11.4		(#33)
	=	18.0		(N= 37 [341 ind], #51)
AFDW	=	17.5	% WW	(N. incisa, Apr., #44)
AFDW	=	69.4	% DW	(N= 42 [429 ind], #14)
	=	70.7 - 82.6		(seasonal var., #15)
	=	78.3		(mean, #15)
	=	69.4		(N= 3, #11)
	=	76.2		(#33)
	=	74.2		(N= 37 [341 ind], #51)

(Nephtys continued)

AFDW	=	80.3	% DW	(N. incisa, Apr., #44)
Joule/mg DW:		17.00		(N. ciliata, N= 3, #13)
		15.50 - 19.75		(seasonal var., #15)
		17.52		(mean, #15)
		15.80 / 17.40		(N. incisa,
				all seasons / Aug - Dec, #44)
Joule/mg AFDW:		19.60 - 25.10		(seasonal var., #15)
		22.17		(mean, #15)

Nereis spp. (N. diversicolor, pelagica)

- N. diversicolor

DW	=	15.7	% WW	(N= 93, #32)
	=	15.5		(N= 47, Aug., #41)
	=	18.1		(Dec., #44)
	=	18.0		(N= 20, Apr., #5)
AFDW	=	14.9	% WW	(Dec., #44)
AFDW	=	75.3 - 88.1	% DW	(seasonal var., #17)
	=	82.8		(mean, #17)
	=	84.4		(#32)
	=	82.3		(Dec., #44)
Joule/mg DW:		21.45		(N= 5, #12)
		16.48 - 19.70		(seasonal var., #17)
		18.04		(mean, #17)
		15.35		(Oct., #41)
Joule/mg AFDW:		23.33		(N= 5, #12)
		20.90 - 22.47		(seasonal var., #17)
		21.76		(mean, #17)

- N. pelagica

DW	=	16.6 / 15.4	% WW	(N= 3 / 3, Nov./Feb., #49)
Joule/mg DW:		24.51 / 21.99		(N= 3 / 3, " , #49)
Joule/mg AFDW:		27.71 / 24.40		(N= 3 / 3, " , #49)

Scoloplos armiger

DW	=	14.9	% WW	(N= 40 [2065 ind], #51)
	=	12.4		(N= 7, #55)
AFDW	=	11.2	% WW	(N= 8 [393 ind], #14)
AFDW	=	56.3	% DW	(N= 22 [1024 ind], #14)
	=	79.3		(N=236 [7803 ind], #51)

2.3.2. Polychaeta sedentaria

Arenicola marina

DW	=	18.2	% WW	(Sep., #44)
AFDW	=	13.7	% WW	(Sep., #44)
AFDW	=	75.3	% DW	(Sep., #44)

Aricidea spp. (A. jeffreysii, suecia)

DW	=	13.6	% WW	(A. suecia, N= 30, Apr.-Oct., #41)
	=	13.7		(A. jeffr., N= 6, #55)
AFDW	=	72.8	% DW	(A. jeffr., N= 19 [190 ind], #14)
	=	80.3		(A. jeffr., N= 40 [115 ind], #51)

Pectinaria spp. (P. hypoboreia, koreni)

DW	=	19.2	% WW	(P. hypo., #13)
	=	6.9 - 11.3		(P. kor., seasonal var., #15)
	=	39.6		(P. kor., N= 17 [43 ind], #51)
AFDW	=	50.5 - 85.7	% DW	(P. kor., seasonal var., #15)
	=	52.0		(P. kor., N=318 [2841 ind], #51)
Joule/mg DW:		14.21		(P. hypo., N= 5, #13)
		13.68 - 19.75		(P. kor., seasonal var., #15)
		16.72		(P. kor., mean, #15)
		13.02		(P. kor., #22)
Joule/mg AFDW:		18.62 - 25.82		(P. kor., seasonal var., #15)
		22.22		(P. kor., mean, #15)

Pygospio elegans

DW	=	24.4	% WW	(N= 20, #32)
AFDW	=	11.1	% WW	(N= 6 [1857 ind], #14)
AFDW	=	61.9	% DW	(N= 23 [7224 ind], #14)

Polydora spp.

DW	=	20.3	% WW	(N= 42 [264 ind], #51)
AFDW	=	83.5	% DW	(N=167 [3645 ind], #51)

Sosane gracilis

DW	=	21.4	% WW	(Feb., #44)
AFDW	=	18.7	% WW	(Feb., #44)
AFDW	=	87.4	% DW	(Feb., #44)

Terebellides stroemi

DW	=	19.6	% WW	(T. spp., #13)
	=	17.0		(N= 4, Nov., #5)
	=	15.5		(N=134, Aug., #41)
	=	13.6		(N= 38 [1850 ind], #51)
	=	13.8		(N= 7, #55)
AFDW	=	60.9	% DW	(N=110 [1099 ind], #51)
Joule/mg DW:		10.47		(T. spp., N= 3, #10)
		17.34		(T. spp., #13)
Joule/mg AFDW:		24.95		(T. spp., N= 3, #10)

2.4. Other Taxa

2.4.1. Turbellaria

DW	=	23.6	% WW	(N= 69, #32)
AFDW	=	78	% DW	(N= 92, Aug., #53)

2.4.2. Nematoda

DW	=	23.0	% WW	(#7)
AFDW	=	80.0	% DW	(#7)
	=	76 - 93		(Aug., #53)

2.4.3. Nemertini

DW	=	20.5	% WW	(Prostoma obscurum, N= 27, #32)
	=	25.5		(N= 15 [39 ind], #51)
	=	12.2		(N= 7, #55)
AFDW	=	22.0	% WW	(#51)
AFDW	=	86.2	% DW	(N= 62 [127 ind]. #51)
Joule/mg DW:	21.3			(P. obscurum, Jul., #26A)
Joule/mg AFDW:	22.5			(P. obscurum, Jul., #26A)

2.4.4. Priapulida

Halicyrptus spinulosus

DW	=	19.0	% WW	(Apr., #7)
	=	9.2		(N= 7, #55)
	=	8.2		(N=139, #41)
	=	7.6		(N=52 [293 ind], #51)
AFDW	=	6.5	% WW	(#51)
AFDW	=	86.1	% DW	(N=368 [1117ind], #51)

Priapulys caudatus

DW	=	8.8	% WW	(#55)
	=	6.2		(N= 36, Apr./Aug., #41)

2.4.5. Oligochaeta

DW	=	18.0	% WW	(Nais elinguis, N=10, Apr., #7)
	=	13.0		(Clitello arenarius, N=10, Apr., #5)
	=	19.0		(N= 13, #32)
Joule/mg DW:		22.79		(Dina microstoma, #42)
		21.51		(Limnodrilus sp., #42)
		22.11		(Tubifex tubifex, #42)
Joule/mg AFDW:		23.23		(T. tubifex, #42)

2.4.6. Insecta

Chironomida Larvae

DW	=	19.5 / 19.0	% WW	(N= 67/34, #32)
	=	21.0		(N= 30, #7)
Joule/mg DW:		22.30		(N= 9, #29)
		22.33		(N= 3, #30)
Joule/mg AFDW:		24.28		(N= 9, #29)
		23.34		(N= 2, #30)

Trichoptera Larvae

DW	=	19.8	% WW	(N= 50, #32)
----	---	------	------	--------------

2.4.7. Echinodermata

Asterioidea (Asterias spp., rubens, vulgaris)

DW	=	24.3	% WW	(A. vulg., #13)
AFDW	=	56.0 / 42.0	% DW	(A. spp., Aug., #10)
	=	50.6		(A. rub., N= 13, #22)
	=	50.7		(A. rub., N= 5, #37)
	=	38.0		(A. rub., N= 21 [39 ind], #51)
Joule/mg DW:		8.55		(A. vulg., #13)
		5.23 - 11.81		(A. rub., #22)
		10.50		(A. rub., N= 5, #37)
		8.92		(A. spp., N= 7, #19)
Joule/mg AFDW:		21.10		(A. rub., N= 5, #37)
		23.60 / 22.12		(A. spp., N=3/3, Aug., #10)

Ophiuroidea (Ophiura albida; Amphiura chiajei, filiformis)

Joule/mg AFDW: 20.94 - 24.83 (3 different species, Aug., #10)

- Ophiura albida

DW = 66.0 / 61.7 % WW (Nov./Apr., #44)
= 55.7 (N= 13 [43 ind], #51)
AFDW = 8.4 / 8.6 % WW (Nov./Apr., #44)
AFDW = 12.7 / 13.9 % DW (Nov./Apr., #44)
= 12.7 (N=230 [1793 ind], #51)

- Amphiura chiajei

DW = 44.9 - 52.9 % WW (seasonal var., #44)
AFDW = 12.2 - 13.5 % WW (seasonal var., #44)
AFDW = 25.0 - 30.1 % DW (seasonal var., #44)

- Amphiura filiformis

DW = 38.6 - 43.6 % WW (seasonal var., #44)
= 40.2 (juv., #44)
AFDW = 13.7 - 17.0 % WW (seasonal var., #44)
= 14.1 (juv., #44)
AFDW = 33.3 - 39.0 % DW (seasonal var., #44)

Echinoidea (Echinocardium; Echinocyamus; Strongylocentrotus)

- Echinocardium cordatum

DW = 17.7 - 21.9 % WW (seasonal var., #44)
AFDW = 2.9 - 3.1 % WW (seasonal var., #44)
AFDW = 14.2 - 16.4 % DW (seasonal var., #44)

- Echinocyamus pusillus

DW = .47.9 % WW (Sep., #44)
AFDW = 7.2 % WW (Sep., #44)
AFDW = 15.0 % DW (Sep., #44)

- Strongylocentrotus spp.

Joule/mg DW:	12.06	(N= 3, #10)
	3.69	(#13)
Joule/mg AFDW:	23.66	(N= 3, #10)

Holothuroidea (Cucumaria)

- Cucumaria spp. (C. elongata, frondosa)

DW	=	47.8	% WW	(C. elong., Jul., #44)
AFDW	=	8.2	% WW	(C. elong., Jul., #44)
AFDW	=	17.2	% DW	(C. elong., Jul., #44)
Joule/mg DW:		12.86		(C. frond. [without CaCO ₃], #13)

3. Length - weight - relationships

Usual log-log-regression form of length-weight-relations:

$$\log s(w) = b * \log s(l) + a ; \quad r, N$$

B = basis of logarithm,

usually 10 unless otherwise stated

l = length

w = weight (WW, SFWW, DW, SFDW, AFDW; see 1.1)

b = slope

a = y-axis intercept

r = correlation coefficient

N = number of values included

an additional number in brackets [] is the number
of individuals included.

Additional parameters included here (if available) are:

- units = units of weight / length,
in mg / mm unless otherwise stated
- range = range of length included in regression
- area and time of sampling

3.1. Mollusca

3.1.1. Bivalvia

species	weight	b	a	r	N	range	area	time	ref.
<u>Abra alba</u> (=Syndosmya alba)	WW	2.94	-0.842	-	-	-	Kiel Bay	-	#15
<u>A. nitida</u>	DW	2.702	-1.403	0.971	196	4 - 13	W.Norway	-	#54
	AFDW	2.786	-2.221	0.934	146	4 - 13	W.Norway	-	#54
<u>A. longicallus</u>	DW	3.008	-1.597	0.956	148	7 - 20	W.Norway	-	#54
	AFDW	3.016	-2.379	0.962	111	7 - 20	W.Norway	-	#54
<u>Arctica islandica</u>	WW	3.08	-0.779	-	-	-	Kiel Bay	-	#15
	SFWW	2.888	-9.590	0.96	-	-	Atlantic	-	#36 loge, g/mm
<u>Astarte borealis</u>	WW	3.034	-0.813	0.996	56	7 - 26	Hanö Bight	-	#41
		3.035	-0.617	0.999	47	-	Kiel Bay	June	#46
<u>A. montagui</u>	WW	3.475	-0.888	0.997	18	-	Kiel Bay	June	#46
<u>A. elliptica</u>	WW	2.878	-0.498	0.998	53	-	Kiel Bay	June	#46
		3.087	-0.835	0.999	57	-	Kiel Bay	June	#46
	DW	2.853	-0.643	0.997	46	-	Kiel Bay	June	#46
	SFDW	2.791	-2.084	0.995	49	-	Kiel Bay	June	#46

Bivalvia (continued)

species	weight	b	a	r	N	range	area	time	ref.	
<u>Cardium edule</u>	SFDW	3.108	-1.972	0.99	29	-	Ythan -	Nov.	#17	shell-height
		3.752	-2.919	0.99	28	-	Estuary	Jan.	"	"
		3.709	-2.907	0.99	21	-	"	Mar.	"	"
		3.412	-2.336	0.99	21	-	"	May	"	"
		3.125	-2.078	0.99	29	-	"	Jul.	"	"
		3.209	-2.090	0.99	24	-	"	Sep.	"	"
	AFDW	2.479	-1.354	0.999	16 [90]	>= 2	Kiel Bay	Jul.	#14	
		2.412	-1.222	0.991	20 [619]		"	Aug.	"	
		2.662	-1.403	0.999	12 [60]		"	Dec.	"	
	AFDW	3.458	-3.854	0.992	18	2 - 6	Skagerrak	Aug.	#34	loge
		3.401	-5.225	0.998	38	4 - 21	"	Nov.	"	"
<u>C. glaucum</u>	WW	3.634	-1.263	0.964	34	5 - 12	Hanö Bight	-	#41	
		2.916	-0.503	0.996	75	5.6-25.5	Askö	Aug.	#5	
	DW	3.655	-1.403	0.945	34	5 - 12	Hanö Bight	-	#41	
		2.848	-0.867	0.987	75	5.6-25.5	Askö	Aug.	#5	
	SFDW	3.274	-2.538	0.967	55	-	Tvärminne	Jul.	#28	
		2.792	-2.032	0.974	58	-	Tvärminne	Aug.	#28	
<u>C. hauniense</u>	WW	2.932	-0.553	0.998	30	1.66-7.9	Askö	Nov.	#5	
	DW	2.760	-0.782	0.993	30	1.66-7.9	Askö	Nov.	#5	

Bivalvia (continued)

species	weight	b	a	r	N	range	area	time	ref.	
<u>Macoma balthica</u>	WW	3.127	-0.964	0.999	15	2 - 21	Hanö Bight	Apr.	#41	
		3.049	-0.814	0.998	137	2 - 19	Askö	-	#7	
		3.049	-6.962	0.995	9	556-1620	Askö	-	#7	µg/µm
	SFWW	2.486	-0.476	0.99	50	2 - 22	Gdansk Bay	-	#39	
	DW	3.165	-1.307	0.996	137	2 - 19	Askö	Aug.	#6	
		3.307	-1.481	0.993	44	5 - 18	Hanö Bight	Apr.	#41	
	SFDW	2.846	-1.534	0.96	126	-	Ythan -	Feb. 9	#17	shell-height
		2.825	-1.565	0.94	116	-	Estuary	Mar. 7	"	"
		2.434	-1.208	0.96	118	-	"	Apr.11	"	"
		2.260	-0.876	0.90	97	-	"	May 4	"	"
		1.839	-0.327	0.92	71	-	"	Jun. 6	"	"
		3.088	-1.540	0.99	66	-	"	Jul.13	"	"
		3.134	-1.892	0.98	72	-	"	Aug.11	"	"
		2.950	-1.452	0.99	66	-	"	Sep. 6	"	"
		2.941	-1.454	0.97	54	-	"	Nov.22	"	"
	SFDW	2.943	-1.928	0.964	97	-	Tvärminne	Jul.	#28	
		2.897	-1.544	0.987	59	-	"	Jul.	"	
		2.897	-1.780	0.990	50	2 - 22	Gdansk Bay	-	#39	
	AFDW	3.192	-2.158	0.997	112	1.5-19.5	Kiel Bay	May	#14	
	SFAFDW	2.847	-1.754	0.99	50	2 - 22	Gdansk Bay	-	#39	

Bivalvia (continued)

species	weight	b	a	r	N	range	area	time	ref.	
<u>Mya arenaria</u>	SFWW	2.993	-1.498	0.992	26	13 - 63	Askö	Apr.	#5	
		2.945	-2.259	0.986	26	13 - 63	Askö	Apr.	#5	
		3.179	-2.635	0.985	66	-	Tvärminne	Aug	#28	
	AFDW	2.134	-3.163	0.951	19	1 - 3	Skagerrak	Jul.	#34	loge
		2.680	-3.804	0.993	30	4 - 10	"	Aug.	"	"
		2.749	-3.553	0.995	42	4 - 25	"	Nov.	"	"
		2.770	-4.425	0.996	18	35 - 77	"	Apr.	"	"
		2.605	-3.741	0.992	15	34 - 75	"	Jun.	"	"
		2.797	-4.410	0.997	16	33 - 62	"	Oct.	"	"
<u>Mytilus edulis</u>	WW	2.328	-0.354	0.986	253	3 - 42	Hanö B.	-	#41	
	DW	2.476	-1.059	0.979	48	9 - 39	Hanö B.	-	#41	
	SFDW	2.468	-1.789	0.949	155	mean 13.7	Tvärminne	-	#28	

Bivalvia (continued)

species	weight	b	a	r	N	range	area	time	ref.
<u>Mytilus edulis</u>	SFDW	2.630	-2.075	0.99	39	-	Askö	Jan. 4 #29	4m depth
		2.709	-2.175	0.99	35	-	"	Feb. 9 "	"
		2.678	-2.147	0.99	39	-	"	Mar. 1 "	"
		2.444	-1.838	0.99	38	-	"	Mar. 21 "	"
		2.636	-1.905	0.99	35	-	"	Apr. 23 "	"
		2.630	-1.925	0.99	35	-	"	May 22 "	"
		2.265	-1.573	0.99	35	-	"	Jun. 28 "	"
		2.300	-1.623	0.99	31	-	"	Aug. 2 "	"
		2.437	-1.843	0.99	33	-	"	Sep. 5 "	"
		2.665	-2.099	0.99	35	-	"	Oct. 9 "	"
		2.562	-1.942	0.99	35	-	"	Nov. 21 "	"
		2.587	-2.001	1.00	32	-	Askö	Mar. 8 "	15m depth
		2.502	-1.663	1.00	35	-	"	Apr. 3 "	"
		2.761	-1.936	0.99	19	-	"	Apr. 17 "	"
		2.534	-1.621	0.98	20	-	"	May 3 "	"
		2.481	-1.641	0.89	20	-	"	May 29 "	"
		2.695	-1.936	0.99	20	-	"	Jun. 6 "	"
		2.064	-1.262	0.99	20	-	"	Jun. 19 "	"
		2.365	-1.731	1.00	35	-	"	Aug. 19 "	"
		2.614	-2.013	0.98	19	-	"	Sep. 2 "	"
		2.571	-1.915	1.00	35	-	"	Dec. 7 "	"

Bivalvia (continued)

species	weight	b	a	r	N	range	area	time	ref.
<u>Pisidium</u> sp.	DW	2.56	-0.81	0.99	-	1 - 6	Bothn.Bay	Jul-Oct	#20

3.1.2. Gastropoda

species	weight	b	a	r	N	range	area	time	ref.
<u>Hydrobia</u> spp.	WW	2.308	-0.543	0.94	30	1.2-5.3	Askö	Apr.	#7
	DW	2.237	-0.660	0.983	30	1.2-5.3	Askö	Apr.	#5
<u>H. ulvae</u>	DW	2.416	-0.750	0.99	54	-	Askö	May	#31
		2.676	-0.908	0.99	54	-	"	Jul.	"
		2.573	-0.841	0.99	93	-	"	May, Jul	"
	SFDW	2.363	-1.587	0.93	32	-	Ythan Est.	Nov. 1	#17
		2.228	-1.457	0.85	29	-	"	Jan. 3	"
		2.174	-1.392	0.86	24	-	"	Mar. 4	"
		1.972	-1.326	0.89	22	-	"	May 6	"
		1.414	-1.108	0.87	23	-	"	Jul. 2	"
		3.353	-1.994	0.88	21	-	"	Sep. 2	"
<u>Littorina litt.</u>	SFDW	3.594	-2.510	0.99	26	-	Ythan Est.	Nov. 1	#17
		3.422	-2.299	0.96	16	-	"	Jan. 3	"
		3.609	-2.518	0.98	23	-	"	Mar. 4	"
		3.058	-1.940	0.96	22	-	"	May 6	"
		4.333	-3.449	0.94	17	-	"	Jul. 2	"
		3.879	-2.799	0.99	25	-	"	Sep. 2	"

Gastropoda (continued)

species	weight	b	a	r	N	range	area	time	ref.
<u>Lymnaea peregra</u>	DW	2.61	-1.02	0.99	-	1 - 20	Bothn. B.	Jul-Oct	#20
<u>Potamopyrgus jenkinsi</u> (Paludestina)	WW	2.594	-0.629	0.985	30	1.45-5.3	Askö	Nov.	#5
<u>Theodoxus fluviat.</u>	DW	2.80	-0.54	0.99	-	2 - 8	Bothn. B.	Jul-Oct	#20
<u>Valvata piscinalis</u>	DW	2.48	-0.40	0.98	-	1 - 5	Bothn. B.	Jul-Oct	#20

3.2. Crustacea

3.2.1. Amphipoda

species	weight	b	a	r	N	range	area	time	ref.
<u>Bathyporeia pilosa</u>	WW	1.788	-0.816	0.971	188	2 - 7	Hanö Bight	Aug.	#41
<u>B. sarsi</u>	DW	2.02	2.86	0.976	6	-	Kiel Bay	-	#3 $\mu\text{g/mm}$, loge
<u>Corophium volutator</u>									
males	WW	2.933	-1.473	0.943	11	5 - 9	Askö	-	#5
females + juv.	WW	2.709	-1.443	0.986	22	2 - 8	"	-	"
fem. + males + juv.	WW	2.989	-1.574	0.984	35	2 - 9	"	-	"
males	DW	3.273	-2.422	0.934	11	5 - 9	Askö	-	#5
females + juv.	DW	2.567	-2.040	0.970	22	2 - 8	"	-	"
fem. + males + juv.	DW	2.938	-2.220	0.973	35	2 - 9	"	-	"
		2.443	-1.709	0.98	41	-	Ythan Est.	Nov.	#17
		2.514	-1.702	0.95	40	-	"	Jan.	"
		3.419	-2.488	0.96	37	-	"	Mar.	"
		3.140	-2.197	0.97	40	-	"	May	"
		2.700	-1.932	0.96	29	-	"	Jul.	"
		2.507	-1.773	0.99	32	-	"	Sep.	"
		2.440	-1.881	0.943	39	3 - 9	Hanö B.	Oct.	#41
<u>C.insidiosum</u>	DW	2.72	1.70	0.997	28	-	Kiel Bay	-	#3 $\mu\text{g/mm}$, loge

Amphipoda (continued)

species	weight	b	a	r	N	range	area	time	ref.	
<u>Gammarus</u> spp.	WW	2.760	-1.447	0.99	43	2 - 24	Askö	Apr.	#7	
		2.219	-0.795	0.962	302	3 - 25	Hanö B.	May-Oct	#41	
	DW	2.140	-1.659	0.993	89	3 - 25	Hanö B.	Aug.	#41	
<u>G. salinus</u>	DW	2.44	0.81	0.98	23	-	Kiel Bay	-	#3	µg/mm, loge
<u>Microdeutopus</u>										
<u>gryllotalpa</u>	DW	2.44	1.95	0.966	15	-	Kiel Bay	-	#3	µg/mm, loge
<u>Pontoporeia affinis</u>	WW	3.060	-1.619	0.997	37	2.9-8.7	Askö	Aug.	#7	
		2.105	-1.123	0.906	81	3 - 9	Hanö B.	May	#41	
		2.116	-1.097	0.973	134	"	"	May	"	
		2.417	-1.268	0.979	302	2 - 8	"	Aug	"	
	DW	2.853	-2.281	-	-	-	Bothnian Sea	Jul.	#1	
		3.269	-2.729	-	-	-	Bothnian Bay	Jul.	#1	
		3.075	-2.258	0.992	37	2.9-8.7	Askö	Aug.	#5	
		3.086	-2.715	0.97	-	-	Askö	Feb. 3	#16	
		3.845	-3.004	0.99	-	-	"	Aug. 8	"	
		2.732	-2.117	0.99	-	-	"	Dec. 4	"	
		2.836	-2.340	0.99	-	-	"	Jan. 17	"	
		2.021	-1.735	0.91	-	-	"	Apr.	"	
		3.006	-2.340	0.99	-	-	"	Jul. 26	"	

(Amphipoda continued)

species	weight	b	a	r	N	range	area	time	ref.
<u>Pontoporeia</u>									
<u>femorata</u>	WW	3.012	-1.598	0.996	22	2.9-9.2	Askö	Aug.	#7
		2.662	-1.218	0.99	50	2 - 11	Gdansk B.	-	#39
		2.342	-1.225	0.982	74	3 - 9	Hanö B.	May	#41
		2.302	-1.073	0.944	55	"	"	Aug.	"
		2.434	-1.135	0.973	154	"	"	Aug.	"
	DW	3.003	-2.281	0.988	22	2.9 -9.2	Askö	Aug.	#5
		2.992	-2.347	0.98	50	2 - 11	Gdansk B.	-	#39
	AFDW	3.042	-2.444	0.98	50	2 - 11	Gdansk B.	-	#39

3.2.2. Other Crustacea

species	weight	b	a	r	N	range	area	time	ref.	
<u>Crangon crangon</u>	DW	2.848	-2.452	0.988	48	10 - 35	Hanö B.	Dec.	#41	g / mm
<u>Diastylis rathkei</u>	WW	3.130	-2.216	-	-	-	Kiel Bay.	-	#15	
		2.973	-2.229	0.974	101	5 - 14	Hanö B.	Aug.	#41	
	DW	3.259	-3.166	0.945	101	5 - 14	Hanö B.	Aug.	#41	
<u>Idothea baltica</u>	DW	2.623	-1.869	0.975	55	6 - 15	Hanö B.	-	#41	
<u>Mesidothea entomon</u>	WW	2.629	-1.430	0.995	42	5 - 56	Hanö B.	May, Aug	#41	
(=Saduria)	DW	2.78	-2.12	0.98	-	5 - 50	Lulea	-	#20	
		2.50	-1.66	-	-	33 - 49	Bothn. B.	-	#28	
<u>Jaera spp.</u>	WW	2.682	-1.249	0.988	24	2 - ?	Hanö B.	-	#41	
females	WW	2.722	-1.271	0.935	17	2.9-4.2	Askö	Apr.	#5	
males	WW	3.399	-1.402	0.895	8	1.5-2.4	Askö	Apr.	#5	
mixed	WW	2.473	-1.126	0.96	25	1.5-4.2	Askö	Apr.	#7	
females	DW	2.911	-1.813	0.970	16	2.9-4.2	Askö	Apr.	#5	
males	DW	2.569	-1.650	0.876	8	1.5-2.4	Askö	Apr.	#5	
mixed	DW	2.868	-1.648	0.981	25	1.5-4.2	Askö	Apr.	#5	
<u>Mysis mixta</u>	WW	3.233	-2.481	0.990	16	11 - 22	Hanö B.	May-Oct	#41	
<u>Neomysis intiger</u>	WW	2.722	-1.873	0.972	16	10 - 18	Hanö B.	Aug-Oct	#41	

3.3. Polychaeta

3.3.1. Polychaeta errantia

species	weight	b	a	r	N	range	area	time	ref.
<u>Harmothoe sarsi</u>	WW	2.672	-1.376	0.96	21	1.9-17.0	Askö	Nov.	#7
		2.870	-1.727	0.976	48	5.0-24.0	Hanö B.	Apr, Aug	#41
		3.333	-1.268	0.996	22	2 - 10	Tvärminne	Aug.	#45 l=breath
	DW	2.415	-2.060	0.96	21	1.9-17.0	Askö	Nov.	#5
		3.254	-1.863	0.994	84	1 - 6	Tvärminne	Aug.	#45 l=breath
	AFDW	3.398	-2.000	0.993	84	"	"	"	"
<u>Nereis diversicolor</u>	WW	2.388	-1.612	0.99	20	9.9-57.0	Askö	Apr.	#7
		2.341	-1.609	0.976	47	5.0-24.0	Hanö B.	Aug.	#41
	DW	2.500	-2.517	0.97	20	9.9-57.0	Askö	Apr.	#5

3.3.2. Polychaeta sedentaria

<u>Aricidea suecia</u>	WW	2.833	-3.060	0.953	134	10 - 21	S.Baltic	-	#41
<u>Terebellides</u> <u>stroemi</u>	WW	2.410	-1.511	0.986	4	6.3-20.0	Askö	Nov.	#5
		2.407	-1.574	0.989	134	7.0-24.0	Hanö B.	Aug.	#41
	DW	2.417	-2.309	0.962	4	6.3-20.0	Askö	Nov.	#5

3.4 Other Taxa

species	weight	b	a	r	N	range	area	time	ref.
---------	--------	---	---	---	---	-------	------	------	------

3.4.1. Priapulida

Halicryptus

<u>spinulosus</u>	adults WW	1.037	-0.480	0.99	13	1.7-39.0	Askö	Apr.	#7	$l = l * \phi^2$
	"	2.972	-1.419	0.968	139	2.0-25.0	Hanö B.	Aug, Oct	#41	
	larvae WW	3.244	-7.778	0.99	3	1200-1300	Askö	Oct.	#7	$\mu g / \mu m$

<u>Priapulus caudatus</u>	WW	2.429	-0.890	0.925	36	2.0-30.0	Hanö B.	Apr-Oct	#41	
---------------------------	----	-------	--------	-------	----	----------	---------	---------	-----	--

3.4.2. Insecta

<u>Chironomus</u>	larvae WW	2.831	-2.350	0.981	30	5.3-16.4	Askö	Apr.	#5	
-------------------	-----------	-------	--------	-------	----	----------	------	------	----	--

3.4.3. Echinodermata

<u>Asterias rubens</u>	DW	0.871	0.786	0.983	68	-	Kiel Bay	Jun-Aug	#4	$l = \phi [cm], \log e$
	AFDW	0.938	0.323	0.978	68	-	"	"	"	"

Appendix II:

References to the data in Appendix I

1. Andersin, A.-B., J. Lassig and H. Sandler (1984): On the biology and production of *Pontoporeia affinis* Lindstr. in the Gulf of Bothnia. *Limnologica* 15: 395-401.
2. Aneer, G. (1979): A two year study of the Baltic herring in the Askö-Lansort area. Contr. Askö Lab., Univ. Stockholm, No 8, 2nd corrected printing.
3. Anger, K. (1979): Untersuchungen zum Lebenszyklus der Amphipoden *Bathyporeia sarsi*, *Microdeutopus gryllotalpa* und *Corophium insidiosum* in der Kieler Bucht. *Mitteilungen Zool. Museum Univ. Kiel* I (3): 1-6.
4. Anger, K., U. Rogal, G. Schriever and C. Valentin (1977): In-situ investigations on the echinoderm *Asterias rubens* as a predator of soft-bottom communities in the western Baltic Sea. *Helgoländer Wiss. Meeresunters.* 29: 439-459.
5. Ankar, S.: Answer to BMB Working Group No 11, unpublished data.
6. Ankar, S. (1980): Growth and production of *Macoma balthica* (L.) in a Northern Baltic soft bottom. *Ophelia Suppl.* 1: 31-48.
7. Ankar, S. and R. Elmgren (1978): The benthic macro- and meio-fauna of the Askö-Landsort area (Northern Baltic Proper). A stratified random sampling survey. Contr. Askö Lab., Univ. Stockholm, No 11, 2nd corrected printing, 115p.
8. Arntz, W.E. and D. Brunswig (1975): An approach to estimating the production of macrobenthos and demersal fish in a Western Baltic *Abra alba* community. *Merentutki. Julk./Havsfors. Skr.* 239: 195-205.
9. Arntz, W.E. and W. Weber (1970): *Cyprina islandica* L. (Mollusca, Bivalvia) als Nahrung von Dorsch und Kliesche in der Kieler Bucht. *Berichte Deutsche Wiss. Komm.* 21: 193-209.
10. Atkinson, E.G. and J.W. Wacasey (1976): Caloric values of zoobenthos and phytobenthos from the Canadian Arctic. Dep. of the Envirom., Fish. and Marine Service, Technical Report No. 632.

11. Banse, K., F.H. Nichols and D.R. May (1971): Oxygen consumption by the seabed III. - On the role of the macrofauna at three stations. *Vie et Milieu* 22 Suppl.: 31-52.
12. Bast, H.-D. and J.-A. v.Oertzen (1976): Zusammenstellung von Energieäquivalenten aquatischer Organismen unter besonderer Berücksichtigung der Ostsee. Teil 2. *Wiss. Zeitschr. Univ. Rostock* 25, Mat.-Nat. Reihe, Heft 3.
13. Brawn, V.M., D.L. Peer and R.J. Bentley (1968): Caloric content of the standing crop of benthic and epibenthic invertebrates of St. Margaret's Bay, Nova Scotia. *J. Fish. Res. Bd. Canada* 25: 1803-1811.
14. Brey, T.: Answer to BMB Working Group No 11, unpublished data.
15. Brunswig, D. (1973): Der Nahrungswert makrobenthischer Fischnährtiere der Kieler Bucht im Jahresgang. Diplom-Thesis, Kiel
16. Cederwall, H. (1975): Production of *Pontoporeia affinis* in the Northern Baltic Proper. Manuscript presented at the 4th BMB-Symposium, Gdansk.
17. Chambers, M.R. and H. Milne (1979): Seasonal variation in the condition of some intertidal invertebrates of the Ythan Estuary, Scotland. *Estuarine Cost. Mar. Sci.* 8: 411-419.
18. Cummins, K.W. and J.C. Wuycheck (1971): Caloric equivalents for investigations in ecological energetics. *Internat. Vereinigung Theoret. Angewandte Limnol., Mitteilung No. 18.*
19. Evans, R. (1977): The shallow water soft bottom benthos in Lindaspolene, Western Norway. 2. Estimation of biomass for some macro-infauna species. *Sarsia* 63: 97-111.
20. Foberg, M. (1976): Produktionsberäkningar för några makrovertebrater i luleå skärgård sommaren 1976. *Zool. Inst. Univ. Stockholm.*
21. Gilbert, M.A. (1973): Growth rate, longevity and maximum shell length of *Macoma balthica* (L.). *Biol. Bull.* 145: 119-126.
22. Gründel, E. (1976): Qualitative und quantitative Untersuchungen an einem Ökosystem "Zostera-Wiese" vor Surendorf (Kieler Bucht, westliche Ostsee). *Univ. Kiel, SFB 95, Report No 18.*
23. Hakala, I. (1979): Seasonal variation in the carbon and energy contents and assimilation of a *Mysis relicta* population in Pääjärvi, Southern Finland. *Ann. Zool. Fennici* 16: 129-137.

24. Hansson, S. (1976): Energetiska studier hos abborre (*Perca fluviatilis*) och skrubbskädda (*Pleuronectes flesus*) från norra Östersjön. Askö Lab., unpubl. manuscript.
25. Hughes, R.H. (1970): An energy budget for a tidal flat population of the bivalve *Scrobicularia plana* (da Costa). J. Anim. Ecol. 39: 357-381.
- 26A Jansson, A.-M. (1974): Community structure, modelling and simulation of the *Cladophora* ecosystem in the Baltic Sea. Contr. Askö Lab., Univ. Stockholm, No 5, 130p.
26. Jansson, B.-O. and F. Wulff (1977): Ecosystem analysis of a shallow sound in the Northern Baltic. A joint study by the Askö group. Contr. Askö Lab., Univ. Stockholm, No 18, 160p.
27. Johnson, M.S. and R.O. Brinkhurst (1971): Production of benthic macroinvertebrates of Bay of Quinte and Lake Ontario. J. Fish. Res. Bd. Canada 28: 1699-1714.
28. Kangas, P.: Answer to BMB Working Group No 11, unpublished data.
29. Kautsky, N. (1981): Quantitative studies on gonad cycle, fecundity, reproductive output and recruitment in a Baltic *Mytilus edulis* L. population. Mar. Biol. 68: 143-160.
30. Kreutzberg, M. and J.-A. v.Oertzen (1973): Zusammenstellung von Energieäquivalenten aquatischer Organismen unter besonderer Berücksichtigung der Ostsee. Wiss. Zeitschr. Univ. Rostock 22, Mat.-Nat. Reihe, Heft 10.
31. Lappalainen, A. (1980): Reproduction, growth and production of *Hydrobia ulvae* (Pennant) in the Northern Baltic Proper. Askö Lab., unpubl. manuscript.
32. Lappalainen, A. and P. Kangas (1975): Littoral benthos of the northern Baltic Sea. 2. Interrelationships of wet, dry and ash-free dry weight of macrofauna in the Tvärminne area. Int. Rev. Ges. Hydrobiol. 60: 297-312.
33. Lie, U. (1968): A quantitative study of benthic infauna in Puget Sound, Washington, USA, in 1963-1964. Fiskeridirektorat. Skrifter Ser. Havunders. 14: 229-556.
34. Möller, P. and R. Rosenberg (1983): Recruitment, abundance and production of *Mya arenaria* and *Cardium edule* in marine shallow waters, Western Sweden. Ophelia 22: 33-55.

35. Møhlenberg, F. and T. Kiørboe (1981): Growth and energetics in *Spisula subtruncata* (da Costa) and the effect of suspended bottom material. *Ophelia* 20: 79-90.
36. Murawski, S.A. and F.M. Serchuk (1979): Shell length - meat weight relationships of Ocean Quahogs, *Arctica islandica*, from the Middle Atlantic shelf. *Proc. National Shellfish. Ass.* 69: 40-46.
37. Norrbin, F. and U. Bamstedt (1984): Energy contents in benthic and planctonic invertebrates of Kosterfjorden, Sweden. A comparison of energetic strategies in marine organism groups. *Ophelia* 23: 47-64.
38. v.Oertzen, J.-A. (1979): Versuch einer kausalökologischen Leistungsanalyse an brackwasserlebenden Tieren. Habil. Diss., Univ. Rostock, 188p.
39. Ostrowski, J. (1977): Biomasa i produkcja skorupiaka *Pontoporeia femorata*/Kroyer/ oraz małza *Macoma balthica*/Linne/ w Zatoce Gdąnski. *Prace MIR Tom 18, seria A.*
40. Petersen, G.H. (1977): The density, biomass and origin of the bivalves of the Central North Sea. *Meddel. Danm. Fisk.- og Havunders. N.S.* 7: 211-273.
41. Persson, E.: Answer to BMB Working Group No 11.
42. Pruus, T. (1970): Calorific value of animals as an element of bioenergetical investigations. *Pol. Arch. Hydrobiol.* 17: 183-199.
43. Rachor, E., W.E. Arntz and H. Rumohr (1982): Seasonal and long-term population fluctuations in *Diastylis rathkei* (Crustacea, Cumacea) of Kiel Bay and German Bight. *Netherl. J. Sea Res.* 16: 141-150.
44. Saebø, B.: unpublished. Mar. Lab. Helsingør.
45. Sarvala, J.: Production of *Harmothoe sarsi* (Polychaeta) at a soft bottom locality near Tvärminne, Southern Finland. Lecture at the 6th BMB Symposium, Gdansk 1975.
46. Schaefer, R.; K. Trutschler and H. Rumohr (1985): Biometric studies on the bivalves *Astarte elliptica*, *A. borealis* and *A. montagui* in Kiel Bay (Western Baltic Sea). *Helgoländer Meeresunters.* 39: 245-253.

47. Semenchenko, V.P. and Z.A. Romanova (1976): Length-weight characteristics and fecundity of *Idotea chelipes* (Crustacea, Isopoda) from the Baltic Sea. *Hydrobiol. J.* 12: 96-99.
48. Slobodkin, L.B. and S. Richman (1961): Calories/g in species of animals. *Nature* 191: 299.
49. Thayer, G.W., W.E. Schaaf, J.W. Angelovic and M.W. LaCroix (1973): Caloric measurements of some estuarine organisms. *Fishery Bull.* 71: 289-296.
50. Tyler, A.V. (1973): Caloric values of some North Atlantic invertebrates. *Mar. Biol.* 19: 258-261.
51. Weigelt, M.: Answer to BMB Working Group No 11, unpublished data.
52. Westin, L.: Answer to BMB Working Group No 11, unpublished data.
53. Widbom, B. (1984): Determination of average individual dry weight and ash-free dry weight in different sieve fractions of meiofauna. *Mar. Biol.* 84: 101-108.
54. Wikander, P.B. (1980): Biometry and behaviour in *Abra nitida* (Müller) and *A. longicallus* (Scacchi) (Bivalvia, Tellinacea). *Sarsia* 65: 255-268.
55. Baltic Sea Environment Proceedings No 9 (1983). Second biological intercalibration workshop, Rønne 1982.

Table of applied methods

refer. No	storage	DW °C / h	AFDW °C / h	calorimeter type	area
1.	?	?	-	-	Baltic, Gulf of Bothnia
2.	Fresh	60/>24	?	-	Baltic, Askö
3.	Formalin	105/ 12	-	-	Baltic, Kiel Bay
4.	Formalin	80/ ?	500/ 3	-	Baltic, Kiel Bay
5.	Fresh	60/>24	-	-	Baltic, Askö
6.	Fresh	60/24	-	-	Baltic, Askö
7.	Fresh	60/24	-	-	Baltic, Askö
8.	Fresh	80/ 24	250/ 1 +550/ 1	Phillipson	Baltic, Kiel Bay
9.	Formalin	-	-	-	Baltic, Kiel Bay
10.	Fresh	100/12-24	500/ 18	Parr	Canadian Arctic
11.	Formalin	?	?	-	Pacific, Seattle
12.	Fresh	60/ 24	500/ 8	MBC-3	Baltic, incl Lit.
13.	Fresh	75/->c	-	Parr	Atlantic, N. Scotia
14.	Formalin	85/ 24	520/ 24	-	Baltic, Kiel Bay
15.	Fresh	80/ 24	250/ 1 +550/ 1	Phillipson	Baltic, Kiel Bay
16.	Fresh	60/336	-	-	N. Baltic Proper
17.	Fresh	freeze- dried		Gallenkamp	North Sea, Scotland
18.					
19.	Formalin/ Alcohol	100/10- 48	550/ 6- 24	-	North Sea, Norway
20.	Fresh	60/>24	525/ 4.3	-	Lulea Archipelago
21.	Fresh	70/ 24	-	Phillipson	Atlantic, Massachu.
22.	Fresh	80/ 24	550/ 24	Phillipson	Baltic, Kiel Bay
23.	?	?	?	?	S. Finland
24.	Fresh	60/>24	?	Gallenkamp	Baltic, Askö
25.	Fresh	50/ 48	500/ 2	Gallenkamp	Irish Sea, N. Wales
26A	Fresh	60/>24	?	Phillipson	Baltic, Askö
26.	Fresh	60/ 24	-	-	Baltic, Tvärminne
27.	Fresh	60/ ?	?	?	Lake Ontario
28.	Fresh or Formalin	70/24	-	-	Baltic, Tvärminne

refer.	storage	DW °C / h	AFDW °C / h	calorimeter type	area
No					
29.	Fresh	60/>24	-	-	Baltic, Askö
30.	-	-	-	-	Baltic, Literature
31.	Fresh or Formalin	60/ 24	-	-	Baltic, Askö
32.	Formalin	70/ 24	350-400/2 +500-550/3	-	Baltic, Tvärminne
33.	Formalin	95/4-8	200-500/8.5 + 500 / 16	-	Pacific, Puget Sound
34.	?	80/to const. weight	550/ 5	-	Skagerrak
35.	?	-	550/ ?	-	Oresund
36.	Fresh	-	-	-	Atlantic
37.	Fresh	60/ 48	500/ ?	Phillipson	Koster Fjord
38.	?	?	?	-	Baltic, Mecklenb.
39.	Fresh	?	?	-	Baltic, Gdansk Bay
40.	Fresh	-	-	-	North Sea
41.	Formalin	100/ ?	-	-	Baltic, Hanö Bight
	or 60/ ?				
42.	-	-	-	-	Literature
43.	Formalin	?	?	Phillipson	Baltic & North Sea
44.	Fresh	105/ 12	550/ 5	-	?
45.	Fresh or Formalin	60/ 24	500/ 3	-	Baltic, Tvärminne
46.	Fresh	80/ 24	-	-	Baltic, Kiel Bay
47.	Formalin	?	-	-	Baltic, Gulf of Finl.
48.	Fresh	?	?	?	?
49.	Fresh	60/ ?	-	Phillipson	Atlantic
	or freeze-dried				
50.	Fresh	60/ 5	-	Parr	Atlantic
51.	Formalin	80/ 24	520/ 24	-	Baltic, Kiel Bay
52.	Fresh	60/>24	?	Gallenkamp	Baltic, Askö
53.	Formalin	60/ 12	550/ 4	-	Gullmar Fjord
54.	Fresh	100/ 24	550/ 24	-	Norway
55.	Formalin	according to BMB-recommend.			Baltic, Bornholm