

## New Records of Marine Algae From the Philippines

JOHN A. WEST and HILCONIDA P. CALUMPONG<sup>1</sup>

*Department of Plant Biology, University of California  
Berkeley, CA, U.S.A. 94720*

**Abstract**—Five species of marine algae, *Acrothamnion preissii* (Sonder) Wollaston (Rhodophyceae, Ceramiales), *Caulacanthus ustulatus* (Mert.) Kütz. (Rhodophyceae, Gigartinales), *Ochrosphaera verrucosa* Schussnig (Prymnesiophyceae, Coccospaerales), *Ostreobium quekettii* Bornet et Flahault (Chlorophyceae, Bryopsidales), and *Rhodosorus marinus* Geitler (Rhodophyceae, Porphyridiales) are recorded for the first time in the Philippines. The sporophytic stage of *Derbesia marina* (Lyngbye) Solier is also reported. *Caulacanthus indicus* Weber-van Bosse and *C. okamurae* Yamada are synonymized with *C. ustulatus* based on morphological and anatomical comparisons of type material with field and culture specimens and successful cross-breeding of Philippine and Australian isolates in culture.

### Introduction

In 1987 and 1988 the authors collected benthic marine algae on several islands in the Philippines. Among the collections of benthic marine algae from these areas were several specimens of unreported species. The plants were collected by hand, from different depths and habitats as indicated. Part of these collections were placed in culture following the methods of West & Calumpong (1988). Sections were made by hand with razor blades. Squash sections were made by first softening the tissue with saturated chloral hydrate solution for 2-5 minutes. These sections were then stained with 0.4% aniline blue-black in 0.1% acetic acid (5% formalin was added to prevent fungal growth in the mounting medium) and mounted in 50% Karo<sup>®</sup> syrup. Coverslips were sealed with clear fingernail varnish. All sections were examined under light microscopy. Type specimens were used for comparison when available. Voucher specimens are deposited in the University Herbarium of the University of California at Berkeley (UC), California, U.S.A. and the Silliman University Herbarium (DGT) at Dumaguete City, Philippines. Voucher cultures are deposited with the Culture Collection of Algae, Department of Botany, University of Texas, Austin TX 78713-7640 U.S.A. Classification follows that of Silva, Meñez & Moe (1987).

Class RHODOPHYCEAE  
Order PORPHYRIDIALES  
Family PORPHYRIDIACEAE  
*Rhodosorus marinus* Geitler

*Rhodosorus marinus* Geitler, 1930: 615–636.

This unicellular red was isolated from a culture of *Laurencia patentiramea* (Montagne) Kützing collected by H. Calumpong from Cang-alwang, Siquijor I., Central Vis-

<sup>1</sup> Present Address: *Department of Biology, Silliman University, Dumaguete City 6200, Philippines*

yas, 10 May 1987 (culture 2789A). The cells are about 5  $\mu\text{m}$  diam and contain numerous starch granules. The chloroplast usually has 3-4 parietal lobes and a central pyrenoid evident in light microscopy. A 3- $\mu\text{m}$  polysaccharide halo surrounds each cell.

The distribution record of *Rhodosorus marinus* is patchy. It was described originally in the Canary Islands (Geitler 1930) and subsequently recorded in France (Giraud 1958), Italy and the Florida Keys (Ott 1967) and Hawaii (West 1969). We believe that the alga is more ubiquitous than records indicate. It may be overlooked in collections because of its very small size.

Order GIGARTINALES

Family RABDONIACEAE

*Caulacanthus ustulatus* (Turner) Kützing

*Fucus acicularis* Wulff var. *ustulatus* Turner, 1809: 143, 144. Type: Cádiz, Spain, collected by S. de R. Clemete (BM!). (Lectotypified by Searles, 43: 46, pl. 10a. 1968.)  
*Caulacanthus ustulatus* (Turner) Kützing, 1843: 395.

TAXONOMIC SYNONYMS:

*Caulacanthus indicus* Weber-van Bosse, 1921: 222-223, fig. 67. Type: Dongala (Paloe Bay), Celebes (Siboga Stat. 86), 18-19 June 1899 (L 939.69 . . . 794!). (Lectotypified herein).

*Caulacanthus okamurae* ("okamurae") Yamada 1933: 277-285, pls. 10-13. Type: Prov. Mikawa, Japan, collected by K. Okamura, 11 June 1899 (SAP!). (Lectotypified at SAP and confirmed herein.)

Taxonomic synonyms as proposed by Searles 1968:

*Laurencia divaricata* Suhr., 1840: 265. Type: "Kaffernküste," South Africa (Holotype: LD 34144!).

*Caulacanthus divaricatus* (Suhr.) Papenfuss, 1943: 86-87.

*Rhodomela? spinella* J. Hooker et Harvey, 1845: 534-535. Type: New Zealand, #236, collected by Colenso (BM!). (Lectotypified herein.)

*Caulacanthus spinellus* (J. Hooker et Harvey) Kützing, 1849: 753.

Misapplied name (*fide* Papenfuss, 1943: 86): *Rhodomela botryocarpa*.

CHARACTERISTICS: Field plants growing together with *Murrayella periclados* (C. Ag.) Schmitz, sterile, consisting of long stolons and uniformly positioned upright axes. Branching pectinate or irregularly pectinate, main branches 145-155  $\mu\text{m}$  diam. No rhizoidal discs seen.

SPECIMENS STUDIED: Culture-2920: Jampason, Initao, Misamis Oriental. 14 May 1988. Collected by Wilfredo Uy.

CULTURE STUDIES: A long stolon from the field collection was cleaned of epiphytes and placed in a deep storage dish in sterile seawater. In less than a month, cut ends regenerated new axes from the medulla. The single apical cell of each branch divided obliquely to form the main axial filament, each cell of which in turn formed two lateral filaments as described by Searles (1968). Rhizoidal discs appeared at points that came into contact with the glass substrate. In two months, the plant produced zonately divided tetrasporangia measuring 55-60  $\mu\text{m}$  diam. Spore germination followed the pattern typical of the order. Gametophytes became reproductive after two months. Female gametophytes were more densely and irregularly branched compared to the males and tetrasporophytes.

Carpogonial branches were typically 3-celled and straight and were embedded in the thallus. Carposporophytes released carpospores that germinated readily.

The Philippine material was cross-bred with *Caulacanthus indicus* Weber-van Bosse (culture 2831) collected from Alma Bay, Magnetic Island, North Queensland, Australia on 6 June 1987 from drift. The Australian plants differed slightly from the Philippine material in size and frequency of branching and rhizoidal discs. The Australian isolate has smaller thalli (125-133  $\mu\text{m}$  diam) that are more densely and irregularly branched (at about 1 mm interval) and have numerous rhizoidal discs (at about 0.35 mm interval). In cross-section, the two have a similar pattern with a large colorless thick-walled central axial cell, two layers of round medullary cells and a layer of subquadrate cortical cells. This is a little different from *C. ustulatus* from Mauritius as described by Børgesen (1950) whereby the cortical cells are more elongated.

Results of the cross showed the two isolates to be interfertile although out-crosses produced fewer cystocarps than self-crosses. In the 2831 self-cross, carpospore formation and release took 17 days. The hybrids released carpospores in 25 days.

Another isolate from Korea, 2596 was collected by In Kyu Lee and identified as *Caulacanthus okamurae* Yamada. This was compared with the other two isolates. *Caulacanthus okamurae* has been grown in 10-12°C since 14 Jan 1982 and has remained vegetative even when exposed to bright light ( $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) and higher temperature (25°C). The axes are the same diameter as the Philippine isolate (140-155  $\mu\text{m}$ ). The cross section resembles the Mauritius plants (Børgesen, 1950) more than the Philippine or the Australian isolates.

**DISCUSSION.** The characters used to separate *Caulacanthus indicus* and *C. okamurae* from *C. ustulatus*: diameter of axes, frequency and density of branching, frequency and density of rhizoidal discs, and dimorphism of sterile and reproductive plants appear to be variable as revealed by culture investigations and may not be reliable. Reproductive female gametophytes assume a more irregular and dense branching as compared with the male gametophyte and tetrasporophytes.

King (1981) lists *Caulacanthus okamurae* for Queensland, Ngan & Price (1979) list *C. ustulatus* for Townsville, and Cribb (1983) lists *C. indicus* for the Great Barrier Reef. All of this indicates a great deal of uncertainty about the species limits. Successful cross-breeding of the Philippine and Australian isolates leads one to conclude that these three are conspecific.

Based on the variability and thereby unreliability of characters separating these three taxa and the fact that the two isolates interbreed in culture, we believe that we are dealing with only one taxon here which under the rules of priority take on the name of *Caulacanthus ustulatus* (Turner) Kützing.

Also in comparing the type specimens of *Caulacanthus indicus* Weber-van Bosse with *C. ustulatus* (Turner) Kützing which were both sterile, no significant morphological and anatomical differences were found. The same is true with *C. okamurae* Yamada *Laurencia divaricata* Suhr, and *Rhodomela spinella* J. Hooker et Harvey.

In considering other closely-related genera, we have compared the vegetative and reproductive anatomy of *Feldmannophycus rayssiae* (J. et G. Feldmann) Augier et Boudouresque with that of *Caulacanthus ustulatus*. The principal features considered by Augier & Boudouresque (1971) and by Feldmann & Feldmann (1961) are sufficiently distinct to warrant the continued separation of the genera.

*Caulacanthus ustulatus* has a tropical to subtropical distribution extending from Japan (Yamada 1933) to Australia (Cribb 1983, King 1981, Ngan & Price 1979), and New Zealand (Searles, 1968) across the Pacific in Baja California (Dawson 1961), in Europe (Italy, Spain: Searles 1968) and Africa (Senegal, South West Africa, South Africa: Searles 1968; Mauritius: Børgesen 1950; Somalia: Sartoni 1986; and Tanzania: Jaasund 1976, Lawson 1980), India (Dixit 1966, Searles 1968). This report represents the first record in the Philippines.

Order CERAMIALES

Family CERAMICEAE

*Acrothamnion preissii* (Sonder) Wollaston

*Callithamnion preissii* Sonder 1845: 52. Type: Western Australia (Holotype: MEL 10260).  
*Antithamnion preissii* (Sonder) De Toni, 1903: 1414–1415.

Taxonomic synonyms as proposed by Wollaston, 1977:

*Callithamnion pulchellum* Harvey, 1855: 561. Syntype localities: Rottnest and Cape Riche, Western Australia.

*Acrothamnion pulchellum* J. Ag., 1892: 25, pl. I, figs. 6–10.

CHARACTERISTICS. Field plants sterile, branching opposite. Main axes 25–30  $\mu\text{m}$  diam, lateral branches of the same diameter as the main axes, generally with terminal gland cell. Axial cells 4 times as long as wide.

SPECIMENS STUDIED. Collected by H. Calumpong from Talisoy, Virac, Catan-duanes Island, eastern Philippines, 14 May 1988 (culture 2885). The plants were growing on shaded limestone in the mouth of a semicave, mixed with other small filamentous reds: *Gymnothamnion elegans* (Schousboe ex C. Ag.) J. Ag., *Haloplegma duperreyi* Montagno and *Dasyphila plumariooides* Yendo.

DISCUSSION. Basson, Silva & Moe (unpubl.) point out that Wollaston failed to indicate the type locality of *Callithamnion pulchellum* when she lectotypified it. They further indicate that *C. pulchellum* Harvey is a later homonym of *C. pulchellum* C. Ag. (1828: 175), hence not priorable, and treat *Acrothamnion pulchellum* as a *nomen novum*.

*Acrothamnion preissii* is found in Western Australia (Wollaston 1977), South Africa (Norris & Aken 1985), Japan (Itono 1977), the Mediterranean (Cinelli & Sartoni 1969) and, now, the Philippines.

Class CHOLOROPHYCEAE

Order BRYOPSIDALES

Family OSTREOBIAZAE

*Ostreobium quekettii* Bornet et Flahault

*Ostreobium quekettii* Bornet et Flahault, 1889: CLXI, pl. IX, fig. 5–8. Type: Luc (Calvados), France, collected by P. A. Dangeard (P!). (Lectotypified herein).

Taxonomic synonym as proposed by Lukas, 1974:

*Ostreobium reineckeii* Bornet in Reinbold, 1896: 269. Type locality: Samoa.

CHARACTERISTICS. Vegetative filaments 3–4  $\mu\text{m}$  minimum diam, extensively branched with many short, often irregularly shaped laterals. Chloroplasts very small, discoid to elongate, (2–7  $\mu\text{m}$ ), appearing yellow-green, lacking pyrenoids and resembling those of the Xanthophyceae, without starch granules, although small (<0.5  $\mu\text{m}$ ) granules that do not stain with IKI are present in the cytoplasm. Colorless, refractive lipid bodies

also common. Small coccoid bacteria are abundant in the vacuoles of filaments. Reproduction is by zoospores and by fragmentation of vegetative filaments. Sporangia arising from apices or short lateral branches, initially appearing subspherical, 25-40  $\mu\text{m}$  diam, becoming multilobed with what may be discharge tubes as they enlarge.

**SPECIMENS STUDIED:** Isolated from specimens of *Acrothamnion preissii* (culture 2885) collected on 14 May 1988, by H. Calumpong on shaded limestone at the mouth of a semicave at Talisoy, Virac, Catanduanes Island, eastern Philippines (culture 2924).

**DISCUSSION.** The above observations were made from plants grown in the laboratory in the absence of calcium carbonate substrate (culture 2824). The small coccoid bacteria which were abundant in the vacuoles of filaments are probably endosymbionts generally common in siphonous greens (Burr & West 1970).

*Ostreobium* usually is "endolithic" in corals, shells and coralline algae so the appearance of free living plants is of interest. The species has been investigated in culture by Kornmann & Sahling (1980). Lukas (1974) recognizes five species and our isolate appears closest to *O. quekettii* Bornet et Flahault. The "type" material of *O. quekettii* is deposited at the Muséum National d'Histoire Naturelle Laboratoire de Cryptogamie at Paris (P) but Bornet & Flahault (1889) did not specify from which of the three oyster shells in that collection they obtained the sample on which they based their description. From the Muséum we received three separate samples of bivalve shell fragments labeled: Le Croisic (September 1887), Brest (October 1886) and Luc (no date—collected by P. A. Dangeard) labeled *Gomontia polyrhiza* Bornet et Flahault. After careful examination of these samples under the steromicroscope pale green 0.5 mm fragments were excised and immersed for 1 hr in 1% HCl for decalcification. Under the compound microscope the green filaments observed were extensively branched, septate and 5-12  $\mu\text{m}$  diam. These appear to be *Ostreobium quekettii* and with reservations we designate the Luc material as the lectotype. .

Although Jeffrey (1968) has identified chlorophyll a and b in *Ostreobium quekettii* from the brain coral *Favia*, this should be reaffirmed with this isolate as well.

*Ostreobium quekettii* is widely distributed in North America (Taylor 1957; Florida Keys: Lukas 1974), Europe (British Isles: Lyle 1929, Wilkinson & Burrows 1972; France: Bornet & Flahault 1889; Netherlands: van den Hoek 1958; North, Baltic, Black and Red Seas: Nadson 1927), the Atlantic (Bermuda: Schroeder 1972), Caribbean (Lukas 1974), Pacific (Caroline and Marshall Islands: Lukas 1973; Samoa: Bornet 1896, Setchell 1924), Indian Ocean (India: Dixit 1966, Krishnamurthy & Joshi 1970, Untawale *et al.* 1983; Maldives: Hackett 1977; Faeröes Islands: Børgesen 1908), and Australia (De Toni & Forti 1923, Lewis 1987).

Class PRYMNESIOPHYCEAE

Order COCCOSPHAERALES

Family HYMENOMONADACEAE

*Ochrosphaera verrucosa* Schussnig

*Ochrosphaera verrucosa* Schussnig, 1940: 317-330.

*Ochrosphaera verrucosa* is widely distributed in the Philippines and occurs in nearly all cultures of macroalgae that we have isolated from there. Because it reproduces so rapidly it is often difficult to eliminate and can be classified as a "pest species." The same

frequency is evident in collections from Hawaii (West 1969), Japan (Inouye & Chihara 1980), Indonesia (Inouye 1988), Australia, Brazil, Mexico, Puerto Rico, Samoa, Guam, Truk, Seychelles, Comoros, Maldives, (West, unpublished). We have isolated one strain from a collection of *Laurencia concinna* Montagne (culture 2773) collected by H. Calumpong from White Beach near the town of Mahatao, Batan Island, Batanes, 22 April 1987. The specimens in culture correspond in size and cytological detail to those described in West (1969) and Inouye & Chihara (1980).

#### Order BRYOPSIDALES

#### Family DERBESIACEAE

#### *Derbesia marina* (Lyngbye) Solier

*Vaucheria marina* Lyngbye, 1819: 79, pl. 22A. Type locality: Kvivig, Strømø, Faeröes.

*Derbesia marina* (Lyngbye) Solier, 1846: 453.

Taxonomic synonyms:

*Gastridium ovale* Lyngbye, 1819: 72, pl. 18B. Syntype localities: various in Faeröes.

*Halicystis ovalis* (Lyngbye) J. Areschoug, 1850: 447.

Only gametophytes (*Halicystis ovalis* stage) of *Derbesia marina* (Lyngbye) Solier were reported in the Philippines (Domantay 1962, Reyes 1978, Saraya & Trono 1980). In a collection of *Laurencia* from Cang-alwang, Siquijor Island, Central Visayas, 8 May 1987, by H. Calumpong (culture 2791D), reproductive filaments of the sporophyte (*Derbesia marina* stage) were seen. The filaments measured 15-30  $\mu\text{m}$  in diameter. Spores germinated into irregular spheres.

#### Acknowledgments

J. West and H. Calumpong acknowledge travel support from NSF International Programs INT-8613341. Dr. E. Meñez (Smithsonian Institution Oceanographic Sorting Center, Washington, D.C.) provided logistical assistance during our trips to the Philippines. Dr. Y. Hara (Tsukuba University, Japan) confirmed the identification of *Rhodosorus marinus*. Dr. Paul Silva (University Herbarium, University of California at Berkeley) provided substantial advice on nomenclature and references. We would like to thank the curators of the following herbaria for lending type material: British Museum (Natural History) at London (BM), Herbarium of Botanical Museum at Lund (LD), the Royal Botanic Gardens and National Herbarium at Melbourne (MEL), Muséum National d'Histoire Naturelle Laboratoire de Cryptogamie at Paris (P), the Herbarium of the Faculty of Science, Hokkaido University at Sapporo (SAP) and Rijksherbarium at Leiden (L).

#### References

Agardh, C. 1820. Species Algarum. Vol. 2, sect. 1, lxxvi+189 pp. Gryphia (Greifswald).

Agardh, J. 1892. Anlecta Algologica. Lunds Univ. Årsskr. . . . Lund. Handlingar 28(6): 182 pp., III pls.

Areschoug, J. E. 1850. *Phycearum quae in maribus Scandinaviae crescunt, enumeratio. Sectio posterior Ulvaceas continens.* Nova Acata R. Soc. Scient. Upsal. 14: 385–454.

Augier, H. & C. F. Boudouresque. 1971. Découverte des cystocarpes de *Feldmannophycus rayssiae* (J. et G. Feld.) nov. gen. (Rhodophycées, Gigartinales). Bull. Soc. Phycol. France 16: 25–30.

Børgesen, F. 1908. The algae-vegetation of the Faeröes coasts with remarks on phytogeography. In *Botany of the Faeröes Based Upon Danish Investigations. Part III.* John Wheldon, London, pp. 683–834.

Børgesen, F. 1950. Some marine algae from Mauritius. Additions to the parts previously published. II. K. danske Vidensk. Selsk. Skr. 18(11): 19–21.

Bornet, E. & C. Flahault. 1889. Sur quelques plantes vivant dans le test calcaire des mollusques. Bull. Soc. Bot. France 36: cxlvii–clxxvi.

Burr, F. & J. West. 1970. Light and electron microscope observations on the vegetative and reproductive structures of *Bryopsis hypnoides*. Phycologia 9: 17–37.

Cinelli, F. & G. Sartoni. 1969. *Acrothamnion* J. Ag. (Rhodophyta, Ceramiaceae): genere algale nuovo per il mare Meditarraneo. Pubbl. Staz. Zool. Napoli. 37: 567–574.

Cribb, A. B. 1983. Marine Algae of the Southern Great Barrier Reef. Part I. Rhodophyta. Australian Coral Reef Society, Handbook No. 2. Watson, Ferguson & Co., Brisbane.

De Toni, G. B. 1903. *Sylloge Algarum . . . Vol. IV. Florideae. Sectio II. Patavii.* pp. 387–776.

De Toni, G. B. & A. Forti. 1923. Alghe di Australia, Tasmania e Nouva Zealand raccolte dal rev. dott. Giuseppe Capra Nel 1908–1909. Mem. del Reale Inst. Venete di Sci., Lett. ed Arti 29(3): 23.

Dixit, S. C. 1966. Species list of Indian marine algae determined by Børgesen. J. Univ. Bombay Sc. 32(54 and 55): 12.

Domantay, J. 1962. An ecological survey of the marine vegetation of the Hundred Islands and vicinity. Phil. J. Sci. 90: 271–295.

Feldmann, J. & G. Feldmann. 1961. Une nouvelle Rhodophycée mediteranéene *Caulacanthus* (?) *rayssiae* sp. nov. Bull. Res. Counc. Israel 10D: 59–65.

Geitler, L. 1930. Ein grunes Filarplasmadium und andere neue Protisten. Arch. Protist. 69: 615–636.

Giraud, G. 1958. Sur le vitesse de croissance d'une Rhodophycée monocellulaire marine, le *Rhodosorus marinus* Geitler. Compte Rendu Acad. Sci. Paris 246: 3501–3504.

Giraud, G. 1962. Les infrastructures de quelques algues et leur physiologie. J. Microscopie 1: 252–254.

Hackett, E. 1977. Marine algae from the Maldives Islands. Atoll Res. Bull. 210: 27.

Harvey, W. H. 1855. Some account of the marine botany of the colony of Western Australia. Trans. R. Ir. Acad. 22 (Science): 525–566.

Hooker, J. D. & W. H. Harvey. 1845. *Algae Novae Zealandiae.* London J. Bot. 4: 534–535.

Inouye, I. 1988. 3. Microalgae of mangroves and characteristics of the community. In

K. Ogino & M. Chihara (eds.), *Biological System of Mangroves. A report on East Indonesian mangrove expeditions, 1986.* pp. 35–43. Ehime University.

Inouye, I. & M. Chihara. 1980. Laboratory culture and taxonomy of *Hymenomonas coronata* and *Ochrosphaera verrucosa* (Class Prymnesiophyceae) from the Northwest Pacific. *Bot. Mag. Tokyo* 93: 195–208.

Itono, H. 1977. Studies on the ceramiaceous algae (Rhodophyta) from southern parts of Japan. *Bibliotheca Phycologia* 35: 366.

Jaasund, E. 1976. *Intertidal Seaweeds in Tanzania.* University of Tromso, Norway. p. 89, fig. 180.

Jeffrey, S. 1968. Pigment composition of Siphonales algae in the brain coral *Favia*. *Biol. Bull.* 135: 141–148.

King, R. 1981. Mangroves and salt marshes plants. In M. Clayton & R. King (eds.), *Marine Botany, An Australian Perspective.* pp. 308–328. Longman Cheshire, Melbourne.

Kornmann, P. & P. Sahling. 1980. *Ostreobium quekettii* (Codiaceae, Chlorophyta). *Helgol. Meeresunters.* 34: 115–122.

Krishnamurthy, V. & H. Joshi. 1970. A check list of Indian marine algae. Central Salt and Marine Chemicals Research Institute. Bhavnagar.

Kützing, F. T. 1843. *Phycologia Generalis oder Anatomie, Physiologie und Systemakunde der Tange.* Leipzig. 458 pp.

Kützing, F. T. 1849. *Species Algarum.* Leipzig. 922 pp.

Lawson, G. W. 1980. *A Checklist of East African Seaweeds (Djibouti to Tanzania).* Lagos, Nigeria: Department of Biological Sciences, University of Lagos. (Mimeo), p. 36.

Lewis, J. 1987. Checklist and bibliography of benthic macroalgae recorded from Northern Australia. III. Chlorophyta. Report MRL-R-1063. Austral. Dept. Defence Materials Res. Lab. Melbourne.

Lukas, K. 1973. Taxonomy and ecology of the endolithic microflora of reef corals with a review of the literature on endolithic microphytes. Ph.D. Thesis, Univ. Rhode Island, 159 pp.

Lukas, K. 1974. Two species of the chlorophyte genus *Ostreobium* from skeletons of Atlantic and Caribbean reef corals. *J. Phycol.* 10: 331–335.

Lyle, L. 1929. Marine algae of some German warships in Scapa Flow and of the neighboring shores. *J. Linn. Soc. (London)* 48: 231–257.

Lyngbye, H. 1819. *Tentamen Hydrophytologie Danicae.* Copenhagen (Hafnia), 248 pp.

Nadson, G. A. 1927. Les algues perforantes, leur distribution et leur rôle dans la nature. *Compte Rendu Acad. Sci.* 184: 1015–1017.

Ngan, Y. & I. R. Price. 1979. The intertidal algae in the mainland coast in the vicinity of Townsville, Queensland. *Atoll Res. Bull.* No. 237: 1–29.

Norris, R. E. & M. E. Aken. 1985. Marine benthic algae new to South Africa. *S. Afr. J. Bot.* 51: 55–65.

Ott, F. 1967. *Rhodosorus marinus* Geitler: a new addition to the marine algal flora of the western hemisphere. *J. Phycol.* 3: 158–159.

Papenfuss, G. F. 1943. Notes on South African marine algae. II. J. S. Afr. Bot. 9: 86-87.

Reinbold, T. 1896. Meeresalgen (Schizophyceae, Chlorophyceae, Phaeophyceae, Rhodophyceae). In F. Reinecke. Die Flora der Samoa-Inseln. Bot. Jahrb. Syst. 23: 266-275.

Reyes, A. 1978. The littoral benthic algae of Siquijor Province. I. Cyanophyta and Chlorophyta. Phil. J. Sci. 105: 133-191.

Saraya, A. & G. Trono. 1980. The marine benthic algae of Santiago Island and adjacent areas in Bolinao, Pangasinan, I. Cyanophyta and Phaeophyta. Nat. Appl. Sci. Bull. (Univ.-Philip. 31: 1-59).

Sartoni, G. 1986. Algal flora and his (sic) vertical distribution on the Gesira Cliff (central-southern Somalia). Webbia 39: 355-377.

Schroeder, J. H. 1972. Calcified filaments of an endolithic alga in Recent Bermuda reefs. Neues Jahrb. Geol. Paleontol. 1972(1): 16-33.

Schussing, B. 1940. Über einige neue Protophyten aus der Adria. Arch. Protistenk. 93: 317-330.

Searles, R. B. 1968. Morphological studies of red algae of the order Gigartinales. Univ. Cal. Publ. Bot. 43: 45-51, figs. 19-21.

Setchell, W. 1924. American Samoa, Part I: Vegetation of Tutuila Island; Part II: Ethnobotany of the Samoans; Part III: Vegetation of Rose Atoll. Carnegie Inst. Wash. Publ. 341: 275 pp.

Silva, P. C., E. G. Meñez & R. L. Moe. Catalog of the Benthic Marine Algae of the Philippines. Smith. Contrib. Mar. Sci. No. 27. Smithsonian Institution Press, Washington, D.C.

Solier, A. J. J. 1846. Sur deux algues zoosporées formant le nouveau genre *Derbesia*. Rev. Bot. 1: 452-454.

Sonder, O. G. 1845. Nova algarum genera et species, quas in itinere ad oras occidentales Nova Hollandiae, collegit L. Preiss, Ph. Dr. Bot. Ztg. 3: 49-57.

Suhr, J. N. 1840. Beiträge zur Algenkunde. Flora 23: 265.

Taylor, W. R. 1957. Marine Algae of the Northeastern Coast of North America. Univ. Mich. Press, Ann Arbor, 227 pp.

Turner, D. 1809. Fuci . . . Vol. II. London, pp. 143-144.

Untawale, A., V. Dhargalkar and V. Agadi. 1983. List of marine algae from India. National Institute of Oceanography, Dona Paula, Goa.

van den Hoek, C. 1958. The algal microvegetation in and on barnacle shells collected along the Dutch and French coasts. Blumea 9(1): 206-214.

Weber-van Bosse, A. 1921. Liste des algues du Siboga. II. Rhodophyceae, premiere partie. Siboga-Exped. Monogr. 59b: 222-223, fig. 67.

West, J. A. 1969. Rare marine microalgae from Hawaii. Phycologia 8: 187-192.

West, J. A. & H. P. Calumpong. 1988. Mixed-phase reproduction of *Bosstrychia* (Ceramiales, Rhodophyta) in culture. I. *B. tenella* (Lam.) J. Ag. Jpn. J. Phycol. (Soruñ) 36: 292-310.

Wilkinson, M. & E. M. Burrows. 1972. An experimental taxonomic study of the algae confused under the name *Gomontia polyrhiza*. J. Mar. Biol. Assoc. (U.K.) N. S. 52(1): 49-57.

Wollaston, E. M. 1968. Morphology and taxonomy of southern Australian genera of *Crouaniae* Schmitz (Ceramiaceae, Rhodophyta). *Aust. J. Bot.* 16: 217-417.

Wollaston, E. M. 1977. Morphology and taxonomy of the genus *Acrothamnion* J. Ag. (Ceramiaceae, Rhodophyta). *Bull. Jpn. Soc. Phycol.* 25(suppl.): 385-393.

Yamada, Y. 1933. Notes on some Japanese algae. *V. J. Fac. Sc. Hokkaido Imp. Univ., ser. 5, 2:* 277-285, pls. 10-13.

