



**Report**

# **Extreme marine environments**



***Objectives and results  
from selected  
MAST projects***



ENERGY, ENVIRONMENT  
AND SUSTAINABLE DEVELOPMENT

EUR 19097 EN

## **Interested in European research?**

**RTD info** is our quarterly magazine keeping you in touch with main developments (results, programmes, events, etc.). It is available in English, French and German. A free sample copy or free subscription can be obtained from:

Directorate-General for Research

Communication Unit

European Commission

Rue de la Loi/Wetstraat 200

B-1049 Brussels

Fax (32-2) 29-58220

E-mail: [rtd-info@cec.eu.int](mailto:rtd-info@cec.eu.int)

Internet: <http://europa.eu.int/comm/dg12/rtdinfo.html>

## **European Commission**

Directorate-General for Research

Unit D.I.3 — Preserving the ecosystem I — Marine ecosystems, infrastructures

*Contact: Mrs Elisabeth Lipiatou*

*Address: European Commission, rue de la Loi/Wetstraat 200 (SDME 7/84),*

*B-1049 Brussels — Tel. (32-2) 29-66286; fax (32-2) 29-63024*

European Commission

# **Extreme marine environments**

**Objectives and results  
from selected  
MAST projects**

*Edited by*  
Anny Asmanidou and Elisabeth Lipiatou

Directorate-General for Research  
Energy, environment and sustainable development programme

## **LEGAL NOTICE**

Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use which might be made of the following information.

### *Acknowledgement*

Ms Christine Vaines is acknowledged for her contribution to the editing of this report.

A great deal of additional information on the European Union is available on the Internet. It can be accessed through the Europa server (<http://europa.eu.int>).

Cataloguing data can be found at the end of this publication.

Luxembourg: Office for Official Publications of the European Communities, 2000

ISBN 92-828-3011-X

© European Communities, 2000

Reproduction is authorised provided the source is acknowledged.

*Printed in Belgium*

PRINTED ON WHITE CHLORINE-FREE PAPER

## **TABLE OF CONTENTS**

	Page
• INTRODUCTION TO EXTREME MARINE ENVIRONMENTS	1
1. Extreme Marine Environments-Definition	1
2. Objectives and Research Tasks of the Marine Science and Technology Programme (MAST III) for the Extreme Marine Environments	1
3. The projects presented and the sub-programme areas of MAST Programme that they represent	1
4. Third European Marine Science and Technology Conference: Section for the Extreme Marine Environments in Europe, Lisbon, 23-27 May 1998	2
5. Workshop on Extreme Marine Environments, GEOMAR Kiel, 19-22 November 1998	2
6. The MAST Advanced Study Courses (ASC) relevant to the Extreme Marine Environments	3
• OVERVIEW OF OBJECTIVES, MAIN RESULTS, AND MAJOR PUBLICATIONS OF THE MAST RELATED PROJECTS	5
• LIST OF PARTICIPANTS OF THE PROJECTS	39
• DOCUMENTATION USED FOR THIS PUBLICATION	49



## **PURPOSE OF THIS PUBLICATION**

The purpose of this publication is to give an overview of the objectives, the main results and the major publications obtained from several MAST-III-funded projects and workshops in the specific field of extreme marine environments from 1996 to 1999.



**ENERGY, ENVIRONMENT AND SUSTAINABLE DEVELOPMENT**





# INTRODUCTION TO THE EXTREME MARINE ENVIRONMENTS

## 1. Definition of Extreme Marine Environments

Extreme conditions are conditions that define the limits of life in our planet; they are found in the deep ocean and at the poles. Short-term conditions can have dramatic consequences for the entire biosphere and evolution of life, as well as for the Earth's climate, because they can alter living conditions or even turn them irreversible. Through adaptation to extreme environments, micro-organisms play a considerable role in global material-cycling. On the above basis, extreme conditions and events are essential components in the assessment of anthropogenic and natural environmental change. The examination of extreme marine environments also contributes to the following. Deliverable of exotic, often economically important minerals, solutes and organics; micro-organisms which function under these conditions change their physical and chemical environment, including environments of commercial interest such as oil and gas reservoirs and sites of waste storage and dumping. Micro-organisms that adapt to extreme environments are extremophiles and will have biotechnological application, for example bio-remediation, waste treatment. Finally, of great importance is the development of deep-sea technologies for observation of extreme marine environments.

## 2. Objectives and Research Tasks of the Marine Science and Technology Programme (MAST III) for the Extreme Marine Environments

Technical areas supported within the MAST III programme (1994-1998) were: Marine Science, Strategic Marine Research, Marine Technology, and Supporting Initiatives (advanced training system, modelling, use of heavy experimental equipment, calibration techniques for marine instrumentation and observational equipment). The overall objective for the study of extreme marine environments was to understand the functioning of any ecosystems under these conditions, and to examine their physical, chemical, biological and geological processes, so as to determine their role in the global environment. The part of research for extreme marine environments included the following disciplinary areas, which the projects of the present study address.

## 3. The MAST III projects on Extreme Marine Environments

### MARINE SCIENCE

The deep sea floor in the North Atlantic and the Mediterranean:

- Azores Mid-Oceanic Ridge Ecosystem Studies: An integrated research programme on deep-sea hydrothermal transfers and fluxes – *AMORES* (MAS3-CT95-0040)
- High resolution temporal and spatial study of the Benthic biology and Geochemistry of a north-eastern Atlantic abyssal Locality – *BENGAL* (MAS3-CT95-0018)

The ice-covered seas in the northern hemisphere:

- European Subpolar Ocean Programme Phase 2: The Thermohaline Circulation in the Greenland Sea – *ESOP II* (MAS3-CT95-0015)

Sedimentary processes in the deep sea, on the continental slope and on the shelf edge:

- **HYdrothermal FLuxes and biological production in the AeGean - AG-HY-FL (MAS3-CT95-0021)**

#### **4. Third European Marine Science and Technology Conference, Lisbon, 23-27 May 1998: Section for the Extreme Marine Environments in Europe: *Extreme Marine Environments in Europe: Scientific and Technological results and challenges***

Some of the outcomes of this section were:

- Determination of the flux of dissolved substances leaving the sediment to return to the bottom water. More specifically, the flux of anthropogenically introduced material across this boundary. Moreover, the deep-sea biota response to the above flux of material has to be examined. Relative results from testing on models, have shown that exchange processes at the deep sea floor play an important role on the oxygen distribution in the oceanic water column and the atmospheric pCO<sub>2</sub>. For a realistic estimation, in situ measurements have to be carried out.
- Microbial oxidation of reduced inorganic compounds by endosymbiotic bacteria is a common feature in hydrothermal vents and cold seeps of subduction zones. It produces a large benthic oxygen demand that affects the local bottom waters. An open field of research is the colonisation of hydrothermal vents by special communities, due to biological fixation, which leads to formation of precipitates, concentration of exotic metals and fractionation of extreme isotopes. Another open research field is the matrix-fluid interaction under extreme temperatures and pressures. More specifically, the phase separation during boiling of hydrothermal fluids, and transport and fractionation of elements.
- Studies on numerous deep brine lakes that have been discovered in the eastern Mediterranean Sea, which are anoxic hypersaline deep basins with the highest dissolved sulfide, magnesium, potassium and helium concentrations ever reported for marine environments. Their origin is not, so far, clearly known.
- Studies on diversity of thermophilic micro-organisms at deep sea hydrothermal vents in Eastern and Southwest Pacific and the mid-Atlantic (at the Guyana's Basin and the MAR) have succeeded in isolating, for the first time, aerobic thermophiles some of them assigned to the genus *Thermus* and several new spore-forming bacilli. Concerning anaerobic hyperthermophiles, investigations have revealed several heterotrophic anaerobic Archaea.
- A recent finding in deep-sea environments is mud volcanoes that consist of pure serpentinite slurries. Furthermore, gas hydrates, a mixture of water, methane and hydrocarbons, carbon dioxide and hydrogen sulfide, might be of great importance as a future energy source and because they may play a role on climate change through their rapid decomposition in shallow waters.

#### **5. Workshop on Extreme Marine Environments, GEOMAR Kiel, 19-22 November 1998: *The deep ocean: A model for processes functioning under extreme environmental conditions***

An international workshop, co-organised by the European Commission, Directorate General XII, Marine Science and Technology Programme, and the German Ministry for Education, Science, Research and Technology. The essential goal was the better understanding of the extreme environmental conditions in the ocean, at the seafloor,

and below the seafloor by European research groups, heading to the explanation of the contribution of such conditions and events in modeling approaches of environmental change. Results of the workshop indicated that *future research must concentrate on: a. identification of the major interacting parameters in complex geo- and bio-topes, b. assessment of frequencies and magnitudes of extreme events, c. modeling of the re-equilibration of environments after perturbation, d. prediction and evaluation of the potential dangers of extreme events, e. identification of feed-back mechanisms, and g. development of new technologies.*

## **6. The MAST Advanced Study Courses (ASC) relevant to the Extreme Marine Environments**

- Deep Benthic Communities based on Chemosynthesis (Hydrothermal Vents and Cold Seeps), FR-Paris, September 1997
- Submarine Volcanism, Hydrothermal Vents, Associated Biota and Mineral Resources, ES- Barcelona, September 1996



# **OVERVIEW OF OBJECTIVES, MAIN RESULTS, AND MAJOR PUBLICATIONS FROM SELECTED MAST PROJECTS**



Title: **AMORES**

**Azores Mid-Oceanic Ridge Ecosystem Studies:** An integrated research programme on deep-sea hydrothermal transfers and fluxes

Contract No: **MAS3-CT95-0040**

Sub-programme area: Extreme Marine Environments - ***The deep sea floor in the North Atlantic and the Mediterranean***

Duration of the project: February 96' - March 99'





**Background:** Three major tectonic plates (European, North American, and African) have their boundaries in the mid-Atlantic ridge system near the Azores, Triple Junction. The Azores region has an unusually shallow position in the North Atlantic. During the MAST 2 MARFLUX/ATJ project, two hydrothermal sites were discovered: the *Lucky Strike* vent field at 37°15', and *Menez Gwen* vent field at 37°45'.

Lucky Strike is one of the largest hydrothermal areas known in the modern oceans, the hydrothermal system of which is controlled by a lava lake. For the depth of 1700 m, the maximum temperature (320° C) is close to the boiling point (340° C). Menez Gwen vent field has a temperature of 281°C, close to the boiling point. The Rainbow hydrothermal site belongs to the AMAR segment and had not been precisely located before the beginning of the project. The site is controlled by a cross cutting fault system, and was located during the MARFLUX/ATJ project from the very high particle content in the plume (German et al. 1996). Both the Lucky Strike and the Menez Gwen vent fields are above the depth at which a typical hydrothermal fluid starts to boil, whereas Rainbow vent field is below this depth.

**Objectives:** The project aimed to identify and quantify the fluxes within and from the hydrothermal vent ecosystem, the interactions between lithosphere and sea-water, and their impact on the ecosystems of the Mid-Atlantic Ridge (MAR), south-west of the Azores Triple Junction (ATJ). AMORES addressed to four themes:

*Theme 1:* The hydrothermal flux of heat and matter from vent-field to segment scale

*Theme 2:* Submersible geological and geochemical studies at the vent field scale

*Theme 3:* Biology of hydrothermal communities at the vent field scale

*Theme 4:* Supporting studies that involve explanation of the occurrence of hydrothermal venting, the mantle's structure and composition, the detailed setting of hydrothermal vent sites, data base management, and model testing.

The project involved seven cruises, that have supported the following study-areas:

- The physical, geochemical and biological dispersion of the neutrally-buoyant hydrothermal plume at a segmental scale;
- High- and low-temperature physical and chemical fluxes, and the internal structure of hydrothermal system at the vent field scale;
- The influence of depth through chemical, physical, and biological processes on hydrothermal vent systems, in the following active segments: a). *AMAR* segment – hydrothermal site Rainbow (2300 m, deep target), b). *FAMOUS* segment (2250-2600 m, deep target), c). *Lucky Strike* vent field (1700 m, shallow target), d). *Menez Gwen* vent field (800 m, shallow target).

**Methodology and Achievements:** Hydrothermal activity was investigated along three different sections of the slow-spreading Mid-Atlantic Ridge. At the scale of the individual study areas, there has been a difference of up to one order of magnitude from the model developed.. Finally, the model developed was partially confirmed by some recent work on the SW Indian Ridge. Methodology and results concern the following (according to themes):

*Theme 1:* This study was a multi-disciplinary investigation of the Rainbow hydrothermal plume (German et al. 1998 & 1999). The dispersing plume was traced and sampled for a distance of >50 km downstream. *In situ* sensors coupled with shipboard and shore-based analyses were used, as well as *BRIDGET*, a deep-towed hydrothermal plume instrument. Combined short-term and long-term physical

oceanographic observations (extensive set of CTD, LADCP, nephelometry profiles, and long-term current meter moorings, respectively) have showed that *the plume is strongly directed to the north along the rift valley wall directly above the vent-site, and then east across a shelf into the adjacent segment.*

The neutrally buoyant plume is dominated by Fe oxy-hydroxides. These oxy-hydroxides are of much lower density than sulfides would be, so the flux of settling particulate material is reduced, compared to other vent-sites, and the total suspended particulate load behaves "conservatively".

There is an unusually high concentration of particle-associated microbial material within the dispersing plume. Although this material is dominated by *Crenarchaeota*, the *Euryarchaeota* (which include halogens and methanogens) are remarkably more abundant than is typical in this chemically-enriched deep-sea environment. Despite the high concentrations of dissolved CH<sub>4</sub> and microbiological activity within the Rainbow hydrothermal plume, there was no evidence for significant larval dispersal. Possible explanations include either a) the occurrence of aggressive predation of such larvae in the high biomass waters overlaying the vent-site or b) discontinuous reproduction of organisms at the vent-site, perhaps responding to seasonal inputs of organic carbon from the overlying water column (see Theme 3, below).

*Theme 2:* The study was based on results from the FLORES cruise, the first submersible investigation of the Rainbow hydrothermal field and the continuation of studies on the Lucky Strike and the Menez Gwen vent fields. Results have focused on the study of geological setting of hydrothermal fields, studies of plume and diffuse flows, hot fluid chemical composition on the seafloor, mineralogical and chemical compositions of sulfide deposits, Pb and Sr isotopic composition of fluids and sulfides, and study of helium isotopes in the hydrothermal fluids. The vent fluids in the Rainbow hydrothermal field, which is a high-temperature site that includes ultramafic rocks, display extremely high dissolved H<sub>2</sub> and CH<sub>4</sub> concentrations, high Fe concentrations, and high Fe/H<sub>2</sub>S ratios

Four types of hydrothermal systems have been distinguished: 1. *Menez Gwen*, characterised by focused discharge, enhanced phase separation, volcanic control, and immature deposit, 2. *Lucky Strike*, characterised by focused discharge, volcanic control, and mature deposit, 3. *Rainbow* field, characterised by tectonic control, ultramafic rocks, and mature deposit, 4. *Saldanha Mound* and *South Lucky Strike*, characterised by regional diffuse discharge related to ultramafic rocks.

The occurrence of active hydrothermal systems in the shallow central topographic high with a high magmatic budget and in the ends of the three investigated segments with ultramafic rocks and high methane anomalies, has been identified. The occurrence in these two distinctly different environments *considerably enlarges the areas where potential hydrothermal vents can be found on slow spreading ridges.* The FLORES results showed that shallow depth near the Azores hot spot connects with both the volcanic style of eruption and the hydrothermal activity. Concerning the volcanic eruption, there is a limit between the effusive and explosive volcanism at 900 m of water. For the hydrothermal activity, the important point is that shallow targets Lucky Strike and Menez Gwen are above the boiling point, and the deep targets Amar and Famous are under the boiling point for a typical hydrothermal fluid. One consequence is a depletion of metals in the fluids and deposits at the shallow Menez Gwen site.

The composition of hydrothermal fluids and mineral deposits, and the associated chemical flux into the ocean vary depending on the geological setting and the underlying rock composition. Hot fluid chemistry is very much affected by

*serpentinisation*, which generates a high flux of  $H_2$  and  $CH_4$  along the axis of the MAR. In these environments, deposits are enriched in Cu, Zn, Co, and Ni.

*Theme 3:* The study has concentrated on the communities that live on the periphery of active vents on three sites of the ATJ area at decreasing depths, so as to find the influence of pressure on the structure, the composition, and the functioning of the system. The theme has particularly addressed: biogeography, microdistribution of organisms, nutritional transfers and the food web, physiological adaptations of marine habitats, reproduction and dispersal of marine habitats.

Rainbow, Lucky Strike, and Menez Gwen were explored and described (Desbruyères et al. submitted). Populations in the two shallower ATJ fields differ from those in other MAR sites, *Bathymodiolus azoricus* being dominant. Composition and structure of communities varied with the depth gradient. Caridean shrimps swarms dominate the deepest site; mussel beds are found at the periphery of sites. Mussel beds are more noticeable in the two shallower sites, and cover the active edifices (structures). The major ecological structuring factor between the three areas is the mineral particle flux. As the ridge gets shallower, non-vent carnivorous animals occupy the vent area feeding on the vent communities. Phase separation processes occurring at the shallowest vents are a major environmental factor, which can explain the basic differences between deep-sea and shallow vents where no specialised fauna are found. Reproduction and dispersal studies suggest discontinuity in reproduction and recruitment likely driven by POM (particulate organic matter) fluxes of photosynthetic origin. The study of the food-web using natural isotope ratios, shows an important connection of chemoautotrophic and photoautotrophic based ecosystems in the ATJ area, which cannot be considered as independent systems.

A deep-sea versatile in situ auto-analyser, ALCHIMIST (AnaLyser CHIMique In SiTu), was adapted and operated by a deep sea submersible. It is based on Flow Injection Analysis and colorimetric detection. It provides real time simultaneous in situ determination of nitrite + nitrate and total sulfide concentrations related to a precise positioning of the sample inlet. The in situ analysis was performed at 1700m depth, over hydrothermal vent biological communities. Further work can concentrate on the extension of the range of parameters that can be analysed in situ.

Chemical differences associated with ultramafic rocks in the hydrothermal sites did not appear to produce major faunal modifications. There is likely to be only one biogeographical province in the hydrothermal vent sites of the MAR, between 15°40' and 38°N. Concerning the homologous habitats, nutrient ( $H_2S$  and  $CH_4$ ) concentrations, metallic ion concentrations, and temperatures hardly varied from one field to another, probably due to the high dilution of the fluids at the organism level.

*Theme 4:* Results have concentrated on the following:

- The *mantle structure studies* have been achieved after sampling across the MAR close to the Azores, but also elsewhere, and by studying the data collected during a cruise. A large asymmetry was confirmed in the upper mantle structure of the Azores and this is the biggest influence of the hotspot on the MAR. To the north, the plate boundary has proved to be regionally orthogonal to the spreading direction, while to the south it is regionally highly oblique (sloping?). So *non-transform discontinuities are more frequent and better developed in the south compared to the north*. Recent trace element and isotopic data analyses have shown that the depleted mantle in the region from 31°N to 41°N is not a homogeneous reservoir.
- The study of comparison of the location of hydrothermal vents on the SW arm of the ATJ with that on the northern and eastern arms have involved independent studies of the AMORES partners away from the Azores region. These studies led to

the production of a preliminary model so as to predict the location of hydrothermal vent sites. Study of the occurrence of hydrothermal venting on the MAR north of the ATJ has been impossible, but combination of knowledge of venting south of the Azores with that from other regions of the Atlantic has led to the following conclusions:

1. *At slow spreading ridges a strong heat supply alone is not enough to induce hydrothermal venting*
2. *Venting is not a simple linear function of spreading rate*
3. *The presence of extensive faulting associated with frequent and well-developed non-transform offsets plays an important role in the control of seafloor hydrothermal activity.*

### **Major Publications:**

Aballéa, M., Radford-Knoery, J., Appriou, P., Bougault, H., Charlou, J. L., Donval, J.P., Etoubleau, J., German, C., Miranda, M., 1998. Manganese distribution in the water column near the Azores Triple Junction along the Mid-Atlantic Ridge and the Azores domain. *Deep-Sea Research I*, 45, 8 (1319-1338).

Chevaldonné, P., Jollivet, D., Feldman, R. A., Desbruyères, D., Lutz, R. A., Vriejenhoek, R. C., 1998. Commensal scale-worms of the genus *Branchipolynoe* (Polychaeta: Polynoidae) at deep-sea hydrothermal vents and cold seeps. *Cahiers de Biologie Marine* 39 (3-4) : 347-350.

Colaço, A., Desbruyères, D., Comtet, T., Alayse, A. M., 1998. Ecology of the Menez Gwen hydrothermal vent field (Mid-Atlantic Ridge/Azores Triple Junction). *Cahiers de Biologie Marine* 39 (3-4) : 237-240.

Comtet, T. and Desbruyères, D., 1998. Population structure and recruitment in mytilid bivalves from the Lucky Strike and Menez Gwen hydrothermal vent fields (37°17'N and 37°50'N on the Mid-Atlantic Ridge). *Marine Ecology Progress Series* 163: 165-177.

Desbruyères, D., Biscoito, M., Caprais, J.-C., Colaço, A., Comtet, T., Crassous, P., Fouquet, Y., Khrpounoff, A., Le Bris, N., Olu, K., Riso, R., Sarradin, P.-M., Segonzac, M., Vangriesheim, A. Variations in deep-sea hydrothermal vent communities when approaching the Azores Plateau *Deep-Sea Research* (submitted July 1999).

Dixon, D. R., Dixon, L. R. J., Pond, D. W., 1998. Recent advances in our understanding of the life history of bresiliid vents shrimps on the MAR. *Cahiers de Biologie Marine* 39 (3-4) ; 383-386.

Douville, E., Charlou, E., Donval, J.P., Hureau, P., Appriou, P., 1999. Le comportement de l'arsenic (As) et de l'antimoine (Sb) dans les fluides provenant de différents systèmes hydrothermaux océaniques. *Comptes Rendus de L'Académie des sciences, C.R.A.S.*, 328 : 97-104.

Fouquet, Y., 1997. Where are the large hydrothermal sulphide deposits in the oceans? *Philosophical Transactions of the Royal Society of London* 355 : 427-441.

Geret, F., Rainglet, F., Cosson, R. P., 1998. Comparison between isolation protocols commonly used for the purification of mollusc metallothioneins. *Marine Environmental Research* 46 (1-5) : 545-550.

German C.R., L.M.Parson and the HEAT Scientific Team. Hydrothermal Exploration at the Azores Triple-Junction: Tectonic control of venting at slow-spreading ridges? *Earth Planet. Sci. Lett.* 138, 93-104, 1996.

German C.R., K.J.Richards, M.D.Rudnicki, M.M.Lam, J.L.Charlou & the FLAME Scientific Party. Topographic control of a dispersing hydrothermal plume. *Earth Planet. Sci. Lett (Express Letter)* 156, 267-273, 1998.

German C.R., K.Richards, M.Lam, A.Thurnherr, J-L.Charlou, J.Radford-Knoery, J-P.Donval, P.Jean-Baptiste, A-M.leClerc, A.Dapoigny, M.Cooper, M.Rudnicki, H.Elderfield, H.Edmonds, D.Green, A.Khipounoff, P.Crassous, A.Vangriesheim, J.D.O'Brien, J.Patching, D.Dixon & P. Herring. A Segment Scale Study of Fluxes Through the Rainbow Hydrothermal Plume, 36N Mid-Atlantic Ridge. *Deep Sea Research (to be submitted, December 1999)*.

Jean-Baptiste, P., Bougault, H., Vangriesheim, A., Charlou, J. L., Radford-Knoery, J., Fouquet, Y., Needham, D., German, C., 1998. Mantle  $^3\text{He}$  in hydrothermal vents and plume of the Lucky Strike site and associated geothermal heat flux. *Earth Planet. Sci. Lett* 157 (1-2) : 69-79.

Pond, D., Dixon, D., Sargent, J., 1997. Wax-ester reserves facilitate dispersal of hydrothermal vent shrimps. *Marine Ecology Progress Series* 146: 289-290.

Pond, D., Dixon, D. R., Bell, M. V., Fallick, A. E., Sargent, J., 1997. Occurrence of 16:2(n-4) and 18:2(n-4) fatty acids in the lipids of the hydrothermal vent shrimps *Pimicaris axiculata* and *Alvinocaris markensis*: nutritional and trophic implications. *Marine Ecology Progress Series* 156: 167-174.

Pond, D., Segonzac, M., Bell, M. V., Dixon, D. R., Fallick, A. E., Sargent, J., 1997. Lipid and lipid carbon stable isotope composition of the hydrothermal vent shrimp *Mirocaris fortunata*: evidence for nutritional dependence on photosynthetically fixed carbon. *Marine Ecology Progress Series* 257 : 221-231.

Radford-Knoery, J., Charlou, J. L., Donval, J. P., Aballea, M., Fouquet, Y., Ondréas, H., 1998. Distribution of dissolved sulphide, methane and manganese near the seafloor at the Lucky Strike and Menez Gwen hydrothermal vent sites on the Mid-Atlantic Ridge. *Deep-Sea Research I* 45 (3) : 367-386.

Sarradin, P. M. and Caprais, J. C., 1996. Analysis of dissolved gases by headspace sampling gas chromatography with column and detector switching. Preliminary results. *Analytical Communications* 33: 371-373.

Truchot, J. P., Lallier, F., 1998. High  $\text{CO}_2$  content in hydrothermal vent water at the Snake Pit area, Mid-Atlantic Ridge. *Cahiers de Biologie Marine* 39 : 153-158.

Valdés, A., Bouchet, P., 1998. Naked in toxic fluids: A nudibranch mollusc from hydrothermal vents. *Deep-Sea Research II* 45 : 319-327.



Title: **BENGAL**

High resolution temporal and spatial study of the **BEN**thic biology  
and **G**eochemistry of a north-eastern **A**tlantic abyssal **L**ocality

Contract No: **MAS3-CT95-0018**

Sub-programme area: Extreme Marine Environments – ***The deep sea floor  
in the North Atlantic and the Mediterranean***

Duration of the project: February 96' – January 99'





**Background:** The site studied was on the Porcupine Abyssal Plain, centered on 48° 50'N: 16°30'W, at a depth of about 4850m, some 270nm to the south-west of Ireland. This locality was also sampled and studied under MAST I and II contracts, in 1989, 1991 and 1994. Those previous studies established that there was significant variability in the sediment chemistry and the abundance, size structure, taxonomic make-up, and behavior of the benthic community at abyssal depths. Moreover, they demonstrated that abyssal benthic ecosystems in the northeast Atlantic were subjected to major seasonal and interannual variations in biogenic fluxes.

Billett et al. (1983) had revealed the deposition of phytodetrital material at the 2000 m deep Porcupine Seabight, south-west of Ireland, using photographic techniques. Subsequently, other observations proved that in the NE Atlantic mass fluxes of fresh organic matter towards the bottom also occur in regions hundreds of km off-shore (Thiel et al, 1990).

**Objectives:** The general objective of the BENGAL project was to identify, quantify and model the principal processes within the abyssal benthic boundary layer (BBL) which affect the permanent sedimentary record of the incoming chemical flux. The project studied how the physics, chemistry and biology of the abyssal benthic boundary layer responded to and modified the chemical input from the overlying surface layers, and how these processes might affect the palaeoceanographic record in the underlying sediment. A highly seasonal abyssal locality in the NE Atlantic, Porcupine Abyssal Plain, was studied over one complete annual cycle. The specific objectives were to:

1. Quantify and characterise the incoming vertical flux of organic and inorganic material, as well as its re-suspension in the near-bottom layers, using sediment traps, transmission-meters, and mid-water cameras
2. Measure the rate of the incorporation of the above flux into the sediment column, and the fluxes across the sediment-water interface, using *in situ* landers, sediment column chemical analysis and sediment profile imaging
3. Study the interaction between the incoming chemical input and the benthic biota, including bioturbation, using conventional sampling and analysis, and *in situ* instrumentation and experimentation
4. Use the resulting data for developing predictive models for the effect of the pulsed incoming flux on the benthic boundary layer and *vice versa*.

The above were incorporated into four work-packages (WP), including 90 Tasks:

- WP1 investigated the primary flux of particulate material to the seafloor, as well as the long-term sedimentary record at the BENGAL site
- WP2 studied the interactions between the particulate flux and the activity of the benthic community
- WP3 investigated the composition and activities of all faunal-size classes in the benthic boundary layer food-web (including microbiota, protozoan and metazoan meiofauna, macrofauna, megafauna, fish and near-bottom zooplankton, micronekton, and scavengers). It also studied the trophic relationships between the various elements of the benthic community
- WP4 aimed at integrating the resulting data from the other work-packages, through modeling and data management.

**Methodology and Achievements:** BENGAL addressed the following specific topics, according to the several work-packages:

**WP1:** One study of this work-package was concentrated on the monitoring of the quantitative and qualitative temporal and spatial variations in particle flux over a full seasonal cycle. Strong seasonal particle fluxes occurred during the summer months at the BENGAL site, as measured by sediment traps at 1000m, 3000m, and 4750m (100m above bottom). Large changes in dry mass flux were measured, particularly at 1000m. The particle flux during the summer contained considerable labile (unstable, but nutritious) organic compounds, such as sterols, fatty acids, proteins, and DNA. Radionuclide data processing indicated that, for 1997, most of the flux was vertical than having arisen from a lateral advection. Resulting data from BENGAL sediment traps, together with results from other sediment trap studies carried out in the same area since 1989, have formed a *very large and long-running data set that establishes a European long-term oceanic monitoring station*.

Another study has involved the monitoring of the hydrodynamic regime within the BBL. The response of the benthic community to the seasonal deposition of organic matter at the PAP site was examined using phytopigments as biomarkers. In 1997, surprisingly, the sediment trap flux maximum was not seen in a similar rise in phytopigments within the sediment. Time-lapse photographs of the seabed did not detect any deposition of phytodetritus either. In the past the seabed had disappeared from view under a green carpet of marine compost, but not in 1997. Long-term measurements of current speed and direction have shown that current speeds are generally low. During geochemical, radionuclide, and nephelometer studies it was proved that re-suspension occurs within 100m of seabed, but there was no proof for transport of material away from the BENGAL site. There is considerable inter-annual variation in the mass deposition of organic material. In September 1996, material was of a relatively fresh character suggesting that a major deposition event had occurred in this year. During the other seasons (March 97', Jul 97', Oct 97'), such fresh material was apparently absent, at least within the seabed. Other BENGAL data suggest that the epibenthic megafauna re-worked the fresh phytodetritus so rapidly that it did not appear in the time-lapse photographs or in the sediment. The gut contents of certain megafauna were full of phytodetritus in July 1997.

Another investigation involved measuring of mass solute fluxes across the sediment/water interface in different seasons. Measurements of fluxes across the sediment/water interface were carried out using three types of lander. Higher mass solute fluxes were expected for the summer months, but, surprisingly there was evidence of higher activity in September 1996. Spatial variability in parameters such as sediment community oxygen consumption (SCOC) was much higher than temporal variability, between September 1996 and March 1998. The very low input of fresh organic matter into the sediment in 1997, must have had an effect on SCOC. In addition, the benthic silica cycle was measured at the PAP site. The downcore profiles of porewater silicic acid and biogenic silica are low. Typical silicic acid profiles increase at the sediment/water interface, showing that dissolution in 1997 mostly occurred in the sediment. This also agrees with the low amounts of organic matter incorporated into the sediment in 1997.

**WP2:** One study studied the temporal kinetics of organic matter (OM) and the relation of OM distribution to the microbiota and meiofauna abundance and distribution. The results gave detailed chemical data for finely sliced sediment core sections that were collected with a multiple corer. This corer causes little disturbance to the critical sediment/water interface during sampling. The analyses included sterols, fatty acids,

proteins and phytopigments. The data for chlorophyll- $\alpha$  and its products showed that there was a weak pulse of fresh organic matter into the sediment in the summer of 1997, in comparison with the measurements for 1996 and 1998. That was confirmed by the finding, through the time-lapse camera observations, that little fresh detritus was incorporated into the sediment in 1997.

**WP3:** One study dealt with a quantitative analytical description of the composition and activities of all benthic faunal-size classes. The dominant lipids of seven species of abyssal holothurian (NE Atlantic Ocean) (*Oneirophanta mutabilis*, *Pseudostichopus villosus*, *Psychropotes longicauda*, *Deima validum*, *Paroriza prouhoi*, *Amperima rosea*, and *Molpadia blakei*) have been examined during this study, concerning their composition and metabolism. Fatty acid compositions are qualitatively similar to those of shallow water holothurians, but relatively higher amounts of unsaturated compounds in the deep-sea animals are found, owing to the adaptation of the latter to maintain membrane fluidity at high pressure and low temperature. Sterol distributions of abyssal holothurians were complex, with mixtures of C<sub>26</sub>-C<sub>30</sub> sterols being present in all animals. *Amperima rosea* is the only species having cholest-7-enol (the dominant product of *de novo* biosynthesis in shallow water animals) as a major component. It also, alone, appears to assimilate 4 $\alpha$ -methylsterols directly, and contains high amounts of steryl-sulphates. The rest of the animals do not appear to biosynthesise their sterols *de novo*. Certainly, inter-species differences in the free sterol distributions reflect their different feeding strategies. *Amperima rosea* increased in abundance by two orders of magnitude during the BENGAL period, compared with sampling before BENGAL. Gut content analyses showed it fed preferentially on phytodetritus. The high abundance and feeding strategy could be one reason why the high organic matter flux recorded by the sediment traps was not seen in the time-lapse photographs, or in changes in sediment chemistry or SCOC.

Another study examined the relation the activity of benthic fauna to the sedimentary record, and assessment of burial dynamics. Studies of the feeding activities of the benthic fauna, together with the organic analyses, showed that only the most recalcitrant organic compounds are buried deep in the sediment. Spatial variability of organic and inorganic carbon recycling rates in PAP sediments was larger than temporal variation (seasonal and inter-annual), during the period 1996-1998. Dissolved organic carbon (DOC) fluxes were comparable to  $\Sigma\text{CO}_2$  fluxes during all BENGAL cruises, implying that organic carbon recycling rates are underestimated if only inorganic carbon and/or oxygen fluxes are measured. Finally, on an annual basis 98% of the particulate organic carbon (POC) and 53% of the particulate inorganic carbon (PIC) deposited in these sediments were recycled to the overlying water. The method used the Göteborg lander for *in situ* sampling of benthic solute fluxes and pore water solutes, in which one of the four experimental chamber modules was exchanged with a gel-peeper module for high vertical resolution (Brünnegård, 1999, in preparation). A polyacrylamide gel was found to give the best results.

**WP4:** A time-dependent model was developed of early diagenesis in sediments. The model can be used to calculate profiles of oxygen, nitrate and organic carbon profiles. The model was developed to simulate the deposition of phytodetritus. Another achievement is a CD-ROM product including geochemical, benthic-biological, and oceanographic data collected by the BENGAL project at the Porcupine Abyssal Plain site. This product will be published as an ISBN registered publication.

## **Major Publications:**

Billett, D.S.M., Lampitt, R.S., Rice, A.L. & Mantoura, R.F.C. 1983. Seasonal sedimentation of phytodetritus to the deep-sea benthos. *Nature, London*, **302**: 520-522.

Rabouille C., Witbaard R., Duineveld G., Wolff G., MacKenzie K. and Khrpounoff A., Temporal variability of sedimentary recycling in the North Atlantic Ocean (BENGAL site) studied with a non-steady-state model. *Progress in Oceanography* (in preparation)

Rice, A. L., BENGAL gets off to a flying start. *Deep-Sea Newsletter* No. 25, 9-10.

Thiel, H., Pfannkuche, O., Schreiver, G., Lochte, K., Gooday, A.J., Hemleben, C., Mantoura, R.F.C., Turley, C.M., Patching, J.P. & Riemann, F. 1990. Phytodetritus on the deep-sea floor in a central oceanic region of the Northeast Atlantic. *Biological Oceanography*, **6**, 203-239

Witbaard R, Duineveld G.C.A., van der Weele J., Berghuis E.M., Reyss J.P., The Benthic response to the seasonal deposition of phyto pigments at the Porcupine Abyssal Plain in the N-E Atlantic.

Ragneneau O., Gallinari M., Hall P., Grandel S., Stahl H., Rickert D., Lampiti R., Witbaard R., Tengberg A., Gooday, A., The benthic silica cycle at the Porcupine Abyssal Plain, N-E Atlantic (submitted to *Deep Sea Research*, in March 99')

*Progress in Oceanography* (in preparation) – BENGAL Special Volume:

Material supply to the abyssal seafloor in the northeast Atlantic R.Lampitt, K.Kiriakoulakis, O.Ragueneau, A. Vangriesheim & G.Wolff

Controls on the organic chemical composition of settling particles in the north-east Atlantic Ocean. K. Kiriakoulakis, E. Stutt, S. Rowland, A. Vangriesheim, R. Lampitt & G. Wolff.

Fluxes of labile organic matter in the NE Atlantic.

M. Fabiano, M. Armeni, R. Danovaro, A. Dell'Anno, A. Pusceddu, & G. Wolff.

Microbial assemblages associated to sinking particles in the Porcupine Abyssal Plain (NE Atlantic). S. Vanucci, R. Danovaro, A. Dell'Anno, M. Fabiano, R. Lampitt & A. Pusceddu.

Aspects of near bottom dynamics and particle fluxes on the PAP site. A. Vangriesheim, B. Springer & R. Witbaard.

Annual and interannual variability of sedimentary recycling in the North Atlantic Ocean (BENGAL site) studied with a non-steady-state model C. Rabouille, R. Witbaard & G. Duineveld.

A lander adapted gel peeper module for the in-situ extraction of sediment pore waters at high vertical resolution using diffusive equilibration. J. Brunnegard, H. Stahl, A. Tengberg, P. Hall, A. Vershinin, S. Hulth & C. Rabouille.

The benthic silica cycle in the North-East Atlantic O. Ragueneau, M. Gallinari, H. Stahl, S. Grandel, A. Tengberg, D. Rickert, R. Witbaard, P. Hall & R. Lampitt.

Imbalance in the carbonate budget of surficial sediments in the North Atlantic Ocean: variations over the last millennium. C. Rabouille, H. Stahl, F. Bassinot, A. Tengberg, J. Brunnegard, P. Hall, K. Kiriakoulakis, J-L. Reyss, L. Dezileau, P. Crassous, P. Roos & R. Lampitt

Biogeochemical processes at the sediment-water interface in a North Eastern Abyssal Locality (BENGAL – Area). S. Varnavas & D. Panagiotaras.

Bacterial Abundance and Activity in deep sea sediments from the Eastern North Atlantic. D. Eardly, M. Carton, J. Gallagher & J. Patching.

*Stainforthia fusiformis* (Williamson) (Protista, Foraminiferida) at the Porcupine Abyssal Plain: a remarkable eurybathyal opportunist. A. Gooday & E. Alve.

Evidence for episodic recruitment in a small opheliid polychaete species from the abyssal NE Atlantic A. Vanreusel, N. Cosson-Sarradin, A. Gooday, G. Paterson, J. Galeron, M. Sibuet & M. Vincx

Are temporal patterns among meiofaunal and macrofaunal taxa (quantitative distribution) at an abyssal NE Atlantic site influenced by sediment lipids chemistry ? J. Galeron, J. Sibuet, M. Vanreusel, A. Mackenzie, K. Gooday, A.J. & Wolff, G.

Impacts of trophic disturbance on benthic fauna at the abyssal NE Atlantic sites (BENGAL and EUMELI): a comparative experimental approach. N. Cosson-Sarradin, M. Sibuet, D. Desbruyères, A. Khrpounoff & P. Crassous.

Temporal and spatial variability in deep-sea megafauna. D. Billett, B. Bett, A. Rice, M. Thurston, M. Sibuet, J. Galeron & G. Wolff.

Lipids in selected abyssal benthopelagic animals: links to the epipelagic? S. Buhring & B. Christiansen.

Temporal variability of megafauna from time lapse photography. B. Bett & B. Wigham.

Food web structure of the benthic community at Porcupine Abyssal Plain (NE Atlantic): a stable isotope analysis. K. Iken, T. Brey, U. Wand & P. Junghans.

What controls the appearance of phytodetrital layers on the abyssal seabed? R. Lampitt, B. Bett & A. Vangriesheim.

Steroid supplies may limit the distribution of large benthic organisms in the deep-sea. M. Ginger, K. Mackenzie, K. Kiriakoulakis, D. Boardman, D. Billett & G. Wolff.

Response of *Oneirophanta mutabilis* (Holothuroidea) to the seasonal deposition of phytopigments at the Porcupine Abyssal Plain in the NE Atlantic. R. Witbaard, G. Duineveld, A. Kok, J. van der Weele & E. Berghuis.

Sediment distribution, enzyme profiles and bacterial activities in the guts of *Oneirophanta mutabilis*, *Psychropotes longicauda* and *Pseudostichopus* sp. - What do they tell us about digestive strategies of abyssal holothurians? D. Roberts, H. Moore, J. Berges, J. Patching, M. Carton & D. Eardley.

Simulation of the impact of a temporal detrital flux on the entire sediment community of an abyssal NE Atlantic site M. Mahaut, D. Billett, J. Galéron, P. Lamont, J. Patching, M. Sibuet & A. Vanreusel.



Title: **ESOP-2**

**European Subpolar Ocean Programme Phase 2: The Thermohaline Circulation in the Greenland Sea**

Contract No: **MAS3-CT95-0015**

Sub-programme area: Extreme Marine Environments – ***The ice-covered seas in the northern hemisphere***

Duration of the project: January 96' – February 99'





**Background:** Nowadays, there is a considerable need for better understanding of the operation of the oceanic circulation of the high latitude regions. These areas are key areas for the oceanic carbon sink and the transfer of heat from low to high latitudes. They also determine the state of the deep sea and the flushing of the world ocean basins. The ESOP-2 project has been a continuation of the *ESOP-1* (MAS2-CT93-0057, Jul 93'- Dec 96'), which in turn was the continuation of the *Greenland Sea Project*. ESOP-1 has made basin-wide studies, so as to determine the large scale role of the ice cover, and meso-scale studies (in the region of the Odden ice tongue in the mid-latitude Greenland Sea), so as to determine the mechanisms by which sea ice is involved in deep convection in winter, and to estimate CO<sub>2</sub> uptake through cooling, convection and biological fixation.

Results from the above projects on the large scale, showed that most of the sea ice in the Greenland Sea enters from the Arctic Basin via Fram Strait, and contributes a fresh water flux, at a range of latitudes, to the surface waters. It also affects ocean-atmosphere energy exchanges. Some sea ice also forms in winter within the Greenland Sea, and the salt flux associated with rapid ice growth is thought to be a critical factor in the onset of deep convection. Deep convection is a vitally important process from a climatic point of view since it leads to carbon dioxide transport into the deep ocean, and to renewal of the deep waters. The region where this interaction occurs has been identified as the Odden ice tongue in the mid-latitude Greenland Sea. In addition, investigations during the Greenland Sea project and ESOP-1 indicated that the initial convection is haline and caused by brine release, but it almost always results in the melting and/or removal of the ice-cover; the final stage of the convection is likely to be thermal.

**Objectives:** The overall objective of the project is to understand the *thermohaline circulation in the Greenland Sea*, its sensitivity and impact on global ocean circulation, using experience gained from ESOP-1 project. In particular, ESOP-2 used intentional release of an anthropogenic tracer (SF<sub>6</sub>), isobaric floats, the observation of standard hydrographic parameters and transient tracers partly from a permanently moored and internally recording CTD, a complete coverage of the carbon cycle, remotely sensed and direct measurements of sea ice fluxes, and an integrated small, meso and large scale modeling. Finally, the combined field and modeling techniques aimed at the understanding of the role of the Greenland Sea in the "Global Conveyor Belt", and to make realistic predictions of climate change.

The specific objectives are. a. to understand the oceanic thermohaline circulation on meso and large scales, and its dependence on surface fluxes of heat, fresh water and momentum, b. to study fresh water fluxes into and out of the central Greenland Sea region, c. to evaluate the role of convection areas in ocean/atmosphere gas exchange, and investigate the role of thermohaline circulation in CO<sub>2</sub> transport, d. to use observations so as to improve and calibrate coupled atmosphere/ocean models. The project has been divided into four workpackages: WP1 involved the tracer experiment and ocean physics, WP2 the ice-ocean interaction, WP3 was devoted to carbon cycle, and WP4 to modeling.

**Methodology and Achievements:** ESOP-2 has resulted in a number of findings, the most important of which the *evidence for climate processes* and the *understanding of circulation of the Greenland Gyre*. At least one unknown mechanism has been detected in addition to deep-water formation from processes earlier proposed. During five cruises, the occurrence of the tracer in the bottom water was studied. In the winter, deep-water forming processes transported water masses labelled with the tracer from 300m to more than 3km depth, causing storms on the seabed. In addition

to the new mechanism of deep-water formation (large-scale movement of water), several local chimney effects have been observed. They are local events of water mixing (thin currents), which take water from the surface straight down to the bottom. One chimney, detected in May 1997, takes surface water down to a depth of 2km. Concerning the circulation of the Greenland Sea Gyre, it was found that the North Atlantic Current transports Atlantic water into the West Spitsbergen Current, further north. Near Fram Strait, it splits into three major branches. The Return Atlantic Water, the western one of the three, forms the eastern side of the East Greenland Current (EGC). Polar Surface Water leaves the Arctic Ocean through Fram Strait, and flows with the EGC along the western periphery of the Greenland Basin. Finally, at about 73° N, the Jan Mayen Current splits eastward from the EGC, carrying both Return Atlantic Water and Polar Surface Water. More specific results of ESOP-2 concentrate on the following:

*Long term trends and interannual variability:* The data analysed in this project are for a circular area from 73.5° to 76.5° N and 9° W to 3° E. This area is sub-divided into smaller circles and rings from the centre of Greenland Sea, each one analysed separately. The central part of the analysed area was examined for temperature and salinity evolution, for the period 1986-1998. Concerning temperature, in the vertical, two regimes were distinguished. Below about 2000m depth, the deep water indicated a general warming which was constant over the period of observation. The average increase was 0.13° C. The layer above 2000m was also under warming, though frequently interrupted by convective events that led to a cooling of the water column. The deepest of those events occurred in the winters of 1987/88 and 1988/89, when mixing was observed to 1950m.

As for salinity evolution, measurements during the late 1980s and during the ESOP-1 and ESOP-2 periods showed that the top 300m of the water column undergo a strong seasonal variability. Overall, there is a deepening of this low salinity layer, from about 150m in the late 1980s to more than 350m at the end of the period of observation. The water column can be divided into three basic layers, which display a different temporal behavior. The top 250m undergo strong seasonal variability, and heating during the summer reaches this depth. Vertical exchanges and complete mixing occur in this top layer almost every year. The layer underneath was sporadically affected by intermediate depth convection during the winter in 1993, 1994, 1996 and 1997. Below 2000m depth, there was no short-term variability, but rather a smooth sinking of the isotherms after 1989. Temperature increase in this deepest part of the water column has been almost constant over time, and was not affected by the smaller convective events of 93/94 and 96/97. The upper boundary of this layer coincides approximately with the depth for which the Greenland Sea Gyre is laterally separated from the adjacent basins.

*Tracer release and spreading:* The tracer release experiment has been the most important result of ESOP-2, an innovative technique (one of the three experiments of this kind, executed in the open ocean) to measure general circulation of the ocean, and to validate ocean modeling in recent years. The tracer, sulfide hexafluoride (SF<sub>6</sub>), is a non-toxic, very inert anthropogenic compound, used in industrial applications since the beginning of the century. It is not abundant in the environment, and destroyed by UV radiation in the upper atmosphere. It is considered as a "marine tracer" because it can be reliably, quickly, and relatively cheaply measured in sea water, with a detection limit of  $< 5 \times 10^{-17}$  mol / kg. The experiment started in August 1996, with the release of 320kg of SF<sub>6</sub> in the centre of Greenland Sea Gyre, on an

isopycnal surface situated at about 300m depth. The tracer was released in streaks, in a rectangle.

The mean vertical profiles in and around the gyre were obtained, as well as the horizontal spread of the tracer, so as to understand vertical mixing processes and estimate large-scale circulation in the Arctic, Nordic Seas, and the northernmost North Atlantic, respectively. Concerning the former, the rates for turbulent mixing are about 100 times higher than those found in the thermocline of the sub-tropical Atlantic. Diapycnal mixing in the Greenland Sea is clearly very rapid compared to typical open ocean values. The amount of water involved in convection, was more precisely defined as smaller for 1997 than that estimated so far, despite the fact of the convective event. That was due to homogenisation of the water column down to the depth of convection. Results for the horizontal spread of the tracer, showed that transmission of the tracer out of the Greenland Sea Gyre has been slow. 25 months after the release, about 1%  $\text{SF}_6$  had reached Fram Strait and was heading north into the Arctic Ocean. Several percent was found in the lower East Greenland Current as far south as the sill of Denmark Strait, which shows considerable though not very large exchange between the EGC and Greenland Sea. About 6% was found in the Norwegian Sea. Overall, about 70% of the tracer remained in the Greenland Sea domain.

*Mesoscale circulation:* During ESOP-2, a large-scale Lagrangian float experiment has been run. Its most important result was that after having been released for periods of 300 days near the centre of the Greenland Gyre, all the floats from the first releases came inside a circle within a 100 nautical miles radius, centered over the deepest part of the Greenland Basin. This indicated that circulation inside the gyre remains quite tight (fixed) for the whole year, without any breaking up during spring or summer time. The above result was confirmed during the second year floats experiment, with 10 out of 12 floats remaining in the Greenland Gyre after one year. This also corresponds to the fact that most of the tracer remained inside the gyre one year later.

Another result came from calculation of float trajectories, and concerns the documentation of an extremely active meso-scale eddy field embedded in a deep mixed patch resulting from deep convection located above the deepest part (> 3500m) of the Greenland Basin. It was mainly in and around this mixed patch that many floats were trapped for about one year. The final result arising from the floats experiment is the documentation of long-lived anticyclonic eddies (>1 year). They consist of a central core set in a solid-body rotation for approximately two-days period, limited to a radius of 5km from the eddy center of rotation and characterized by a strong negative relative vorticity (vortical movement).

*Sea ice convection:* The first systematic study on ice kinematics in the Greenland Sea region was done during the ESOP-1 project and covered the years 1993 and 1994; a high variability of the sea ice flux through Fram Strait was measured. Total mean ice transport for these two years was estimated at 1700km<sup>3</sup>/year. There is a discrepancy between this estimation and other estimates based on numerical modeling, and it is due to the uncertainties in knowledge of ice thickness distribution. A dynamic-thermodynamic ice model was used to expand the time series of sea ice flux further to the past. Results showed a high interannual and seasonal variability. The average seasonal cycle indicated that the highest fluxes are reached in the winter months with maximum flow in December, and minimum in the summer months (August).

The relation of sea ice dynamics and the salinity of the underlying ocean was also investigated. Ice dynamics were studied by releasing ice drifters, brine drainage rates by direct sampling of frazil and pancake ice, and ice type distribution by shipborne validation of satellite remote sensing. In addition, a salt flux model was generated, so as to estimate the resultant salt fluxes from the above observations. Salt flux was measured for the winter 96'-97'; the flux was strongly negative in the region that faces the southern edge of Odden, and strongly positive along the inner edge of Odden and in Nordbukta. After a number of years without an Odden sea ice feature in the Greenland Sea, it can be estimated that ice transport from the Arctic Ocean through Fram Strait is negatively correlated with the size and duration of the Odden sea ice feature.

*Carbon cycle and carbon fluxes:* During the project, it has been shown that the Greenland Sea is a year round sink of CO<sub>2</sub>, but with strong interannual and seasonal contrasts. The uptake of atmospheric CO<sub>2</sub> is positive during all year, in the order of 69±16gC/m<sup>2</sup>y or 20±4×10<sup>12</sup>gC/y, if integrated over the ice-free part of the Greenland Sea. Anthropogenic carbon is removed by deep-water production (below 1500m) in a rate 10 times less than the air/sea flux. As 2/3 of the production of intermediate waters (200-1500m) probably comes from convection in the Greenland and Iceland Sea, anthropogenic carbon removal by deep-water production increases by a factor of 10, therefore is equal to the atmospheric uptake of carbon.

The uptake of carbon in the Greenland Sea is controlled equally by biological primary production and heat loss. The combined effect of the uptake by these processes is that surface water leaving Greenland Sea is of higher concentration of total dissolved inorganic carbon than the inflowing, warmer surface water. A significant fraction of the outflowing surface water flows to the south into the Iceland Sea, where it may mix down into deep water that flows into the deep North Atlantic.

**Dissemination of results:** Two ESOP-2 CD-ROMs are available: a. one prepared at ICES, containing cruise specific data, and b. one produced at the Danish Meteorological Institute, containing model outputs from the ESOP-2 models.

### **Major Publications:**

Anderson, L., Olsson, K., Chierici, M., 1998. A Carbon budget for the Atlantic Ocean, *Global Biogeochemical Cycles*, 12, 455-465.

Anderson, L. G., Chierici, M., Fransson, A., Olsson, K., Jones, E. P., 1998. Anthropogenic carbon dioxide in the Arctic Ocean: Inventory and sinks. *J. Geophys. Res.*, 103, 27, 707-27, 716.

Backhaus, J. O., Harms, I. H., Kämpf, J., Rubino, A., 1997. Formation of Dense Bottom Water in Arctic Shelf Seas. *ICES Journal of Marine Science*, 23 pp.

Backhaus, J. O., Fohrmann, H., Kämpf, J., Rubino, A., 1997. Formation and export of water masses produced in Arctic shelf polynyas - process studies of oceanic convection. *ICES Journal of Marine Science*, 54, 366-382.

Kämpf, J., Backhaus, J. O., 1997. Shallow, brine driven free convection in polar oceans: Nonhydrostatic numerical process studies. *J. Geophys. Res.*, 103, 5577-5593.

Lherminier, P., Gascard, J. C., 1998. Drifting isobaric float response to deep convective activity in the Greenland Sea. C. R. Acad. Sci. Paris, Earth and Planetary Sciences, 326, 341-346.

Vinje, T., Nordlund, N., Kvambekk, A., 1998. Monitoring ice thickness in Fram Strait. Journal of Geophys. Res. Vol 103, 10,437-10,449.



Title: **AG-HY-FL**

Hydrothermal fluxes and biological production in the Aegean

Contract No: **MAS3-CT95-0021**

Sub-programme area: Marine Systems Research - ***Sedimentary processes  
in the deep sea, on the continental slope and on the shelf edge***

Duration of the project: February 96' - January 99'





**Background:** Major hydrothermal systems have been found along the Hellenic Volcanic Arc at Methana, Sousaki, Milos, Santorini, Kos, Yaki and Nisiros. Geothermal areas have also been found in many coastal regions and islands around the edge of the Aegean Sea, including the regions around Antemus, Volvi-Langada, Strymon, Nestos-Xanthi and Alexandroupoli basins, in the north, Lesbos in the east, and the Gulf of Maliachos and Sperchios basins in the west. The areas studied in this project were mainly Milos, but also Methana, Santorini, Kos, and Lesbos, which are shown in *Figure 1*. Previous studies in those areas have recently been reviewed by Dando et al. (1999, in press).

The project examined the biogeochemical processes and fluxes in relation to hydrothermalism in different time and space scales, in the back arc area of the Aegean Sea, the Sea of Crete. Hydrothermal areas around Milos were examined as part of previous EC-funded, MAST-1 and MAST-2 projects that looked at areas of high sulfide and methane concentrations and at interfaces between oxic and anoxic conditions. These studies showed that 35 km<sup>2</sup> of seabed around Milos were hydrothermally active. Areas of seabed were influenced by hydrothermal brine with a composition similar to that in the reservoir 1 km below. Free CO<sub>2</sub> flux was estimated to be 1-4% of the global mid-ocean ridge flux. Layers of alga-mineral-bacterial mats, containing sulphur-oxidising bacteria, up to 100m across, overlay the seeps. It was also found that bacterial H<sub>2</sub>S oxidation lowered the pH to 2 and released high concentrations of phosphate. Additional phosphate, ammonia, silicate, and other nutrients were released by the venting of hydrothermal fluids and gases. Finally, episodic events were verified, including earthquakes, white smoker activity, and release of brines, resulting in a bottom-held (hugging) submarine "fog".

**Objectives:** The overall objective was the assessment of the effect of hydrothermalism on the chemistry, production and biodiversity of the oligotrophic Sea of Crete. More specifically, the project involved four objectives:

1. Determination of the extent of submarine venting, measurement of bacterial production and export, quantification of geochemical fluxes and their modification by bacterial growth. Finally, establishment of the periodicity of venting, the role of episodic events due to storms, and seismicity.
2. Estimation of the effects of naturally occurring excess nutrients and CO<sub>2</sub> on the photosynthetic rate of phytoplankton, benthic diatoms and cyanobacteria, and on chemosynthesis by sulphur-oxidising bacteria.
3. Comparison of the diversity of bacteria and epifauna in venting and non-venting areas, estimation of the importance of chemosynthesis in comparison with photosynthesis in the food-web.
4. Isolation, identification, and characterization of bacteria from vent sites (hyperthermophiles and halophiles) with bio-product potential.

**Methodology and Achievements:** The major achievements of the project are the following, according to the 8 main tasks into which the project was divided:

*Task 1* has involved fieldwork organisation and logistics. There were four major field trips to the submarine hydrothermal field offshore Milos, on June 96', September 96', June 97', and June 98'. There was also a field trip to Kos, in September 97'. Additionally, several cruises have taken place. During the *Meteor* cruise of December 97' to Milos, Santorini, and the western part of Kos, important achievements were: a. two hot GKG cores obtained from 110m depth, one with gas bubbling that lasted for

24h after retrieval, b. a hot multicore from Santorini Caldera and the discovery of high methane concentrations outside the Caldera, c. the discovery of a new hydrothermal field south of Kos and the detailed mapping of Milos vents and hydrothermal plumes.

*Task 2* involved the study of seismic records (microseismicity) and of the relation of the records with venting periodicity. A seismic network of five three-component, triggering mode digital stations, was installed on Milos Island in the summer 96', and recorded 400 local micro-earthquakes and 500 signals. The analysis indicated low seismicity, probably associated with hydrothermal activity. The sources producing the recorded micro-earthquakes were typical sources of volcanic activity, most probably related to the existing volcanoes of Fyriplaka and Trachilas. The longer period signals may be related to venting procedures, and the micro-earthquake trains may originate from very small sources which act as microfractures in a self-organised critical system. This process possibly connects to changes in pore pressure. A data base of seismic records has been established.

*Task 3* examined the influence of water pressure changes on venting activities, the measurement of water currents and levels in the area near the vents, and aimed at explaining, by a mathematical model, the water dynamics in Palaeochori Bay, Milos. Gaso-hydrothermal vent positions were located by echo-sounders and accurately mapped using DGPS, allowing detailed vent maps to be produced for the Milos hydrothermal fields (*Figure*). A schematic geological section through Palaeochori Bay, from the shore southwards, is presented in *Figure*. The basement consists of a fractured metamorphic rock with block movements along the faults. It is overlain by marine and volcanoclastic sediments of various thickness, consisting predominantly of sand with variable mud and gravel components. Permeability is locally highly variable; it ranges from the nearly impermeable, due to the formation of hydrothermal precipitates, to the highly permeable. Three fluid sources could be distinguished by their acidity, chlorinity and enrichment of solute contents, compared to ambient sea water. a. a magmatic source indicated by the presence of a mantle-derived He component, b. fluids showing seawater-rock interactions in the deeper crust, c. fluids displaying fluid-sediment-bacterial interactions in the upper few dm of the sediment.

*Task 4* had the objective to determine chemical and bacterial fluxes from hydrothermal fields into the Aegean Sea. Around Milos Island, diffuse but widespread hydrothermal venting in shallow water (intertidal-150m) occurs. Despite its diffuse nature, hydrothermal activity leads to slight but measurable changes in the water column. Vent water composition varied a lot, due to mixing of hydrothermal reservoir fluids, vapor condensate and seawater, altered by interactions of fluid-sediment-bacteria in different proportions, in the gasohydrothermal vents. Compositions ranged from nearly seawater with only slightly reduced pH, to higher or lower salinity fluids with a pH as low as 3 and with large enrichments in heavy and trace metals. Phase separation was a common feature at these shallow vents. The dry gas phase was mainly CO<sub>2</sub>, with significant amounts of H<sub>2</sub>S, CH<sub>4</sub> and H<sub>2</sub>. *In-situ* measurements of temperature, pH, red-ox potential, H<sub>2</sub>S and O<sub>2</sub> confirmed that the shallow water vent sites were extreme environments with low pH, high sulfide content, and high temperature. Repeated measurements during three fieldtrips to the same sites have demonstrated the stability of some of the vents over a two years period.

Gas composition, as well as flow rate, varied both between sites and between vents. The collected gases contained up to 95% CO<sub>2</sub>, 9% H<sub>2</sub>, 3% H<sub>2</sub>S, and 1.8% CH<sub>4</sub>. Calculations of the number of vents per area, the average fluids flow rate, and their free and dissolved gas composition can lead to an estimation of the global

importance of the Aegean vents in geochemical budgets. Results of the project show that hydrothermal gas fluxes coming from the Aegean vents play a very important role, considering that a vent may consist of thousands of single outlets clustered together and that there are tens of vent fields off Milos and more in the rest of the Aegean Arc.

Higher values of particulate material as well as phytoplankton and zooplankton, were found during bottom water analyses, in June 97' than in September 96'. Bottom water samples were enriched in particulate organic carbon and nitrogen, total particulates and chlorophyll southwards from the active vent areas, and these components may be exported from the Bay in the observed bottom water current. The suspended particulate material (SPM) from venting waters, collected by SCUBA divers, was analysed for Fe, Mn, Cu, Pb, Al, Ba, Sr, Ca, Si and Li. Sediment trap moorings located along a transect from the active vent zone to an area free of any major vent influence showed during a 3-month summer period that mass fluxes were on average 7 to 9 times higher at the vent site than at the other stations and had a completely different pattern over time. At the vent zone, fluxes reached a maximum of  $5.9 \text{ g m}^{-2} \text{ d}^{-1}$  compared to  $1.0$  and  $1.1 \text{ g m}^{-2} \text{ d}^{-1}$  at the other sites. Organic C and N fluxes showed similar trends with two periods of pulsed sedimentation of organic material observed only at the vent site (POC flux up to  $1.4 \text{ g m}^{-2} \text{ d}^{-1}$  at the vent against  $0.1 \text{ g POC m}^{-2} \text{ d}^{-1}$  at the other stations). Particles collected at all stations were largely of pelagic origin and not directly related to phytoplankton production; large brown fecal material was important at the vent site in September. The presence of hyperthermophilic Archaea and the chemical signature of the sedimenting particles suggest that the high organic fluxes at the vent area are associated to the hydrothermal activity.

*Task 5* aimed at measurements of the fluid, heat, chemical and bacterial fluxes from individual hydrothermal vent sites, and the determination of their periodicity. A lander system for *in situ* long-term measuring and sampling of diffuse submarine hydrothermal vents at shallow water depths (200-300m) has been developed and tested. It consists of a long-time measuring cell – IPGVENT (Institute for Petrography and Geochemistry VENT system), a long-term sampling system LOVESA (Long-term Vent Sampler), an ultra low flow meter, an energy pack, and a funnel for collecting hydrothermal fluids. Thermo-chains were also set up with the lander, to measure the heat flux in a wider area. The lander was be used as an underwater short- or long-term measuring unit, giving data and water samples for a few days or for up to half a year. The data measured were pH, conductivity, temperature,  $\text{O}_2$  and  $\text{H}_2\text{S}$  contents, and water samples are of 100ml volume and are tested for major and trace element analysis.

*Task 6* was devoted to the study of biodiversity and with the culture of organisms with bio-product potential, by comparing the diversity of organisms at vent and non-vent sites, and explaining the differences in terms of physical and chemical environment and the food-web. The diversity of the microbial species at the vent sites studied was large, and a high percentage of new taxa were found. The high biodiversity of Bacteria, Archaea and epifaunal species surrounding the vents, can be explained by the complex conditions in the region of the vents. Some of the new taxa of microbes found are expected to be of biotechnological potential. Epifaunal diversity was unexpectedly high (over 200 species recorded at the shallower Milos vents), although no vent-specific species were found. Hydrothermalism was shown to increase species richness by creating greater spatial heterogeneity, by operating as an

"intermediate disturbance" (mortality due to toxic fluids) and by inducing higher recruitment. Other effects of vent activity on epibenthic communities include the reduction of biological cover (only at a very local scale), modified trophic condition for suspension-feeders and enhanced deposition of carbonates by bioconstructional organisms. The discovery of several exotic warm water species around the vents suggests that the biota at vent sites in the Mediterranean can be used to predict changes in the ecosystem, which may occur as a result of rising sea temperatures.

*Task 7* had the overall objective to determine the effects of hydrothermalism on benthic and pelagic primary production. The water column was always dominated by respiring rather than photosynthesising organisms. No significant difference was found between plankton production at venting and non-venting sites, despite a large range of inorganic carbon concentration. Small changes in ambient temperature had, on the contrary, a considerable effect on plankton respiration and a less effect on plankton photosynthesis. Photosynthetic rates in the water column did not appear to be enhanced by the venting. However, in two vent sites, high such rates were measured on bottom water samples and may be due to re-suspension of an active benthic algal mat. Much of the photosynthetic productivity, at least in the shallow water sites south of Milos, may be due to the sea-grass meadows, and carbon from this source may be a major input to the heterotrophic bacteria surrounding the vents. Particulate organic matter fluxes to the bottom were exceptionally high for short periods of 10-20 days in the vent area compared to the control area, due to high pelagic activity and the fallout of fecal faecal pellets. The sedimenting material had a vent signature ( presence of thermophilic Archaea, stable isotope composition). However, the contribution of bacterial productivity at the vents to zooplankton production is currently unclear.

*Task 8* involved the data management of the project. All the data coming from the different partners are collected from Consiglio Nazionale delle Ricerche, Istituto per lo Studio della Dinamica delle Grandi Masse Stazione Oceanografica, and are stored, creating inventories. Finally, *technology implementation plans* have been set up by GEOMAR (DE) and University of Kiel, for the exploitation of the following that resulted through the project. a. A low-velocity thermistor flow meter, b. a submarine gas flow meter, c. cultures of 53 strains of *hyper-thermophilic Archaea* isolated from the hydrothermal sites at a temperature of 90 °C.

### **Major Publications:**

Aliani, S., Amici, L., Dando, P. R., & Meloni, R. Time series of water pressure and bottom temperature in a marine shallow water hydrothermal vent off Milos Island (Aegean Volcanic Arc): preliminary results. *Rapports et Procès Verbaux de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée* **35**: 46-47

Aliani, S., Bianchi, C. N., Cocito, S., Dando, P. R., Meloni, R., Morri, C., Niemeyer, A., Peirano, A., & Ziebis, W. 1998. A map of seagrass meadows in Palaeochori Bay (Milos Island, Greece), a marine area with hydrothermal activity. *Rapports et Procès Verbaux de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée* **35**: 512-513

Bianchi, C. N., Aliani, S., Balduzzi, A., Boyer, M., Cocito, S., De Biasi, A. M., Ferdeghini, F., Morri, C., Pansini, M., Peirano, A. & Pestarino, M. 1999. Biodiversità dell'epifauna marina sessile in un'isola del Mar Egeo: Milos, Cicladi. *Biologia Marina Mediterranea* **6**: 179-183.

Böttcher, M. E., Sievert, S. M. & Küver, J. 1999. Sulfur isotope fractionation during dissimilatory reduction of sulfate by a thermophilic gram-negative bacterium at 60°C *Archives of Microbiology* **172**: 125-128.

Brinkhoff, T., Sievert, S. M., Kuever, J. & Muyzer, G. Distribution and diversity of sulfur-oxidizing *Thiomicrospira* spp. at a shallow-water hydrothermal vent in the Aegean Sea (Milos, Greece). *Applied and Environmental Microbiology* **65**:3843-3849.

Dando, P. R., Thomm, M., Arab, H., Brehmer, M., Hooper, L. E., Jochimsen, B., Schlesner, H., Stöhr, R., Miquel, J.-C. & Fowler, S. W. 1998. Microbiology of shallow hydrothermal sites off Palaeochori Bay, Milos (Hellenic Volcanic Arc). *Cahiers de Biologie Marine* **39**: 369-372

Jochimsen, B., Peinemann-Simon, S., Völker, H., Stüben, D., Botz, R., Stoffers, P., Dando, P. R. & Thomm, M. 1997. *Stetteria hydrogenophila*, gen. nov. and sp. nov., a novel mixotrophic sulfur-dependent crenarchaeote isolated from Milos, Greece. *Extremophiles* **1** (2): 67-73

Miquel, J. C., Fowler, S. W., La Rosa, J., Aliani, S. & Meloni, R. Particulate and organic fluxes in a coastal hydrothermal area off Milos, Aegean Sea. 1998. *Rapports et Procès Verbaux de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée* **35**: 276-277

Morri, C., Bianchi, C. N. Hydroids (Cnidaria: Hydrozoa) from the Aegean Sea, mostly epiphytic on algae. *Cahiers de Biologie Marine* **40**: 283-291.

Robinson, C., Ziebis, W., Müller, S., Eichstaedt, K., Dando, P., Linke, P., Varnavas, S., Megalovasilis, P. & Panagiotaras, D. 1997. *In situ* investigations of shallow water hydrothermal vent systems, Palaeochori Bay, Milos, Aegean Sea. *Proceedings of the Fourth Underwater Science Symposium of the Society for Underwater Technology, Newcastle upon Tyne* pp. 85-100

Sievert S. M., Brinkhoff, T., Muyzer, G., Ziebis, W. and Kuever, J.. 1999. Spatial heterogeneity of bacterial populations along an environmental gradient at a shallow submarine hydrothermal vent near Milos island (Greece). *Applied and Environmental Microbiology* **65**: 3834-3842.

Stueben, D., Koelbl, R., Haushahn, P. & Schaupp, P. 1998. Measuring and sampling diffuse submarine hydrothermal vents. *International Ocean Systems Design*, **2** (2): 6-12

Aliani, S. & Bergamasco, A. Coastal current variability in a hydrothermal vents area off Milos Island (Aegean Sea) *Atti XXI Congresso dell'Associazione Italiana di Oceanologia e Limnologia*, Ancona, 27-30 Settembre 1998

Cocito, S., Bianchi, C. N., Morri, C. & Peirano, A. First survey of sessile communities on subtidal rocks in an area with hydrothermal vents: Milos Island, Aegean Sea. *Hydrobiologia*

Dando, P. R., Aliani, S., Arab, H., Bianchi, C. N., Brehmer, M., Cocito, S., Fowler, S. W., Gundersen, J., Hooper, L. E., Kölbl, R., Kuever, J., Linke, P., Makropoulos, K. C., Meloni, R., Miquel, J.-C., Morri, C., Müller, S., Robinson, C., Schlesner, H., Sievert, S., Stöhr, R., Stüben, D., Thomm, M., Varnavas, S. P. & Ziebis, W. Hydrothermal studies in the Aegean Sea. *Physics and Chemistry of the Earth*

Dando, P. R., Stüben, D. & Varnavas, S. P. Hydrothermalism in the Mediterranean Sea. *Progress in Oceanography*

Morri, C., Bianchi, C. N., Cocito, S., Peirano, A., De Biasi, A. M., Aliani, S., Pansini, M., Boyer, M., Ferdeghini, F., Pestarino, M. & Dando, P. Biodiversity of marine sessile epifauna in an Aegean island: Milos, Eastern Mediterranean Sea. *Marine Biology*

Sartoni, G. & De Biasi, A. M. A survey of the marine algae of Milos Island, Greece. *Cryptogamie et Algologie*

Sievert S. M., Heidorn, T., Kuever, J., 1999. *Halothiobacillus kellyi* sp. nov., a mesophilic obligately chemolithoautotrophic sulfur-oxidizing bacterium isolated from a shallow-water hydrothermal vent in the Aegean Sea and emended description of the genus *Halothiobacillus*. *International Journal of Systematic Bacteriology*

Varnavas, S. P., Panagiotaras, D., Megalovasilis, P., Dando, P., Aliani, S. & Meloni, R. Compositional characterization of suspended particulate matter in Hellenic Volcanic Arc hydrothermal centres. *Physics and Chemistry of the Earth*

# **LIST OF PARTICIPANTS OF THE PROJECTS**





## **1. AMORES (MAS3-CT95-0040)**

Prime Contractor:

IFREMER - Institut Français de Recherche pour l'Exploitation de la Mer,  
Département Environnement Profond, Centre de Brest, Plouzane, BRETAGNE,  
FRANCE

Contact Person : DESBRUYERES, Daniel  
Tel: +33-98-224301  
Fax: +33-98-224547  
Email: ddesbruy@ifremer.fr

Other Contractors

- University of Durham, Department of Geological Sciences, Durham, UNITED KINGDOM

Contact Person : SEARLE, Roger Clive

- University of Bristol, Department of Geology, Bristol, UNITED KINGDOM

Contact Person : PALMER, Martin

- Centre for Applied Microbiology and Research (CAMR), Microbial Technology Laboratory, Salisbury, UNITED KINGDOM

Contact Person : SHARP, RICHARD JAMES

- University Wales - Bangor, School of Ocean Sciences, Gwynedd, WALES, UNITED KINGDOM

Contact Person : DANDO, Paul

- University of Cambridge, Department of Earth Sciences, Cambridge, UNITED KINGDOM

Contact Person : ELDERFIELD, Henry

- University of Southampton, Department of Oceanography, Southampton, UNITED KINGDOM

Contact Person : TYLER, Paul Alan

- Centre National de la Recherche Scientifique, UPR 9042 - Station Biologique de Roscoff SBR, Roscoff, BRETAGNE, BRETAGNE

Contact Person : LALLIER, François

- Université de Bretagne Occidentale, Laboratoire de Géochimie Isotopique du Groupement de Recherche, Plouzane, BRETAGNE, FRANCE

Contact Person : DOSSO, Laure

- Université de Bretagne Occidentale, Département de Chimie-Faculté des Sciences, Brest, BRETAGNE, FRANCE

Contact Person : APPRIOU, Pierre

- NERC Centre of Coastal and Marine Sciences, Plymouth, CORNWALL, DEVON, UNITED KINGDOM

Contact Person : DIXON, David

- National University of Ireland, Galway, Department of Marine Microbiology - The Martin Ryan Marine Science Institute, Galway, IRELAND

Contact Person : PATCHING, John W.

- Université Pierre et Marie Curie - Paris VI, Laboratoire de Pétrologie, Minéralogie, Magnatologie, Métallogénie, Paris, FRANCE

Contact Person : MEVEL, Catherine

- Era-Maptec Ltd., Dublin, IRELAND

Contact Person : COLLIER, David

- Vrije Universiteit Brussel, Afdeling Analytische Chemie - Faculteit van de Wetenschappen, Bruxelles, BELGIUM

Contact Person : DEHAIRS, Frank

- Commissariat à l'Energie Atomique (CEA), : Laboratoire de Modélisation du Climat et de l'Environnement, Direction des Sciences de la Matière, Gif-sur-Yvette, FRANCE

Contact Person : JEAN-BAPTISTE, Philippe

- Institut Français de Recherche pour l'Exploitation de la Mer (IFREMER), Département des Géosciences Marines-DRO, Plouzane, BRETAGNE, FRANCE

Contact Person : FOUQUET, Yves

- Instituto do Mar, Laboratório Marítimo da Guia, Cascais, LISBOA E VALE DO TEJO, PORTUGAL

Contact Person : SALDANHA, Luiz

- NERC Southampton Oceanography Centre, Southampton, UNITED KINGDOM

Contact Person : RUSSELL, Christopher

## **2. BENGAL (MAS3-CT95-0018)**

Prime Contractor

NERC Southampton Oceanography Centre, Southampton, UNITED KINGDOM

Contact Person : Name: RICE, Anthony Leonard

Tel: +44-1428-684141

Fax: +44-1428-683066

Email: pht@ua.nwo.ac.uk

Other Contractors

- University of Plymouth, Plymouth, UNITED KINGDOM
- Universiteit Gent, Vakgroep Morfologie, Systematiek en Ecologie, Faculteit Wetenschappen, Gent, BELGIUM

Contact Person : VINCX, Magda Margareta Mari

- University of Patras, Department of Geology - School of Natural Sciences, Patras, GREECE

Contact Person : VARNAVAS, Soterios

- University of Liverpool, Department of Earth Sciences, Oceanography Laboratories, Liverpool, UNITED KINGDOM

Contact Person : WOLFF, George Andrew

- Commissariat à l'Energie Atomique (CEA), Centre des Faibles Radioactivités, Direction des Sciences de la Matière, Gif-sur-Yvette, FRANCE

Contact Person : RABOUILLE, Christophe

- IFREMER - Institut Français de Recherche pour l'Exploitation de la Mer, Département Environnement Profond, Centre de Brest, Plouzane, BRETAGNE, FRANCE

Contact Person : SIBUET, Myrian

- University College Galway, Department of Marine Microbiology - The Martin Ryan Marine Science Institute, Galway, IRELAND
- Alfred-Wegener-Institut für Polar- und Meeresforschung, Abteilung Biologie 1 (Zoologie) - Fachbereich Biologie, Bremerhaven, GERMANY

Contact Person : BREY, Thomas

- Queen's University of Belfast, Medical Biology Centre, School of Biology and Biochemistry, Belfast, NORTHERN IRELAND, UNITED KINGDOM

- Christian-Albrechts Universität Kiel, Abteilung Marine Umweltgeologie, Forschungszentrum für Marine Geowissenschaften, Kiel, GERMANY

- Netherlands Institute for Sea Research, Department of Marine Ecology, Texel, Den Burg, NETHERLANDS

Contact Person : DUINEVELD, Gerard C.A.

- Università degli Studi di Ancona, Centro di Spesa Cattedra di Biologia Marina, Ancona, ITALY

Contact Person : MAURO, Fabiano

- Centre National de la Recherche Scientifique, URA 1513 - Flux de Matière et Réponses du Vivant, Brest, BRETAGNE, FRANCE

Contact Person : TREGUER, Paul

- University of Göteborg, Department of Analytical and Marine Chemistry - Faculty of Mathematics and Natural Sciences, Göteborg, SWEDEN

Contact Person : HALL, Per O.J.

- Universität Hamburg, Institut für Hydrobiologie und Fischereiwissenschaft, Hamburg, GERMANY

Contact Person : WEIKERT, Horst

- Institut Océanographique 'Fondation Albert I Prince de Monaco', Laboratoire de Physiologie des Etres Marins, Paris FRANCE

Contact Person : LAUBIER, Lucien

- Scottish Association of Marine Science, Oban, SCOTLAND, UNITED KINGDOM

Contact Person : GAGE, John

- INTERNATIONAL MARINE DREDGING CONSULTA, Antwerpen, BELGIUM

### **3. ESOP-2 (MAS3-CT95-0015)**

#### **Prime Contractor**

Universitetet i Bergen, Centre for Environment and Resource Studies, Bergen,  
NORWAY

Contact Person : Name: JANSEN, EYSTEIN

Tel: +47-55213491

Fax: +47-55324416

#### **Other Contractors**

- University of Cambridge, Scott Polar Research Institute, Cambridge, UNITED KINGDOM
- University of Cambridge, Department of Applied Mathematics and Theoretical Physics, Cambridge, UNITED KINGDOM

Contact Person : USCINSKI, Barry

- Danmarks Tekniske Universitet, Department of Electromagnetic Systems, Lyngby, DENMARK

Contact Person : TOUDAL, Leif

- Nansen Environmental and Remote Sensing Center, Solheimsviken, NORWAY

Contact Person : HAUGAN, Peter Mosby

- Université de Paris VI (Université Pierre et Marie Curie), Laboratoire d'Océanographie Dynamique et de Climatologie, Paris, FRANCE

Contact Person : GASCARD, Jean-Claude

- University of Göteborg, Department of Analytical and Marine Chemistry - Faculty of Mathematics and Natural Sciences, Göteborg, SWEDEN

Contact Person : ANDERSON, Leif G.

- Norges Teknisk-Naturvitenskapelige Universitet, Department of Biotechnology - Faculty of Chemistry and Chemical Technology, Norges Tekniske Høgskole, Trondheim, NORWAY

Contact Person : BOERSHEIM, Knut Yngve

- The Marine Research Institute of Iceland, Division of Oceanography and Ecology, Reykjavik, ICELAND

Contact Person : OLAFSSON, Jon

- International Council for the Exploration of the Sea, København, DENMARK

Contact Person : DOOLEY, Harry

- Meteorological Office, Department of Ocean Applications, Bracknell, UNITED KINGDOM

Contact Person : CATTLE, Howard

- Icelandic Meteorological Office, Sea Ice Research - Department of Research and Processing, Reykjavik, ICELAND

Contact Person : JAKOBSSON, Thor

- Dansk Hydraulisk Institut, Hørsholm, DENMARK

Contact Person : RASMUSSEN, Erland Bo

- Alfred-Wegener-Institut für Polar- und Meeresforschung, Abteilung Meeresphysik und Messwesen - Fachbereich Physik des Ozeans und der Atmosphäre, Bremerhaven, GERMANY

Contact Person : LEMKE, Peter

- Havforskningsinstituttet, Division for Chemical Oceanography, Bergen-Nordnes, NORWAY

Contact Person : REY, Francisco

- Centre National de la Recherche Scientifique (CNRS), Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse (CSNSM). (UPR 6412), Institut National de Physique Nucléaire et de Physique des Particules, Orsay, FRANCE

Contact Person : RAISBECK, Grant, Miller

- Universität Hamburg, Institut für Meereskunde, Hamburg, GERMANY

Contact Person : BACKHAUS, Jan

- Danish Meteorological Institute, Research and Development Department, København Ø, DENMARK

Contact Person : CHRISTENSEN, Jens Hesselbjerg

- Consiglio Nazionale delle Ricerche (CNR), Istituto per lo Studio delle Metodologie Geofisiche Ambientali, Modena, ITALY

Contact Person : PARMIGGIANI, Flavio Fiorigi

- Norwegian Polar Institute of the Ministry of Environment, Oslo, NORWAY

Contact Person : VINJE, Torgny

#### **4. AG-HY-FL (MAS3-CT95-0021)**

##### **Prime Contractor**

University of Wales, Bangor, School of Ocean Sciences, Menai Bridge, WALES,  
UNITED KINGDOM

Contact Person : Name: DANDO, Paul

Tel: +44-1248-351151

Fax: +44-12482620

Email: OSS109@SOS.BANGOR.AC

##### **Other Contractors**

- University of Patras, Department of Geology - School of Natural Sciences, Patras, GREECE

Contact Person : VARNAVAS, Soterios

- Consiglio Nazionale delle Ricerche (CNR), Stazione Oceanografica di La Spezia, Istituto per lo Studio della Dinamica delle Grandi Masse, La Spezia, ITALY

Contact Person : MELONI, Roberto

- Århus Universitet, Department of Microbial Ecology, Institute of Biological Sciences  
Århus C, DENMARK

Contact Person : GUNDERSEN, Jens Kristian

- National and Kapodistrian University of Athens, Department of Geophysics and Geothermy, Athens, GREECE

Contact Person : MAKROPOLOUS, Kostas

- Ente per le Nuove Tecnologie l'Energia e l'Ambiente (ENEA), Centro Ricerche Ambiente Marino, Lerici, ITALY

Contact Person : BIANCHI, Carlo Nike

- Christian-Albrechts Universität Kiel, Abteilung Marine Umweltgeologie, Forschungszentrum für Marine Geowissenschaften Kiel, GERMANY

Contact Person : LINKE, Peter

- Christian-Albrechts Universität Kiel, Institut für Allgemeine Mikrobiologie, Kiel, GERMANY

Contact Person : THOMM, Michael

- Max - Planck Institut for Marine Microbiology, Bremen, GERMANY

Contact Person : JOERGENSEN, Bo Barker

- Universität Karlsruhe (Technische Hochschule), Institut für Petrographie und Geochemie - Fakultät für Bio- und Geowissenschaften, Karlsruhe, GERMANY

Contact Person : STUBEN, Doris



## **DOCUMENTATION USED FOR THIS PUBLICATION**



## **AMORES (MAS3-CT95-0040)**

1. Final AMORES Report: "Azores Mid-Oceanic Ridge Ecosystem Studies : integrated research programme on deep-sea hydrothermal transfers and fluxes", April 1999
2. "Third European Marine Science & Technology Conference", Lisbon, 23-27 May 1998: Conference Proceedings, Project Synopses
3. German C.R., K.J.Richards, M.D.Rudnicki, M.M.Lam, J.L.Charlou & the FLAME Scientific Party. Topographic control of a dispersing hydrothermal plume. *Earth Planet. Sci. Lett (Express Letter)* 156, 267-273, 1998.
4. Pond, D., Dixon, D., Sargent, J., 1997. Wax-ester reserves facilitate dispersal of hydrothermal vent shrimps. *Marine Ecology Progress Series* 146: 289-290.
5. Pond, D., Dixon, D. R., Bell, M. V., Fallick, A. E., Sargent, J., 1997. Occurrence of 16:2(n-4) and 18:2(n-4) fatty acids in the lipids of the hydrothermal vent shrimps *Pimicaris axoculata* and *Alvinocaris markensis*: nutritional and trophic implications. *Marine Ecology Progress Series* 156: 167-174.
6. Pond, D., Segonzac, M., Bell, M. V., Dixon, D. R., Fallick, A. E., Sargent, J., 1997. Lipid and lipid carbon stable isotope composition of the hydrothermal vent shrimp *Mirocaris fortunata*: evidence for nutritional dependence on photosynthetically fixed carbon. *Marine Ecology Progress Series* 257 : 221-231.

## **BENGAL (MAS3-CT95-0018)**

1. Final BENGAL Report: "High resolution temporal and spatial study of the Benthic biology and Geochemistry of a north-eastern Atlantic abyssal Locality", April 1999
2. "Third European Marine Science & Technology Conference", Lisbon, 23-27 May 1998: Conference Proceedings, Project Synopses
3. Rabouille C., Witbaard R., Duineveld G., Wolff G., MacKenzie K. and Khripounoff A., Temporal variability of sedimentary recycling in the North Atlantic Ocean (BENGAL site) studied with a non-steady-state model. *Progress in Oceanography* (in preparation)
4. Witbaard R, Duineveld G.C.A., van der Weele J., Berghuis E.M., Reyss J.P., The Benthic response to the seasonal deposition of phyto pigments at the Porcupine Abyssal Plain in the N-E Atlantic.
5. Ragneneau O., Gallinari M., Hall P., Grandel S., Stahl H., Rickert D., Lampiti R., Witbaard R., Tengberg A., Gooday, A., The benthic silica cycle at the Porcupine Abyssal Plain, N-E Atlantic (submitted to *Deep Sea Research*, in March 99)
6. Ståhl, H. J., Brunnegård, J., Hall, P. O. J., Roos, P., Tengberg, A., 1998. Input, recycling and burial of organic and inorganic carbon in sediments of the Porcupine Abyssal Plain, NE Atlantic. *EU Meetings on Extreme Marine Environments at GEOMAR, Kiel, Nov. 18-21, 1998*
7. Ginger, M. L., Santos, Vera, L. C. S., Wolff, G. A., (submitted). The lipids of abyssal holothurians from the north-east Atlantic Ocean. *Journal of the Marine Association, Newsletter no. 11/9, p 41-43.*

## ESOP-2 (MAS3-CT95-0015)

1. Final ESOP-II Report "European Subpolar Ocean Programme Phase 2: The Thermohaline Circulation in the Greenland Sea", August 1999.
2. *ESOP-2 Newsletter*, September 1996.
3. *ESOP-2 Newsletter*, February 1997. Truls Johannessen, Leif Toudal, Gereon Budeus.
4. *ESOP-2 Newsletter*, September 1997. Stephanie Wolf, Henning Wehde, Blandine L'Heveder and Detlef Quadfasel (Photos: T. Johannessen).

## AG-HY-FL (MAS3-CT95-0021)

1. Dando, P. R., Aliani, S., Arab, H., Bianchi, C. N., Brehmer, M., Cocito, S., Fowler, S. W., Gundersen, J., Hooper, L. E., Kölbl, R., Kuever, J., Linke, P., Makropoulos, K. C., Meloni, R., Miquel, J.-C., Morri, C., Müller, S., Robinson, C., Schlesner, H., Sievert, S., Stöhr, R., Stüben, D., Thomm, M., Varnavas, S. P. & Ziebis, W. Hydrothermal studies in the Aegean Sea. *Physics and Chemistry of the Earth (in press)*.
1. Dando, P. R., Stüben, D. & Varnavas, S. P. Hydrothermalism in the Mediterranean Sea. *Progress in Oceanography (in press)*.
2. Kölbe, R. H., Stüben, D., 1998. Plume characteristics at a shallow water hydrothermal system of Milos, Greece. *Physics and Chemistry of the Earth (part of the Final Report for the AG-HY-FL project)*.
3. Makropoulos, K., Kouskouna, K., Karnassopoulou, A., Dando, P., Varnavas, S. P. Seismicity in the Aegean hydrothermal systems in relation to biogeochemical parameters *Physics and Chemistry of the Earth. (part of the Final Report for the AG-HY-FL project)*.
4. Robinson, C., Ziebis, W., Müller, S., Eichstaedt, K., Dando, P., Linke, P., Varnavas, S., Megalovasilis, P. & Panagiotaras, D. 1997. *In situ* investigations of shallow water hydrothermal vent systems, Palaeochori Bay, Milos, Aegean Sea. *Proceedings of the Fourth Underwater Science Symposium of the Society for Underwater Technology, Newcastle upon Tyne* pp. 85-100
5. Sartoni, G. & De Biasi, A. M. A survey of the marine algae of Milos Island, Greece. *Cryptogamie et Algologie (part of the Final Report for the AG-HY-FL project)*.
6. Stueben, D., Koelbl, R., Haushahn, P. & Schaupp, P. 1998. Measuring and sampling diffuse submarine hydrothermal vents. *International Ocean Systems Design*, 2 (2): 6-12
7. Ziebis, W., Böttcher, M. E., Weter, A., Miquel, J. C., Sievert, S. M., Linke, P., Deep water secondary productivity and intake sulphate reduction in a hypersulphidic basin in the Mediterranean Sea: Implications for the sulphur isotope record (*submitted to Nature, part of the Final Report for the AG-HY-FL project*).

European Commission

**EUR 19097 — Extreme marine environments  
Objectives and results from selected MAST projects**

*Edited by Anny Asmanidou and Elisabeth Lipiatou*

Luxembourg: Office for Official Publications of the European Communities

2000 — V, 52 pp. — 17.6 x 25 cm

ISBN 92-828-3011-X

Price (excluding VAT) in Luxembourg: EUR 7



## BELGIQUE/BELGIË

**Jean De Lannoy**  
Avenue du Roi 202/Koningslaan 202  
B-1190 Bruxelles/Brussel  
Tél. (32-2) 538 43 08  
Fax (32-2) 538 08 41  
E-mail: jean.de.lannoy@infoboard.be  
URL: <http://www.jean-de-lannoy.be>

## La librairie européenne/De Europese Boekhandel

Rue de la Loi 244/Wetstraat 244  
B-1040 Bruxelles/Brussel  
Tél. (32-2) 295 36 39  
Fax (32-2) 295 08 60  
E-mail: mail@libeurop.be  
URL: <http://www.libeurop.be>

## Moniteur belge/Belgisch Staatsblad

Rue de Louvain 40-42/Leuvenseweg 40-42  
B-1000 Bruxelles/Brussel  
Tél. (32-2) 552 22 11  
Fax (32-2) 511 01 84

## DANMARK

### J. H. Schultz Information A/S

Herstedvang 12  
DK-2620 Albertslund  
Tlf. (45) 43 63 23 00  
Fax (45) 43 63 19 69  
E-mail: schultz@schultz.dk  
URL: <http://www.schultz.dk>

## DEUTSCHLAND

### Bundesanzeiger Verlag GmbH

Vertriebsabteilung  
Amsterdamer Straße 192  
D-50735 Köln  
Tél. (49-221) 97 66 80  
Fax (49-221) 97 66 82 78  
E-Mail: Vertrieb@bundesanzeiger.de  
URL: <http://www.bundesanzeiger.de>

## ΕΛΛΑΔΑ/GREECE

### G. C. Eleftheroudakis SA

International Bookstore  
Panepistimiou 17  
GR-10564 Athina  
Tél. (30-1) 331 41 80/112/3/4/5  
Fax (30-1) 323 98 21  
E-mail: elebooks@netor.gr

## ESPAÑA

### Boletín Oficial del Estado

Tratfalgar, 27  
E-28071 Madrid  
Tél. (34) 915 38 21 11 (Libros),  
Fax (34) 913 84 17 15 (Suscrip.),  
Fax (34) 915 38 21 21 (Libros),  
Fax (34) 913 84 17 14 (Suscrip.),  
E-mail: clientes@com.boe.es  
URL: <http://www.boe.es>

### Mundt Prensa Libros, SA

Castelló, 37  
E-28001 Madrid  
Tél. (34) 914 36 37 00  
Fax (34) 915 75 39 98  
E-mail: librena@mundtprensa.es  
URL: <http://www.mundtprensa.com>

## FRANCE

### Journal officiel

Service des publications des CE  
26, rue Desaix  
F-75727 Paris Cedex 15  
Tél. (33) 140 58 77 31  
Fax (33) 140 58 77 00  
E-mail: europublications@journal-officiel.gouv.fr  
URL: <http://www.journal-officiel.gouv.fr>

## IRELAND

### Government Supplies Agency

Publications Section  
4-5 Harcourt Road  
Dublin 2  
Tél. (353-1) 661 31 11  
Fax (353-1) 475 27 60  
E-mail: opw@iol.ie

## ITALIA

### Licosa SpA

Via Duca di Calabria, 1/1  
Casella postale 552  
I-50125 Firenze  
Tél. (39) 055 64 83 1  
Fax (39) 055 64 12 57  
E-mail: licosa@licosa.com  
URL: <http://www.licosa.com>

## LUXEMBOURG

### Messageries du livre SARL

5, rue Raiffaissen  
L-2411 Luxembourg  
Tél. (352) 40 10 20  
Fax (352) 49 06 61  
E-mail: mail@mdl.lu  
URL: <http://www.mdl.lu>

## NETHERLAND

### SDU Servicecentrum Uitgevers

Christoffel Plantijnstraat 2  
Postbus 20014  
2500 EA Den Haag  
Tél. (31-70) 378 98 80  
Fax (31-70) 378 97 83  
E-mail: sdu@sdul.nl  
URL: <http://www.sdu.nl>

## ÖSTERREICH

### Manz'sche Verlags- und Universitätsbuchhandlung GmbH

Kohlmarkt 16  
A-1014 Wien  
Tél. (43-1) 53 16 11 00  
Fax (43-1) 53 16 11 67  
E-Mail: bestellen@manz.co.at  
URL: <http://www.manz.at>

## PORTUGAL

### Distribuidora de Livros Bertrand L.d.ª

Grupo Bertrand, SA  
Rua das Terras dos Vales, 4-A  
Apartado 60037  
P-2700 Amadora  
Tél. (351) 214 95 87 87  
Fax (351) 214 96 02 55  
E-mail: dlb@ip.pt

### Imprensa Nacional-Casa da Moeda, SA

Rua da Escola Politécnica nº 135  
P-1250 -100 Lisboa Codex  
Tél. (351) 213 94 57 00  
Fax (351) 213 94 57 50  
E-mail: spcoe@incm.pt  
URL: <http://www.incml.pt>

## SUOMI/FINLAND

### Akateeminen Kirjakauppa/

Akademiska Bokhandeln  
Keskuskatu 1/Centralgatan 1  
FIN-00101 Helsinki/Helsingfors  
P/fn (358-9) 121 44 18  
F/fax (358-9) 121 44 35  
Sähköposti: sps@akateeminen.com  
URL: <http://www.akateeminen.com>

## SVERIGE

### BTJ AB

Traktorvägen 11  
S-221 82 Lund  
Tlf. (46-46) 18 00 00  
Fax (46-46) 30 79 47  
E-post: btjeu-pub@btj.se  
URL: <http://www.btj.se>

## UNITED KINGDOM

### The Stationery Office Ltd

Orders Department  
PO Box 276  
London SW9 5DT  
Tél. (44-171) 870 60 05-522  
Fax (44-171) 870 60 05-533  
E-mail: book.orders@hso.co.uk  
URL: <http://www.tsonline.co.uk>

## ISLAND

### Bokabud Larusar Blöndal

Skólavörðustíg, 2  
IS-101 Reykjavík  
Tél. (354) 552 55 40  
Fax (354) 552 55 60  
E-mail: bokabud@simnet.is

## NORGE

### Swets Norge AS

Østenjoveien 18  
Boks 6512 Etterstad  
N-0606 Oslo  
Tél. (47-22) 97 45 00  
Fax (47-22) 97 45 45  
E-mail: kyterid@swets.nl

## SCHWEIZ/SUISSE/SVIZZERA

### Euro Info Center Schweiz

c/o OSEC  
Stampfenbachstraße 85  
PF 492  
CH-8003 Zürich  
Tél. (41-1) 365 53 15  
Fax (41-1) 365 54 11  
E-mail: eics@osec.ch  
URL: <http://www.osec.ch/eics>

## BĂLGARIA

### Europressa Euromedia Ltd

59, blvd Vitoshka  
BG-1000 Sofia  
Tél. (359-2) 980 37 66  
Fax (359-2) 980 42 30  
E-mail: Milena@mbos.cit.bg

## ČESKÁ REPUBLIKA

### ÚSIS

NIS-prodejna  
Havelskou 22  
CZ-130 00 Praha 3  
Tél. (420-2) 24 23 14 86  
Fax (420-2) 24 23 11 14  
E-mail: volanova@usiscr.cz  
URL: <http://usiscr.cz>

## CYPRUS

### Cyprus Chamber of Commerce and Industry

PO Box 1455  
CY-1509 Nicosia  
Tél. (357-2) 68 95 00  
Fax (357-2) 66 10 44  
E-mail: demetrap@ccci.org.cy

## EESTI

### Eesti Kaubandus-Tööstuskoda

(Estonian Chamber of Commerce and Industry)  
Toom-Kooli 17  
EE-0001 Tallinn  
Tél. (372) 646 02 44  
Fax (372) 646 02 45  
E-mail: einfo@koda.ee  
URL: <http://www.koda.ee>

## HRVATSKA

### Mediatrade Ltd

Pavia Hatza 1  
HR-10000 Zagreb  
Tél. (385-1) 481 94 11  
Fax (385-1) 481 94 11

## MAGYARORSZÁG

### Euro Info Service

Hungexpo Európa Ház  
PO Box 44  
H-1441 Budapest  
Tél. (36-1) 264 82 70  
Fax (36-1) 264 82 75  
E-mail: euroinfo@euroinfo.hu  
URL: <http://www.euroinfo.hu>

## MALTA

### Miller Distributors Ltd

Malta International Airport  
PO Box 25  
Luqa LQA 05  
Tél. (356) 66 44 88  
Fax (356) 67 67 99  
E-mail: gwirth@usa.net

## POLSKA

### Ars Polona

Krakowskie Przedmiescie 7  
Slrk, pocztowa 1001  
PL-00-950 Warszawa  
Tél. (48-22) 826 12 01  
Fax (48-22) 826 62 40  
E-mail: books119@arspolona.com.pl

## ROMÂNIA

### Euromedia

Strada Franceza Nr 44 sector 3  
RO-70749 Bucuresti  
Tél. (40-1) 315 44 03  
Fax (40-1) 315 44 03  
E-mail: mnediciu@pcnet.pcnet.ro

## ROSSIYA

### CCEC

60-letiya Oktyabrya Av. 9  
173131 Moscow  
Tél. (7-095) 135 52 27  
Fax (7-095) 135 52 27

## SLOVAKIA

### Centrum VTI SR

Nám. Slobody, 19  
SK-81223 Bratislava  
Tél. (421-7) 54 41 83 64  
Fax (421-7) 54 41 83 64  
E-mail: europ@tbl1.sltk.stuba.sk  
URL: <http://www.sltk.stuba.sk>

## SLOVENIJA

### Gospodarski Vestnik

Dunajska cesta 5  
SLO-1000 Ljubljana  
Tél. (386) 613 09 16 40  
Fax (386) 613 09 16 45  
E-mail: europ@gvestnik.si  
URL: <http://www.gvestnik.si>

## TÜRKIYE

### Dünya Infotel AS

100, Yil Mahallesi 34440  
TR-80050 Bagcilar-Istanbul  
Tél. (90-212) 629 46 89  
Fax (90-212) 629 46 27  
E-mail: infotel@dunya-gazete.com.tr

## AUSTRALIA

### Hunter Publications

PO Box 404  
3067 Abbotsford, Victoria  
Tél. (61-3) 94 17 53 61  
Fax (61-3) 94 19 71 54  
E-mail: jpdavies@ozemail.com.au

## CANADA

### Les éditions La Liberté Inc.

3020, chemin Sainte-Foy  
G1X 3V6 Sainte-Foy, Québec  
Tél. (1-418) 658 37 63  
Fax (1-800) 567 54 49  
E-mail: liberte@mediom.qc.ca

### Renouf Publishing Co. Ltd

5369 Chemin Canolek Road Unit 1  
K1J 9J3 Ottawa, Ontario  
Tél. (1-613) 745 26 65  
Fax (1-613) 745 76 60  
E-mail: order.dept@renoufbooks.com  
URL: <http://www.renoufbooks.com>

## EGYPT

### The Middle East Observer

41 Sherif Street  
Cairo  
Tél. (20-2) 392 69 19  
Fax (20-2) 393 97 32  
E-mail: inquiry@meobserver.com  
URL: <http://www.meobserver.com>

## INDIA

### EBIC India

3rd Floor, Y. B. Chavan Centre  
Gen. J. Bhosale Marg.  
400 021 Mumbai  
Tél. (91-22) 282 60 64  
Fax (91-22) 285 45 64  
E-mail: ebic@qasbm01.vsnl.net.in  
URL: <http://www.ebicindia.com>

## JAPAN

### PSI-Japan

Asahi Sanbancho Plaza #206  
7-1 Sanbancho, Chiyoda-ku  
Tokyo 102  
Tél. (81-3) 32 34 69 21  
Fax (81-3) 32 34 69 15  
E-mail: books@psi-japan.co.jp  
URL: <http://www.psi-japan.co.jp>

## MALAYSIA

### EBIC Malaysia

Level 7, Wisma Hong Leong  
18 Jalan Perak  
50450 Kuala Lumpur  
Tél. (60-3) 21 62 62 98  
Fax (60-3) 21 62 61 98  
E-mail: ebic-kl@mol.net.my

## MÉXICO

### Mundi Prensa Mexico, SA de CV

Río Pánuco No 141  
Colonia Cuauhtémoc  
MX-06500 Mexico, DF  
Tél. (52-5) 536 56 58  
Fax (52-5) 514 67 99  
E-mail: 101545.2361@compuserve.com

## PHILIPPINES

### EBIC Philippines

19th Floor, PS Bank Tower  
Sen. Gil J. Puyat Ave. cor. Tindalo St.  
Makati City  
Metro Manila  
Tél. (63-2) 759 66 80  
Fax (63-2) 759 66 90  
E-mail: eccpcom@globe.com.ph  
URL: <http://www.eccp.com>

## SOUTH AFRICA

### Eurochamber of Commerce in South Africa

PO Box 781738  
2146 Sandton  
Tél. (27-11) 884 39 52  
Fax (27-11) 883 55 13  
E-mail: info@eurochamber.co.za

## SOUTH KOREA

### The European Union Chamber of Commerce in Korea

5th Fl, The Shilla Hotel  
202, Jangchung-dong 2 Ga, Chung-ku  
100-392 Seoul  
Tél. (82-2) 22 53-5631/4  
Fax (82-2) 22 53-5635/6  
E-mail: eucock@eucock.org  
URL: <http://www.eucock.org>

## SRI LANKA

### EBIC Sri Lanka

Trans Asia Hotel  
115 Sir Chittampalam  
A. Gardiner Mawatha  
Colombo 2  
Tél. (94-1) 074 71 50 78  
Fax (94-1) 44 87 79  
E-mail: ebicsl@itlmin.com

## THAILAND

### EBIC Thailand

29 Vanissa Building, 8th Floor  
Soi Chidlom  
Floenchit  
10330 Bangkok  
Tél. (66-2) 655 06 27  
Fax (66-2) 655 06 28  
E-mail: ebicbkk@ksc15.th.com  
URL: <http://www.ebicbkk.org>

## UNITED STATES OF AMERICA

### Bernan Associates

4611-F Assembly Drive  
Lanham MD20706  
Tél. (1-800) 274 44 47 (toll free telephone)  
Fax (1-800) 865 34 50 (toll free fax)  
E-mail: query@bernan.com  
URL: <http://www.bernan.com>

## ANDERE LÄNDER/OTHER COUNTRIES/

### AUTRES PAYS

### Bitte wenden Sie sich an ein Büro Ihrer Wahl/ Please contact the sales office of your choice/ Veuillez vous adresser au bureau de vente de votre choix

### Office for Official Publications of the European Communities

2, rue Mercier  
L-2985 Luxembourg  
Tél. (352) 29 29-4245  
Fax (352) 29 29-42758  
E-mail: info.info@cec.eu.int  
URL: <http://eur-op.eu.int>

This publication provides an overview of the objectives, the main results, and the major publications of MAST-III-funded projects in the specific field of extreme marine environments. Extreme environments are defined as having conditions at the edge of the limit of life on our planet, and are found mainly in the deep ocean and at the poles. Extreme conditions are essential components in the assessment of anthropogenic and natural environmental change.

14

15

CG-NA-19097-EN-C

Price (excluding VAT) in Luxembourg: EUR 7



OFFICE FOR OFFICIAL PUBLICATIONS  
OF THE EUROPEAN COMMUNITIES

L-2985 Luxembourg

ISBN 92-828-3011-X



9 789282 830116 >