

## SKIN DISEASES IN CETACEANS

Marie-Françoise Van Bressem<sup>1</sup>, Koen Van Waerebeek<sup>1</sup>, Leonardo Flach<sup>2</sup>, Julio César Reyes<sup>3</sup>, Marcos César de Oliveira Santos<sup>4</sup>, Salvatore Siciliano<sup>5</sup>, Monica Echegaray<sup>3</sup>, Francisco Viddi<sup>6</sup>, Fernando Felix<sup>7</sup>, Enrique Crespo<sup>8</sup>, Gian Paolo Sanino<sup>9</sup>, Isabel Cristina Avila<sup>10</sup>, Natalia Fraijia<sup>11</sup> and Cristina Castro<sup>12</sup>

<sup>1</sup>Cetacean Conservation Medicine Group (CMED), CEPEC, Museo de Delfines, Pucusana, Peru; <sup>2</sup>Projeto Boto Cinza, Rua Sta Terezinha, 531 – Vila Muriqui, 23860-000 - Mangaratiba-RJ, Brazil; <sup>3</sup>Areas Costeras y Recursos Marinos – ACOREMA; <sup>4</sup>Projeto Atlantis, Laboratório de Biologia da Conservação de Cetáceos, Departamento de Zoologia, Instituto de Biociências, Universidade Estadual Paulista "Júlio de Mesquita Filho" (UNESP), Campus Rio Claro, Brasil, 13506-900; <sup>5</sup>Grupo de Estudios de Mamíferos Marinhos da Região dos Lagos (GEMM-Lagos), Departamento de Endemias Samuel Pessoa, Escola Nacional de Saúde Pública/FIOCRUZ. Rua Leopoldo Bulhões, 1480-6o. andar, Manguinhos, Rio de Janeiro, RJ 21041-210, Brazil; <sup>6</sup>Marine Mammal Research Group, Graduate School of the Environment, Macquarie University, NSW 2109 Sydney, Australia and Centro Ballena Azul, Valdivia, Chile; <sup>7</sup>Fundación Ecuatoriana para el Estudio de Mamíferos Marinos (FEMM), PO Box 09-01-11905, Guayaquil, Ecuador; <sup>8</sup>Centro Nacional Patagónico (CONICET), Boulevard Brown 3600, 9120 Puerto Madryn, Chubut, Argentina; <sup>9</sup>Centre for Marine Mammals Research LEVIATHAN, 7640392 Chile; <sup>10</sup>Universidad del Valle, Departamento de Biología, Grupo de Investigación en Ecología Animal, Cali, Colombia; <sup>11</sup>Universidad Jorge Tadeo Lozano, Bogota, Colombia; <sup>12</sup>Pacific Whale Foundation, Ecuador.

### ABSTRACT

Micro-organisms that are known or suspected to cause skin diseases in cetaceans are briefly reviewed. Viruses belonging to four families i.e. *Caliciviridae*, *Herpesviridae*, *Papillomaviridae* and *Poxviridae* were detected by electron microscopy, histology and molecular techniques in vesicular skin lesions, black dots perceptible by the touch, warts and tattoos in several species of odontocetes and mysticetes. Herpesviruses, poxviruses and likely a cutaneous papillomavirus are cetacean specific. Among bacteria, *Dermatophilus* spp., *Erysipelothrix rhusiopathiae*, *Mycobacterium marinum*, *Pseudomonas* spp., *Streptococcus iniae* and *Vibrio* spp. were isolated from ulcerative dermatitis, pyogranulomatous dermatitis and panniculitis, diamond skin disease and slow-healing ulcers and abscesses. *Aeromonas* spp., *Mycobacterium marinum*, *Pseudomonas* spp. and *Vibrio* spp. are normally present in the marine environment while *Erysipelothrix rhusiopathiae* and *Streptococcus iniae* are fish pathogens that may also infect captive dolphins. Most seem to be opportunistic pathogens, exploiting some break-down in the host's defenses to initiate an infection. Selection of antibiotic-resistant bacteria through the prophylactic use of antibiotics in aquaculture is suggested to be a growing problem in South America and may account for the emergence of unusual cutaneous conditions. At least four groups of fungi i.e. *Candida albicans*, *Fusarium* spp., *Trichophyton* spp. and *Lacazia loboi* cause skin diseases. Candidiasis occurs predominantly in captive odontocetes. The lesions are often localized around the body orifices and may become extensive, granulating and ulcerated. Fusariosis is characterized by firm, erythematous, cutaneous nodules. *Trichophyton* spp. was isolated from widespread superficial nodules in an Atlantic *T. truncatus* kept in captivity in Japan. Lobomycosis or lacaziosis is distinguished by grayish, whitish to slightly pink, verrucous lesions, often in pronounced relief that may ulcerate. While initially described only in *Tursiops truncatus* and *Sotalia guianensis* from the Americas, lobomycosis seems to be expanding to other continents. The role of ballast water in transporting fungi worldwide should be investigated. Finally, ciliated protozoans, likely *Kyaroikeus cetarius*, caused invasive dermatitis in small cetaceans from the USA and Korea. The aquatic environment of cetaceans is naturally home to bacteria and fungi but cetacean skin has several mechanisms to impede invasion. Chemical contaminants may affect natural skin barriers and depress the immune system. Wounds and specific viral infection (poxvirus, herpesvirus) may provide routes of entry.

### INTRODUCTION

Skin diseases in cetaceans have been reported at least since the 1950s (Simpson *et al.*, 1958; Slijper, 1962; Greenwood *et al.*, 1974). Caldwell *et al.* (1975) provided a comprehensive description of lobomycosis in common bottlenose dolphins *Tursiops truncatus* from the Northwest Atlantic. Geraci *et al.* (1979) gave an insight into the clinical aspects and histopathology of tattoo skin disease (TSD) in captive and free-ranging *T. truncatus* and Atlantic white-sided dolphin *Lagenorhynchus acutus*. More recently the epidemiology and ecology of skin diseases including TSD, lobomycosis and

lobomycosis-like disease (LLD) were investigated in some detail in free-ranging dolphins worldwide (Reif *et al.*, 2006; Van Bresseem and Van Waerebeek, 1996; Van Bresseem *et al.*, 2003a,b, 2007a, 2008a,b). Skin lesions were also recently reported in mysticetes (Pettis *et al.*, 2004; Brownell *et al.*, 2007; Bertelotti *et al.*, 2008; Mouton *et al.*, 2008). The role of natural and anthropogenic factors in the ecology of cutaneous illnesses of known and unknown aetiology has been thoroughly examined and the data gathered suggest that their emergence as well as their high prevalence in some populations is linked to a degraded environment, though water temperature and salinity may play a role too (Harzen and Brunnick, 1997; Wilson *et al.*, 1999; Van Bresseem *et al.*, 2003a, 2007a, 2008a,b; Reif *et al.*, 2006; Klaich *et al.*, 2008). Skin lesions also often reflect the health status of cetaceans (Pettis *et al.*, 2004; Van Bresseem *et al.*, 2008a).

Here we review the micro-organisms that are known or suspected to cause skin diseases in cetaceans.

## VIRUSES

### Caliciviruses

A strain of vesicular exanthema of swine virus (VESV), the cetacean calicivirus (CCV Tur-1) belongs to the genus *Vesivirus* of the family *Caliciviridae* (ICTV 2006). It was isolated from vesicular skin lesions that developed on a tattoo and old scars in two Atlantic bottlenose dolphins. The vesicles quickly eroded, leaving shallow ulcers in one of the dolphins (Smith *et al.*, 1983). Though CCV Tur-1 was isolated in the early 1980s, there are no published reports of further isolates and very little is known about its distribution and epidemiology. The role of vesiviruses in the aetiology of cutaneous vesicles observed in several cetaceans in South American coastal waters is suspected (Brownell *et al.*, 2007; Van Bresseem *et al.*, 2007a; Flach *et al.*, 2008).

Neutralizing antibodies to several other marine vesiviruses were detected by serology in mysticetes from the North Pacific (Smith and Latham, 1978, Smith and Boyt, 1990) and Alaska (Smith *et al.*, 1987, O'Hara *et al.*, 1998). California sea lions and one of their prey, the opaleye fish *Girella nigricans* (Dailey, 1970), may be primarily involved in the maintenance of these viruses in the North Pacific Ocean (Smith and Boyt, 1990). Transmission of the virus to other marine mammals is probably linked to contacts between species, migratory pathways and vectors (Smith *et al.*, 1980a,b, Smith and Boyt, 1990). Metazoan parasites like *Zalophotrema* sp. and *Parafilaroides decorus* may act as mechanical vectors (Smith *et al.*, 1980a,b).

### Herpesviruses

Herpes-like virus particles were detected by electron microscopy (EM) in epithelial cells from skin lesions of free-ranging and captive beluga whales *Delphinapterus leucas* from the St. Lawrence estuary and the Churchill river, Canada (Martineau *et al.*, 1988, Barr *et al.*, 1989) and of two dusky dolphins caught off central Peru in 1991 (Fig. 1) (Van Bresseem *et al.*, 1994). The skin lesions in the belugas were paler than the normal skin, circular or elliptical and slightly in relief (Martineau *et al.*, 1988, Barr *et al.*, 1989). In the St. Lawrence estuary individual the marks showed a dark small center surrounded by a narrow ill-defined dark rim (Martineau *et al.*, 1988). They were either numerous, generalized and up to 2cm in diameter (St. Lawrence beluga), or few, localized and up to 20-30 cm in diameter (Churchill river's individual). The infection did not seem to have affected the general body condition of the captive beluga (Barr *et al.*, 1989). Histologically, the lesions were characterized by epithelial necrosis and intranuclear inclusions. Intracellular oedema with micro-vesicle formation in superficial epithelium was also observed (Martineau *et al.*, 1988, Barr *et al.*, 1989). In the dusky dolphins, black dots, perceptible by the touch, were present on the head and body (Fig. 1). The infection did not seem to be more than mildly pathogenic (Van Bresseem *et al.*, 1994). Herpesvirus-like lesions were also observed in Burmeister's porpoises *Phocoena spinipinnis* and long-beaked common dolphins *Delphinus capensis* from the central coast of Peru in 1991-1995 (Van Bresseem *et al.*, 2006; Van Bresseem and Van Waerebeek, unpublished observations). In Peruvian small cetaceans they seem to mostly affect immature animals (Van Bresseem *et al.*, 1994, 2006; Van Bresseem, Van Waerebeek, Garcia-Godos, unpublished observations).

Previously unknown members of the subfamily *Alphaherpesvirinae* were detected by polymerase chain reaction (PCR) in skin lesions from two Atlantic *T. truncatus*, closely resembling those described in the dusky dolphins (Manire *et al.*, 2006; Smolarek Benson *et al.*, 2006). Though related to the simplexviruses of humans, they are cetacean-specific and probably co-evolved with their hosts for thousands of years (Smolarek Benson *et al.*, 2006). Identical or very similar alphaherpesviruses may be responsible for both cutaneous and fatal systemic infections in Atlantic *T. truncatus* (Smolarek Benson *et al.*, 2006). However, much remains to be learned on the ecology of these viruses as well as on their impact on survival.

### Papillomaviruses

Intranuclear virus-like particles were described in hyperplastic epithelial lesions observed on the skin of a harbour porpoise *Phocoena phocoena* incidentally taken in the Bay of Fundy (Geraci *et al.*, 1987) and of a captive killer whale (*Orcinus orca*) of Icelandic origin (Bossart *et al.*, 1996). Papillomavirus infection was demonstrated by

immunohistochemistry in cutaneous warts from a male *P. phocoena* (Fig. 2) stranded alive in Germany in December 1993 (Van Bresse *et al.*, 1999). Cutaneous warts are infrequently observed on the skin of North Atlantic harbour porpoises and were never seen in Peruvian small cetaceans in the period 1993-1994 though genital warts were highly prevalent in these species (Van Bresse *et al.*, 1996). The low prevalence of cutaneous PVs in small cetaceans may be explained by i) a high rate of unapparent or subclinical infections perhaps related to the high turnover of epidermis; ii) an early regression of the warts; iii) the development of a long lasting protective immunity; and iv) weak contagious nature of the disease (Van Bresse *et al.*, 1999). The impact on survival of cutaneous PVs is supposedly low or nihil.

### **Poxviruses**

Newly recognized members of the family Poxviridae possibly belonging to a new genus of *Chordopoxvirinae*, caused tattoo skin disease (TSD) characterised by very typical, irregular, grey, black or yellowish, stippled lesions in cetaceans worldwide (Fig.3) (Bracht *et al.*, 2006, Pearce *et al.*, 2008). These viruses are cetacean-specific and likely sub-order and family-specific (Bracht *et al.*, 2006, Pearce *et al.*, 2008). The epidemiology and ecology of TSD has been thoroughly investigated. Results of a recent study suggest that the prevalence and severity of the disease is the highest in coastal species and populations living in a contaminated environment (Van Bresse *et al.*, 2003a, 2008a). Thus, TSD may be considered as a general health indicator for cetaceans and their habitat. Tattoos may provide a route of entry to other viruses, bacteria and fungi that may increase the severity of the lesions (Smith *et al.*, 1983; Frasca *et al.*, 1996; Flach *et al.*, 2008; Van Bresse *et al.*, 2008c).

### **BACTERIA**

Several bacteria were isolated from skin lesions in cetaceans. Most are opportunistic, exploiting some break in the host defenses to initiate an infection. Some may have developed antibiotic-resistance due to the uncontrolled use of antibiotics in aquaculture in some countries (Cabello, 2004, 2006) that may render them more difficult to fight.

#### ***Aeromonas* spp.**

*A. hydrophila* caused ulcerative dermatitis and pneumonia in a *T. truncatus* from North Carolina, USA in the 1970s (Cusick and Bullock, 1973). *Aeromonas* species thrive in the aquatic environment and have been recorded in the proximity of salmon farms in Chile (Mirand and Zemelman, 2002). In humans they may cause severe skin lesions including necrotizing fasciitis (Lehane and Rawlin, 2000; Hiransuthikul *et al.*, 2005; Tsai *et al.*, 2007). They are likely candidates in the aetiology of skin diseases in cetaceans from SA (Fig. 4).

#### **Dermatophilosis**

*Dermatophilus* spp. are actinomycetes of the family *Dermatophilaceae*. *Dermatophilus*-like bacteria were observed in slightly depressed, round cutaneous areas with a dull surface and pale grey colour in six belugas stranded along the shores of the St Lawrence estuary (Mikaelian *et al.*, 2001). The lesions measured 0.5-4 cm in diameter and, in five of the six cetaceans, were numerous, covering about 5% of the surface of the whole body. Attempts to isolate *D. congolensis* were unsuccessful, possibly because another species of *Dermatophilus* was involved (Mikaelian *et al.*, 2001). The morphology of *Dermatophilus* spp. is so unique that a strong presumptive diagnosis can be made on examination of stained smears alone (Quinn *et al.*, 1994).

#### ***Erysipelothrix rhusiopathiae***

*E. rhusiopathiae* belongs to the family *Erysipelotrichaceae*. It may cause a sub-acute to chronic dermatological disease and septicemia in captive cetaceans (Geraci *et al.*, 1966). The skin disease is characterized by dermal infarction that results in sloughing of the epidermis. The formation of micro-infarcts may result in the rhomboid areas of cutaneous necrosis, known as diamond skin disease and characterized by grayish, raised, irregular in relief patches that may ulcerate (Geraci *et al.*, 1966). Animals with this form usually recover with timely antibiotic treatment. The septicemic form is usually acute or peracute, with the animal found moribund or dead. *E. rhusiopathiae* is a common contaminant of fishes. Ingestion of contaminated fish or injuries from the teeth of other cetaceans is presumed to be the route of infection for this disease.

#### ***Mycobacterium marinum***

*M. marinum* belongs to the *Mycobacteriaceae* family and is ubiquitous to fresh and marine sediments. It is an opportunistic, infrequent pathogen of homeotherms, including humans and is usually acquired through contamination of wounds but aerosol transmission also occurs (Bowenkamp *et al.*, 2001). *M. marinum* was isolated from multicentric 1-3 cm diameter ulcerative skin lesions that developed on the thorax and abdomen and near the genital folds of a captive beluga. On cut section they were epidermal ulcers with cores of purulent material that extended deep into the blubber.

Histologically, the lesions consisted of pyogranulomatous dermatitis and panniculitis (Bowenkamp *et al.*, 2001). There is no other report of skin disease related to this bacterium in cetaceans. However, it has been suggested that captive *T. truncatus* may harbour the bacteria asymptotically and transmit the disease to humans through biting (Flowers, 1970).

#### ***Pseudomonas* spp.**

Belonging to the family *Pseudomonadaceae*, *P. aeruginosa* is an opportunistic pathogen widely present in the terrestrial and fresh water environments as well as in the open ocean. It may be found in a biofilm, attached to some surface or substrate, or in a planktonic form, as a unicellular organism. It may cause disease and mortality in plants, humans and animals (Khan *et al.*, 2006). Most *Pseudomonas* infections are both invasive and toxinogenic. The ultimate *Pseudomonas* infection may be seen as composed of three distinct stages: i) bacterial attachment and colonization; ii) local invasion; iii) disseminated systemic disease (Todar, 2008). *Pseudomonas aeruginosa* caused fatal bronchopneumonia and extensive dermatitis in an Atlantic *T. truncatus* caught off Florida in the 1970s (Diamond *et al.*, 1979). The dolphin died 70 days after being caught after displaying dyspnea and anorexia. It had hard, round, and raised dermal nodules with necrotic centers all over its body surface. Similar skin lesions were observed in a female *L. obscurus* caught off Peru in 1994: it presented several nodules (some bleeding) on the back and tailstock (Van Bresse and Van Waerebeek, unpublished observations).

#### ***Staphylococcus delphini***

A member of the Family *Streptococcaceae*, *S. delphini* was isolated from multiple suppurating skin lesions that responded well to antibiotic treatment in captive dolphins (Varaldo *et al.*, 1988). *Staphylococcus* spp. have also been isolated from the respiratory tract and the genitals of a franciscana *Pontoporia blainvillei* and a southern right whale *Eubalaena australis* from Brazil, respectively (S. Siciliano, pers. observations).

#### ***Streptococcus iniae***

*Streptococcus iniae* is a member of the family *Streptococcaceae*. It is a serious pathogen of fishes that causes high losses in farmed fishes (Agnew and Barnes, 2007). It has a zoonotic potential with human infection linked to fish manipulation. It triggers 'golf ball disease' in captive bottas (*Inia geoffrensis*). The disorder is characterised by slow growing, multiple subcutaneous abscesses that eventually rupture (Bonar and Wagner, 2003). The disease is apparently linked to the fishes used to feed the dolphins and environmental conditions (Bonar *et al.*, 2007). It may also occur in free-ranging *I. geoffrensis* from the Brazilian Amazon Basin (da Silva *et al.*, 2008).

#### ***Vibrio* spp.**

Members of the *Vibrionaceae* Family, *Vibrio* spp. thrive in the aquatic environment. Their presence and numbers are influenced by factors such as temperature, salinity and algal density. *V. damsela*, *V. alginolyticus*, *V. parahaemolyticus*, *V. vulnificus* and *V. fluvialis* were isolated from slow-healing ulcers and abscesses of captive *T. truncatus* as well as from miscellaneous wounds sampled in free-ranging cetaceans from Brazil (Schroeder *et al.*, 1985; Fujioka *et al.*, 1988; Pereira *et al.*, 2007). Traumatized in coastal dolphins living in biologically contaminated environment likely provide a portal of entry to these bacteria.

## **FUNGI**

At least four groups of fungi have been isolated from skin lesions in cetaceans. Recently reports of mycotic diseases in free-ranging cetaceans (Fig. 5) have dramatically increased worldwide (Reif *et al.*, 2006; Van Bresse *et al.*, 2007; Kiska and Van Bresse, 2008; Moreno *et al.*, 2008; Mouton *et al.*, 2008; Shirakihara *et al.*, 2008; Siciliano *et al.*, 2008; V. Peddemors, P. Le Noury and Van Bresse, unpublished data). *Lacazia loboi* seems to be the more widespread fungal pathogen in free-ranging cetaceans. However, further histological and EM studies are required to confirm this assumption.

#### **Candidiasis**

Belonging to the Order Saccharomycetales, *C. albicans* lives under normal circumstances in 80% of the human population with no harmful effects, although overgrowth results in candidiasis. Candidiasis is often observed in immunocompromised individuals. Candidiasis is relatively common in captive cetaceans and occurs secondary to stress, unbalanced water disinfection with chlorines, or indiscriminate antibiotic therapy. The lesions usually may be localized around the body orifices. However, *C. albicans* has also caused extensive, granulating and sometimes ulcerated skin lesions and oesophago-gastric ulcerations. Cases of disseminated candidiasis caused the death of *T. truncatus*, *P. phocoena* and a long-finned pilot whale (*Globicephala melas*) (Nakeeb *et al.*, 1977; Dunn *et al.*, 1982).

### **Fusariosis**

*Fusarium* species are common soil saprophytes and plant pathogens. However, in recent years, they have been reported with increasing frequency as causes of opportunistic infections in humans and in animals, including reptiles, turtles, pinnipeds and dolphins (Frasca *et al.*, 1996; Cabañes *et al.*, 1997). *F. oxysporum*, *F. solani* and *F. verticillioides* are also serious emerging pathogens of humans with a broad resistance to the available anti-fungal drugs. They may cause lethal, invasive, fungal infections in immuno-compromised patients (Ortoneda *et al.*, 2004). *F. oxysporum* is broadly used in South America to destroy coca (*Erythroxylum coca*) plantations. *Fusarium* spp. caused raised, firm, erythematous, 2 to 5 mm cutaneous nodules that were most prominent on the head, trunk, and the caudal portion of an Atlantic white-sided dolphin and a pygmy sperm whale *Kogia breviceps* stranded along the northeast coast of the USA in 1991 (Frasca *et al.*, 1996) as well as in a captive beluga (Bowenkamp *et al.*, 2001). The yeast isolated from the *L. acutus* and *K. breviceps* appeared most like *F. oxysporum* (Frasca *et al.*, 1996).

### **Lobomycosis or lacaziosis**

*Lacazia loboi* is an uncultivated pathogen that belongs with the other dimorphic fungal pathogens to the order Onygenales (Herr *et al.*, 2001). *L. loboi* naturally affects humans and cetaceans from the Americas causing lobomycosis (Caldwell *et al.*, 1975; Reif *et al.*, 2006; Van Bresseem *et al.*, 2007a). *L. loboi* cells found in *T. truncatus* infected tissues are significantly smaller than those found in humans, suggesting that the organism may not be identical in the two hosts (Haubold *et al.*, 2000). Lobomycosis in dolphins is characterized by grayish, whitish to slightly pink, verrucous lesions, often in pronounced relief that may ulcerate (Migaki *et al.*, 1971). Though the disease evolves slowly it may eventually lead to death (Simões-Lopes *et al.*, 1993; Van Bresseem *et al.*, 2007a; Bermúdez-Villapol *et al.*, 2008). The clinical symptoms, histology and epidemiology of the disease are reviewed in detail in Reif *et al.*, 2006; Van Bresseem *et al.*, 2007a; Moreno *et al.*, 2008; Paniz-Mondolfi *et al.*, 2008; Siciliano *et al.*, 2008. In South America, several cases highly reminiscent of lobomycosis were observed in free-ranging coastal *T. truncatus* and *S. guianensis* (Fig. 6). In the absence of a histological diagnosis, the disease was called lobomycosis-like disease (Van Bresseem *et al.*, 2007a).

### **Trichophyton spp.**

*Trichophyton* spp. was isolated from widespread superficial nodules on the trunk of a captive Atlantic *T. truncatus* kept in captivity in Japan (Hoshina *et al.*, 1956).

## **PROTOZOANS**

Dermatitis with invasive ciliates have been reported in *T. truncatus*, spotted dolphins *Stenella attenuata*, a common dolphin *Delphinus delphis*, a pygmy sperm whale, a Fraser's dolphin *Lagenodelphis hosei* and a killer whale *Orcinus orca* from the United States as well as in a "*Tursiops gilli*" kept in captivity in Seoul Grand Park, Korea (Schulman and Lipscomb, 1987; Choi *et al.*, 2003). The disease was characterized by discrete ulcers with subjacent dermal, and often subcutaneous, necrosis and inflammation in several *T. truncatus* that died during a morbillivirus epizootic along the Atlantic coast of the US in 1987-1988 (Schulman and Lipscomb, 1999). Prevalence of this condition was high (19%; N=95) during the epizootic in comparison with other years, probably because of the immune-suppression caused by the viral infection. The ciliates observed in the skin lesions were morphologically identical to *Kyaroikeus cetarius* isolated from the blowhole of captive and free-ranging cetaceans (Sniezek *et al.*, 1995; Poynton *et al.*, 2001). They may be opportunistic invaders taking advantage of skin traumas (Schulman and Lipscomb, 1999). In the *T. gilli* the lesions initially developed on a traumatic skin injury on the neck area and then spread throughout the body surface (Choi *et al.*, 2003). The protozoans detected in the *T. gilli* were also morphologically similar to *K. cetarius* (Choi *et al.*, 2003). Protozoan-like organisms were also observed by transmission electron microscopy in rounded marks spread over the entire body surface of a male *D. capensis* and in small, dark, rounded marks, often punctured, present in an immature female *L. obscurus* (Van Bresseem *et al.*, 2006; unpublished observations).

## **CONCLUSIONS**

Commensal bacteria, fungi and micro-algae are naturally present in marine waters. The nanorough surface of the dolphin skin smoothed by a shear-resistant gel at the epidermal surface likely prevent their adhesion (Meyer *et al.*, 2008). The presence of the antibiotic, cationic enzyme lysozyme and antibiotic peptide group of Defensins in the epidermis may also serve as non-specific defense against these micro-organisms (Meyers and Seegers, 2004). Chemical contaminants may affect natural skin barriers while wounds and specific viral infection (poxvirus, herpesvirus) may provide routes of entry to opportunistic pathogens. By depressing the immune response of infected animals, morbilliviruses may favour protozoan infection (Schulman and Lipscomb, 1999).

The apparent emergence of skin diseases in cetaceans worldwide has generated an increased interest in these conditions and the role of environmental factors. Previously unrecognised cetacean viruses have been isolated, shedding more light into the phylogeny of several virus families. However, in a large number of cases the aetiology of these diseases is unknown, often because the affected cetaceans are free-ranging and sampling unrealistic. Photo-identification provides a good way of following the development of the illness, examine its general characteristics and study its epidemiology. When feasible, samples should be taken from stranded or by-caught specimens for isolation, electron microscopy and molecular studies. The remoteness of some study areas and lack of financial support sometimes prevents that such studies are undertaken.

The association between coastal waters that have been anthropogenically degraded and the health of its marine inhabitants has been clearly demonstrated in fishes years ago (Murchelano, 1990). Unless, drastic measures are taken to improve treatment of run-off waters, reduce chemical contamination, decontaminate water ballast and stop or at least reduce the heavily use prophylactic antibiotics in aquaculture, coastal environments will become inhabitable and unsuitable for human recreation purposes.

## ACKNOWLEDGEMENTS

The work of M-F Van Bresseem was funded by WDCS and CSI. A COREMA received support from the Gesellschaft zur Rettung der Delphine (GRD -Germany) and the Cetacean Society International (CSI-USA). Dolphin research in the Paracas National Reserve was carried out under permits issued by the Instituto Nacional de Recursos Naturales (INRENA). CEPEC research and data analysis has benefitted from sustained IFAW support. Van Waerebeek attends the IWC/60 SC meeting as head of scientific delegation for Federal Public Service, Public health, Food Chain security and Environment, Brussels, Belgium. We kindly thank Drs Vic Peddemors and Ron Kastelein and Pamela Le Noury, for providing some of the photographs used in this article. S. Siciliano is supported by fellowships provided by FIOCRUZ/FAPERJ and PROJETO PIATAM OCEANO – CENPES/PETROBRAS, UFF, UFPA and MPEG.

## LITERATURE CITED

- Agnew, W. and Barnes, A.C. 2007. *Streptococcus iniae*: an aquatic pathogen of global veterinary significance and a challenging candidate for reliable vaccination. *Vet Microb* 122: 1-15
- Barr, B., Dunn, J.L., Daniel, M.D. and Banford, A. 1989. Herpes-like viral dermatitis in a beluga whale (*Delphinapterus leucas*). *J Wildl Dis* 25: 608-611
- Bermúdez-Villapol, L.A., Sayegh, A.J. and Van Bresseem, M-F. 2008. Lobomycosis-like disease in a bottlenose dolphin (*Tursiops truncatus*) from Venezuela Paper SC/60/DW7 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]
- Bertellotti, M., Varisco, A., Aguado, G. and Francisco, E. 2008. Skin lesions in Southern right whales off the coast of Valdes Peninsula, Argentina. Paper SC/60/DW14 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]
- Bonar, C.J. and Wagner, R.A. 2003. A third report of "golf ball disease" in an Amazon River dolphin (*Inia geoffrensis*) associated with *Streptococcus iniae*. *J Zoo Wildl Med*: 34, 296-301.
- Bonar, C.J., Boede, E.O., García Hartmann, M.G., Lowenstein-Whaley, J., Mujica-Jorquera, E., Parish, S.V., Parish, J.V., Garner, M.M. and Stadler, C.K. 2007. A retrospective study of pathologic findings in the Amazon and Orinoco dolphin (*Inia geoffrensis*) in captivity. *J Zoo Wildl Med* 38: 177-191.
- Bossart, G. D., Cray, C., Solorzano, J. L., Decker, S. J., Cornell, L. H. and Altman, N. H. 1996. Cutaneous papovaviral-like papillomatosis in a killer whale (*Orcinus orca*). *Mar Mamm Sc* 12: 274-281.
- Bowenkamp, K., Frasca, S. Jr, Draghi II, Tsongalis, G.J., Koerting, C. Hinckley, L., De Guise, S., Montali, R.J., Goertz, C.J., St Aubin, D. J. and Dunn, L. 2001. Mycobacterium dermatitis and chronic pleuritis in a captive white whale (*Delphinapterus leucas*) with aortic rupture. *J Vet Diagn Invest* 13: 524-530.
- Bracht, A.J., Brudek, R.L., Ewing, R.Y., Manire, C.A., Burek, K.A., Rosa, C., Beckmen, K.B., Maruniak, J.E and Romero, C.H. 2006. Genetic identification of novel poxviruses of cetaceans and pinnipeds. *Arch Virol* 151: 423-438.
- Brownell, R. L. Jr., Carlson, C.A., Galletti Vernazzani, B. and Cabrera, E. 2007. Skin lesions on blue whales off Southern Chile: possible conservation implications? Paper SC/59/SH21 presented to the IWC Scientific Committee, May 2007 (unpublished).
- Cabañes, F.J., Alonso, J.M., Castellá, G., Alegre, F., Domingo, M. and Pont, S. 1997. Cutaneous hyalophomycosis caused by *Fusarium solani* in a loggerhead sea turtle (*Caretta caretta* L.). *J Clin Microb* 35:3343-3345.
- Cabello, F.C. 2004. Antibioticos y acuicultura en Chile: consecuencias para la salud humana y animal. *Rev Med Chile* 132: 1001-1006.

- Cabello, F.C. 2006. Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environm Microb* 8: 1137-1144.
- Caldwell, D.K., Caldwell, M.C., Woodard, J.C., Ajello, L., Kaplan, W. and McLure, H.M. 1975 Lobomycosis as a disease of the Atlantic bottle-nosed dolphin (*Tursiops truncatus* Montagu, 1821). *Am J Tropic Med* 24: 105-114.
- Choi, Y-K., Kang, M-S., Kim, D-Y. and Sohn, H-R. 2003. Disseminated ciliated protozoan infection in a Pacific dolphin (*Tursiops gilli*). *Vet Rec* 153: 714-715.
- Cusick, P.K. and Bullock, B.C. 1973. Ulcerative dermatitis and pneumonia associated with *Aeromonas hydrophila* infection in the bottle-nosed dolphin. *JAVMA* 163: 578-579.
- Dailey, M.D. 1970. The transmission of *Parafilaroides decorus* (Nematoda: Metastrongyloidea) in the California sea lion (*Zalophus californianus*). *Proc Helminthol Soc Wash* 37:215-222
- da Silva, V.M.F., Martin, A.R. and Mikesch, E. 2008. Skin disease and lesions of the boto *Inia geoffrensis* in the Central Amazon. Paper SC/60/DW11 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal].
- Diamond, S.S. Ewing, D.E. and Caldwell, G.A. 1979. Fatal Bronchopneumonia and dermatitis caused by *Pseudomonas aeruginosa* *JAVMA* 175: 984-987.
- Dunn, J.L., Buck, J.D. and Spotte, S. 1982. Candidiasis in captive cetaceans 1982. *JAVMA* 181:1316
- Flach, L., Van Bresse, M-F., Reyes, J.C., Echegaray, M., Siciliano, S., Santos, M., Viddi, F., Crespo, E., Klaich, J., Moreno, I., Tavares, M., Felix, F. and Van Waerebeek, K. 2008. Miscellaneous skin lesions of unknown aetiology in small cetaceans from South America. Paper SC/60/DW4 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal].
- Flowers, D.J. 1970. Human infection due to *Mycobacterium marinum* after a dolphin bite. *J Clin Pathol* 23: 475-477.
- Frasca, S. Jr., Dunn, J.L., Cooke J.C. and Buck J.D. 1996. Mycotic dermatitis in an Atlantic white-sided dolphin, a pygmy sperm whale, and two harbor seals. *JAVMA* 208:727-729.
- Fujioka, R.S., Greco, S.B., Cates, M.B. and Schroeder, J.P. 1988. *Vibrio damsela* from wounds in bottlenose dolphins *Tursiops truncatus*. *DAO* 4:1-8.
- Geraci, J.R., Sauer, R.M. and Medway, W. 1966. Erysipelas in dolphins. *Am J Vet Research* 27: 597-606.
- Geraci, J.R., Hicks, B.D. and St Aubin, D.J. 1979. Dolphin pox: a skin disease of cetaceans. *Canadian Journal of Comparative Medicine* 43:399-404.
- Geraci, J.R., Palmer, N.C. and St Aubin, D.I. 1987. Tumors in cetaceans: analysis and new findings. *Can J Fish Aquat Sc* 44, 1289-1300
- Greenwood, A.G., Harrison, R.J. and Whitting, H.W. 1974. Functional and pathological aspects of the skin of marine mammals. In: R.J. Harrison (ed) *Functional anatomy of marine mammals*. Academic Press, New York, pp 73-110.
- Harzen, S. and Brunnick, B. 1997. Skin disorders in bottlenose dolphins (*Tursiops truncatus*), resident in the Sado estuary, Portugal. *Aquat Mamm* 23: 59-68
- Haubold, E.M., Cooper, C.R. Jr, Wen, J.W., McGinnis, M.R. and Cowan, D.F. 2000. Comparative morphology of *Lacazia loboi* (syn. *Loboa loboi*) in dolphins and humans. *Med Mycol* 38: 9-14.
- Herr, R.A., Tarcha, E.J., Taborda, P.R., Taylor, J.W., Ajello, L. and Mendoza, L. 2001. Phylogenetic analysis of *Lacazia loboi* places this previously uncharacterized pathogen within the Dimorphic Onygenales. *J Clin Microb* 39: 309-114
- Hiransuthikul N., Tantisiriwat W., Lertutsahakul K., Vibhagool A. and Boonma P. 2005. Skin and soft-tissue infections among tsunami survivors in southern Thailand. *Clin Infect Dis* 41: 889-890.
- Hoshina, T. and Sigiura, Y. 1956. On a skin disease and nematode parasite of a dolphin *Tursiops truncatus* (Montagu, 1821). *Scient Whale Res Inst* (Tokyo) 11:133-137.
- ICTV.2006. Caliciviridae. In ICTVdB – The Universal Virus Database, version 4. Edited by C. Buchen-Osmond. New York: Columbia University. <http://www.ncbi.nlm.nih.gov/ICTVdb/Ictv/index.htm>
- Khan, N.H., Ishii, Y., Kimata-Kino, N., Esaki, H., Nishino, T., Nishimura, M., and Kogure, K. 2006. Isolation of *Pseudomonas aeruginosa* from open ocean and comparison with freshwater, clinical, and animal isolates. *Microb Ecol* DOI: 10.1007/s00248-006-9059-3.
- Lehane, L. and Rawlin GT. 2000. Topically acquired zoonosis from fish: a review. *Med J Aust* 174: 480-481
- Manire C.A., Smolarek K.A., Romero C.H., Kinsel M.J., Clauss T.M. and Byrd, L. 2006. Proliferative dermatitis associated with a novel alpha herpesvirus in an Atlantic bottlenose dolphin (*Tursiops truncatus*). *J Zoo Wildl Med.* 37:174-81.
- Martineau, D., Lagace, A., Beland, P., Higgins, R., Armstrong, D. and Shugart, L.R. 1988. Pathology of stranded beluga whales (*Delphinapterus leucas*) from the St. Lawrence Estuary, Quebec. *Can J Comp Pathol* 98:287-3
- Meyer, W. and Seegers, U. 2004. A preliminary approach to epidermal antimicrobial defense in the Delphinidae. *Mar Biol* 144: 841-844
- Meyer, W., Kloepper, J.E. and Fleischer, L-G. 2008. Demonstration of  $\beta$ -glucan receptors in the skin of aquatic mammals—a preliminary report. *Eur J Wildl Res* DOI 10.1007/s10344-008-0173.

- Migaki, G., Valerio, M.G., Irvine, B.A and Graner, F.M. 1971. Lobo's disease in an Atlantic bottle-nosed dolphin. *JAVMA* 159: 578-582.
- Mikaelian, I., Lapointe, J.-M., Labelle, P, Higgins, R., Paradis, M. and Martineau, D. 2001. Dermatophilus-like infection in beluga whales, *Delphinapterus leucas*, from the St Lawrence estuary. *Vet Dermatol* 12: 59-62.
- Mirand, C.D. and Zemelman, R. 2002. Antimicrobial multiresistance in bacteria isolated from freshwater Chilean salmon farms. *Sci Total Environ* 293:207-18.
- Moreno, I.G., Ott, P.H., Tavares, M., Oliveira, L.R., Borba, M.R., Driemeier, D., Nakashima, S.B., Heinzmann, L.S., Siciliano, S. and Van Bresse, M-F. 2008. Mycotic dermatitis in common bottlenose dolphins (*Tursiops truncatus*) from southern Brazil, with a confirmed record of lobomycosis disease. Paper SC/60/DW1 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal].
- Mouton, M., Reeb, D., Thornton, M., Botha, A. and Best, P.B. 2008. Cutaneous fungi on southern right whales (*Eubalaena australis*) from South Africa as a possible indicator of health risks. Paper SC/60/DW17 presented to the IWC Scientific Committee, May 2008 (unpublished).
- Murchelano, R.A. 1990. Fish health and environmental health. *Environm Health Perspect* 86: 257-259.
- Nakeeb, S., Targowski, S.P. and Spotte, S. 1977. Chronic cutaneous candidiasis in bottle-nosed dolphins. *JAVMA* 171: 961-965.
- Ortoneda, M., Guarro, J., Madrid, M.P., Caracuel, Z., Roncero, I.G., Mayayo, E. and Di Pietro, A. 2004. *Fusarium oxysporum* as a multihost model for the genetic dissection of fungal virulence in plants and mammals. *Infect Imm* 72: 1760-1766.
- O'Hara, T.M., House, C., House, J.A., Suydam, R.S. and George, J.C. 1998. Viral serologic survey of bowhead whales in Alaska. *J Wildl Dis* 34:39-
- Paniz-Mondolfi, A.E., Hoffman, L.S., Reyes Jaimes, O. And Pérez Álvarez, A.. 2008. Epidemiology of lobomycosis in humans and its possible emergenc in other animal species. Paper SC/60/DW9 presented to the IWC Scientific Committee, May 2008 (unpublished). 9pp. [Available from the Office of this Journal].
- Pearce, G., Blacklaws, B.A., Gajda, A.M., Jepson, P., Deaville, R. and Van Bresse, M-F. 2008. Molecular identification and phylogenetic relationships in poxviruses from cetacean skin lesions. Conference of the European Cetacean Society, The Netherlands, 10-12 March 2008.
- Pereira, C.S., Amorim, S.D. Santos, A.F.M., Siciliano, S., Moreno, I.M.B., Ott, P.H. and Rodrigues, D.P. 2007. *Vibrio* spp. isolados de mamíferos marinhos capturados na região litorânea do sudeste ao sul do Brasil. *Pesq Vet Bras* 27:81-83.
- Pettis, H.M., Rolland, R.M., Hamilton, P.K., Brault, S., Knowlton, A.R. and Kraus, S.D. 2004. Visual health assessment of North Atlantic right whales (*Eubalaena glacialis*) using photographs. *Can J Zool* 82: 8-19
- Poynton, S. L., Whitaker, B.R. and Heinrich, A.B. 2001. A novel trypanoplasm-like flagellate *Jarrellia atramenti* n. g., n. sp. (*Kinetoplastida: Bodonidae*) and ciliates from the blowhole of a stranded pygmy sperm whale *Kogia breviceps* (Physeteridae): morphology, life cycle and potential pathogenicity. *Dis Aquat Org* 44: 191-201.
- Quinn, P. J., Carter, M. E., Markey, B. K. and Carter, G.R. 1994. Clinical Veterinary Microbiology. Wolfe Publishing, Mosby-Year Book Europe Ltd, London, UK.
- Reif, J.S., Mazzoil, M.S., McCulloch, S.D., Varela, R., Goldstein, J.D., Fair, P. and Bossart, G.D. 2006. Lobomycosis in Atlantic bottlenose dolphins from the Indian River Lagoon, Florida. *JAVMA* 228: 104-108.
- Siciliano, S., Van Bresse, M-F., Moreno, I.B., Ott, P.H., Tavares, M., Flores, P.A.C., Flach, L., Cesar Reyes, J., Echegaray, M., de Oliveira Santos, M.C., Viddi, F., Crespo, E., Klaich, J., Felix, F., Sanino, G.P. and Van Waerebeek, K. 2008. Review of lobomycosis and lobomycosis-like disease in Cetacea from South America. Paper SC/60/DW13 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal].
- Schulman, F.Y. and Lipscomb, T.P. 1999. Dermatitis with Invasive Ciliated Protozoa in Dolphins That Died During the 1987-1988 Atlantic Bottlenose Dolphin Morbilliviral Epizootic. *Vet Pathol* 36:171-174
- Schroeder, J.P., Wallace, J.G. , Cates, M.B., Greco, S.B. and Moore, P. W. B. 1985. An infection by *Vibrio alginolyticus* in an Atlantic bottlenose dolphin housed in an open ocean pen. *J Wildl Dis* 21: 437-438
- Simões-Lopes, P.C., Paula, G.S., Xavier, F.M. and Scaramelo, A.C. 1993. First case of lobomycosis in a bottlenose dolphin on Southern Brazil. *Mar Mamm Sc* 9: 329 - 331
- Simpson, C.F., Wood, F.G. and Young F. 1958. Cutaneous lesions on a porpoise with Erysipelas. *JAVMA* 133: 558-560.
- Slijper, E.J. 1962. Whales. Hutchinson, London
- Sniezek, J.H., Coats, D.W. and Small, E.B.. 1995. *Kyaroikeus cetarius* N. G., N. Sp.: A Parasitic Ciliate from the Respiratory Tract of Odonticete Cetacea. *J Eukar Microb* 42: 260-268
- Smith, A.W. and Lalham, A.B. 1978. Prevalence of vesicular exanthema of swine antibodies among feral mammals associated with the southern California coastal zones. *Am J Vet Res* 39:291-296
- Smith A.W. and Boyt, P.M. 1990. Calicivirus of ocean origin: a review. *J Zoo Wildl Med* 21:3-23
- Smith, A.W., Skilling, D.E., Dardiri, A.H. and Latham, A.B. 1980a Calicivirus pathogenic for swine: a new serotype isolated from opaleye *Girella nigricans*, an ocean fish. *Science* 209: 940-941

- Smith, A.W., Skilling, D.E. and Brown, R.J. 1980b. Preliminary investigation of a possible lung worm (*Parafilaroides decorus*), fish (*Girella nigricans*) and marine mammal (*Callorhinus ursinus*) cycle for San Miguel sea lion virus type 5. *Am J Vet Res* 41:1846-1850
- Smith, A.W., Skilling, D.E. and Ridgway, S.H. 1983. Calicivirus-induced vesicular disease in cetaceans and probable interspecies transmission. *JAVMA* 11: 1223-1225
- Smith, A.W., Skilling, D.E., Bernirschke, K., Albert, F.T. and Barlough, J.E. 1987. Serology and virology of the bowhead whale (*Balaena mysticetus* L.). *J Wildl Dis* 23:92-98
- Smolarek Benson, K.A., Manire, C.A., Ewing, R.Y., Saliki, J.T., Townsend, F.I., Ehlers, B. and Romero, C.H. 2006. Identification of novel alpha- and gammaherpesviruses from cutaneous and mucosal lesions of dolphins and whales. *J Virol Methods* 136:261-6.
- Todar, K. 2008. Todar's Online Textbook of Bacteriology. <http://www.textbookofbacteriology.net/>
- Tsai, Y.H., Hsu, R.W., Huang, T.J., Hsu, W.H., Huang, K.C., Li, Y.Y. and Peng, K.T. 2007. Necrotizing soft-tissue infections and sepsis caused by *Vibrio vulnificus* compared with those caused by *Aeromonas* species. *J Bone Joint Surg Am* 89: 631-636
- Van Bresseem, M.-F. and Van Waerebeek, K. 1996. Epidemiology of poxvirus in small cetaceans from the Eastern South Pacific. *Mar Mamm Sc* 12: 371-382.
- Van Bresseem, M.-F., Van Waerebeek, K., Garcia-Godos, A., Dekegel, D. and Pastoret, P.-P. 1994. Herpes-like virus in dusky dolphins, *Lagenorhynchus obscurus*, from coastal Peru. *Mar Mamm Sc* 10: 354-359.
- Van Bresseem, M.-F., Van Waerebeek, K., Piérard, G. and Desaintes, C. 1996. Genital and lingual warts in small cetaceans from coastal Peru. *Dis Aquat Org* 26: 1-10
- Van Bresseem, M.-F., Kastelein, R. A., Flamant, P. and Orth, G. 1999. Cutaneous papillomavirus infection in a harbour porpoise (*Phocoena phocoena*) from the North Sea. *Vet Rec* 144: 592-593.
- Van Bresseem, M.-F., Van Waerebeek, K., Raga, J.A., Gaspar, R., Di Benedetto, A.P., Ramos, R. and Siebert, U. 2003a. Tattoo disease of odontocetes as a potential indicator of a degrading or stressful environment: a preliminary report. Document SC/55/E1, International Whaling Commission, Berlin, Germany.
- Van Bresseem, M.-F., Gaspar, R. and Aznar, J. 2003b. Epidemiology of tattoo skin disease in bottlenose dolphins (*Tursiops truncatus*) from the Sado estuary, Portugal. *Dis Aquat Org* 56: 171-179.
- Van Bresseem, M.-F., Van Waerebeek, K., Montes, D., Kennedy, S., Reyes J.C., Garcia-Godos, A.I., Onton-Silva, K. and Alfaro-Shigueto, J. 2006. Diseases, lesions and malformations in the long-beaked common dolphin (*Delphinus capensis*) from the Southeast Pacific. *Dis Aquat Org* 68: 149-165.
- Van Bresseem, M.-F., Van Waerebeek, K., Reyes, J.C., Félix, F., Echegaray, M., Siciliano, S., Di Benedetto, A.P., Flach, L., Viddi, F., Avila, I.C., Herrera, J.C., Tobón, I.C., Bolaños, J., Moreno, I.B., Ott, P.H., Sanino, G.P., Castineira, E., Montes, D., Crespo, E., Flores, PAC, Haase, B., Mendonça de Souza, S.M.F., Laeta, M. and Fragoso, A.B. 2007a. A preliminary overview of skin and skeletal diseases and traumata in small cetaceans from South American waters. *LAJAM* 6 (in press)
- Van Bresseem, M.-F., Cassonnet, P., Rector, A., Desaintes, C., Van Waerebeek, K., Alfaro-Shigueto, J., Van Ranst, M and Orth, G. 2007b. Genital warts in Burmeister's porpoises: characterization of *Phocoena spinipinnis* papillomavirus type 1 (PsPV-1) and evidence for a second, distantly related PSPV. *J Gen Virol* 88: 1928-1933.
- Van Bresseem, M.-F., Van Waerebeek, K., Raga, J.A., Jepson, P.D., Duignan, P., Deaville, R., Flach, L., Viddi, F., Baker, J.R., Di Benedetto, A.P., Echegaray, M., Genov, T., Reyes, J.C., Felix, F., Gaspar, R., Ramos, R., Peddemors, V. and Siebert, U. 2008a. Epidemiology of tattoo skin disease worldwide: a general health indicator for cetaceans and their environment? *Dis Aquat Org* (submitted).
- Van Bresseem, M.-F., Santos M, Faria Oshima, J. 2008b. Skin diseases in marine tucuxis (*Sotalia guianensis*) from the Parana estuary, Brazil: A possible indicator of a compromised marine environment. *Mar Environm Res* (submitted).
- Van Bresseem, M.-F., Raga, J.A., di Guardo, G., Jepson, P., Duignan, P., Siebert, U., Barrett, T., Santos, M.C.d.O, Moreno, I., Siciliano, S., Aguilar, A. and Van Waerebeek, K. 2008c. Emerging and recurring diseases in cetaceans worldwide and the role of environmental stressors. Paper SC/60/DW5 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal].
- Varaldo, P.E., Kilpper-Baltz, R., Biaviaco, F., Satta, G. and Schleifer, K.H. 1988. *Staphylococcus delphini* sp. nov. A coagulase positive species isolated from dolphins. *Int J Syst Bacteriol* 38: 436-439.
- Wilson, B., Arnold, H., Berazi, G., Fortuna, C.M., Gaspar, R., Ingram, S., Liret, C., Pribanic, S., Read, A.J., Ridoux, V., Schneider, K., Unrian, K.W., Wells, R.S., Wood, C., Thompson, P.M. and Hammond, P.S. 1999. Epidermal lesions in bottlenose dolphins: impacts of natural and anthropogenic factors. *Proc R Soc Lond B*. 266: 1077-1083

FIGURES



Figure 1. Black dots, perceptible by the touch in a *Lagenorhynchus obscurus* by-caught in December 1993 off central Peru.




Figure 2. Papillomavirus skin infection in a *Phocoena phocoena* stranded in Germany in December 1993.



Figure 3. Tattoo lesions in a *Lagenorhynchus obscurus* by-caught in June 1994 off central Peru.



Figure 4. Whitish, velvety lesions in a *Tursiops truncatus* from the Bay of Paracas, Peru, in July 2006.



Figure 5. Mycotic disease in a *Tursiops aduncus* from Plettenberg Bay, South Africa in December 2004.



Figure 6. Lobomycosis-like disease in a *Sotalia guianensis* from the Paraná estuary, Brazil in February 2007.