

By-catch in a tropical shrimp fishery: are TEDs and BRDs effective in excluding elasmobranchs?


Tomas Willems^{1,2}, Kris Hostens¹ & Magda Vincx²

¹Institute for Agricultural and Fisheries Research - Biological Environmental Research

²Ghent University, Department of Biology, Marine Biology Section

INTRODUCTION

TROPICAL SHRIMP FISHERIES are highly wasteful. Using fine-meshed demersal trawl nets, they generate a third of the world's fisheries by-catch, mostly thrown back overboard dead or dying. Technical gear adaptations are often applied to tackle this problem. A **TED** (Turtle Excluder Device) excludes large organisms, mainly seaturtles, from the net while a **BRD** (Bycatch Reduction Device) reduces by-catch of small fish. **ELASMOBRANCHS** (Chondrichthyes) are cartilaginous fish including rays and sharks. In general, their reproductive biology makes them highly vulnerable to overfishing: many species occurring on tropical shrimp fishing grounds are **IUCN red listed**. A **CASE STUDY** on trawling for Atlantic seabob-shrimp *Xiphopenaeus kroyeri* in Suriname is presented here. We assessed the effectiveness of existing gear adaptations (TED and BRD) in reducing elasmobranch by-catch.




Trawl codend with gear modifications: TED and BRD



By-catch of rays

DATA COLLECTION

- Commercial shrimp trawler
- Offshore Suriname
- 34 comparative hauls of 1 h
- February – July 2012
- All elasmobranchs identified and measured




MAJOR FINDINGS

8 SPECIES of rays and sharks occur in by-catch. Overall, TED and BRD cause a **40%-REDUCTION** in elasmobranch by-catch. This reduction is mainly caused by the exclusion of **3 LARGE RAY SPECIES** which are commonly caught (83% of by-catch). Hence, by-catch is the result of the **TED** and **NOT BRD**, through which only small-bodied species could possibly escape. **ESCAPE THROUGH THE TED** is highest for *D. geijskesi* and lowest for *G. micrura*. Escape through the TED is size-dependent.

Species	By-catch with TED and BRD
All species	- 40.0 % *
<i>Dasyatis guttata</i>	- 51.5 % *
<i>Dasyatis geijskesi</i>	- 69.7 % *
<i>Gymnura micrura</i>	- 38.6 % *

* Exact binomial test, $p < 0.001$

COMPOSITION OF ELASMOBRANCH BY-CATCH




CONCLUSIONS

A TED prevents **LARGE-BODIED RAYS** from being captured. It is least effective, however, for the most abundant ray in by-catch, *G. micrura*.

SURVIVAL AFTER ESCAPE through TED remains completely unknown.

TEDs nor BRDs exclude **SMALL-BODIED ELASMOBRANCHS**. Development of new devices reducing by-catch of small and juvenile elasmobranchs is desirable, since they make up a major part of the total by-catch in the Suriname seabob trawl fishery.



Exclusion rates at different sizes for three ray species : output of logistic regression model on body width measurements. \times 's on x-axis indicate individual measurements (n = 548, 86 and 1033 respectively, Av. = average body width)