

# APPARENT OPTICAL PROPERTIES OF THE AEROSOLS FOR MERIS/OLCI-S3 - SELECTION OF BEST LUTS FOR ATMOSPHERIC CORRECTION OVER OCEAN -

Francis Zagolski<sup>(1,2)</sup>, Richard Santer<sup>(1,4)</sup>, and Ouahid Aznay<sup>(1,3)</sup>

<sup>(1)</sup> ADRINORD, Association pour le Développement de la Recherche et de l'Innovation dans le bassin du Nord-Pas-de-Calais, 2 rue des Canoniers, F59000, Lille - FRANCE.  
E-mail: Santer.Richard@yahoo.fr

<sup>(2)</sup> PARBLEU Technologies Inc., 79 Veilleux street, St Jean-sur-Richelieu (QC), J3B-3W7 - CANADA.  
E-mail: Francis\_Zagolski@yahoo.ca

<sup>(3)</sup> C-S Systèmes d'Information, ZAC de la Grande Plaine, F31506, Toulouse - FRANCE.  
E-mail: Ouahid.Aznay@c-s.fr

<sup>(4)</sup> Université du Littoral Côte d'Opale, MREN, 32 Avenue Foch, F62930, Wimereux - FRANCE.  
E-mail: Santer@univ-littoral.fr

## INTRODUCTION

In the aerosol remote sensing algorithm over ocean, the models are deduced from the spectral dependence of the path radiances in the near-infrared (NIR) region. Based on these derived optical properties, the atmospheric signal is then extrapolated from the NIR to the shorter wavelengths in the visible (VIS) domain. To perform such a classical atmospheric correction scheme requires a well defined climatology representative of the aerosols encountered in remote areas of oceans.

Different climatologies are candidate to be implemented as auxiliary data (*i.e.*, look-up tables) including radiative properties of the aerosols. In the MERIS (Medium Resolution Imaging Spectrometer) atmospheric correction algorithm over ocean, we will consider: (i) the standard aerosol models (SAMs) [1] used in the 3<sup>rd</sup> MERIS reprocessing, (ii) the new set of AERONET (Aerosol Robotic Network) derived models [2] characterized by their micro-physical properties, and (iii) the new set of AERONET derived models by WOPAE [3] described through their inherent optical properties (IOPs) [4]. All sets are processed with the MERIS Auxiliary data Tool (MERISAT) for generating these LUTs for the MEGS (MERIS-ESA Ground Segment) prototype.

By using ODESA (Optical Data processor of ESA), the MERIS level-1 data extracted from MERMAID (MERIS Matchup In-situ Database) will be processed to get the level-2 data. AAOT (Acqua Alta Oceanographic Tower, Venice - Italy) being a reference station for MERIS Cal/Val activities, this site will be then selected to evaluate the MERIS level-2 aerosol product as well as the retrieved marine reflectance, obtained for each of the three sets of aerosol LUTs.

## THE AEROSOL INHERENT OPTICAL PROPERTIES

The IOPs, required in 13 MERIS bands, are the extinction coefficient, the single scattering albedo (SSA) and phase functions  $P_p$ . For the SAMs, the IOPs are computed with the Mie's theory using the micro physical properties of the aerosols. For all the other IOPs, coming from AERONET, they are known in three spectral bands (440, 675 and 870 nm). This first step for the AAOT site is fully described in a companion poster [5].

These IOPs are then distributed in 16 classes which are organized around the values of the Angstrom exponent ( $\alpha$ ) computed in the NIR region between 675 and 870 nm. The first class starts at  $\alpha=0$  and the last one is centred at  $\alpha=2.25$ , by using a step of 0.15 to get the others. For each class, we computed the mean, the minimum and the maximum values of at one sigma. A spectral interpolation has been finally achieved to get these IOPs at the 15 nominal MERIS wavelengths.

At the end, we get 8 sets of IOPs. Both for the SAMs [1] and the IOPA [4], these 2 sets are at a global scale. For the AAOT site, we refer the sets of IOPs as provided by AERONET [5] as AERO\_mean, AERO\_min and AERO\_max, and for the sets of IOPs derived by WOPAE we call them as IOPA\_mean, IOPA\_min and IOPA\_max.




Fig.1: Aerosol phase function at 870 nm versus the scattering angle for the 8 aerosol models and the class #11.

The performance of the atmospheric correction (AC) has not to be judged on the retrieval of the AOT but on the possibility to predict  $\rho_p(\lambda, \lambda_0) = \frac{\log(\rho_p(\lambda)/\rho_p(\lambda_0))}{\log(\lambda/\lambda_0)}$

the spectral dependency of  $\rho_p$  from the NIR to the blue region. More exactly, we extract the aerosol reflectance  $\rho_p$  in the NIR and we want to know it in the blue. In the primary scattering approximation,  $\rho_p$  is proportional to  $\tau_p P_p$ . Figure 2 gives the multiplicative factor we need to apply to  $\rho_p$  in B13 to get  $\rho_p$  in B2. Clearly the R70 model over estimates the atmospheric correction while the models derived from AERONET well compare.




Fig.2: Ratio of MERIS B2/B13 versus the scattering angle for the 8 aerosol models.

## THE AOP IN THE MERIS AUXILIARY DATA FILE(ADF)

The successive order of scattering (SO) code has been used to produce the aerosol apparent optical properties (AOPs): path radiances and transmittances for all of the 8 aerosol families in the MERIS bands for a set of 17 AOT550's (from 0 to 0.8 by a step of 0.05). The SO outputs have then been reformatted as auxiliary data files (ADFs) compatible with MEGS and ODESA.

## THE ODESA -MERMAID PROCESS

The MERIS-RR L1 products are available in MERMAID. They correspond to the standard extraction of 5x5 pixels window around the AAOT. ODESA was applied on this L1 data set with the 8 new aerosol ADFs and without vicarious adjustment. The L2 outputs were merged with the MERMAID data to produce a new file which includes the *in situ* data and replaces the L2 products and break points by the ODESA outputs. This procedure was applied on all the 25 pixels of the window.

We then averaged the L1 and L2 on the 25 pixels keeping the flag setting used for the validation: no land, no cloud, no ice-haze, no white scatters, no glint, no PCD 13, no PCD 19. We also computed the rms values.

At the end, 126 matchups were selected. Table 1 displays a comparison between AOT measurements and MERIS MEGS-v8.0 retrieved values. First, the size of the aerosols is representative of coastal aerosols and not maritime ones. The spectral dependency in NIR is comparable because the MERIS aerosol reflectance does not depend much on the aerosol model. The dispersion on  $\alpha$  is higher with AERONET because of its inaccurate determination at small AOTs. MERIS overestimates the AOT.

	AERONET	MERIS
	AOT_870	ALP(670,870) TAU_AER_13 ALPHA
mean	0.079	1.438 0.119 1.426
rms	0.050	0.863 0.061 0.343

Table 1: Comparison between AERONET and MEGS: AOT at 870 nm and  $\alpha$ : in mean and rms.

## Acknowledgments

We thank G. Zibordi for the use of the AAOT AERONET data, the MERMAID and ODESA team. This work was partially supported by the INTERREG 2 Seas program.



## EVALUATION OF THE AEROSOL PRODUCT

The aerosol product is the AOT at 865 nm and  $\alpha(779,865)$ . As a break point, the AOT550 is proposed. We generated the L2 product for the SAMs, IOPAs and the 3 regional AEROs (mean, min, max). The 3 regional IOPAs are similar to the AEROs.

The AERONET measurements correspond to the AOT at 440, 675 and 870 nm for 2 sets of measurements which bracket the time of the MERIS overpass. The AERONET data is matched with the MERIS AOT with: (i) a time interpolation at the time of MERIS overpass, (ii) a spectral interpolation on the AOT between 675 nm and 870 nm gives the AOT778 and AOT865, and (iii) a spectral interpolation on the AOT between 440 and 675 nm yields to the AOT550.




Fig.3: AOT at 3 wavelengths. Comparison between in situ and MEGS-v8.0 for the SAMs.

	SAM	IOPA global	AERO mean	AERO min	AERO max
N	113	113	113	113	113
mean	1.02	1.37	1.38	1.35	1.45
min	1.00	0.71	0.68	0.68	0.67
max	1.04	1.68	1.68	1.68	1.68
ratio	1.12	2.48	2.13	2.13	2.13
sigma	0.13	0.14	0.23	0.23	0.23

Table 2: Number of matchups, mean ratio and sigma on the AOT between MERIS and AERONET at 865 nm (upper set) and 440 nm (lower set). A total of 5 different sets of aerosol ADFs are reported.

Clearly, with the SAMs MEGS overestimate the AOT both in the NIR and blue regions. Results are better with the aerosol models derived from AERONET both on the mean and on the dispersion, even if they are far to be perfect.

## THE SPATIAL VARIABILITY OF THE AEROSOL PRODUCT

One indicator of the quality of the selected aerosol family is its ability to do not amplify the natural spatial variability of the signal. The inputs to aerosol remote sensing module are the aerosol reflectances at 779 and 865 nm. These values, obtained after gaseous correction and the BPAC, are independent on the aerosol ADF. The optical continuity of the aerosol models in a given family results in that the relative dispersion on the AOT should be comparable to the dispersion of the aerosol reflectance in MERIS B12 and B13.




Fig.4: Relative variation in the (5x5) pixels window of the AOT versus the relative variation of the aerosol reflectance in B13 for the SAMs and for the IOPAs.

## EVALUATION OF THE WATER REFLECTANCE RETRIEVAL

The water reflectance is measured in 5 spectral bands centred at 412, 443, 490, 560 and 665 nm. We compare the 5 MERIS outputs in these bands.




Fig.5: Water reflectance at 2 wavelengths. Comparison between in situ and MEGS-v8.0 for the SAMs.

The first comparison between *in situ* and MERIS water reflectances is achieved in B2 and B5 with MEGS and SAMs without vicarious adjustment.

The MERIS water reflectance is above the *in-situ* one in B2 and very scattered. Results in B5 appear as more acceptable.

The choice of the criterion to intercompare the results is controversial. For each of the ADF, we report in this table, the number of matchups, the mean and the rms of the MERIS / *in-situ* ratio of the water reflectance in B2 and B5. We also display the difference *in situ* - MERIS. The salient point is the high dispersion which illustrates the difficulty to make the AC in the coastal region as well to interpret these results.

	ratio	difference
	B2	B5
SAM	1.097	0.943
129	0.354	0.333
IOPA	0.878	1.073
129	0.358	0.459
AERO_mean	1.645	0.778
123	0.527	0.411
AERO_min	1.646	0.732
134	0.540	0.748
AERO_max	1.617	0.767
124	0.523	0.396

## Conclusion

We reported here for the AAOT data set a preliminary study on the influence of the aerosol models on some of the MERIS L2 products. What we clearly see for all the ADF candidates is a large dispersion on the AOT between *in situ* and MERIS. This dispersion is explained by the natural variability of the aerosol IOPs [4] but clearly this dispersion is reduced when the aerosol models are defined locally from their IOPs as for the AERO models. The spatial homogeneity is driven by the optical continuity of the aerosol models versus  $\alpha$ , and the SAMs are the worst. For the water reflectance retrieval, we have no firm conclusion because of large error bars. With the algorithm as it is, we need to better identify the aerosol model and the only small hope we have is to use the relative humidity. Alternatively a combined retrieval of the aerosols and of the water colour is suggested.

This study needs to be consolidated and extended, starting by the AERONET Ocean Colour network. We need to process a large amount of matchups to be able to conduct an analysis for each type of aerosols (oceanic and coastal) and for different scattering angle domains. It is clear than compared to MERIS, OLCI will go in the backscattering, a difficult region in terms of the variability of the IOPs.

## References

- [1] Antoine, D., and A. Morel, 1999, "A multiple scattering algorithm for atmospheric correction of remotely sensed ocean color (MERIS instrument): Principle and implementation for atmospheres carrying various aerosols including absorbing ones", *International Journal of Remote Sensing*, 20, 1875-1916.
- [2] Dubovik, O., and M. D. King, 2000, "A flexible inversion algorithm for retrieval of aerosol optical properties from Sun and sky radiance measurements", *Journal of Geophysical Research*, 105: 20673-20696.
- [3] Santer, R., F. Zagolski, and O. Aznay, 2009, "Aerosol phase function derived from CIMEL measurements", *International Journal of Remote Sensing*, 31 (4), 969-992.
- [4] Zagolski, F., R. Santer, and O. Aznay, 2007, "A new climatology for atmospheric correction based on the aerosol inherent optical properties", *Journal of Geophysical Research*, vol. 112, no. D14.
- [5] Santer, R., F. Zagolski, and O. Aznay, 2012, "Inherent optical properties of the aerosols for MERIS/OLCI-S3: From micro-physical properties to optical properties", *Proceedings of MERIS/ATSR & OLCI/SLSTR Preparatory Workshop*, Frascati (Italy), 15-19 October, 2012.