

Check-list of Cnidaria and Ctenophora from the coasts of Turkey

Melih Ertan ÇINAR^{1*}, Mehmet Baki YOKES², Şermin AÇIK³, Ahmet Kerem BAKIR¹

¹Department of Hydrobiology, Faculty of Fisheries, Ege University, Bornova, İzmir, Turkey

²Department of Molecular Biology and Genetics, Haliç University, Şişli, İstanbul, Turkey

³Institute of Marine Sciences and Technology, Dokuz Eylül University, İnciraltı, İzmir, Turkey

Received: 28.05.2014

Accepted: 13.08.2014

Published Online: 00.00.2013

Printed: 00.00.2013

Abstract: This paper presents the actual status of species diversity of the phyla Cnidaria and Ctenophora along the Turkish coasts of the Black Sea, the Sea of Marmara, the Aegean Sea, and the Levantine Sea. A total of 195 cnidarian species belonging to 5 classes (Hydrozoa, Cubozoa, Scyphozoa, Staurozoa, and Anthozoa) have been determined in these regions. Eight anthozoan species (*Arachnanthus oligopodus*, *Bunodactis rubripunctata*, *Bunodeopsis strumosa*, *Corynactis viridis*, *Halcampoides purpureus*, *Sagartiogeton lacerates*, *Sagartiogeton undatus*, and *Pachycerianthus multiplicatus*) are reported for the first time as elements of the Turkish marine fauna in the present study. The highest number of cnidarian species (121 species) was reported from the Aegean Sea, while the lowest (17 species) was reported from the Black Sea. The hot spot areas for cnidarian diversity are the Prince Islands, İstanbul Strait, İzmir Bay, and Datça Peninsula, where relatively intensive scientific efforts have been carried out. Regarding ctenophores, 7 species are distributed along the Turkish coasts, 5 of which were reported from the Black Sea. A total of 16 alien cnidarian and 2 ctenophore species were determined in the regions. Two species (*Sagartiogeton laceratus* and *Pachycerianthus multiplicatus*) are new alien species for the Mediterranean Sea and could have been introduced to the northern part of the Sea of Marmara and İskenderun Bay, areas from which these species are recorded, by ships from the North-East Atlantic.

Key words: Species diversity, cnidarian, ctenophore, alien species, protected species, Mediterranean Sea, Black Sea

1. Introduction

Cnidaria is a diverse phylum containing 10,211 species worldwide and 752 species in the Mediterranean Sea, inhabiting exclusively aquatic and mostly marine environments (Coll et al., 2010; Appeltans et al., 2012). Many cnidarians are distributed in shallow waters as they are in a symbiotic relationship with endosymbiotic algae for food and calcification (Pearse and Muscatine, 1971; Muscatine and Porter, 1977). They form reefs in tropical waters, constituting a highly productive ecosystem (Hatcher, 1997). Blooms of jellyfish suddenly appearing in the anthropogenically deteriorated areas are known to be an indication of greatly altered ecosystem function (Parsons and Lalli, 2002; Purcell et al., 2007). They have also direct effects on human enterprise, particularly obscuring tourism by stinging swimmers (Öztürk and İşinibilir, 2010; Çınar et al., 2011), fishing and aquaculture by clogging nets and feeding zooplankton and ichthyoplankton (Richardson et al., 2009; Dong et al., 2010), and power plants by clogging seawater-intake pipes (Galil, 2007).

The jellyfish bloom along the coast of Turkey was first reported by Antipa (1892) in İzmir Bay for *Drymonema*

dalmatinum (cited as *D. cordelio*). Afterwards, blooms of alien (*Rhopilema nomadica*) and native (*Aurelia aurita*, *Chrysaora hysoscella*, and *Rhizostoma pulmo*) jellyfish have been frequently observed along the coasts of Turkey (Balık, 1972; Bingel et al., 1991; Mutlu et al., 1994; Alparslan, 2001; Çınar et al., 2011). More recently, blooms of *Liriope tetraphylla*, a hydromedusa species, caused a temporal regime shift from a crustacean- to a jellyfish-controlled system in the Sea of Marmara (Yilmaz, 2014).

The naturalist and diplomat Luigi Ferdinando Marsili, who provided the first scientific understanding of the dynamics of the exchange currents through the Turkish Straits System, was the first to observe and draw a coral species (probably *Savalia savaglia*) in the Sea of Marmara in 1691 (Özsoy and Pinardi, 2010). Scientifically, the first reports of anthozoan species were given by Forbes (1844), who said: "The only corals met with of any size were *Cladocora caespitosa* and *Porites daedalea*. The former is extremely abundant near water-mark on the coast of Asia Minor, where it forms elegant cauliflower-like patches of bright orange, from the hue of the animals, adhering to the rocks". The species *Porites daedalea* was then considered

* Correspondence: melih.cinar@ege.edu.tr

as a misidentification of *Madracis pharensis* by Zibrowius (1980). Colombo (1885) and Ostroumoff (1896) listed 5 and 15 anthozoan species in the Çanakkale Strait and Sea of Marmara, respectively. Demir (1952) identified 14 species among the material collected from the İstanbul Strait and Prince Islands. Tixier-Durivault (1961) described a new species (*Parerythropodium bosphorense*) in the İstanbul Strait. This group of animals was first specifically studied by Coşar (1974), who reported 11 species in İzmir Bay. More recently, Topçu and Öztürk (2013) identified 9 octocoral species from Balıkcı Island (Sea of Marmara). Other records of anthozoan species were given in some general ecological works (i.e. Çinar et al., 2006b) or a species atlas (i.e. Gökalp, 2011).

The phylum Ctenophora, also known as comb jellies, is mainly characterized by bearing 8 rows of comb-like bands of cilia used for swimming. This phylum includes almost 190 species worldwide and 30 species in the Mediterranean Sea (Coll et al., 2010; Appeltans et al., 2012). The first record of ctenophores from the Turkish coast was given by Demir (1952), who reported *Pleurobrachia rhodopsis* around the İstanbul Strait and Prince Islands. This species was later found off Trabzon (Demir, 1954). The well-known ctenophore species in the region is *Mnemiopsis leidyi*, which was unintentionally introduced to the Black Sea from the northwestern Atlantic by ballast water of ships (Vinogradov et al., 1989). This zooplanktonic predator reached an enormous biomass level ($>1 \text{ kg m}^{-2}$) in some areas in the summer of 1989, which was estimated to be greater than that of all other zooplankton combined, leading to a sharp decrease in anchovy production in the region (Kideys and Niermann, 1994; Kideys, 2002). After the arrival of the other alien species *Beroe ovata*, which feeds on other ctenophores, the sharpest drop was estimated for biomass and density of *Mnemiopsis leidyi* in 1999 compared with late summer data during the previous 10 years (Shiganova et al., 2001).

The present study gives a check-list of cnidarian and ctenophore species reported from the coasts of Turkey and presents some anthozoan species new to the Turkish marine fauna. Finally, the hot spot diversity areas for these major taxonomic groups are presented.

2. Materials and methods

The check-list was prepared by compiling all existing scientific papers about Cnidaria and Ctenophora species along the coasts of Turkey. In addition, new records of anthozoan species, which were identified in recent research projects but have not been cited in any papers, are presented here. The stations where new records of species were found are indicated in Figure 1. The station numbers are superscripted over “PS” in the Table. The first reports of all species and their depth and habitat distributions are given in the Table. Notes regarding the recorded species names cited in the literature considering the Turkish marina fauna are also provided in the Table.

In order to assess the diversity hot spots for the taxonomic groups considered in this study and to identify the areas where different degrees of research efforts — a major factor apart from environmental characteristics regulating diversity patterns — have been performed to date (gap analysis), the coasts of Turkey were divided into grids with squares of $15 \times 15 \text{ km}$. All data of species distribution were entered into an Excel file and then imported and digitized with ArcGIS 9.3 software.

3. Results and discussion

According to previous data and new data presented in this paper, the phylum Cnidaria along the coasts of Turkey is represented by 195 species belonging to 5 classes (Hydrozoa, Cubozoa, Scyphozoa, Staurozoa, and Anthozoa) and 76 families. The phylum Ctenophora along the coasts of Turkey is represented by 7 species belonging to 2 classes (Nuda and Tentaculata) and 4




Figure 1. Map of the stations where new records of anthozoan species were found.

Table. Species list of the phyla Cnidaria and Ctenophora from the coasts of Turkey *: Alien species; BS: Black Sea; SM: Sea of Marmara; AS: Aegean Sea; LS: Levantine Sea; DR: depth range (I: 0–10 m; II: 11–50 m; III: 51–100 m; IV: 101–200 m; V: 201–400 m; VI: 401–600 m; VII: >600 m); H: habitat (Hs: hard substratum – including algae, sponge, mussels, etc.; Ss: soft substratum – including all phanerogams; P: pelagic); PS: present study (superscripted numbers indicate station numbers as shown in Figure 1).

Group/Species	BS	SM	AS	LS	DR	H	Remarks
Phylum: Cnidaria							
Class: Hydrozoa							
Subclass: Trachylinae							
Family: Geryoniidae							
<i>Geryonia proboscidalis</i> (Forsskål, 1775)	-	-	70	1		P	
<i>Liriope tetrphylla</i> (Chamisso & Eysenhardt, 1821)	-	55	-	-		P	
Family: Rhopalonematidae							
<i>Rhopalonema velatum</i> Gegenbaur, 1857	-	-	13	-		P	
<i>Aglaura hemistoma</i> Péron & Lesueur, 1810	-	5	61	16	II, III	P	
Family: Olindiidae							
<i>Olindias phosphorica</i> (Delle Chiaje, 1841)	-	-	58	-	I	P	
Family: Aeginidae							
<i>Solmundella bitentaculata</i> (Quoy & Gaimard, 1833)	-	55	-	-		P	
Subclass: Hydroidolina							
Family: Forskaliidae							
<i>Forskalia contorta</i> (Milne-Edwards, 1841)	-	-	-	1		P	
Family: Abylidae							
<i>Abylopsis tetragona</i> (Otto, 1823)	-	-	1	1		P	
Family: Diphyidae							
<i>Muggiaea kochii</i> (Will, 1844)	-	5	-	-	I–VI	Hs	
<i>Lensia conoidea</i> (Keferstein & Ehlers, 1860)	10	-	40	-	II	P	
<i>Lensia subtiloides</i> (Lens & Van Riemsdijk, 1908)	-	-	16	16		P	
<i>Eudoxoides spiralis</i> (Bigelow, 1911)	-	-	13	16		P	
Family: Cladonematidae							
<i>Cladonema radiatum</i> Dujardin, 1843	-	4	-	-	II	Hs	
Family: Pennariidae							
<i>Pennaria disticha</i> Goldfuss, 1820	-	-	-	20	I	Hs	
Family: Porpitidae							
<i>Porpita porpita</i> (Linnaeus, 1758)	-	-	-	1		P	= <i>Porpita glandifera</i>
Family: Cladocorynidae							
<i>Cladocoryne floccosa</i> Rotch, 1871	-	-	19	-	I	Hs	
Family: Corynidae							
<i>Sarsia tubulosa</i> (M. Sars, 1835)	-	8	-	-	I	Hs	= <i>Sarsia sarsii</i> = <i>Syncoryne sarsii</i>
* <i>Coryne eximia</i> Allman, 1859	-	8	-	-	I	Hs	
<i>Coryne muscoides</i> (Linnaeus, 1761)	-	-	19	-	I	Hs	

Table 1. (Continued).

Family: Tubulariidae							
<i>Tubularia indivisa</i> Linnaeus, 1758	-	4	-	-	II	Hs, Ss	
<i>Ectopleura crocea</i> (Agassiz, 1862)	-	-	15	-	I	Hs	= <i>Tubularia</i>
<i>Ectopleura larynx</i> (Ellis & Solander, 1786)	-	8	19	-	I	Hs	<i>mesembryanthemum</i>
Family: Cordylophoridae							
* <i>Cordylophora caspia</i> (Pallas, 1771)	-	-	52	-	I	Hs	
Family: Bougainvilliidae							
<i>Bougainvillia muscus</i> (Allman, 1863)	14	2	19	20	I–III	Hs	= <i>Eudendrium ramosum</i> Van Beneden, 1844 = <i>Bougainvillia fruticosa</i>
<i>Rhizorhagium arenosum</i> (Alder, 1862)	-	8	-	-	I	Hs	
<i>Garveia grisea</i> (Motz-Kossowska, 1905)	-	-	20	-		Hs	
<i>Garveia nutans</i> Wright, 1859	14	-	-	-		Hs	
<i>Dicoryne conferta</i> (Alder, 1856)	-	24	-	-	I	Hs	
Family: Hydractiniidae							
<i>Hydractinia carnea</i> (M. Sars, 1846)	-	8	-	-	I	Hs	
<i>Hydractinia echinata</i> (Fleming, 1828)	-	5	-	-	I, II	Hs	
<i>Podocoryna carnea</i> M. Sars, 1846	-	5	-	-	II	Hs	
<i>Stylactaria fucicola</i> (M. Sars, 1857)	-	-	20	-		Hs	
Family: Eudendriidae							
<i>Eudendrium armstrongi</i> Stechow, 1909	-	5	-	-	II	Hs	= <i>Eudendrium ramosum</i> Armstrong, 1879
<i>Eudendrium capillare</i> Alder, 1856	-	8	19	-	I–III	Hs	
<i>Eudendrium glomeratum</i> Picard, 1952	-	-	20	-		Hs	
* <i>Eudendrium merulum</i> Watson, 1985	-	39	39	-	I	Hs	
<i>Eudendrium racemosum</i> (Cavolini, 1785)	-	-	20	-	I, II	Hs	
<i>Eudendrium rameum</i> (Pallas, 1766)	-	8	-	-	II	Hs	
<i>Eudendrium simplex</i> Pieper, 1884	-	-	20	-		Hs	= <i>Eudendrium</i> <i>motzkossowskiae</i>
Family: Pandeidae							
<i>Leuckartiara octona</i> (Fleming, 1823)	-	8	-	-	I–III	Hs	= <i>Perigonimus repens</i>
<i>Neoturris pileata</i> (Forsskal, 1775)	-	55	-	-		P	
Family: Halopterididae							
<i>Halopteris catharina</i> (Johnston, 1833)	-	-	20	-	I, II	Hs	
<i>Halopteris diaphana</i> (Heller, 1868)	-	-	20	-		Hs	
<i>Antennella secundaria</i> (Gmelin, 1791)	-	5	20	-	II, III	?Hs	
Family: Plumulariidae							
<i>Plumularia obliqua</i> (Johnston, 1847)	-	-	24	-	I	Hs	
<i>Plumularia setacea</i> (Linnaeus, 1758)	-	-	20	-		Hs	
<i>Plumularia syriaca</i> Billard, 1931	-	24	-	6	II, III	Hs	
<i>Nemertesia antennina</i> (Linnaeus, 1758)	-	2	24	-	I–IV	Hs	= <i>Nemertesia pentasticha</i>
<i>Nemertesia ramosa</i> (Lamarck, 1816)	-	2	-	-	III	Hs	

Table 1. (Continued).

<i>Nemertesia tetrasticha</i> Meneghini, 1845	-	5	-	-	III	Hs	= <i>Plumularia disticha</i>
Family: Aglaopheniidae							
<i>Aglaophenia dichotoma</i> Kirchenpauer, 1872	-	11	-	-	II	Hs	
<i>Aglaophenia elongata</i> Meneghini, 1845	-	5	20	-	II, III	Hs	
<i>Aglaophenia octodonta</i> Heller, 1868	-	11	20	-	II	Hs	
<i>Aglaophenia picardi</i> Svoboda, 1979	28	-	-	-	II	Hs	
<i>Aglaophenia pluma</i> (Linnaeus, 1758)	14	4	19	-	I, II	Hs	= <i>Plumularia cristata</i>
* <i>Macrorhynchia philippina</i> Kirchenpauer, 1872	-	-	-	51	I	Hs	
<i>Lytocarpia myriophyllum</i> (Linnaeus, 1758)	-	24	24	-	II-IV	Hs	
Family: Kirchenpaueriidae							
<i>Kirchenpaueria halecioides</i> (Alder, 1859)	-	8	24	-	I	Hs	
<i>Kirchenpaueria pinnata</i> (Linnaeus, 1758)	-	-	19	6	I	Hs	= <i>Kirchenpaueria echinulata</i>
Family: Haleciidae							
<i>Halecium beanii</i> (Johnston, 1838)	-	24	24	-	I-III	Hs	
<i>Halecium halecinum</i> (Linnaeus, 1758)	-	11	24	-	I-IV	Hs	= <i>Halecium halecium</i>
<i>Halecium labrosum</i> Alder, 1859	-	37	-	-	III	Hs	
<i>Halecium lankesterii</i> (Bourne, 1890)	-	-	20	-		Hs	= <i>Halecium lankesteri</i>
<i>Halecium pusillum</i> Sars, 1856	-	-	19	-	I	Hs	
<i>Halecium mediterraneum</i> Weismann, 1883	-	-	20	-		Hs	
<i>Halecium nanum</i> Alder, 1859	-	-	20	-		Hs	
<i>Hydrodendron mirabile</i> (Hincks, 1866)	-	-	19	-	I	Hs	= <i>Ophiodissa mirabilis</i>
Family: Sertulariidae							
<i>Sertularia distans</i> (Lamouroux, 1816)	-	-	24	-	II	Hs	= <i>Sertularella fusiformis</i>
* <i>Sertularia marginata</i> (Kirchenpauer, 1864)	-	-	20	-		Hs	
<i>Sertularia perpusilla</i> Stechow, 1919	-	-	20	-		Hs	
<i>Sertularella crassicaulis</i> (Heller, 1868)	-	4	-	-	II-IV	Hs	
<i>Sertularella ellisii</i> (Deshayes & Milne Edwards, 1836)	-	4	19	-	I, II	Hs	
<i>Sertularella mediterranea</i> Hartlaub, 1901	-	-	19	6	I	Hs	
<i>Sertularella polyzonias</i> (Linnaeus, 1758)	7	5	20	-	I-III	Hs	
<i>Sertularella simplex</i> (Hutton, 1873)	-	-	24	-	I	Hs	= <i>Sertularella fusiformis</i>
<i>Salacia desmoides</i> (Torrey, 1902)	-	-	20	-		Hs	
<i>Dynamena disticha</i> (Bosc, 1802)	-	-	20	20		Hs	
<i>Amphisbetia operculata</i> (Linnaeus, 1758)	-	-	24	-	I	Hs	= <i>Dynamena operculata</i>
Family: Aequoreidae							
* <i>Aequorea globosa</i> Eschscholtz, 1829	-	-	-	65	II	P	
<i>Aequorea forskalea</i> Péron & Lesueur, 1810	-	-	-	71	I, II	P	
Family: Phialellidae							
<i>Phialella quadrata</i> (Forbes, 1848)	-	8	-	-	I	Hs	= <i>Campanulina repens</i>
Family: Campanulariidae							

Table 1. (Continued).

<i>Campanularia hincksii</i> Alder, 1856	-	-	20	-		Hs	= <i>Orthopyxis alta</i>
<i>Clytia gracilis</i> (Sars, 1850)	-	8	24	-	I, II	Hs	
<i>Clytia hemisphaerica</i> (Linnaeus, 1767)	-	4	19	20	I, II	Hs	= <i>Clytia johnstoni</i>
* <i>Clytia linearis</i> (Thorneley, 1900)	-	-	20	-		Hs	= <i>Clytia gravieri</i>
<i>Clytia paulensis</i> (Vanhöffen, 1910)	-	-	20	-		Hs	
<i>Obelia bidentata</i> Clark, 1875	-	8	24	-	I–III	Hs	= <i>Obelia biscuspidata</i>
<i>Obelia dichotoma</i> (Linnaeus, 1758)	-	8	36	6	I, II	Hs	
<i>Obelia geniculata</i> (Linnaeus, 1758)	-	-	19	-	I	Hs	
<i>Obelia longissima</i> (Pallas, 1766)	-	24	-	-	I	Hs	= <i>Obelia flabellata</i>
<i>Orthopyxis everta</i> (Clark, 1876)	-	-	20	-		Hs	= <i>Orthopyxis crenata</i>
<i>Orthopyxis integra</i> (MacGillivray, 1842)	-	24	20	-	I, II	Hs	= <i>Orthopyxis caliculata</i>
<i>Hartlaubella gelatinosa</i> (Pallas, 1766)	14	8	-	-	I, II	Hs	
<i>Gonothyraea loveni</i> (Allman, 1859)		8	-	-	I, II	Hs	= <i>Gonothyraea hyalina</i>
<i>Laomedea angulata</i> Hincks, 1861	14	8	24	-	I, II	Hs, Ss	
<i>Laomedea calceolifera</i> (Hincks, 1871)	-	38	20	-	I	Hs	
<i>Laomedea exigua</i> M. Sars, 1857	-	24	24	-	I, II	Hs	
<i>Laomedea flexuosa</i> Alder, 1857	-	5	24	-	I–III	Hs	
Family: Lafoeidae							
<i>Lafoea dumosa</i> (Fleming, 1820)	-	8	20	-	I–IV	Hs	= <i>Lafoea gracillima</i> = <i>Scandia pocillum</i>
<i>Filellum serpens</i> (Hassall, 1848)	-	24	-	-	II	Hs	
* <i>Filellum serratum</i> (Clarke, 1879)	-	24	20	-	II	Hs	
<i>Acryptolaria conferta</i> (Allman, 1877)	-	24	-	-	IV	Hs	
Family: Hebellidae							
<i>Hebella scandens</i> (Bale, 1888)	-	-	20	-		Hs	
<i>Anthohebella parasitica</i> (Ciamician, 1880)	-	5	20	-	III	Hs	
Family: Tiarannidae							
<i>Modeeria rotunda</i> (Quoy & Gaimard, 1827)	-	24	-	-	II–IV	Hs	= <i>Stegopoma fastigiata</i>
Family: Laodiceidae							
<i>Laodicea undulata</i> (Forbes & Goodsir, 1853)	-	-	20	-		Hs	
Class: Cubozoa							
Family: Carybdeidae							
<i>Carybdea marsupialis</i> (Linnaeus, 1758)	-	-	18	-	?	P	
Class: Scyphozoa							
Subclass: Discomedusae							
Family: Ulmaridae							
<i>Aurelia aurita</i> (Linnaeus, 1758)	9	8	13	60	I, II	P	
Family: Pelagiidae							
<i>Chrysaora hysoscella</i> (Linnaeus, 1767)	-	45	13	27	I, II	P	
<i>Pelagia noctiluca</i> (Forsskal, 1775)	-	-	13	60	I, II	P	

Table 1. (Continued).

Family: Cyaneidae						
<i>Drymonema dalmatinum</i> Haeckel, 1880	-	-	3	-	I, II	P
Family: Cassiopeidae						
* <i>Cassiopea andromeda</i> (Forsskål, 1775)	-	-	60	44	I	Ss, P
Family: Cepheidae						
<i>Cotylorhiza tuberculata</i> (Macri, 1778)	-	-	2	-	I, II	P
Family: Mastigiidae						
* <i>Phyllorhiza punctata</i> Lendenfeld, 1884	-	-	69	57	I	P
Family: Rhizostomatidae						
<i>Rhizostoma pulmo</i> (Macri, 1778)	35	53	1	27	I, II	P
* <i>Rhopilema nomadica</i> Galil, Spanier & Ferguson, 1990	-	-	66	33	I, II	P
Family: Periphyllidae						
<i>Nausithoe punctata</i> Kölliker, 1853	-	-	59	-	II	Hs
<i>Periphylla periphylla</i> (Péron & Lesueur, 1810)	-	5	-	-	I, II	P
Family: Paraphyllinidae						
<i>Paraphyllina ransoni</i> Russell, 1956	-	55	-	-	I, II	P
Class: Staurozoa						
Family: Kishinouyeidae						
<i>Lucernariopsis campanulata</i> (Lamouroux, 1815)	47	8	-	-	I	?Hs
Class: Anthozoa						
Subclass: Hexacorallia						
Family: Actiniidae						
<i>Actinia cari</i> Delle Chiaje, 1822	-	-	17	54	I–III	Hs
<i>Actinia equina</i> (Linnaeus, 1758)	41	8	15	48	I	Hs
<i>Anemonia viridis</i> (Forskål, 1775)	-	8	15	48	I, II	Hs = <i>Anemonia sulcata</i>
<i>Aulactinia verrucosa</i> (Pennant, 1777)	-	-	17	PS ⁶	I	Hs
<i>Bunodactis rubripunctata</i> (Grube, 1840)	-	-	PS ⁴	-	I	Hs
<i>Condylactis aurantiaca</i> (Delle Chiaje, 1825)	-	60	59	PS ¹⁰	I, II	Ss
Family: Actinostolidae						
<i>Paranthus rugosus</i> Andrès, 1880	-	-	50	-	I	Hs
Family: Aiptasiidae						
<i>Aiptasia diaphana</i> (Rapp, 1829)	-	-	17	PS ⁶	I	Hs
<i>Aiptasia mutabilis</i> (Gravenhorst, 1831)	-	60	56	PS ⁶	I, II	Hs
<i>Aiptasiogeton pellucidus</i> (Hollard, 1848)	-	PS ¹	50	-	I	Hs = <i>Aiptasiogeton comatus</i>
Family: Aliciidae						
<i>Alicia mirabilis</i> Johnson, 1861	-	-	59	60	I, II	Hs
Family: Arachnactidae						
<i>Arachnanthus oligopodus</i> (Cerfontaine, 1891)	-	-	-	PS ⁶	II	Ss
Family: Boloceroididae						

Table 1. (Continued).

<i>Bunodeopsis strumosa</i> Andrès, 1881	-	PS ³	PS ²	PS ⁶	I	Hs	
Family: Cerianthidae							
<i>Cerianthus lloydii</i> (Gosse, 1859)	-	59	-	-	I	Ss	Questionable
<i>Cerianthus membranaceus</i> (Spallanzani, 1784)	-	5	30	48	I–V	Ss	
<i>Pachycerianthus solitarius</i> (Rapp, 1829)	23	8	PS ⁵	PS ⁶	II	Ss	
* <i>Pachycerianthus multiplicatus</i> Carlgren, 1912	-	-	-	PS ¹⁰	I	Ss	
Family: Corallimorphidae							
<i>Corynactis viridis</i> Allman, 1846	-	PS ¹	-	-	II	Hs	
Family: Diadumenidae							
* <i>Diadumene lineata</i> (Verrill, 1869)	-	-	43	-	I	Hs	
* <i>Diadumene cincta</i> Stephenson, 1925	-	59	-	-	I	Hs	Questionable
Family: Edwardsiidae							
<i>Edwardsia clapedii</i> (Panceri, 1869)	23	-	50	67	I–III	Ss	
Family: Halcampoididae							
<i>Halcampoides purpureus</i> (Studer, 1879)	-	-	-	PS ⁶	I, II	Ss	
Family: Hormathiidae							
<i>Adamsia palliata</i> (Fabricius, 1779)	-	-	17	PS ^{6,8}	I, II	Hs	
<i>Calliactis parasitica</i> (Couch, 1842)	-	-	15	60	I, II	Hs	
Family: Epizoanthidae							
<i>Epizoanthus couchii</i> Johnston in Couch, 1844	-	59	-	-	II	Hs	
Family: Isophellidae							
<i>Telmatactis cricoides</i> (Duchassaing, 1850)	-	-	60	32	I	Hs	
<i>Telmatactis forskalii</i> (Ehrenberg, 1834)	-	-	19	PS ⁶	I	Hs	= <i>Phellia limicola</i>
Family: Haloclavidae				-			
<i>Peachia cylindrica</i> (Reid, 1848)	14	46	17	-	I, II	Ss	= <i>Peachia hastata</i>
Family: Sagartiidae							
<i>Actinothoe clavata</i> (Ilmoni, 1830)	23	-	-	-	III	Ss	
<i>Cereus pedunculatus</i> (Pennant, 1777)	-	59	17	-	I, II	Ss	
<i>Sagartia elegans</i> (Dalyell, 1848)	-	60	17	-	II	Hs	= <i>Sagartia rhododactylos</i>
* <i>Sagartiogeton laceratus</i> (Dalyell, 1848)	-	PS ¹	-	-	II	Hs	
<i>Sagartiogeton undatus</i> (Müller, 1778)	-	PS ¹	-	-	II	Hs	
<i>Sagartiogeton viduatus</i> (Müller, 1776)	-	5	-	-	II	Ss	
Family: Schizopathidae							
<i>Parantipathes larix</i> (Esper, 1788)	-	5	-	-	V–VII	Ss	= <i>Anthipathes larix</i>
Family: Parazoanthidae							
<i>Parazoanthus axinellae</i> (Schmidt, 1862)	-	5	48	48	II	Hs	
<i>Savalia savaglia</i> (Bertoloni, 1819)	-	25	PS ²	-	II, III	Hs	= <i>Gerardia savaglia</i>
Family: Dendrophyllidae							
<i>Balanophyllia</i> (<i>Balanophyllia</i>) <i>europaea</i> (Risso, 1826)	-	5	21	60	I–III	Hs, Ss	= <i>Balanophyllia italica</i>

Table 1. (Continued).

<i>Dendrophyllia ramea</i> (Linnaeus, 1758)	-	5	-	-	II	Ss	
<i>Leptopsammia pruvoti</i> Lacaze-Duthiers, 1897	-	-	59	60	II	Hs	
Family: Caryophyllidae							
<i>Caryophyllia (Caryophyllia) cyathus</i> (Ellis & Solander, 1786)	-	2	-	-	II, III	Hs	
<i>Caryophyllia (Caryophyllia) inornata</i> (Duncan, 1878)	-	-	21	60	I, II	Hs	
<i>Caryophyllia (Caryophyllia) smithii</i> Stokes & Broderip, 1828	-	5	48	48	II, III	Hs, Ss	= <i>Caryophyllia clava</i>
<i>Coenocyathus anthophyllites</i> Milne Edwards & Haime, 1848-	-	5	-	-	II	Ss	
<i>Desmophyllum dianthus</i> (Esper, 1794)	-	64	-	-	VII	Hs	
<i>Hoplangia durotrix</i> Gosse, 1860	-	-	59	-	I	Hs	
<i>Paracyathus pulchellus</i> (Philippi, 1842)	-	8	-	-	II	Hs	
<i>Phyllangia mouchezii</i> (Lacaze-Duthiers, 1897)	-	-	22	22	II	Hs	
<i>Polycyathus muelleriae</i> (Abel, 1959)	-	-	21	PS ⁷	I	Hs	
Family: Faviidae							
<i>Cladocora caespitosa</i> (Linnaeus, 1767)	-	62	1	48	I, II	Hs	
Family: Oculinidae							
* <i>Oculina patagonica</i> De Angelis, 1908	-	-	-	51	I	Hs	
Family: Pocilloporidae							
<i>Madracis pharensis</i> (Heller, 1868)	-	-	1	21	II	Hs	= <i>Porites daedalea</i>
Subclass: Octocorallia							
Family: Funiculinidae							
<i>Funiculina quadrangularis</i> (Pallas, 1766)	-	5	-	-	II-IV	Ss	
Family: Alcyoniidae							
<i>Alcyonium acaule</i> Marion, 1878	-	72	60	-	II	Hs	
<i>Alcyonium coralloides</i> (Pallas, 1766)	-	4	PS ²	-	II, III	Hs, Ss	
<i>Alcyonium palmatum</i> Pallas, 1766	-	2	15	-	II-V	Ss	
<i>Parerythropodium bosphorense</i> Tixier-Durivault, 1961	-	12	-	-	II	Ss	
Family: Clavulariidae							
<i>Clavularia crassa</i> (Milne Edwards, 1848)	-	8	-	-	I	Hs	
<i>Sarcodictyon roseum</i> (Philippi, 1842)	-	4	-	-	III	Ss	= <i>Rhizoxenia rosea</i>
Family: Plexauridae							
<i>Bebryce mollis</i> Philippi, 1842	-	5	-	-	III	Ss	
<i>Paramuricea clavata</i> (Risso, 1826)	-	25	48	-	II	Hs	
<i>Paramuricea macrospina</i> (Koch, 1882)	-	5	-	-	II, III	Hs, Ss	= <i>Muricea macrospina</i>
<i>Spinimuricea klavereni</i> (Carpine & Grasshoff, 1975)	-	5	-	-	II, III	Hs, Ss	= <i>Muricea echinata</i>
Family: Gorgoniidae							
<i>Eunicella cavolini</i> (Koch, 1887)	-	48	48	-	II	Hs	
<i>Eunicella singularis</i> (Esper, 1791)	-	48	48	-	II	Hs	
<i>Eunicella verrucosa</i> (Pallas, 1766)	-	8	-	-	II, III	Hs	
Family: Veretillidae							

Table 1. (Continued).

<i>Veretillum cynomorium</i> (Pallas, 1766)	-	4	15	-	II-IV	Ss	
Family: Paralyconiidae							
<i>Paralcyonium spinulosum</i> Delle Chiaje, 1822	-	68	-	-	II	Hs	
Family: Pennatulidae							
<i>Pennatula phosphorea</i> Linnaeus, 1758	-	2	-	67	II, III	Ss	
<i>Pennatula rubra</i> (Ellis, 1761)	-	49	-	54	II, III	Ss	
<i>Pteroeides griseum</i> (Linnaeus, 1767)	-	8	-	PS ⁸	?	Ss	
<i>Pteroeides spinosum</i> (Ellis, 1764)	-	2	-	PS ⁹	II	Ss	
Family: Virgulariidae							
<i>Virgularia mirabilis</i> (Müller, 1776)	7	46	-	-	II, III	Ss	
Family: Cornulariidae							
<i>Cornularia cornucopiae</i> (Pallas, 1766)	-	46	-	-	II	Hs	
Family: Kophoblemnidae							
<i>Kophobelemnon stelliferum</i> (Müller, 1776)	-	8	-	-	II	Ss	= <i>Cophobelemnon leuckarti</i>
Phylum: Ctenophora							
Class: Nuda							
Family: Beroidae							
* <i>Beroe ovata</i> Bruguière, 1789	42	34	-	-	I, II	P	
Class: Tentaculata							
Subclass: Cyclocoela							
Family: Bolinopsidae							
<i>Bolinopsis vitrea</i> (L. Agassiz, 1860)	63	-	13	PS	I-III	P	
<i>Bolinopsis infundibulum</i> (O.F. Müller, 1776)	-	-	60	-	II	P	
* <i>Mnemiopsis leidyi</i> A. Agassiz, 1865	35	26	31	29	I, II	P	
Family: Cestidae							
<i>Cestum veneris</i> Lesueur, 1813	-	-	-	60	I	P	
Subclass: Typhlocoela							
Family: Pleurobrachiidae							
<i>Pleurobrachia pileus</i> (O. F. Müller, 1776)	35	34	-	-	I, II	P	
<i>Pleurobrachia rhodopsis</i> Chun, 1879	9	8	13	-	I	P	

1. Forbes, 1844; 2. Colombo, 1885; 3. Antipa, 1892; 4. Ostroumoff, 1894; 5. Ostroumoff, 1896; 6. Billard, 1931; 7. Jakubova, 1948; 8. Demir, 1952; 9. Demir, 1954; 10. Einarsson and Gürtürk, 1959; 11. Tortonese, 1959; 12. Tixier-Durivault, 1961; 13. Ergen, 1967; 14. Băcescu et al., 1971; 15. Geldiay and Kocataş, 1972; 16. Gökalp, 1972; 17. Coşar, 1974; 18. Geldiay and Balık, 1977; 19. Kocataş, 1978; 20. Marinopoulos, 1979; 21. Zibrowius, 1979a; 22. Zibrowius, 1980; 23. Kiseleva, 1981; 24. Ünsal, 1981; 25. Öztürk and Bourguet, 1990; 26. Artüz, 1991; 27. Bingel et al., 1991; 28. Svoboda and Cornelius, 1991; 29. Uysal and Mutlu, 1993; 30. Ergen et al., 1994; 31. Kideys and Niermann, 1994; 32. Hartog, 1995; 33. Kideys and Gücü, 1995; 34. Shiganova et al., 1995; 35. Shiganova et al., 1998; 36. Koçak et al., 1999; 37. Albayrak and Balkis, 2000; 38. Koçak and Küçüksezgin, 2000; 39. Marques et al., 2000; 40. Tarkan, 2000; 41. Bat et al., 2001; 42. Kideys and Romanova, 2001; 43. Sarı et al., 2001; 44. Bilecenoglu, 2002; 45. İnanmaz et al., 2002; 46. Uysal et al., 2002; 47. Gönlgür Demirci, 2004; 48. Öztürk et al., 2004; 49. Topaloğlu et al., 2004; 50. Çinar et al., 2006b; 51. Çinar et al., 2006a; 52. Çinar et al., 2008; 53. Mavili, 2008; 54. Mutlu and Ergev, 2008; 55. İşinibilir et al., 2010; 56. Aslan Cihangir et al., 2011; 57. Çevik et al., 2011; 58. Eleftheriou et al., 2011; 59. Gökalp, 2011; 60. Gözcelioğlu, 2011; 61. Mavili, 2011; 62. Özalp and Alparslan, 2011; 63. Öztürk et al., 2011; 64. Taviani et al., 2011; 65. Turan et al., 2011; 66. Gülşahin and Tarkan, 2011a; 67. Çinar et al., 2012; 68. Evirgen and Ekşiyan, 2012; 69. Gülşahin and Tarkan, 2012; 70. Gülşahin et al., 2013; 71. Gürlek et al., 2013; 72. Topçu and Öztürk, 2013.

families (Table; Figure 2). Seven anthozoan species (*Arachnanthus oligopodus*, *Bunodactis rubripunctata*, *Bunodeopsis strumosa*, *Corynactis viridis*, *Halcampoides purpureus*, *Sagartiogeton lacerates*, and *Sagartiogeton undatus*) constitute new records for the Turkish marine fauna. A brief description and distribution of these species is presented below.

Arachnanthus oligopodus (Cerfontaine, 1891)

(Figure 2A)

Notes: Column up to 80 mm (usually up to 25 mm) long. Thin, not sclerotised, granular tube-living relatively loose in substrata; often u-shaped; tearing easily. Up to 20 marginal or labial tentacles each in a pseudocycle; marginal tentacles short; usually brown-grey banded, but uniformly brown or transparent; siphonoglyph very wide (Figure 2A). Active at night, rolls its tentacles immediately if touched (Hofrichter, 2003).

Distribution: This species was encountered on sand at 26 m depth at station 6 (2 specimens), and in a *Cymodocea nodosa* bed at 16 m depth at station 7 (1 specimen). It is a species endemic to the Mediterranean (Vafidis, 2010).

Bunodactis rubripunctata (Grube, 1840)

(Figure 2B)

Notes: Basal disc almost 35 mm; column 40 mm high, 30 mm base; mouth disc 45 mm. Column brownish red or green mottling with fine red, irregular; 48 vertical rows of red dotted verrucae; significant fosse; short capitulum. Limbus lighter than wall sheet, clearly notched; margin with 96 alternating small and large, composite (pseudo) acrorhagi. Ninety-six tentacles slightly longer than oral disc diameter; hyaline brown grey, light grey, reddish grey with yellowish longitudinal stripes on surface, conical (Figure 2B). Inner mouth brown field greenish or reddish, peripheral half golden yellow, olive green border, mesenteric insertion red. Two distinct siphonoglyphs (Hofrichter, 2003).

Distribution: This species was only found on hard substrate (1 specimen) at 1.5 m depth at station 4 (Poyraz Limanı-Karaada). It is an Atlanto-Mediterranean species (Vafidis, 2010).

Bunodeopsis strumosa Andrès, 1881

(Figure 2C)

Notes: Basal disc almost 4 mm in diameter; column up to 6 mm high, column brownish green with large golden yellow, with bubble-like protuberances. Oral disc small; 26 conical tentacles, almost 4 times longer than oral disc diameter; mottling with brownish stripes (Figure 2C). No significant siphonoglyphs.

Distribution: This species was found on *Posidonia oceanica* and *Cymodocea nodosa* leaves (many specimens) at 2–3 m depth at stations 2, 3, and 6. It is a species endemic to the Mediterranean Sea (Vafidis, 2010).

Corynactis viridis Allman, 1846

(Figure 2D)

Notes: Column usually short and squat, usually overlapped by oral disc. Tentacles short to moderate, arranged in radial rows increasing in size towards the margin; each with a spherical knob (acrosphere) at tip (Figure 2D). Disc about 10 mm diameter; span of tentacles to 25 mm. Colours varied, with disc, tentacles, and often acrospheres in contrasting colours.

Distribution: Two colonies of this species (each >100 individuals) were found on rocky wall at 28–30 m depths at station 1 (Yassiada and Sivriada). Both colonies observed were colourless, without zooxanthellae. It is an Atlanto-Mediterranean species and was previously reported from the western Mediterranean Sea, Adriatic Sea, and Aegean Sea (Vafidis, 2010).

Halcampoides purpureus (Studer, 1879)

(Figure 2E)

Notes: Column elongated, not divided into distinct regions, its aboral end rounded and provided with cinclides; periderm is absent. Disc small, with a slight hypostome; tentacles very long, even in contraction being much longer than the disc diameter, in extension tapering to fine points (Figure 2E). Length of column when not buried up to 100 mm. Disc and tentacles translucent grey-brown or pinkish, disc with reddish lines on mesenteric insertions and often with small white spots on the exocoels, close to the mouth (Manuel, 1988).

Distribution: This species was observed on sand at 5–12 m depth (6 specimens) at station 6. It is distributed in the East Atlantic Ocean and Mediterranean Sea (Vafidis, 2010).

Sagartiogeton laceratus (Dalyell, 1848)

(Figure 2F)

Notes: Base broad and closely adherent, its outline usually ragged due to basal laceration; column tall and pillar-like extension, 30 mm high, 15 mm base diameter; disc twice diameter of column; tentacles long, arranged irregularly, owing to habitual pedal laceration, up to about 200. Column buff, dull orange or reddish brown, pale cream stripes running upwards from limbus to about middle; disc translucent cream or greyish, mesenteric insertions being indicated by fine orange lines; flanked by a pair of opaque cream ones, lining of actinopharynx orange (Figure 2F); acontia not readily emitted (Manuel, 1988).

Distribution: This species was found on rocky substrate at 26–35 m depth at station 1 (many specimens). It is distributed along the north-eastern Atlantic coasts from Sweden to Portugal (Manuel, 1988). This species has not been recorded in the Mediterranean Sea before and its presence in the Sea of Marmara could be attributed to transport by ships.




Figure 2. A) *Arachnanthus oligopodus* at 26 m depth at station 6, B) *Bunodactis rubripunctata* at 1.5 m depth at station 4, C) *Bunodeopsis strumosa* at 2 m depth at station 2, D) *Corynactis viridis* at 28 m depth at station 1, E) *Halcampoides purpureus* at 12 m depth at station 6, F) *Sagartiogeton lacerates* at 30 m depth at station 1, G) *Sagartiogeton undatus* at 30 m depth at station 1, H) *Pachycerianthus multiplicatus* at 8 m depth at station 10 (only C and H photographed by ME Çinar, the others by B Yokeş).

Sagartiogeton undatus (Müller, 1778)

(Figure 2G)

Notes: Column up to 120 mm high and 20 mm in diameter, becoming very low in contraction. Oral disc wider than column, with an unusually wide mouth; tentacles up to 200, long. Column grey or buff, with regular vertical stripes of brown flecks of variable intensity; acontia not readily emitted (Figure 2G). Disc and tentacles translucent grey or buff, with a pattern of dark markings and cream spots; often with irregular dark radial wedges; tentacles with 2 dark longitudinal stripes (Manuel, 1988).

Distribution: Many specimens of this species were observed on rocky substrate at 25–35 m depths at station 1. It is an Atlanto-Mediterranean species (Vafidis, 2010).

Pachycerianthus multiplicatus Carlgren, 1912

(Figure 2H)

Notes: Marginal tentacles arranged in 4 pseudocycles, numbering almost 120; diameter of tentacles ca. 150 mm. Marginal tentacles pale yellowish to white at tips, banded with brown; labial tentacles pale yellowish (Figure 2H). Tips of marginal tentacles slender, bending towards direction of prevailing current.

Distribution: This species was found only on sandy mud bottom (3 specimens) at 8 m depth at station 10 (Yumurtalık). It is known from the coasts of the United Kingdom and Scandinavia (Manuel, 1988). This is the first time this species is being reported from the Mediterranean Sea. This species could have been introduced to the area (near the Baku–Tbilisi–Ceyhan Oil Terminal) by ballast waters of ships.

Three anthozoan species were excluded from the list: 1) *Phellia elongata* (Delle Chiaje, 1825) was reported on alga *Ellisolandia elongata* (cited as *Corallina mediterranea*) in İzmir Bay by Kocataş (1978), but this species was considered *nomen dubium* by van der Land and den Hartog (2001); 2) based on the specimens collected at 4 stations (depth range: 73–496 m) in the Sea of Marmara, Ostroumoff (1896) presented *Palythoe conchilega* as a new species together with a small note, but as the species was not adequately described or figured, it is considered here as *nomen nudum*; 3) Demir (1952) identified the species *Gorgonia flabellum* Linnaeus, 1758 (cited as *Rhipidigorgia flabellum*) among the Sea of Marmara material deposited at the Fisheries Institution in İstanbul. It is in fact a western Atlantic (Antilles) species (Bayer, 1953) and its report from the Sea of Marmara is probably a misidentification since the species has no other records from the Turkish coasts or the whole Mediterranean Sea. Finally, 2 anthozoan species, namely *Cerianthus lloydii* and *Diadumene cincta*, are regarded as questionable as no morphologic/anatomic characters specific to these species were provided by Gökalp (2011).

The total number of Cnidaria and Ctenophora species along the coasts of Turkey was estimated to be 202 species, of which 195 species belonged to Cnidaria and 7 species to Ctenophora (Figure 3). The highest number of cnidarian species was determined in the Aegean Sea (121 species) and the Sea of Marmara (115 species), with the lowest (17 species) in the Black Sea. Unlike cnidarians, the highest number of ctenophore species (5 species) was encountered in the Black Sea, with the lowest (3 species) in the Levantine Sea.

Among the classes of Cnidaria and Ctenophora, Hydrozoa ranked first in terms of the number of species (106 species), followed by Anthozoa (75 species) and Scyphozoa (12 species) (Figure 3). Scyphozoa included 10 species in the Aegean Sea, 7 species in the Levantine Sea, and 2 species (*Aurelia aurita* and *Rhizostoma pulmo*) in the Black Sea. The Sea of Marmara possessed roughly half the number of hydrozoan species (57 species) and the majority of anthozoans (52 species) known to occur along the coasts of Turkey. This sea also contained 24 anthozoan species (e.g., *Corynactis viridis*, *Sagartiogeton* spp., *Dendrophyllia ramea*, *Desmophyllum dianthus*, *Parerythropodium bosphorense*, *Spinimuricea klavereni*, and *Paramuricea macrospina*) that have not been reported from any other seas surrounding Turkey. Only 1 Cubozoa (*Carybdea marsupialis*) and 1 Staurozoa (*Lucernariopsis campanulata*) species were reported along the coasts of Turkey (Demir, 1952; Geldiay and Balık, 1977; Gönülğür Demirci, 2004). The number of ctenophore species varied between 3 (Levantine Sea) and 5 (Black Sea) in the region.

The numbers of cnidarian and ctenophore species along the coasts of Turkey are shown in Figure 4. The hot spot areas for the cnidarian species are the Prince Islands, İstanbul Straits, İzmir Bay, and Datça Peninsula (Figure 4A). Only 9 grid squares (15 × 15 km) have more than 18 cnidarian species in the region. Cnidarians along the Black Sea coast of Turkey have been poorly studied to date, with the presence of 2 jellyfish (*Aurelia aurita* and *Rhizostoma pulmo*), 8 hydrozoan, and 6 anthozoan species, most of which were reported from the pre-Bosphorus area (i.e. Jakubova, 1948; Kiseleva, 1981). The diversity and population densities of ctenophores have been relatively well studied in the Black Sea and the Sea of Marmara. Every grid in these seas has at least one ctenophore species (Figure 4B). The less studied regions in terms of the ctenophore diversity are the Aegean Sea and Levantine Sea, where the limited data regarding this group are only available from Fethiye Bay, İzmir Bay, and Gökçeada.

The numbers of hydrozoan species attained the highest levels (max. 32 species) in areas such as the Datça Peninsula, Prince Islands, and İstanbul Strait, where relatively strong scientific efforts have been made, particularly thanks to Marinopoulos (1979) and Ünsal




Figure 3. The number of species of the phyla Cnidaria and Ctenophora along the coasts of Turkey. HYD: Hydrozoa, CUB: Cubozoa, SCY: Scyphozoa, STA: Staurozoa, ANT: Anthozoa, NUD: Nuda, TEN: Tentaculata. ΣS indicates the total number of species of two phyla.

(1981) (Figure 5A). Only 3 grid squares in the Aegean Sea and 5 grid squares in the Sea of Marmara have more than 10 species. Only grid squares located near the Black Sea entrance of the İstanbul Strait possessed the hydrozoan species (totally 8 species) in the Black Sea, where almost 26 species are known (Zaitsev and Mamaev, 1997). There are also some sporadic data on hydrozoan fauna along the Levantine coast of Turkey. The scyphozoan species were relatively well studied in İzmir Bay (Ergen, 1967; Balık, 1972) and around Gökçeada (Öztürk and Topaloğlu, 2011), where more than 4 species were reported (Figure

5B). *Aurelia aurita* is a common species along the coast of the Black Sea (Kideys and Romanova, 2001), *Cotylorhiza tuberculata* in Gökova Bay (Gülşahin and Tarkan, 2011b), and *Rhopilema nomadica* in Mersin Bay and İskenderun Bay (Avşar, 1999). The highest number of anthozoan species (>12 species) was found around the Prince Islands and Kaş, where relatively intensive research has been performed (Figure 5C). Some species seem to have a narrow geographic distribution (i.e. confined to a specific area). For instance, the alien invasive coral species *Oculina patagonica* was only encountered in Akkuyu (Çinar et al.,




Figure 4. The distribution of the number of cnidarian (A) and ctenophore (B) species along the coasts of Turkey. Each grid square has dimensions of 15×15 km.

2006a), *Desmophyllum dianthus* in deep water (Çınarcık Basin) of the Sea of Marmara (Taviani et al., 2011), and *Parerythropodium bosphorense* in the İstanbul Strait (Tixier-Durivault, 1961). Data regarding the anthozoan fauna of the Black Sea of Turkey are only available from the pre-Bosphorus region and Sinop Peninsula.

Only 34 species were found in the pelagic domain and the remaining in benthic assemblages. The majority of the species (139) were encountered on hard substrata and 33 species were from soft substrata. Seven species occurred on both hard and soft substrata. A total of 155 species were reported from 0–50 m depths and only 5 species (*Alcyonium palmatum*, *Desmophyllum dianthus*, *Parantipathes larix*, *Cerianthus membranaceus*, and *Muggiaea kochii*) inhabited depths of about 200 m or downwards.

The alien Cnidaria and Ctenophora species inhabiting the coasts of Turkey were evaluated by Çınar et al. (2011), who listed 8 species (2 Hydrozoa, 3 Scyphozoa, 1 Anthozoa, and 2 Ctenophora species) in the region. The number of alien species (18) presented in this study is more than the double that. There are 3 reasons for this

increase: 1) 5 alien hydrozoan (*Coryne eximia*, *Eudendrium merulum*, *Sertularia marginata*, *Clytia linearis*, and *Filellum serratum*) and 1 anthozoan (*Diadumene lineata*) species escaped the attention of Çınar et al. (2011); 2) 2 species (*Aequorea globosa* and *Diadumene cincta*) were recently identified in the area (Gökalp, 2011; Turan et al., 2011); and, finally, 3) 2 species (*Sagartiogeton laceratus* and *Pachycerianthus multiplicatus*) are newly reported as components of the Turkish marine fauna in the present study. The xenodiversity (i.e. alien diversity) regarding the phylum Cnidaria is 16 species (8 hydrozoan, 3 scyphozoan, 5 anthozoan); for the phylum Ctenophora, it is 2 species. The majority of alien species should be considered as Lessepsian migrants. The presence of the cnidarian species *Diadumene* spp., *S. laceratus*, *P. multiplicatus*, and *Oculina patagonica* and the ctenophores (*Mnemiopsis leidyi* and *Beroe ovata*) along the Turkish coasts should probably be attributed to introduction by shipping. Finally, *Cordylophora caspia* has been considered as a cryptogenic species (Zenetos et al., 2010). The distribution of the number of alien species along the coast of Turkey is shown in Figure 6. Two alien ctenophore species (*Mnemiopsis*




Figure 5. The distribution of the number of species for the major Cnidaria classes along the coasts of Turkey: A) Hydrozoa, B) Scyphozoa, and C) Anthozoa. Each grid square has dimensions of 15 × 15 km.

leidy and *Beroe ovata*) are very common in the Black Sea and the Sea of Marmara. Five grid squares in the Sea of Marmara and 3 in the Levantine Sea (İskenderun Bay) have maximally 3 alien species, whereas only 2 grid squares have 2 alien species in the Aegean Sea. The alien jellyfish species (*Cassiopea andromeda*, *Phyllorhiza punctata*, and *Rhopilema nomadica*) occurred only along the Aegean and Levantine coasts of Turkey.

In the IUCN Red List, the gorgonian *Eunicella verrucosa* is classified as Vulnerable, *Madracis pharensis* as

Least Concern, and *Balanophyllia europaea* and *Cladocora caespitosa* as Data Deficient (IUCN, 2014). *Oculina patagonica*, which is an invasive alien species in the Mediterranean Sea, might have been erroneously included in the list with wrong information (i.e. endemic species for the Mediterranean Sea). According to the Barcelona and Bern Conventions, *Savalia savaglia* is recognised as an endangered or threatened species. In the national legislation (Official Notice for Regulating Commercial Fisheries of Water Resources, No. 2012/65), fisheries of




Figure 6. The distribution of the number of alien cnidarian and ctenophore species along the coasts of Turkey. Each grid square has dimensions of 15×15 km.

Corallium rubrum and *Savalia savaglia* are prohibited along the coasts of Turkey. *Savalia savaglia* is known to occur in the Sea of Marmara (Öztürk and Bourguet, 1990) and the North Aegean Sea (present study), but no report has been made about the occurrence of *C. rubrum* along the Turkish coast. Therefore, the presence of this species in the list lacks scientific basis. However, the presence of this species in the Aegean Sea is known since the classical work of Forbes (1844), who reported it as “very numerous minute specimens occurring of *Corallium rubrum*, for instance, but none being met with of sufficient size as to render them of value in commerce”. This species was then encountered near Meis (Kastellorizon) Island at 128–274 m depth (Zibrowius, 1979b) and in the North Aegean Sea (Chintiroglou et al., 1989) including Midilli Island at 50–91 m depths (Vafidis et al., 1994).

The species reports of Cnidaria and Ctenophora sharply increased after the 1940s in all seas except for the Black Sea, where a relatively gradual increase was observed in the course of time (Figure 7). The authors that made the major contributions to the understanding of the cnidarian diversity in the region were Ostroumoff (1896) with 23 new records, Demir (1952) with 26 new records, Marinopoulos (1979) with 25 new records, and Ünsal (1981) with 13 new records. In the last 14 years, new species records have been given by different authors with almost equal contributions.

This paper compiled all existing species data regarding Cnidaria and Ctenophora along the coasts of Turkey and provides a database for future studies. The paper also indicates the urgent need of precise taxonomic studies to be focused on certain classes of these phyla (i.e. Anthozoa) that have not been adequately studied in the area yet. The




Figure 7. Changes over time in the number of new records of Cnidaria and Ctenophora along the coasts of Turkey (T: Turkey, AS: Aegean Sea, SM: Sea of Marmara, LS: Levantine Sea, BS: Black Sea).

lack of faunistic data in many marine areas of Turkey (including deep seas) and the requisite of expanding faunistic studies from some specific areas (e.g., İzmir Bay) where marine stations are located towards other unexplored areas of the Turkish coasts are also emphasised.

References

- Albayrak S, Balkis N (2000). Hydroid polyps of the Bosphorus. *Turk J Mar Sci* 6: 41–53.
- Alparslan M (2001). Succession of Scyphozoa-Ctenophora in the Harbour of Çanakkale. *Turk J Mar Sci* 7: 59–68.
- Antipa GR (1892). Eine neue Art von *Drymonema*. *Jen Z Naturwiss* 27: 337–339 (in German).
- Appeltans W, Ah Yong ST, Anderson G, Angel MV, Artois T, Bailly N, Bamber R, Barber A, Batsch I, Berta A et al. (2012). The magnitude of global marine species diversity. *Current Biology* 22: 2189–2202.
- Artüz MI (1991). Ctenophora invasion in the Marmara Sea. *J Sci Tech* 9: 9–11 (in Turkish).
- Aslan Cihangir H, Pancucci Papadopoulou MA, Can Yılmaz E (2011). First record of *Aiptasia mutabilis* (Cnidaria: Anthozoa) from Turkish Seas. *Turk J Zool* 53: 447–450.
- Avşar D (1999). Physico-chemical characteristics of the eastern Mediterranean in relation to distribution of the new Scyphomedusae (*Rhopilema nomadica*). *Tr J Zool* 23: 605–616 (in Turkish with English abstract).
- Băcescu MC, Müller GI, Gomoiu MT (1971). *Ecologie Marina. Cercătări de ecologie bentală în Marea Neagră-Analiza cantitativă, calitativă și comparată a faunei bentale pontice*. Edit Acad Rep Soc Rom 4: 1–357 (in Romanian).
- Balık S (1972). İzmir Körfezi ve civarında bulunan deniz analarının “Scyphozoa, Coelenterata” taksonomik ve ekolojik özellikleri üzerine araştırmalar. MSc, Ege University, İzmir, Turkey (in Turkish).
- Bat L, Akbulut M, Sezgin M, Çulha M (2001). Effects of sewage pollution the structure of the community of *Ulva lactuca*, *Enteromorpha linza* and rocky macrofauna in Dışlman of Sinop. *Turk J Biol* 25: 93–102.
- Bayer FM (1953). Zoogeography and evolution in the octocorallian family Gorgoniidae. *Bull Mar Sci Gulf Carib* 3: 100–119.
- Bilecenoglu M (2002). Türkiye kıyılarında ilk kez rastlanan bir denizanası türü: *Cassiopea andromeda* (Forsskal, 1775) (Cassiopeidae, Scyphozoa). *Sualtı Dünyası Dergisi* 72: 42–43 (in Turkish).
- Billard A (1931). Hydroïdes des Côtes de Syrie. In: Gruvel A, editor. *Les Etats de Syrie. Richesses marines et fluviales. Exploitation actuelle. Avenir*. Paris, France: Societe d’Edition Geographiques, Maritimes et Coloniales, pp. 389–395.
- Bingel F, Avşar D, Gücü AC (1991). Occurrence of jellyfish in Mersin Bay. In: UNEP: Jellyfish Blooms in the Mediterranean. Proceedings of the 2nd Workshop on Jellyfish in the Mediterranean Sea. MAP Tech Rep Ser 47: 65–71.
- Çevik C, Derici OB, Çevik F, Cavas L (2011). First record of *Phyllorhiza punctata* von Lendenfeld, 1884 (Scyphozoa: Rhizostomeae: Mastigiidae) from Turkey. *Aquat Inv* 6: 27–28.
- Chintiroglou H, Dounas C, Koukouras A (1989). The presence of *Corallium rubrum* (Linnaeus, 1758) in the eastern Mediterranean Sea. *Mitt Zool Mus Berl* 65: 145–149.
- Çinar ME, Bilecenoglu M, Öztürk B, Can A (2006a). New records of alien species on the Levantine coast of Turkey. *Aquat Inv* 1: 84–90.
- Çinar ME, Bilecenoglu M, Öztürk B, Katakın T, Yokeş MB, Aysel V, Dağlı E, Açıık S, Özcan T, Erdoğan H (2011). An updated review of alien species on the coasts of Turkey. *Medit Mar Sci* 12: 257–315.
- Çinar ME, Katagan T, Koçak F, Öztürk B, Ergen Z, Kocatas A, Önen M, Kirkim F, Bakir K, Kurt G et al. (2008). Faunal assemblages of the mussel *Mytilus galloprovincialis* in and around Alsancak Harbour (İzmir Bay, eastern Mediterranean) with special emphasis on alien species. *J Mar Sys* 71: 1–17.
- Çinar ME, Katagan T, Öztürk B, Dağlı E, Açıık S, Bitlis B, Bakir K, Dogan A (2012). Spatio-temporal distributions of zoobenthos in Mersin Bay (Levantine Sea, eastern Mediterranean) and the importance of alien species in benthic communities. *Mar Biol Res* 8: 954–968.
- Çinar ME, Katagan T, Öztürk B, Egemen Ö, Ergen Z, Kocatas A, Önen M, Kirkim F, Bakir K, Kurt G et al. (2006b). Temporal changes of soft-bottom zoobenthic communities in and around Alsancak Harbour (İzmir Bay, Aegean Sea), with special attention to the autecology of exotic species. *Mar Ecol* 27: 229–246.
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Lasram FBR, Aguzzi J, Ballesteros E, Bianchi CN, Corbera J, Dailianis T et al. (2010). The biodiversity of the Mediterranean Sea: estimates, patterns and threats. *PLoS One* 5: 11842.
- Colombo A (1885). Raccolte Zoologiche eseguite del R. Piroscopa Washington nelle campagna-abissale talassografica dell’anno 1885. *Riv Marit*: 1–34 (in Italian).
- Coşar G (1974). İzmir Körfezi ve civarında yaşayan deniz şakayıkları (Actiniaria, Anthozoa) üzerine taksonomik ve ekolojik araştırmalar. MSc, Ege University, İzmir, Turkey (in Turkish).

- Demir M (1952). Boğazlar ve adalar sahillerinin omurgasız dip hayvanları. Ist Univ Rac Sci Hydrobiol Res Ins Publ 3: 1–615 (in Turkish).
- Demir M (1954). Report on the plankton of the south eastern coast of the Black Sea. Ist Univ Rac Sci Hydrobiol Res Ins Publ Ser B 1: 284–286.
- Dong Z, Liu D, Keesing JK (2010). Jellyfish blooms in China: dominant species, causes and consequences. Mar Poll Bull 60: 954–963.
- Einarsson H, Gürtürk N (1959). On plankton communities in the Black Sea. Et Bal Kur Balık Araş Mer Rap 1: 1–28.
- Eleftheriou A, Anagnostopoulou-Visilia E, Anastasopoulou E, Ateş SA, Bachari NEI, Cavas L, Cevik C, Çulha M, Cevik F, Delos AL et al. (2011). New Mediterranean biodiversity records (December 2011). Medit Mar Sci 12: 491–508.
- Ergen Z (1967). İzmir Körfezi'nde tespit edilen başlıca planktonik organizmalar. Sci Rep Fac Sci Ege Univ 47: 1–47 (in Turkish).
- Ergen Z, Kocataş A, Kayağan T, Çınar ME (1994). Gencelli Limanı (Aliağa-İzmir) bentik faunası. Ege Üniv J Fac Sci Ser B (Suppl. 16): 1047–1059 (in Turkish).
- Evirgen A, Ekşiyan S (2012). Marmara'da hayat var şimdilik. İstanbul, Turkey: Adalı Press (in Turkish).
- Forbes E (1844). Report on the Mollusca and Radiata of the Aegean Sea, and on their distribution, considering as bearing on geology. In: Report of the 13th Meeting of the British Association for the Advancement of Science 13: 130–193.
- Galil BS (2007). Seeing red: alien species along the Mediterranean coast of Israel. Aquat Inv 4: 281–312.
- Geldiay R, Balık S (1977). Ege Sahilinden tesbit edilen yeni bir scyphozoid türü hakkında. Ege Univ J Fac Sci Ser B 1: 5–12 (in Turkish).
- Geldiay R, Kocataş A (1972). Note préliminaire sur les peuplements benthiques du golfe d'Izmir. Sci Monog Fac Sci Ege Univ 12: 3–33 (in French).
- Gökalp M (2011). Doğa Rehberi (Deniz Canlıları). İstanbul, Turkey: Inkılâp Press (in Turkish).
- Gökalp N (1972). Edremit, Bodrum ve İskenderun Körfezlerinin plankton durumunun karşılaştırmalı incelenmesi. J Hydrobiol 3: 1–33 (in Turkish).
- Gönlüğü Demirci G (2004). A new record for the Turkish waters from Sinop coast of Black Sea: *Lucernariopsis campanulata* (Lamouroux, 1815) (Stauromedusae, Scyphozoa). Ege Univ J Fish Aquat Sci 21: 331–333 (in Turkish with English abstract).
- Gözcüoğlu B (2011). Denizlerimizizin Sakinleri. İstanbul, Turkey: Gökçe Ofset Basım Yayın Sanayi (in Turkish).
- Gülşahin N, Tarkan AN (2011a). The first confirmed record of the alien jellyfish *Rhopilema nomadica* Galil, 1990 from the southern Aegean Coast of Turkey. Aquat Inv 6: 95–97.
- Gülşahin N, Tarkan AN (2011b). A familiar organism in Muğla Region, the Aegean Sea *Cotylorhiza tuberculata* (Maori, 1778). In: Turan C, Öztürk B, editors. First National Workshop on Jellyfish and Other Gelatinous Species in Turkish Marine Waters; 20–21 May 2011; Muğla, Turkey. İstanbul, Turkey: Turkish Marine Research Foundation, pp. 53–57.
- Gülşahin N, Tarkan AN (2012). The first record of *Phyllorhiza punctata* von Lendenfeld, 1884 from the southern Aegean Coast of Turkey. BioInv Rec 1: 41–44.
- Gülşahin N, Tarkan AN, Bilge G (2013). The hydrozoan *Geryonia proboscoidalis* (Forsk., 1775), new for Turkey (Hydrozoa). Zool Middle East 59: 93–94.
- Gürlek M, Yağhoğlu D, Ergüden D, Turan C (2013). A new jellyfish species in the Turkish coastal waters *Aequorea forskalea* Péron and Lesueur, 1810 (Cnidaria: Hydrozoa). J Black Sea/Medit Environ 19: 380–384.
- Hartog JC den (1995). The genus *Telmatactis* Gravier, 1916 (Actiniaria: Acontinaria: Isophelliidae) in Greece and the eastern Mediterranean. Zool Med Leiden 69: 153–176.
- Hatcher BG (1997). Coral reef ecosystems: how much greater is the whole than the sum of the parts? Coral Reefs 16: 77–91.
- Hofrichter R (2003). Das Mittelmeer. Fauna, Flora, Ökologie, Band II/1: Bestimmungsführer Procaryota, Protista, Fungi, Algae, Plantae, Animalia (bis Nemertea). Heidelberg, Germany: Spektrum-Verlag (in German).
- İnanmaz ÖE, Bekbolet M, Kideys AE (2002). A new scyphozoan species in the Sea of Marmara: *Chrysaora hysoscella* (Linné, 1766). In: Yılmaz A, editor. Oceanography of the Eastern Mediterranean and Black Sea. Ankara, Turkey: TÜBİTAK, pp. 857–859.
- İşinbilir M, Yılmaz İN, Piraino S (2010). New contributions to the jellyfish fauna of the Marmara Sea. Italian J Zool 77: 179–185.
- IUCN (2013). The IUCN Red List of Threatened Species. Version 2013.2. Gland, Switzerland: IUCN.
- Jakubova LI (1948). Osobennosti biologii pribosforskogo uchastka Chernogo Morya. Trudy SBS 6: 274–285 (in Russian).
- Kideys AE (2002). Fall and rise of the Black Sea ecosystem. Science 297: 1482–1484.
- Kideys AE, Niermann U (1994). Occurrence of *Mnemiopsis* along the Turkish coasts (from northeastern Mediterranean to Istanbul). ICES J Mar Sci 51:423–427.
- Kideys AE, Gücü AC (1995). *Rhopilema nomadica*: A Lessepsian scyphomedusan new to the Mediterranean coast of Turkey. Israel J Zool 41: 615–617.
- Kideys AE, Romanova Z (2001). Distribution of gelatinous macrozooplankton in the southern Black Sea during 1996–1999. Mar Biol 139: 535–547.
- Kiseleva MI (1981). Benthos Ryixlyix Gruntov Chernogo Morya. Kiev, Ukraine: Naukova Dumka (in Russian).
- Koçak F, Ergen Z, Çınar ME (1999). Fouling organisms and their developments in a polluted and an unpolluted marina in the Aegean Sea (Turkey). Ophelia 50: 1–20.

- Koçak F, Küçüksezgin F (2000). Sessile fouling organisms and environmental parameters in the marinas of the Turkish Aegean coast. *Indian J Mar Sci* 29: 149–157.
- Kocataş A (1978). İzmir Körfezi kayalık sahillerinin bentik formları üzerinde kalitatif ve kantitatif araştırmalar. EÜ Fac Sci Monogr Ser 12: 1–93 (in Turkish).
- Manuel RL (1988). British Anthozoa (Coelenterata: Octocorallia and Hexacorallia). Keys and Notes for the Identification of the Species. 2nd ed. Synopses of the British Fauna (New Series) 18. Devon, UK: NHBS.
- Marinopoulos J (1979). Biological survey of the eastern Mediterranean Sea: hydroids. *Rapp Comm Int Mer Medit* 25/26: 119–120.
- Marques AC, Peña Cantero AL, Vervoort W (2000). Mediterranean species of *Eudendrium* Ehrenberg, 1834 (Hydrozoa, Anthomedusae, Eudendriidae) with the description of a new species. *Zool Soc London* 252: 197–213.
- Mavili S (2008). Kuzey Ege ve Marmara Denizi Kıyısız Yüzey Sularında Scyphomedusae ve Ctenophora Üzerine Yerinde Gözlemler. *Ege Univ J Fish Aquat Sci* 25: 141–146 (in Turkish).
- Mavili S (2011). İzmir Körfezi (Ege Denizi) yüzey tabakalarında (0-10 m) Hydromedusae popülasyonu ve *Aglaura hemistoma* Peron ve Lesueur, 1810 (Cnidaria, Hydrozoa)'nın dağılımı. *Ege Univ J Fish Aquat Sci* 28: 25–28 (in Turkish).
- Muscatine L, Porter JW (1977). Mutualistic symbioses adapted to nutrient-poor environments. *BioScience* 27: 454–460.
- Mutlu E, Bingel F, Gucu AC, Melnikov VV, Niermann U, Ostr NA, Zaika VE (1994). Distribution of the new invader *Mnemiopsis* sp. and the resident *Aurelia aurita* and *Pleurobrachia pileus* populations in the Black Sea in the years 1991-1993. *ICES J Mar Sci* 51: 407–421.
- Mutlu E, Ergev MB (2008). Spatio-temporal distribution of soft-bottom epibenthic fauna on the Cilician shelf (Turkey), Mediterranean Sea. *Rev Biol Trop* 56: 1919–1946.
- Ostroumoff A (1894). Dal'neishie materialy k estestvennoi istoriyi Bosfora. *Bull Acad Imp Sci Saint Petersburg* 74: 1–46 (in Russian).
- Ostroumoff A (1896). Otchet o dragirovka i planktonnykh ulovax ekspeditsii "Selyanika". *Bull Acad Imp Sci Saint Petersburg* 5: 33–92 (in Russian).
- Özalp HB, Alparslan M (2011). The first record of *Cladocora caespitosa* (Linnaeus, 1767) (Anthozoa, Scleractinia) from the Marmara Sea. *Turk J Zool* 35: 701–705.
- Özsoy E, Pınardı N (2010). Marsili in İstanbul notes, maps and drawings. In: Marmara Sea 2010 Symposium, 25–26 September 2010, İstanbul, Turkey, pp. 25–32.
- Öztürk B, Aktan Y, Topaloğlu B, Keskin Ç, Karakulak S, Öztürk AA, Dede A, Türkozan O (2004). Marine Life of Turkey in the Aegean and Mediterranean Seas. İstanbul, Turkey: Turkish Marine Research Foundation.
- Öztürk B, Bourguet JP (1990). Données préliminaires sur le corail noir de la Mer de Marmara (Turquie) *Gerardia savaglia* (Bertolini, 1819). *Ist Uni J Aquat Prod* 4: 45–48 (in French).
- Öztürk B, İsinibilir M (2010). An alien jellyfish *Rhopilema nomadica* and its impacts to the eastern Mediterranean part of Turkey. *J Black Sea/Med Environ* 16: 149–156.
- Öztürk B, Mihneva V, Shiganova T (2011). First records of *Bolinopsis vitrea* (L. Agassiz, 1860) (Ctenophora: Lobata) in the Black Sea. *Aquat Inv* 6: 355–360.
- Öztürk B, Topaloğlu B (2011). A short note on the jellyfish at the Gökçeada Island, the North Aegean Sea, Turkey. In: Turan C, Öztürk B, editors. First National Workshop on Jellyfish and Other Gelatinous Species in Turkish Marine Waters; 20–21 May 2011; Muğla, Turkey. İstanbul, Turkey: Turkish Marine Research Foundation, pp. 86–89.
- Parsons TR, Lalli CM (2002). Jellyfish population explosions: revisiting a hypothesis of possible causes. *La Mer* 40: 111–121.
- Pearse VB, Muscatine L (1971). Role of symbiotic algae (Zooxanthellae) in coral calcification. *Biol Bull* 141: 350–363.
- Purcell JE, Uye S, Lo WT (2007). Anthropogenic causes of jellyfish blooms and their direct consequences for humans: a review. *Mar Ecol Prog Ser* 350: 153–174.
- Richardson AJ, Bakun A, Hays GC, Gibbons MJ (2009). The jellyfish joyride: causes, consequences and management responses to a more gelatinous future. *Trend Ecol Evol* 24: 312–322.
- Sarı HM, Balık S, Özbek M, Aygen C (2001). Bafa Gölü'nün (Ayдын) makro ve meiobentik omurgasız faunası. *Anadolu Univ J Sci Tech* 2: 285–291 (in Turkish).
- Shiganova T, Tarkan AN, Dede A, Cebeci M (1995). Distribution of the ichthyoplankton *Mnemiopsis leidyi* (Agassiz, 1865) in the Marmara Sea (October 1992). *Turk J Mar Sci* 1: 3–12.
- Shiganova TA, Bulgakova YV, Volovik SP, Mirzoyan ZA, Dudkin SI (2001). The new invader *Beroe ovata* Mayer 1912 and its effect on the ecosystem in the northeastern Black Sea. *Hydrobiologia* 451: 187–197.
- Shiganova TA, Kideys AE, Gücü AS, Niermann U, Khoroshilov VS (1998). Changes of species diversity and their abundance in the main components of pelagic community during last decades. In: Ivanov LI, Oguz T, editors. Ecosystem Modeling as a Management Tool for the Black Sea. Boston, MA, USA: Kluwer Academic Publishers.
- Svoboda A, Cornelius PFS (1991). The European and Mediterranean species of *Aglaophenia* (Cnidaria: Hydrozoa). *Zool Verhandl* 274: 1–72.
- Tarkan AN (2000). Abundance and distribution of zooplankton in coastal area of Gökçeada Island (Northern Aegean Sea). *Turk J Mar Sci* 6: 201–214.
- Taviani M, Vertino A, López Correa M, Savini A, De Mol B, Remia A, Montagna P, Angeletti L, Zibrowius H, Alves T et al. (2011). Pleistocene to Recent scleractinian deep-water corals and coral facies in the eastern Mediterranean. *Facies* 57: 579–603.
- Tixier-Durivault A (1961). Sur un nouvel Alcyonaire: *Parerythropodium bosphorense*. *Bull Mus Nat Hist Natur* 33: 322–325 (in French).

- Topaloğlu B, Öztürk B, Karakulak FS (2004). The macrozoobenthic invertebrate fauna in the Marmara Sea. *Rapp Comm Int Mer Médit* 37: 554.
- Topçu EN, Öztürk B (2013). Octocoral diversity of Balıkçı Island, the Marmara Sea. *J Black Sea/Medit Environ* 19: 46–57.
- Tortonese E (1959). Osservazioni sul bentos del Mar di Marmara e del Bosforo. *Natura Milano* 50: 18–26 (in Turkish).
- Turan C, Gürlek M, Yağlıoğlu D, Seyhan D (2011). A new alien jellyfish species in the Mediterranean Sea–*Aequorea globosa* Eschscholtz, 1829 (Cnidaria: Hydrozoa). *J Black Sea/Medit Environ* 17: 282–286.
- Ünsal I (1981) Ege ve Marmara denizlerinin hidroid polipleri. Docent thesis, İstanbul University, İstanbul, Turkey (in Turkish).
- Uysal A, Yüksek A, Okuş E, Yılmaz N (2002). Benthic community structure of the Bosphorus and surrounding area. *Water Sci Tech* 46: 37–44.
- Uysal Z, Mutlu E (1993). Preliminary note on the occurrence and biometry of Ctenophoran *Mnemiopsis leidyi* finally invaded Mersin Bay. *Doğa Turk J Zool* 12: 229–236.
- Vafidis D (2010). Checklist of class Anthozoa (Phylum Cnidaria). In: Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F, Aguzzi J, Ballesteros E, Bianchi CN, Corbera J, Dailianis T et al., editors. *The Biodiversity of the Mediterranean Sea: Estimates, Patterns, and Threats*. PLoS ONE 5: 36.
- Vafidis D, Koukouras A, Voultsiadou-Koukoura E (1994). Octocoral fauna of the Aegean Sea with a check list of the Mediterranean species: new information, faunal comparisons. *Ann Inst Océanogr Paris* 70: 217–229.
- van der Land J, den Hartog JH (2001). Actiniaria. In: Costello MJ, Emblow C, White, RJ, editors. *European Register of Marine Species: A Check-List of the Marine Species in Europe and a Bibliography of Guides to Their Identification*. Collection Patrimoines Naturels 50: 106–109
- Vinogradov ME, Shushkina EA, Musaeva EI, Sorokin PY (1989). Ctenophora *Mnemiopsis leidyi* (A. Agassiz) (Ctenophora: Lobata)-new settler in the Black Sea. *Oceanology* 29: 293–298.
- Yılmaz IN (2014). Collapse of zooplankton stocks during *Liriope tetraphylla* (Hydromedusa) blooms and dense mucilaginous aggregations in a thermohaline stratified basin. *Mar Ecol* (in press).
- Zaitsev Y, Mamaev V (1997). *Marine Biological Diversity in the Black Sea. A Study of Change and Decline*. New York, NY, USA: United Nations Publications.
- Zenetos A, Gofas S, Verlaque M, Çinar ME, García Raso JE, Bianchi CN, Morri C, Azzurro E, Bilecenoglu M, Frogia C et al. (2010). Alien species in the Mediterranean by 2010. A contribution of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Med Mar Sci* 11: 381–493.
- Zibrowius H (1979a). Campagne de la Calypso en Méditerranée nord-orientale (1955, 1956, 1960, 1964). 7. Scléractiniaires. *Ann Inst Océanogr Paris* 55: 7–28 (in French).
- Zibrowius H (1979b). À propos du corail rouge en Méditerranée orientale. *Rapp Comm int Mer Médit* 25/26: 4 (in French).
- Zibrowius H (1980). Les Scléractiniaires de la Méditerranée et de l'Atlantique nord-oriental. *Mem Inst Océanog Monaco* 11: 1–227 (in French).