



OPEN

Acidification reduced growth rate but not swimming speed of larval sea urchins

SUBJECT AREAS:

CLIMATE-CHANGE
ECOLOGY

ECOLOGY

Received
22 September 2014Accepted
16 March 2015Published
15 May 2015Correspondence and
requests for materials
should be addressed to
K.Y.K.C. (karenchan@
ust.hk)Kit Yu Karen Chan^{1,2}, Eliseba García³ & Sam Dupont⁴

¹Biology & Applied Ocean Physics and Engineering, Woods Hole Oceanographic Institution, Woods Hole, MA, U.S.A, ²Division of Life Science, Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong, ³Departamento de Biología Animal (Ciencias Marinas), Universidad de La Laguna, La Laguna, Tenerife, Islas Canarias, Spain, ⁴Department of Biological and Environmental Sciences, University of Gothenburg, The Sven Lovén Centre for Marine Sciences – Kristineberg, Fiskebäckskil, Sweden.

Swimming behaviors of planktonic larvae impact dispersal and population dynamics of many benthic marine invertebrates. This key ecological function is modulated by larval development dynamics, biomechanics of the resulting morphology, and behavioral choices. Studies on ocean acidification effects on larval stages have yet to address this important interaction between development and swimming under environmentally-relevant flow conditions. Our video motion analysis revealed that pH covering present and future natural variability (pH 8.0, 7.6 and 7.2) did not affect age-specific swimming of larval green urchin *Strongylocentrotus droebachiensis* in still water nor in shear, despite acidified individuals being significantly smaller in size (reduced growth rate). This maintenance of speed and stability in shear was accompanied by an overall change in size-corrected shape, implying changes in swimming biomechanics. Our observations highlight strong evolutionary pressure to maintain swimming in a varying environment and the plasticity in larval responses to environmental change.

Many marine benthic organisms have limited mobility as adults and rely on a planktonic larval phase for dispersal. Planktonic larvae thus play a significant role in determining population abundance, genetic diversity, distribution, and their resilience to disturbances^{1,2}. Larval survival, transport, and successful settlement are affected by environmental variables (e.g., prey and predator concentrations, current direction and speed) which is in turn modulated by larval swimming behaviors^{3–5}. Larval swimming is bounded by the physiological and biomechanical limits of the larval form, and modulated through behavioral choices^{6–8}.

Global stressors such as ocean acidification (OA) can affect larval growth and development. These stressors can lead to changes in organisms' morphology⁹ and modify dynamics of development¹⁰. Larval ability to carry out ecological functions are tightly coupled with their development and morphology. For example, changes in length and orientation of the ciliated arms of larval urchins affect their abilities to filter particles for food¹¹, swim in still water^{12,13}, and maintain directed movement (stability) in flow¹⁴. This tight coupling suggests that any changes associated with environmental stressors can have substantial consequences.

To date, very few studies have investigated how observed effects of OA on larval development and morphology can impact swimming behavior, and thus, potential consequences for successful dispersal. Non-invasive video motion analysis has been applied to investigate swimming of larval sand dollars *Dendraster excentricus* in still water¹⁵. Their results suggested that OA condition did not affect larval swimming speeds. The authors hypothesized that a coordinated change in shape could help maintain swimming performance. Planktonic larvae, however, are rarely found in completely still water in nature. They rather experience moving water, and thus, it is important to explore larval swimming performance in flow^{16,17}.

The aim of this paper was to test the effect of OA on the relationship between developmental dynamics, larval morphology (with an emphasis on overall shape using geometric morphometrics, Fig. 1d), and swimming behavior (in still and moving water using non-invasive video motion analysis, Fig. 2). We exposed our focal organism, the larval green urchin *Strongylocentrotus droebachiensis*, to different rearing pHs (see Suppl. Table 1 for detailed carbonate chemistry) and compared their overall shapes and swimming behaviors in still water and in shear over time.

pH affected growth but not mortality rate. Relative mortality rate (RMR in day⁻¹) was calculated as the coefficient of the significant linear regression between relative density and time (days post fertilization). All regressions were statistically significant ($p < 0.001$) and had r^2 values ranged from 0.59 to 0.85 (Suppl. Table 2).




Figure 1 | Micrographs of larval *Strongylocentrotus droebachiensis* (10 days post fertilization) reared at three nominal pH levels (a). Geometric morphometric techniques were used to analyze shape change in these larvae through identification of landmarks (b), followed by computation and comparison of Procrustes coordinates on transformation grids (c).

RMR were $0.036 \pm 0.0045 \text{ day}^{-1}$ for pH 8.0, $0.039 \pm 0.0043 \text{ day}^{-1}$ for pH 7.6, and $0.027 \pm 0.0097 \text{ day}^{-1}$ for pH 7.2. Neither pH nor replicate had a significant effect on larval mortality ($p = 0.21$ and $p = 0.28$ respectively, Fig. 3a).

A logarithmic regression between total body length (mm) and days post-fertilization (hereafter larval age) was used to determine larval growth rate. The coefficient of regression between body length and the natural logarithm of larval age denotes the growth rate in a unit of $\text{mm ln}(\text{day})^{-1}$. All regressions were statistically significant ($p < 0.001$) and had r^2 values ranged from 0.77 to 0.9 (Fig. 3b, Suppl. Table 3). Growth rates were $0.105 \pm 0.0035 \text{ mm ln}(\text{day})^{-1}$ for pH 8.0, $0.103 \pm 0.0017 \text{ mm ln}(\text{day})^{-1}$ for pH 7.6, and $0.079 \pm 0.0046 \text{ mm ln}(\text{day})^{-1}$ for pH 7.2. pH had a significant effect on larval growth ($p < 0.001$) and post-hoc test showed that growth rates were significantly different between the three pH treatments ($p < 0.01$). Replicate also had a significant effect on larval growth rate ($p < 0.001$) and a post-hoc test showed that at pH 7.6 growth rate in one of the replicates (number 3) was significant different from the other two and at pH 7.2 larval urchins in all three replicates had different growth rates.

Acidification induced overall shape change. Based on the canonical variate analysis, larval urchins reared under the three pHs differed significantly in overall shapes (Fig. 4, $p < 0.0001$, Suppl. Table 4). Canonical variate 1 (CV1) and CV2 accounted for 65.4% and 20.7% of the total variance in the population. CV1 mainly varied with the ratio of body size and arm length; individuals with a lower CV1 scores had shorter total arm length but relatively longer arms per unit length of the body at a given size. CV2 varied with the arm lengths (overall height) and the separation distances between arms; individuals with a lower CV2 scores had shorter arms that are less spread out in the horizontal direction (Suppl. Table 4, Fig. 5). In general, larval urchins in the low pH treatments had a more negative CV2 score at a given CV1 score.

Our previous analysis showed that larval urchins had different growth rates when reared under different pH, therefore, we compared growth patterns of the three pH treatments by regressing the CV1 and CV2 scores against Procrustes centroid sizes. Regressions for

CV1 were significant and had r^2 value over 0.62 (Suppl. Table 6). Such allometric relationships did not hold for CV2 and the variance explained by the regression were low ($r^2 > 0.05$). In other words, larvae that were larger in size did not necessary have longer overall height, arm lengths, or shorter separation distances between arms (CV2) (Suppl. Table 6).

Larval swimming in shear is not affected by pH. Net movement and swimming metrics of larvae were computed after flow subtraction. For upward swimming larvae, average vertical swimming velocities differed between larval age and flow conditions ($p < 0.0001$, Suppl. Table 7, Fig. 5). At a given age, upward swimming larvae exposed to shear condition had higher vertical velocities. Maximum upward swimming speed was observed 10 days post fertilization and subsequently decreased with age. pH alone did not have a significant effect on vertical velocities ($p = 0.67$). There were also no significant interactions between pH and flow ($p = 0.10$) Net horizontal velocity was not affected by the three factors studied i.e., flow, age, and pH ($p \geq 0.12$, Suppl. Table 7, Fig. 5).

For downward swimming larvae, average vertical and horizontal velocities differed between larval age and flow ($p < 0.001$, Suppl. Table 7, Fig. 6). Vertical velocities became less negative as larvae aged. Horizontal velocities were higher in shear than in still condition for larval urchins at a given age, i.e., there is more horizontal movement towards downward moving flow when exposed to shear. Larval urchins in all three pH treatments had higher horizontal velocities at 7 and 10 days post-fertilization. However, pH alone did not have a significant effect on vertical velocities ($p = 0.24$). There were also no significant interactions between pH and flow ($p = 0.26$, Suppl. Table 6, Fig. 6).

At a same day of age, larval urchins from the different pH treatments had different total body lengths. Such delay in growth could have potential implications for larval swimming (Fig. 3b). However, statistical comparison on the effect of flow and pH with size as a covariate could not be performed because there was very little overlap in total body length between the pH treatments when the video observations were performed. Larvae reared in pH 7.3 did not catch up in size over the 14 days of observations (Suppl. Fig. 2).

Discussion and conclusion

Swimming of planktonic larvae has significant consequences for population dynamics and is tightly coupled with larval physiology and biomechanics⁷. When reared under decreased pH conditions, larval green urchins had reduced growth rates, changed overall shapes but were able to maintain their age-specific swimming performances.

Larvae reared under decreased pH grew slower and in a different shape. Within the observed pH range (8.0–7.2), reduction in pH had no significant effect on larval survivorship over 14 days (Fig. 3a). This observation supports previous observation on the same species which identified pH 7.0 as the physiological tipping point¹⁸. The rate and pattern of growth were significantly impacted by pH (Fig. 3b). Reduced growth rate could prolong pelagic larval duration and reduce the number of settlers due to the high mortality in the plankton^{10,19,20}. These slower growing individuals might also settle at smaller sizes²¹ or less discriminately to the chemical cues²² which could have negative implications on juvenile survival in the wild.

In addition to differences in rate of growth, overall shapes of the acidified larval urchins also differed from those of the control (Fig. 4). Previous studies on pH impact on larval morphology focus on linear measurement and have shown contrasting results: in some observations, including that on larval green urchins, ratio between body length and arm length/stomach length were affected by pH^{18,23,24} but not in others^{25,26}. In the latter cases, the pH effect became




Figure 2 | Example of background flow field in shear and swimming behaviors observed in two flow conditions (still and shear). Shear refers to a velocity gradient in the horizontal direction (a, b) and the shear level generated is similar to that in coastal water c). Red indicates upward moving water and blue indicates downward moving water. Swimming trajectories observed over five minutes of larvae reared at pH_T 8.0 were overlaid on these flow fields, with the red circles indicating the beginning of the paths (d, e). Under shear condition, larval urchins experienced horizontal transport across flow lines.

non-significant when correcting for the reduced growth rate by including body length as a covariate in the analysis. In our data set, CV1 scores correlated significantly with Procrustes centroid sizes such that the relative ratio between larval body and arms varied in an allometric manner. However, the lack of correlation between Procrustes centroid size and CV2 scores suggest that the change in length of arms and separation distances between them do not allometrically increase with overall size (Suppl. Table 6). The differences between different aspects of shape change (CV scores) relative to overall size may help reconcile the difference in previous reports.

Observed shape changes under decreased pH may have significant biomechanical implications for larval movement. Larvae with the

lower CV1 and 2 scores appeared to have an overall more “squat” morphology. This decrease in arm extension without the corresponding change in body length (decrease in CV1 score) would increase the distance between larval centers of buoyancy and gravity, and hence, the torque exerted when tilted in moving water would be decreased, thereby conferring stability^{7,27}. In addition, the reduction in separation distance between arms (decrease in CV score) present a narrower width exposed to the shear gradient, which would result in less difference in water velocity across the larval body, less tilting of the larva^{8,13}. Future modeling effort on fluid interactions of larvae should consider this kind of coordinated morphological variation in “armed larvae”.




Figure 3 | Average relative mortality rate (inset of a) calculated from linear regression of the change in larval density over time did not differ between pH treatments (a). However, pH has a significant effect on larval growth rate computed by logarithmic regression of increase in total body length over time (b). Closed symbols and solid block line represent pH 8.0 replicates, gray symbols and dash line represent pH 7.6, and open symbols and dotted line represent pH 7.2.

Decreased pH did not impact larval swimming. Contrary to the hypothesis that the change in shape would affect swimming, larval urchins raised under decreased pH had different age-specific morphology but maintained the same swimming performance in both still and moving water (Fig. 5, 6).

While pH had no significant effect on age-specific larval swimming velocities, this performance metric significantly varied with larval age. For example, older larvae had higher horizontal velocities

and may be more likely entrained in downward moving waters by crossing flow lines. This could translate into differential transport through ontogeny²⁸. It is remarkable that this age-specific pattern was maintained within our tested pH range despite significant differences in growth rates. This could be a consequence of the observed shape change allowing the smaller larvae raised in decreased pH to maintain stability. Other non-mutually exclusive mechanisms could also allow larval urchins to maintain their swimming performances




Figure 4 | Larval urchins reared at different pHs had significantly different overall shape after accounting for size difference using landmark analysis. The transformation grids render the space between the landmarks with thin plate splines to aid visualization. On these grids each lollipop shows the original starting point of a landmark with a filled circle and the shift of landmark to the target shape is indicated by a line. Each dot represents the canonical variate scores of an individual observed in each of the pH treatments, color of the dots indicate different pH treatment. CV1 and CV2 together account for over 85% of the total variance observed and their transformation grids with the corresponding transformation grids are shown as insets.




Figure 5 | Vertical velocities of both upward and downward swimming larval urchins varied with age and background flow conditions but pH had no effect on larval swimming. Averages and standard errors of larval swimming velocities from each pH treatment are shown.

despite changes in overall shape. First, larvae raised in decreased pH are smaller in size (this study) and could potentially be less calcified²⁹. These morphological differences could in turn reduce the neutral weight of an individual, and hence, a relatively smaller lifting force is required to propel the larva^{13,30}. Second, change in pH is reported to affect ciliary activities of the ciliate *Paramecium caudatum*, and a decrease in pH reduced the amount of ciliary reversals and sensitivity to KCl stimulation³¹. In sea urchins, ciliary bands play a key role in swimming and swimming speeds could be maintained through reducing number of reversal, such that more of the work done by the cilia would be directed to forward propulsion. Pluteus of other sea urchins (*Hemicentrotus pulcherrimus* and *Anthocidaris crassispina*) are able to swim at the same speed regardless of compass direction, implying a compensation for variation in gravitational pull³². Both this and our observation that plutei maintain their age-specific swimming abilities despite slower growth suggest that propulsive activities of larval urchins are under strong selective pressure, actively modulated, and highly plastic.

Ecological implications for plasticity in swimming behaviors.

Larval urchins use their ciliary bands for both food capture and swimming suggesting potential tradeoff between the two functions⁸. Volume of water filtered per unit time and hence food capture is dependent on total length of the ciliated band and its leakiness. Changes in larval size, shape, and weight may affect particle capture³³. Also, if ciliary reversal is affected by changes in pH, particle capture rate would also decrease³⁴. However, previous incubation experiment showed increased feeding rates in green urchin plutei raised under decreased pH³⁵. This was interpreted as a compensatory mechanism following a decrease in digestion efficiency.

Maintaining age-specific swimming performance under ocean acidification could also have implications for larval transport. Earlier modeling and observation on pluteus larvae of sand dollars suggested that the changes in morphology and swimming through ontogeny affect selective transport such that older larvae were less stable (higher horizontal velocity in shear) and are more likely to be transported downwards²⁸. This prediction also matches vertical distribution of larval sand dollars in the field; older larvae are found in deeper waters³⁶. If this is the case, the observed maintenance of age-specific swimming behavior regardless of developmental stage implies larvae raised under decreased pH might be transported to deeper water prematurely, e.g. before being able to settle (competency). This mismatch between behavior and development could potentially decouple competent window from approaching of settlement sites.

In conclusion, larval *S. droebachiensis* demonstrated strong phenotypic plasticity in the face of decreased pH. The changes in overall shape did not negatively impact on larval swimming ability in still and moving water. On the contrary, this shape change to a more “squat form” might have helped provide stability. Maintained performance of swimming under OA condition reinforces the importance of this ecological function to early life stages. While larval urchins demonstrated capacity to cope with decreased pH in the lab, it is important for future studies to evaluate the ecological consequences of these functional impacts acting in concert. Together, these small, sub-lethal changes could negatively impact survivorship and transport, and ultimately the distribution and abundance of populations. However, this observation also illustrates the importance to include relevant functional parameters as endpoint and to avoid over-interpretation of *change* (e.g. in size) as negative consequences.




Figure 6 | Horizontal velocities (Average \pm S.E.) of downward swimming larval urchins varied with age and background flow conditions (c,d). Downward swimming individuals (d) had higher horizontal velocities than upward swimming individuals (b) in shear i.e., larvae were crossing flow lines towards downward moving background flow (grey area). For both groups of larvae, pH had no effect on larval horizontal swimming velocities.

Methods

Larval rearing and seawater chemistry. Adult sea urchins *Strongylocentrotus droebachiensis* were collected from Droebak, Norway. Two males and two females were spawned and gametes were fertilized with protocol previously described. Larvae were kept at 10 individuals mL⁻¹ at 9°C and fed a constant diet of 150 µg C L⁻¹ *Rhodomonas* sp. (~ 6000 cells mL⁻¹) starting five days post fertilization when their mouths opened. Three pH treatments (pH_T 8.0, 7.6, and 7.2) were used, with three replicate jars for each pH level. These pH levels correspond to average surface ocean today (8.0), predicted average value for 2100 and present-day extreme (7.6), and extreme prediction for 2100 (7.2)¹⁸. pH levels were maintained through CO₂ bubbling and monitored with pH electrode and alkalinity titration¹⁸.

Larval mortality and growth. Duplicate 10 mL subsamples were taken and fixed from each replicate daily. The number of larvae was counted under a dissecting microscope and the change in relative density was used to calculate mortality rate through linear regression against time. Micrographs of twenty individuals were taken and the total body lengths (Fig. 1) were measured at different times. A logarithmic regression i.e. change in body length (mm) per the natural log of unit time (days) was used to compute the growth rate (mm ln(day)⁻¹).

Geometric morphometrics analysis. Eleven landmarks were identified including tip of the larval body, tips and bases of posterodorsal arm and preoral arms (Fig. 1). These landmarks were digitized for 10 randomly selected individuals from each replicate at four time points 4, 7, 10, and 13 days post fertilization using the imaging software Fiji³⁷. The coordinates extracted were imported into the software MorphoJ for General Procrustes Analysis³⁸. To reduce the dimensionality of the data set, a canonical variate analysis was performed and changes in overall shape were visualized with transformation grids.

Video motion analysis. Swimming behaviors were observed at four different time points (4, 7, 10, and 13 days post fertilization). Larvae were observed under no induced flow (still) and in moving water with a linear vertical velocity gradient (shear). Shear is a realistic simplification of turbulence in the natural environment for small larvae, as argued in previous studies²⁸. The same shear tank from these studies was used to create shear by enclosing a narrow inner chamber with two flanking outer chambers held at different temperatures (Suppl. Fig. 1). Observations were made at the larval rearing pH and the two flow conditions were compared: in still condition

where the temperature of both outer chambers were held at 10°C, and in shear where the temperature of the outer chambers were 0 and 20°C respectively. The temperature in the middle of the inner chamber was therefore held close to the rearing temperature at 10°C ($\pm 1^\circ\text{C}$) and checked at the beginning of each observation.

Trajectories of larval urchins (200 larvae per trial) and those of algal tracer particles (~6000 cells mL⁻¹) were recorded using a modified HD webcam (Logitech Pro Cam 7200) equipped with a CCTV lens at 7.5 fps under fiber optic lights illumination. Two, five minutes long video clips were first collected in still water and two additional clips in shear after a ten minute acclimation time. Video was processed and analyzed using Avidemux2.4 and Tracker3D^{15,39}. Smoothing splines were applied to each trajectory to remove frame rate noise and to differentiate the overall direction of travel (axis) from the helical path (path). Swimming speed was computed based on the path, net vertical and horizontal velocity based on the axis. Flow subtraction is applied to all velocities calculation because the observed larval movement is the sum of both the passive transport by the moving water and the individual swimming behavior⁴⁰. The background flow was computed by binning the algal trajectories from 1500 frames into 13 \times 17 spatial bins, and the flow velocity in each bin was the median of all the trajectories passing through each bin. The 20°C temperature difference used in the observation chamber resulted in shear ranging from -0.6 to 0.25 s⁻¹. Given the horizontal velocity generated was negligible, only the vertical component of the larval speed was corrected. Data were divided into up-swimming and down-swimming individuals based on overall displacement for statistical analysis, such that up-swimmers are those with their final positions above their starting position (i.e. a positive change z-coordinates) and down-swimmers are those with the final positions below their starting position (i.e. a negative change in z-coordinates). In this current scheme, the down-swimmers include both individuals that are sinking through ciliary arrest and those that are actively swimming downward through ciliary reversal.

Statistical Analysis. The statistical analyses were performed using SAS or PASW software. Statistical assumption of normality and homogeneity of variance were tested with Shapiro-Wilk test and Levene's test. For ANCOVAs, additional test of homogeneity of regression slope and linearity in relation between covariate and dependent variable were performed with scattered plot and linear regressions. pH and replicate effect on carbonate chemistry were tested using an ANOVA. pH and replicate effect on mortality and survival were tested using an ANCOVA with age as covariate. pH effect on shape (CV scores) was tested using an ANOVA and the scores were also linearly regressed against centroid sizes. Effect of pH, flow, and their



interactions on larval swimming was tested using an ANCOVA with age as covariate. Post-hoc Tukey's tests were performed where appropriate. All significant levels were set at $p < 0.05$.

- Grosberg, R. K. & Levitan, D. R. For adults only? Supply-side ecology and the history of larval biology. *Trends in Ecology & Evolution* **7**, 130–133 (1992).
- Gaines, S. & Roughgarden, J. Larval settlement rate: a leading determinant of structure in an ecological community of the marine intertidal zone. *Proceedings of the National Academy of Sciences* **82**, 3707–3711 (1985).
- Metaxas, A. Behaviour in flow: perspectives on the distribution and dispersion of meroplanktonic larvae in the water column. *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 86–98 (2001).
- North, E. *et al.* Vertical swimming behavior influences the dispersal of simulated oyster larvae in a coupled particle-tracking and hydrodynamic model of Chesapeake Bay. *Marine Ecology Progress Series* **359**, 99 (2008).
- Levin, L. A. Recent progress in understanding larval dispersal: new directions and digressions. *Integrative and Comparative Biology* **46**, 282–297 (2006).
- Emlet, R. B. Functional Constraints on the evolution of larval forms of marine-invertebrates - Experimental and comparative evidence. *American Zoologist* **31**, 707–725 (1991).
- Chan, K. Y. K. Biomechanics of larval morphology affect swimming: Insights from the sand dollars. *Dendraster excentricus*. *Integrative and Comparative Biology* **52**, 458–469 (2012).
- Strathmann, R. R. & Grünbaum, D. Good eaters, poor swimmers: compromises in larval form. *Integrative and Comparative Biology* **46**, 312–322, doi:10.1093/icb/ijc031 (2006).
- Sheridan, J. A. & Bickford, D. Shrinking body size as an ecological response to climate change. *Nature Climate Change* **1**, 401–406 (2011).
- Dupont, S., Dorey, N. & Thorndyke, M. What meta-analysis can tell us about vulnerability of marine biodiversity to ocean acidification? *Estuarine, Coastal and Shelf Science* **89**, 182–185 (2010).
- Hart, M. W. & Strathmann, R. R. Functional consequences of phenotypic plasticity in echinoid larvae. *Biological Bulletin* **186**, 291 (1994).
- Emlet, R. B. Locomotion, drag, and the rigid skeleton of larval echinoderms. *Biological Bulletin* **164**, 433–445 (1983).
- Grünbaum, D. & Strathmann, R. R. Form, performance and trade-offs in swimming and stability of armed larvae. *Journal of Marine Research* **61**, 659–691 (2003).
- Clay, T. W. & Grünbaum, D. Swimming performance as a constraint on larval morphology in plutei. *Marine Ecology Progress Series* **423**, 185–196 (2011).
- Chan, K. Y. K., Grünbaum, D. & O'Donnell, M. J. Effects of ocean-acidification-induced morphological changes on larval swimming and feeding. *Journal of Experimental Biology* **214**, 3857–3867 (2011).
- Pepper, R. E., Variano, E. & Koehl, M. Turbulent flow from a microscopic organism's perspective: What does it feel like to be tiny in the ocean? *Bulletin of the American Physical Society* **58** (2013).
- Koehl, M. & Hadfield, M. G. Hydrodynamics of larval settlement from a larva's point of view. *Integrative and Comparative Biology* **50**, 539–551 (2010).
- Dorey, N., Lançon, P., Thorndyke, M. & Dupont, S. Assessing physiological tipping point of sea urchin larvae exposed to a broad range of pH. *Global Change Biology* **19**, 3355–3367 (2013).
- Vaughn, D. & Allen, J. D. The peril of the plankton. *Integrative and Comparative Biology* **50**, 552–570 (2010).
- Lamare, M. D. & Barker, M. F. In situ estimates of larval development and mortality in the New Zealand sea urchin *Evechinus chloroticus* (Echinodermata: Echinoidea). *Marine Ecology Progress Series* **180**, 197–211 (1999).
- Ebert, T. A. Growth and survival of post-settlement sea urchins. *Developments in Aquaculture and Fisheries Science* **32**, 79–102 (2001).
- Botello, G. & Krug, P. J. Desperate larvae revisited: age, energy and experience affect sensitivity to settlement cues in larvae of the gastropod. *Alderia* sp. *Marine Ecology Progress Series* **312**, 149–159 (2006).
- Matson, P. G., Pauline, C. Y., Sewell, M. A. & Hofmann, G. E. Development under elevated pCO₂ conditions does not affect lipid utilization and protein content in early life-history stages of the purple sea urchin, *Strongylocentrotus purpuratus*. *Biological Bulletin* **223**, 312–327 (2012).
- Yu, P. C. *et al.* Growth attenuation with developmental schedule progression in embryos and early larvae of *Sterechinus neumayeri* raised under elevated CO₂. *PLoS ONE* **8**, e52448, doi:10.1371/journal.pone.0052448 (2013).
- Gonzalez-Bernat, M. J., Lamare, M., Uthicke, S. & Byrne, M. Fertilisation, embryogenesis and larval development in the tropical intertidal sand dollar *Arachnoides placenta* in response to reduced seawater pH. *Marine Biology* **160**, 1927–1941 (2013).
- Dupont, S., Dorey, N., Stumpp, M., Melzner, F. & Thorndyke, M. Long-term and trans-life-cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus droebachiensis*. *Marine Biology* **160**, 1835–1843 (2013).
- Mogami, Y., Ishii, J. & Baba, S. A. Theoretical and experimental dissection of gravity-dependent mechanical orientation in gravitactic microorganisms. *Biological Bulletin* **201**, 26–33 (2001).
- Clay, T. & Grünbaum, D. Morphology–flow interactions lead to stage-selective vertical transport of larval sand dollars in shear flow. *Journal of Experimental Biology* **213**, 1281–1292 (2010).
- Clark, D., Lamare, M. & Barker, M. Response of sea urchin pluteus larvae (Echinodermata: Echinoidea) to reduced seawater pH: a comparison among a tropical, temperate, and a polar species. *Marine Biology* **156**, 1125–1137 (2009).
- McDonald, K. A. & Grünbaum, D. Swimming performance in early development and the “other” consequences of egg size for ciliated planktonic larvae. *Integrative and Comparative Biology* **50**, 589–605 (2010).
- Doughty, M. J. Effect of extracellular pH on motility and K⁺-induced ciliary reversal in *Paramecium caudatum*. *Journal of Protozoology* **33**, 435–441, doi:10.1111/j.1550-7408.1986.tb05635.x (1986).
- Mogami, Y., Oobayashi, C., Yamaguchi, T., Ogiso, Y. & Baba, S. A. Negative geotaxis in sea urchin larvae: A possible role of mechanoreception in the late stages of development. *Journal of Experimental Biology* **137**, 141–156 (1988).
- Emlet, R. & Strathman, R. Gravity, drag, and feeding currents of small zooplankton. *Science* **228**, 1016 (1985).
- Hart, M. W. Particle captures and the method of suspension feeding by echinoderm larvae. *Biological Bulletin* **180**, 12–27 (1991).
- Stumpp, M. *et al.* Digestion in sea urchin larvae impaired under ocean acidification. *Nat. Clim. Change* **3**, 1044–1049 (2013).
- Pennington, J. T. & Emlet, R. B. Ontogenetic and diel vertical migration of a planktonic echinoid larva, *Dendraster excentricus* (Eschscholtz): Occurrence, causes, and probable consequences. *Journal of Experimental Marine Biology and Ecology* **104**, 69–95 (1986).
- Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).
- Klingenberg, C. P. Morpho: an integrated software package for geometric morphometrics. *Molecular Ecology Resources* **11**, 353–357 (2011).
- Chan, K. Y. K. & Grünbaum, D. Effects of temperature and diet on larval swimming behaviors of sand dollar *Dendraster excentricus*. *Marine Ecology Progress Series* **419**, 419–429 (2010).
- Wheeler, J. *et al.* Upward swimming of competent oyster larvae *Crassostrea virginica* persists in highly turbulent flow as detected by PIV flow subtraction. *Marine Ecology Progress Series* **488**, 171–185 (2013).

Acknowledgements:

We thank J. Pineda, V. Starzack, and J. Wheeler for their input on data interpretation; N. Dorey, I. Casties, B. Lundve for technical assistance. K.C. was supported by the Postdoctoral Scholar Program at the Woods Hole Oceanographic Institution (WHOI), with funding provided by the Coastal Ocean Institute, the Croucher Foundation, and the Royal Swedish Academy of Sciences. S.D. was financially supported by the Linnaeus Centre for Marine Evolutionary Biology at the University of Gothenburg (<http://www.cemeb.science.gu.se/>) and a Linnaeus grant from the Swedish Research Councils VR and Formas. Additional funding was provided from the European Seventh Framework Programme under grant agreement 265847.

Author contributions

K.C. and S.D. conceived and designed the experiment. K.C., E.G., S.D. conducted the experiment. K.C. and S.D. analyzed the data and prepared the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/scientificreports>

Competing Financial Interests: The authors declare no competing financial interests.

How to cite this article: Chan, K.Y.K., Garcia, E. & Dupont, S. Acidification reduced growth rate but not swimming speeds of larval sea urchins. *Sci. Rep.* **5**, 9764; DOI:10.1038/srep09764 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>