



Not to be cited without prior reference to the authors

Correlation analysis of interannual variation in cephalopod landings from European waters

G J Pierce¹, M.A. Collins¹, M.M. Cunha², A. Guerra³, U. Piatkowski⁴, F. Porteiro⁵ & J.P. Robin⁶

¹ Department of Zoology, University of Aberdeen, Tillydrone Avenue, Aberdeen AB9 2TN, UK

² Instituto Portugués de Investigação Marítima, Ministério do Mar, Avenida de Brasília, 1400 Lisboa, Portugal

³ Instituto de Investigaciones Marinas, Consejo Superior de Investigaciones Científicas, r/r Eduardo Cabello, 6, Vigo-36208, Espania

⁴ Institut für Meereskunde, Universität Kiel, Dösterbrookweg 20, D-24105 Kiel, Germany

⁵ Departamento de Oceanografia e Pescas, Universidade dos Açores, Caís de St Cruz, PT-9900 Horta, Faial, Açores, Portugal

⁶ Laboratoire de Biologie et Biotechnologies Marines, Université de Caen, Esplanade de la Paix, 14032 Caen Cedex, France

Abstract

The correspondence of interannual trends in cephalopod catches across European waters is analyzed. Data on cephalopod landings and, where available, fishing effort, by European countries were compiled and entered into a correlation analysis.

Relevant fishing effort data could be obtained only for the UK and Portugal. In the UK data, CPUE was strongly correlated with landings, suggesting that landings could be used as an index of abundance. For both Scotland and Portugal there was reasonable correspondence between interannual trends in catches by different fishing gears. However, it cannot be assumed that landings are always a realistic index of abundance.

In the Northeast Atlantic and the Mediterranean, interannual trends in landings of certain cephalopod categories in adjacent countries tended to be similar both across species and across areas. There was some evidence of corresponding trends over longer time periods and on a larger spatial scale, particularly for *Loligo forbesi*, pointing to the possible role of large-scale climatic or oceanographic factors in determining cephalopod abundance.

Introduction

Cephalopods are an increasingly important fishery resource in European waters and a number of species of squid, cuttlefish and octopus are landed commercially. Recent research has focused mainly on squid. In European waters, the squid species of greatest importance to commercial fisheries are the so-called common squids (Cephalopoda: Loliginidae), *Loligo forbesi* (veined squid) and *L. vulgaris* (European squid). Both are widely distributed in Northeast Atlantic and Mediterranean waters, although *L. forbesi* has a more northerly distribution and is scarce in southern Portuguese and Mediterranean waters, while *L. vulgaris* has a more southerly distribution and is rarely found in Scottish waters. An isolated, genetically distinct, population of *L. forbesi* is found in the Azores (Martins 1982, Brierley *et al.*, 1993, 1995; Pierce *et al.*, 1994a) while a sub-species of *L. vulgaris*, *L.v. reynaudii*, occurs on the southern and western coasts of Africa.

Landings of common squids (*Loligo* spp.) in European countries vary widely from year to year, apparently unrelated to fishing effort in those countries which record fishing effort (Pierce *et al.*, 1994b). Since

1990, annual landings in several European countries have decreased. However, this apparent trend has not been subject to rigorous analysis and previous data suggest that there may be cyclic variation in abundance (Pierce *et al.*, 1994b). The short (annual) life-cycle (Coelho *et al.*, 1994; Guerra & Rocha, 1994; Moreno *et al.*, 1994; Pierce *et al.*, 1994c; Porteiro & Martins, 1994; Collins *et al.*, 1995) may in part account for the high variability in abundance, since the populations are not buffered against fluctuations in recruitment or excessive mortality of breeding adults (Pierce & Guerra, 1994).

It is commonly thought that cephalopods such as *Loligo* spp. may be highly sensitive to variation in hydrographic conditions (Caddy, 1983; Forsythe, 1993). Indirect evidence that this is the case may be obtained if common trends in abundance are apparent across a wide geographical area.

The present paper examines correlations between interannual trends in squid and other cephalopod landings in different EC countries.

Materials and Methods

ICES supplied data on squid landings 1973-94 by European countries fishing in the Northeast Atlantic and data on total annual cephalopod landings in EC countries 1965-92 from the Northeast Atlantic (FAO area 27) and Mediterranean (FAO area 37) were assembled from FAO Yearbooks of Fishery Statistics, Catches and Landings (Food & Agriculture Organization, Rome). Long-finned (lolinid) squid landings from the NE Atlantic were also compiled by direct access to national databases in England and Wales (1980-94), Scotland (1904-94), France (1960-94), Spain (1983-94), Portugal (1960-93) and the Azores (1948-93). Where possible the data were subdivided by area and/or gear. Where effort data were available, overall CPUE was derived (total landings divided by total effort).

Similarities in inter-annual trends for different data categories were evaluated by non-parametric correlation analysis (BMDP statistical software). Due to the very large number of possible comparisons, the analysis is restricted to answering specific questions. The first three questions relate to the utility of landings data as an abundance index:

- (a) Are the same trends seen in landings and CPUE?
 - (b) Are the same trends seen for different gear types?
 - (c) Are the same trends seen in catches by different countries fishing in the same area?
- Two further questions relate to the trends over large areas:
- (d) Are the same trends seen for different cephalopod categories within the same area?
 - (e) Are the same trends seen for a particular species in different areas?

Comments on the categorization of cephalopod landings in published data

Cephalopods are classified into a series of categories by FAO, the use of which has been inconsistent, and not all of which are mutually exclusive. Some cephalopod landings are recorded as "Cephalopods not elsewhere identified". Many EC countries recorded squid landings from area 27 as "Squid (Lolinidae or Ommastrephidae)" up to approx. 1980, after which landings were categorized as "*Loligo* spp.". For Scotland, Northern Ireland, Ireland and Portugal it is reasonable to assume that, prior to 1980, the squid landed was usually *Loligo*. The category "Squid (Lolinidae or Ommastrephidae)" continues to appear until 1985 in data for England & Wales, possibly reflecting participation in the *Todarodes sagittatus* fishery in the early 1980s. France recorded squid landings as "*Loligo* spp." until 1980 and as "Squid (Lolinidae or Ommastrephidae)" thereafter. However, landings are probably still mainly of *Loligo forbesi* and/or *L. vulgaris*. Spain has switched from always recording "*Loligo* spp.", "*Illex illecebrosus*" and "*Todarodes sagittatus*" separately to recording an increasing proportion of landings as "Squid (Lolinidae or Ommastrephidae)". Because Spain has a fishery for ommastrephids, this category is certain to include a mixture of species, including the ommastrephid *Todaropsis eblanae* as well as the above mentioned species. Similar problems of categorization arise less frequently for cuttlefish and octopus, but Spain and Portugal have both switched between using "Sepiidae or Sepiolidae" and "*Sepia officinalis*" for all cuttlefish landings. Also, Spain normally records octopus landings as "Octopodidae" but during 1973-6 recorded all octopus landed as "*Eledone*". France recorded landings of "*Octopus vulgaris*" in 1975-76 but normally recorded "Octopodidae". Note that the *Illex* species landed by Spain is more likely to be *Illex coindetii* than *I. illecebrosus*, since the latter is a Northwest Atlantic species.

Data for FAO area 37 (Mediterranean) present fewer problems, but France and Greece both switched (in 1981) from recording "Sepiidae or Sepiolidae" to recording "*Sepia officinalis*". Spain and Greece recorded

landings of *Illex coindetii*, although Greece switched in 1981 to recording "Squid (Loliginidae or Ommastrephidae)". Greece also started recording *Octopus vulgaris* in 1983, as well as Octopodidae.

Results

(a) Landings and CPUE for *Loligo*

This analysis used data extracted directly from National databases. In data on *Loligo* landings from Scotland (Fig. 1) and England + Wales, landings and CPUE were significantly correlated, for individual gear-types and overall (Table 1). For Portugal, annual *Loligo* landings from the trawl fishery (1988-1992) were negatively correlated with CPUE but the trend is not statistically significant. Data for other countries were not available.

(b) Comparisons between gears for *Loligo* landings and CPUE

This analysis used data extracted directly from National databases. Landings of *Loligo* in Scotland by different gear categories were correlated, except for seine nets versus heavy trawling gear and the same was true of CPUE (Fig. 2, Table 2). Landings of *Loligo* in England by different gears were generally not correlated, except for heavy trawls versus seine, and this also applies for CPUE data. Landings of loliginids in Portugal from the artisanal fishery were correlated with trawl and with seine landings.

(c) Comparisons between landings of *Loligo* by different countries fishing in the same area

This analysis used data provided by ICES. The statistical power of the analysis is restricted by incomplete data. Possibly many missing values actually represent zero catches but it was necessary to assume they were missing data. Interrannual trends in landings of *Loligo* by different countries fishing in the same ICES fishery subdivision were well correlated in some important squid fishery areas, particularly the west coast of Scotland (VIa; Fig. 3) and the Irish Sea areas (VIIa) but not in other areas (Table 3). In the case of area VIa, it is possible that landings by Ireland are poorly correlated with those of other countries because landings from VIb were included with those from VIa.

(d) Different cephalopod species within the same area

This analysis used data published by FAO. For all countries except Spain, the different FAO categories of squid, cuttlefish and octopus in area 27 were combined for analysis, but the squid can be assumed to have been mostly *Loligo* spp.

For area 27 as a whole, there were no significant correlations between landings of the three main taxa, but positive correlations were seen in landings by Spain, except for *Todarodes* (Fig. 4), and France, also between squid and cuttlefish for England and Wales (Table 4).

More categories of cephalopods are distinguished in Mediterranean (area 37). Overall, there were positive correlations between many of these categories (Table 5). For individual countries (Italy, Greece, Malta and Cyprus), some positive correlations between species were seen, although *Illex* landings in Greece were negatively correlated with those of other species, while *Sepia* and *Todarodes* landings in Italy were negatively correlated (Fig. 5). Squid and octopus landings in Spain were also negatively correlated.

(e) Trends across large areas

ICES data were available for many countries: the two examples chosen here are for countries fishing across different, but overlapping, areas. ICES landings data for England & Wales showed comparable interannual trends in *Loligo* landings for many fishery areas, with only landings from the northern North Sea (IVa) being negatively correlated with landings from elsewhere (Table 6). Spain fishes squid from a wider area, and positive correlations are seen between landings in Spain from the west coasts of Scotland (VI) and Ireland (VIIbc), the Celtic Sea (VIIg-k), the Bay of Biscay (VIII) and west Portugal (IX) (Fig. 6, Table 6).

The FAO data show some correspondence between landings of *Loligo* from area 27 by different European countries, and this was also the case for cuttlefish (Table 7). In area 37, landings of these cephalopod categories by different countries were as often negatively correlated as positively correlated (Table 8), although landings of octopods by different countries were mostly positively correlated. A difficulty with interpreting these trends is that the fishing areas of different countries overlap extensively.

Comparisons between landings from FAO areas 27 and 37 are possible using area totals and for the two countries, France and Spain, which routinely fish for cephalopods in both areas. Squid landings from the two areas tended to be positively correlated, octopus landings negatively correlated. Overall cuttlefish landings from the two areas were correlated but landings of cuttlefish in France from the two areas were negatively related (Table 9).

Longer time series of data for long-finned squid (*Loligo*) landings were obtained directly from National databases held in the Azores, Portugal and Scotland. Additional data series were obtained for France and Spain but covered shorter periods. Previous comments about the FAO data notwithstanding, it appeared from the new Spanish data (1983-94) that published (area 27) totals for FAO categories "Loligo" and "Squid (Loliginidae or Ommastrephidae)" should be added to approximate to total loliginid landings. These new FAO totals were used for pre-1983 data and combined with the new 1983-94 data to create a new data series. The new French data (1980-94) were similarly grafted onto earlier FAO data to create a new data series. Landings by Scotland were correlated with those in France and the Azores, while landings in Spain were negatively correlated with those in the Azores (Table 10). The pattern of landings since 1980 appears strikingly similar for Scotland, the Azores, Portugal and France (Fig. 7). However, statistically, only the correlation between Scotland and the Azores is significant over this period (Table 10).

Discussion

This kind of analysis will always be limited by the quality of the data available, but collection and reporting of data on cephalopods, which are not quota species in EC waters, is particularly poor (Anon., 1993, 1994; Donoso-Perez & Forest, 1993; Pierce & Guerra, 1994). The main problems are:

- (a) Limited objectives of data collection by National Governments. Many countries record total cephalopod landings but not all record associated data on location or fishing effort.
- (b) Low accuracy: there is ample scope for human error in the way data are collected. In the UK fishery we are aware of examples of large amounts of squid going unrecorded due to categorization as "other species" or being landed by small boats, also of quota species being mis-reported as squid.
- (c) Inconsistent categorization of cephalopods: continuity of data over time is compromised by splitting and lumping of categories (see above).
- (d) Incomplete reporting to International bodies.

Additionally, interpreting trends in such data series is difficult. When a large number of variables is being analysed, correlation analysis will inevitably generate some spurious correlations, and correlations may not indicate any underlying causal relationship. Temporal autocorrelation may also be a problem, although for short-lived species with non-overlapping generations, abundances in two adjacent years may probably be regarded as independent, because the only direct link is through recruitment. In any case, the approach used in the present paper represents one of the few feasible means of addressing the questions raised.

Pierce *et al.* (1994b) argued that, in the Scottish fishery, landings of *Loligo* were a reasonable index of abundance because they were strongly correlated with CPUE. Even though squid may be targeted by Scottish fishermen in years of high abundance, the impact of this on overall CPUE (by the entire fleet) appears to have been negligible. This conclusion is supported by the present analysis for Scottish and English data but not for Portuguese data. Thus, for countries which report landings but not effort there must remain doubt as to whether landings are related to abundance.

The UK data also show that landings and CPUE recorded for different gear-types do not always record the same picture. Similarly, trends in landings by different countries fishing in the same ICES fishery subdivision did not always correspond. This is likely to reflect differences in fishing effort or methods rather than abundance, so it is generally preferable to treat data from different countries separately.

Analyses of both FAO and ICES datasets showed the existence of parallel trends both across species and across areas: there are certainly more significant positive correlations than would have been expected by chance alone. At present however we can only provisionally suggest that this might, to some extent, reflect a common response to changing environmental conditions.

The analysis of data obtained by direct access to National databases was particularly interesting, revealing striking correspondence between trends in landings of *Loligo* in Scotland and the Azores over the last 15 years. Landings in France and Portugal showed somewhat similar trends, while the pattern of Spanish landings was entirely different. It may be relevant that Scotland and the Azores are unique in landing only *Loligo forbesi*, albeit possibly different subspecies, whereas the other countries land a mixture of *Loligo forbesi* and *L. vulgaris*. If this correspondence between trends in midAtlantic and Continental shelf landings can be shown to reflect trends in abundance rather than a common response of fisheries to changing technology and

market forces - and in the case of Scotland we believe it can - this represents the best evidence yet available that squid abundance could be controlled by large-scale climatic or oceanographic factors. In a complementary analysis, Pierce (1995) showed that squid landings in Scotland could be well-predicted from sea surface temperature data.

Acknowledgements

UK landings data were supplied by SOAFD and MAFF. French data were supplied by the Affaires Maritimes. Additional information on squid landings was supplied by ICES and we would like to thank Maria Zarecki for her assistance with extracting these data. The analysis was funded by a grant from the European Commission, (project no. AIR 1 CT92 0573) and we are grateful to many colleagues working on squid and associated with this project for discussions and contributions to these ideas.

References

- Anon., 1993. Report of the Study Group on Cephalopod Biology. Kiel, 21-22 September 1992. *International Council for the Exploration of the Sea. C.M. 1993/K:66.*
- Anon., 1994. Report of the Study Group on The Life History and Assessment of Cephalopods. *International Council for the Exploration of the Sea. CM 1994/K:7*
- Brierley, A.S., Thorpe, J.P., Clarke, M.R., Martins, H.R., 1993. A preliminary biochemical genetic investigation of the population structure of *Loligo forbesi* Steenstrup, 1856 from the British Isles and the Azores. In Okutani, T., O'Dor, R.K. and Kubodera, T. (eds.) *Recent advances in cephalopod fisheries biology*. Tokai University Press, Tokyo, 61-69.
- Brierley, A., Thorpe, J., Pierce, G.J., Clarke, M.R., Boyle, P.R., 1995. Genetic variation in *Loligo forbesi* in the Northeast Atlantic. *Mar. Biol.*, **121**, 501-508.
- Caddy, J.F., 1983. The cephalopods: factors relevant to their population dynamics and to the assessment and management of stocks. In *Advances in assessment of world cephalopod resources*, edited by J.F. Caddy, FAO Fish. Tech. Pap., **231**: 416-449.
- Coelho, M.L., J. Quintela, V. Bettencourt, G. Olavo and H. Villa, 1994. Population structure, maturation patterns and fecundity of the squid *Loligo vulgaris* from southern Portugal. *Fish. Res.*, **21**, 87-102.
- Collins, M.A., Burnell, G.M. & Rodhouse, P.G., 1995. Reproductive strategies of male and female *Loligo forbesi* (Cephalopoda: Loliginidae). *J. Mar. Biol. Ass. U.K.*, **75**, 621-634.
- Donoso-Perez, R. & Forest, A., 1993. Analyse des pêcheries françaises de céphalopodes dans le golfe de Gascogne (division VIII du CIEM). ICES C.M. 1993/K:51
- Forsythe, J.W., 1993. A working hypothesis of how seasonal temperature change may impact the field growth of young cephalopods. In: Okutani, T., O'Dor, R. and Kubodera, T. (eds), *Recent advances in Cephalopod Fisheries Biology*, Tokai University Press, Tokyo, pp. 133-143.
- Guerra, A. and F. Rocha, 1994. The life history of *Loligo vulgaris* and *Loligo forbesi* (Cephalopoda: Loliginidae) in Galician waters (NW Spain). *Fish. Res.*, **21**, 43-70.
- Martins, H.R. 1982. Biological studies of the exploited stock of *Loligo forbesi* (Mollusca: Cephalopoda) in the Azores. *J. Mar. Biol. Ass. U.K.*, **62**, 799-808.
- Moreno, A., M.M. Cunha and J.M.F. Pereira, 1994. Population biology of veined squid (*Loligo forbesi*) and European Squid (*Loligo vulgaris*) from the Portuguese coast. *Fish. Res.*, **21**, 87-102.
- Pierce, G.J., 1995. Stock assessment with a thermometer: correlations between sea surface temperature and landings of squid (*Loligo forbesi*) in Scotland. *International Council for the Exploration of the Sea. C.M. 1995/K:21.*
- Pierce, G.J., Boyle, P.R., Bailey, N. & Hastie, L.C., 1994. Distribution and abundance of the fished population of *Loligo forbesi* in UK waters: analysis of research cruise data. *International Council for the Exploration of the Sea. C.M. 1994/K:22*
- Pierce, G.J., Boyle, P.R., Hastie, L.C. & Key, L., 1994c. The life history of *Loligo forbesi* in Scottish waters. *Fish. Res.*, **21**, 17-41
- Pierce, G.J., Boyle, P.R., Hastie, L.C. & Shanks, A., 1994b. Distribution and abundance of the fished population of *Loligo forbesi* in UK waters: analysis of fishery data. *Fish. Res.*, **21**, 193-216.
- Pierce, G.J. & Guerra, A., 1994. Stock assessment methods used for cephalopod fisheries. *Fish. Res.* **21**, 255-285.
- Pierce, G.J., Thorpe, R.S., Hastie, L.C., Brierley, A.S., Boyle, P.R., Guerra, A., Jamieson, R. & Avila, P., 1994a. Geographic variation in *Loligo forbesi* in the Northeast Atlantic: analysis of morphometric data and tests of causal hypotheses. *Mar. Biol.*, **119**, 541-548.
- Porteiro F. and H. R. Martins, 1994. Population biology of *Loligo forbesi* Steenstrup, 1856 (Mollusca: Cephalopoda) in the Azores: sample composition and maturation of squid caught by jigging. *Fish. Res.*, **21**, 103-114.

Table 1

Correlations between *Loligo* landings and CPUE

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. Data for individual gear-types are from 1980-94 (i.e. N=15). The all-gear data for Scotland goes back to 1974 (N=21).

Gear	Scotland	England
Heavy trawl	0.946	0.957
Light trawl	0.986	0.950
Pair trawl	0.986	0.975
Seine	0.954	0.807
All	0.979	0.589

Table 2

Correlations between gear-types for *Loligo* landings and CPUE.

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. Data for individual gear-types are from 1980-94 (N=15) in Scotland and England + Wales, and 1986-93 (N=8) in Portugal. Heavy = heavy otter trawl, Light = light otter trawl, Pair = pair trawl, Seine = demersal seines, Art. = artisanal fishery.

Landings, Scotland				CPUE, Scotland					
Gear	Heavy	Light	Pair	Seine	Gear	Heavy	Light	Pair	Seine
Light	0.557				Light	0.696			
Pair	0.511	0.864			Pair	0.579	0.782		
Seine	0.050	0.664	0.711		Seine	0.025	0.564	0.611	
All	0.664	0.982	0.857	0.632	All	0.725	0.989	0.775	0.557

Landings, England & Wales				CPUE, England & Wales					
Gear	Heavy	Light	Pair	Seine	Gear	Heavy	Light	Pair	Seine
Light	-0.186				Light	-0.196			
Pair	-0.393	-0.036			Pair	-0.325	0.139		
Seine	0.554	0.014	0.150		Seine	0.679	0.104	-0.264	
All	-0.486	0.271	0.700	0.079	All	-0.214	0.032	0.896	-0.225

Landings, Portugal			
Gear	Trawl	Art.	Seine
Art.	0.810		
Seine	0.527	0.898	
All	0.952	0.857	0.647

Table 3

Correlations between landings of squid* by different countries fishing in the same area.

Significant correlations are indicated by bold type. The analysis is based on ICES data (1973-94). *Landings of "Loligo" and "Squid not elsewhere included" were summed because the use of these categories was ambiguous. For most countries, the squid can be assumed to be *Loligo*. Scot = Scotland, N. Ire = Northern Ireland, I. Man = Isle of Man, Engl = England & Wales, Belg = Belgium, Fran = France, Port = Portugal. IVa = Northern North Sea, IVb = central North Sea, IVc = Southern North Sea, Vla = west of Scotland, Vlb = Rockall, VIIa = Irish Sea, VIIbc = West of Ireland, VIIId = Eastern English Channel, VIIe = Western English Channel, VIIIf = Bristol Channel, VIIg-k = Celtic Sea, VIII = Bay of Biscay, IX = West Portugal.

IVa (N=9a, 14b, 15c)		IVb (N=17a, 20b)		IVc (N=13a, 15b, 7)						
Engl	Fran	Engl	Fran	Engl	Fran					
Fran	0.210 ^b	-0.258 ^c	Engl	0.397 ^a						
Scot	0.018 ^b		Fran	0.279 ^a	0.509 ^a					
Vla (N=13a, 16a, 19a, 10a, 11a, 14a, 17a, 8a, 12a)	Engl	Fran	Ireland	Scot						
Fran	0.495 ^a		Engl	0.205 ^a						
Ireland	-0.242 ^a	-0.219 ^a	Fran	0.060 ^a	0.300 ^a					
Scot	0.409 ^a	0.255 ^a	Scot	0.071 ^a	-0.316 ^a					
Spain	0.571 ^a	0.618 ^a	Spain		-0.400 ^a					
VIIa (N=19a, 11a, 12a, 16a, 14a, 18a, 21a, 8a)	Belg	Engl	Fran	I Man	Ireland	N. Ire	VIIbc (N=5a, 9a, 14a)	Engl	Fran	Ireland
Engl	0.215 ^a		Fran	-0.158 ^a						
Franc	0.132 ^b	0.557 ^a	Ireland	0.418 ^a	0.359 ^a					
I. Man	0.046 ^a	0.817 ^a	Spain	0.892 ^a	0.170 ^a					
Ireland	0.235 ^a	0.255 ^a			-0.299 ^a					
N. Ire	-0.119 ^a	0.652 ^a								
Scot	0.020 ^a	0.738 ^a								
VIIId (N=21a, 16a)	Belg	Engl	VIIe (N=19a, 13a)	Belg	Engl	VIIIf (N=18a, 15a, 13a)	Engl	Belg	Engl	
Engl	0.055 ^a		Engl	0.115 ^a		Engl	-0.127 ^a			
Fran	0.010 ^a	0.590 ^a	Fran	0.232 ^a	0.658 ^a	Fran	-0.203 ^a	0.577 ^a		
VIIg k (N=21a, 14a, 17a, 16a, 12a)	Belg	Engl	Fran			Engl	-0.048 ^a			
Engl	-0.047 ^a		Fran			Fran	0.500 ^a	-0.200 ^a		
Fran	0.459 ^a	0.205 ^a	Spain			Spain	-0.500 ^a	0.500 ^a		
Ireland	-0.088 ^a	0.759 ^a					-0.220 ^a			
Spain	-0.188 ^a	-0.196 ^a						0.117		

Table 4

Correlations between landings of different cephalopod categories: FAO area 27.

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. The analysis is based on FAO data for landings from area 27, 1965-1992. Only countries recording more than one category* of cephalopod are included. *Categories have been combined where appropriate (see text). Cuttle = "*Sepia officinalis*" + "Sepiidae or Sepiolidae", Oct = "Octopodidae" or "*Octopus vulgaris*" or "Eledone spp.", Squid = "*Loligo* spp." or "Squid (Loliginidae or Ommastrephidae)" or "*Illex illecebrosus*" or "*Todarodes sagittatus*". Tod. = "*Todarodes sagittatus*".

England & Wales (N=10)		Channel Islands (N=9)		Ireland (N=6)	
Cuttle	Squid	Cuttle	Squid	Cuttle	Squid
0.675		0.098			
Oct	-0.119	-0.480			
France (N=19 ^a , 15 ^a)		Spain (N=20 ^a , 8 ^a , 27, 28 ^a , 19 ^a)		Area 27 totals (N=28)	
Cuttle	Squid	Cuttle	Squid	Cuttle	Squid
0.691^a		0.447^a		0.136	
Oct	0.644^a	0.716^a		Oct	0.413^a
Area 27 totals (N=28)					
Cuttle	Squid	Cuttle	Squid	Cuttle	Squid

Table 5

Correlations between landings of different cephalopod categories: FAO area 37.

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. The analysis is based on FAO data for landings from area 37, 1965-1992. Only countries recording more than one category* of cephalopod are included. *Categories have been combined where appropriate (see text). Cuttle = "*Sepia officinalis*" + "Sepiidae or Sepiolidae", Oct = "Octopodidae" or "*Octopus vulgaris*" or "Eledone spp.", Squid = "*Loligo* spp." or "Squid (Loliginidae or Ommastrephidae)" or "*Illex coindetii*" or "*Todarodes sagittatus*". Tod. = "*Todarodes sagittatus*", O.vul = "*Octopus vulgaris*", Eled. = "Eledone spp."

Spain (N=20 ^a , 11 ^a , 28 ^a)				France (N=19)			
Lol.	Illex	Squid	Cuttle	Lol.	Illex	Squid	Cuttle
0.477^b				0.388			
Squid	-			O. vul	-0.060	0.037	
Cuttle	-0.012 ^a			0.058^a			
Oct	-0.701^a	-0.037^a	-0.740^a	-0.381^a			
Italy (N=28)				Malta (N=28)			
Lol.	Tod.	Sepia	O. vul	Lol.	Tod.	Sepia	O. vul
0.260				0.753			
Sepia	0.086	-0.439		0.769			
O. vul	0.457	0.796	-0.232				
Eled.	0.334	0.397	-0.199	0.536			
Greece (N=20 ^a , 8 ^a , 10 ^a)				Cyprus (N=20)			
Lol.	Illex	Squid	Cuttle	Lol.	Illex	Squid	Cuttle
-0.696 ^a				0.869			
Squid	-						
Cuttle	0.877^a	-0.823^a	0.823^a				
O. vul	0.891^a	-	0.891^a	0.952^a			
Oct	0.791^a	-0.609 ^a	0.829^a	0.845^a	0.709^a		
Area 37 totals (N=28 or N=15 ^a)							
Lol.	Tod.	Illex	Squid	O. vul	Eled.	Cuttle	
0.204							
Illex	-0.338 ^a	0.205^a					
Squid	0.772	0.705	-0.010 ^a				
O. vul	0.199	0.827	-0.007 ^a	0.593			
Eled.	-0.027	0.338	0.171 ^a	0.196	0.458		
Cuttle	0.550	0.581	-0.096 ^a	0.694	0.744	0.287	
Oct	0.127	0.878	0.254 ^a	0.594	0.894	0.546	0.639

Table 8

Correlations between landings by different countries: FAO area 37.

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. Codes for ICES Fisheries Subdivisions: IVa = Northern North Sea, IVb = central North Sea, IVc = Southern North Sea, VIa = west of Scotland, VIb = Rockall, VIIa = Irish Sea, VIIb = West of Ireland, VIIId = Eastern English Channel, VIIe = Western English Channel, VIIIf = Bristol Channel, VIIg k = Celtic Sea, VIII = Bay of Biscay, IX = West Portugal.

												<i>Top right: Spain (N=5^a, 12^b, 8^c, 15^d)</i>			
IVa	IVb	IVc	VIa	VIb	VIIa	VIIbc	VIIId	VIIe	VIIIf	VIIg k	VIII	IX			
IVa	-0.257 ^a														
IVb		0.596 ^b													
IVc	0.350 ^b	0.049 ^c	0.460 ^c		0.700 ^a		0.284 ^b				0.590 ^b	0.526 ^b	0.756 ^b		
VIa	0.260 ^c	0.049 ^c	0.460 ^c		0.057 ^d		0.024 ^c				0.228 ^c	0.476 ^c	0.000 ^f		
VIb	0.242 ^c	-0.109 ^d	0.264 ^d		0.057 ^d										
VIIa	-0.362 ^c	0.167 ^b	0.469 ^b	0.449 ^b	0.216 ^c						0.846 ^d	-0.323 ^d	0.492 ^d		
VIIb	-0.567 ^c	0.147 ^b	0.009 ^c	0.368 ^b	-0.538 ^c	0.028 ^b									
VIIId	-0.437 ^c	0.448 ^b	0.248 ^b	0.387 ^b	0.177 ^c	0.557 ^b	0.390 ^b								
VIIe	-0.316 ^c	0.319 ^b	0.723 ^b	0.617 ^b	0.116 ^c	0.708 ^b	0.349 ^b	0.559 ^b							
VIIIf	0.009 ^c	0.043 ^c	0.340 ^c	0.526 ^b	0.161 ^c	0.541 ^b	0.396 ^b	0.481 ^b	0.598 ^b						
VIIg k	-0.496 ^c	0.615 ^b	0.764 ^b	0.048 ^c	-0.172 ^c	0.460 ^b	0.681 ^b	0.227 ^c	0.532 ^c	0.290 ^c	-0.121 ^d	0.551 ^c			
VIII	-0.691 ^c	-0.097 ^b	-0.639 ^c	0.280 ^b	-0.207 ^b	0.225 ^b	0.185 ^b	0.620 ^b	0.346 ^b	0.122 ^b	0.274 ^b	0.333 ^c			
IX															
<i>Bottom left: England & Wales (N=13^a, 11^b, 14^c, 8^d, 18^e, 16^f, 20^g, 10^h, 12ⁱ, 1P, 9^j, 15^k, 19^l, 17^m, 21ⁿ)</i>															

Table 6

Correlations between landings from different ICES areas.

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. The analysis is based on FAO data for landings from area 27, 1965-1992. Fran = France, Gree = Greece, Ital = Italy, Cypr = Cyprus, Ts = *Todarodes sagittatus*, Ov = *Octopus vulgaris*, El = *Eledone* spp..

<i>"Loligo spp."</i> (N=28 ^a , 21 ^b , 8 ^c)				<i>Illex</i> (N=4 ^a , 11 ^b , 8 ^c)			
Fran	Spain	Italy	Gree	Fran	Spain	Italy	Gree
Spain	-0.326 ^a						
Ital	-0.320 ^a	-0.147 ^b					
Gree	-0.257 ^a	-0.234 ^b	0.010 ^c				
Malta	0.347 ^a	-0.322 ^b	0.147 ^b	-0.485 ^b			

<i>Cuttlefish</i> (N=19 ^a , 21 ^b , 28 ^c)					
Fran	Spain	Italy	Gree		
Spain	-0.137 ^a				
Ital	0.535 ^a	-0.070 ^b			
Gree	-0.791 ^a	-0.035 ^b	-0.518 ^b		
Malta	0.426 ^a	-0.166 ^b	-0.153 ^b	-0.241 ^b	
Cypr	-0.510 ^a	-0.034 ^b	-0.296 ^b	0.702 ^b	-0.469 ^b

<i>Octopodidae</i> (N=19 ^a , 28 ^b , 20 ^c)						
Fran	Spain	Ital	Gree			
Spain	-0.409 ^a					
Ov Ital	-0.053 ^a	0.675 ^b				
El Ital	-0.327 ^a	0.620 ^b	0.536 ^b			
Gree	-0.230 ^a	0.724 ^b	0.716 ^b	0.588 ^b		
Malta	-0.118 ^a	0.234 ^b	0.114 ^b	0.214 ^b	-0.789 ^b	
Cypr	-0.092 ^a	0.583 ^b	0.501 ^b	0.603 ^b	0.787 ^b	-0.573 ^b

Table 7

Correlations between landings by different countries: FAO area 27.

Spearman's rank correlation coefficients. Significant correlations are indicated by bold type. The analysis is based on FAO data for landings from area 27, 1965-1992. Scot = Scotland, N. Ire = Northern Ireland, Ire = Ireland, I. Man = Isle of Man, Engl = England & Wales, Belg = Belgium, Ch Isl = Channel Islands, Fran = France, Port = Portugal, Germ = Germany, Norw = Norway, Icel = Iceland, Denm = Denmark, Holl = Holland. The first tabulation, for "Loligo spp." also includes "Squid (Loliginidae or Ommastrephidae)" for those countries which either normally land only *Loligo* or record other Ommastrephids separately (see text).

<i>"Loligo spp."</i> (or "Squid (Loliginidae or Ommastrephidae)") (N=26 ^a , 17 ^b , 12 ^c , 28 ^d , 20 ^e , 27 ^f , 15 ^g , 9 ^h , 18 ⁱ , 16 ^j , 25 ^k , 10 ^l , 12 ^m , 1P, 9 ⁿ , 29 ^o , 19 ^p)											
Scot	N. Ire	Ire	I. Man	Engl	Belg	Ch Isl	Fran	Spain			
0.266 ^a											
0.440^a	0.016 ^b										
0.438 ^a	0.857^b	0.219 ^c									
0.577^a	0.767^b	0.268^b	0.565^b								
0.331^a	0.237^b	0.453^b	-0.112^c	0.412^b							
0.212 ^a	-0.147 ^c	0.025^b	-0.275^b	0.394^b	0.474^b						
0.564^a	-0.066 ^c	0.197^b	-0.287^b	0.138^b	0.322^b	0.271^b					
-0.191 ^a	-0.475 ^b	-0.379 ^b	-0.397 ^b	-0.540^b	-0.232^b	-0.430^b	0.019 ^b				
0.099 ^a	0.114 ^b	0.109 ^b	0.298 ^b	0.208 ^b	0.066 ^b	0.537^b	-0.132 ^b	-0.255 ^b			

<i>Illex</i> (N=6 ^a , 8 ^b)				<i>Todarodes</i> (N=7 ^a , 4 ^b)			
Spain	Port	Spain	Norw	Spain	Norw	Ital	Port
0.406 ^a	-0.796 ^b	0.000 ^b					
Germ							
Holl							
Engl							
Ch Isl							
Fran							
Spain							
Port							

<i>Cuttlefish</i> (N=10 ^a , 9 ^b , 17 ^c , 25 ^d , 12 ^e , 11 ^f , 8 ^g , 18 ^h , 19 ⁱ , 28 ^j)							
Denm	Holl	Engl	Ch Isl	Fran	Spain		
0.138 ^a							
0.829^a							
0.854^a			0.748 ^b				
0.303 ^a	0.661 ^b	0.842^b	0.891^b				
-0.722 ^a	-0.133 ^b	0.728^b	0.466^b	0.221^b			
0.287 ^a	-0.336 ^b	0.127 ^b	0.241^b	0.558^b	-0.303 ^b		

<i>Octopodidae</i> (N=5 ^a , 6 ^b , 10 ^c , 15 ^d , 27 ^e)				
Irل	Engل	Fran	Spain	
-0.216 ^a				
-0.746^a	0.549 ^b			
-0.795^a	0.085 ^b	0.335 ^b		
-0.500 ^a	0.596^b	0.601^b	-0.390 ^b	

Table 10

Correlations between landings of long-finned squid from the NE Atlantic by different European countries.

<i>Top right: 1980-1994 (N=15^a, 14^b)</i>				
Scotland	Fran	Spain	Portugal	Azores
	0.379 ^a	-0.400 ^b	0.284 ^b	0.508 ^b
0.439^a		-0.579^b	-0.002 ^b	0.279 ^b
-0.216 ^a	0.090 ^b		0.103 ^b	-0.248 ^b
0.060 ^a	0.022 ^b	0.186^b		-0.231 ^b
0.270 ^a	-0.208 ^b	-0.750 ^b	-0.109 ^b	

Bottom left: 1948-1994 (N=30^a, 34^b, 45^c, 29^d, 28^e, 33^f)

Figure 1




Figure 2





Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

Figure legends

Figure 1. Landings of *Loligo* in Scotland, hours fishing and overall CPUE for *Loligo*, 1980-94.

Figure 2. Landings of *Loligo* in Scotland classified by gear-type, 1980-94.

Figure 3. Landings of squid (presumed to be *Loligo*) from ICES fishery subdivision Vla (west coast of Scotland) by different European countries, 1973-94.

Figure 4. Landings of different cephalopods from FAO area 27 (NE Atlantic) by Spain, 1965-92.

Figure 5. Landings of different cephalopods from FAO area 37 (Mediterranean) by Italy, 1965-92.

Figure 6. Landings of squid from different ICES fishery subdivisions by Spain, 1973-94. Note that, for several of these years, no data were reported to ICES.

Figure 7. Landings of long-finned squid (*Loligo*) from the NE Atlantic by Scotland, France, Spain, Portugal and the Azores, 1948-94. Only the first and last of these datasets extend back to 1948.