

SCIENTIFIC REPORTS

OPEN

Ocean acidification increases copper toxicity differentially in two key marine invertebrates with distinct acid-base responses

Received: 15 September 2015

Accepted: 25 January 2016

Published: 22 February 2016

Ceri Lewis, Robert P. Ellis, Emily Vernon, Katie Elliot, Sam Newbatt & Rod W. Wilson

Ocean acidification (OA) is expected to indirectly impact biota living in contaminated coastal environments by altering the bioavailability and potentially toxicity of many pH-sensitive metals. Here, we show that OA (pH 7.71; $p\text{CO}_2$ 1480 μatm) significantly increases the toxicity responses to a global coastal contaminant (copper $\sim 0.1 \mu\text{M}$) in two keystone benthic species; mussels (*Mytilus edulis*) and purple sea urchins (*Paracentrotus lividus*). Mussels showed an extracellular acidosis in response to OA and copper individually which was enhanced during combined exposure. In contrast, urchins maintained extracellular fluid pH under OA by accumulating bicarbonate but exhibited a slight alkalosis in response to copper either alone or with OA. Importantly, copper-induced damage to DNA and lipids was significantly greater under OA compared to control conditions (pH 8.14; $p\text{CO}_2$ 470 μatm) for both species. However, this increase in DNA-damage was four times lower in urchins than mussels, suggesting that internal acid-base regulation in urchins may substantially moderate the magnitude of this OA-induced copper toxicity effect. Thus, changes in metal toxicity under OA may not purely be driven by metal speciation in seawater and may be far more diverse than either single-stressor or single-species studies indicate. This has important implications for future environmental management strategies.

Ocean acidification (OA), the drop in ocean pH associated with increasing levels of carbon dioxide in the atmosphere and hence the oceans, is now widely considered to be one of the most pervasive human impacts on global marine biodiversity^{1,2}. Recent projections published as part of the Representative Concentration Pathways (RCP) database suggest that atmospheric $p\text{CO}_2$ levels may exceed 1000 μatm early in the next century (RCP 8.5)³ causing the average pH of the world's surface waters to drop by as much as 0.43 units to around 7.73^{3,4}. There is now a wealth of evidence that this change in ocean carbonate chemistry has the potential to impact upon the health and physiology of a wide range of marine invertebrate species^{5–9} with 63% of echinoderms and 51.6% of molluscs tested so far showing negative impacts of near-future OA, making them amongst the most sensitive phyla¹⁰.

Whilst the physiological impacts of OA for many marine biota have been widely studied, the potential for OA to interact with additional environmental stressors remains poorly understood. Changes in ocean carbonate chemistry are happening against a background of additional anthropogenically driven changes such as warming, sea level rise, increasing hypoxic and anoxic zones and chronic coastal pollution. The urgent need for 'multi-stressor' studies is now widely acknowledged by the OA community¹¹ but to date such studies have tended to focus on combining OA with either temperature, salinity or hypoxia^{12–14}. Of particular concern for environmental monitoring purposes is the potential for the predicted changes in ocean pH to alter the behaviour and bioavailability of historical and chronic coastal contaminants, such as metals^{15–17}.

Metals are one of the most common types of coastal contaminant globally and are found in high concentrations in the waters and sediments of many coastal and estuarine systems^{18,19}. For example, concentrations of total dissolved copper in U.K. coastal and estuarine waters can range from chronic low levels of 0.004 μM ²⁰ to as high as 1.61 μM in highly contaminated habitats²¹. OA is expected to alter the bioavailability of waterborne metals¹⁵ as a result of changes in their speciation in seawater, driven by the declining pH. The toxic free-ion concentration of copper (Cu^{2+}) is predicted to increase by 115% in coastal waters in the next 100 years due to reduced pH^{22,23},

College of Life and Environmental Sciences, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter, EX4 4QD, UK. Correspondence and requests for materials should be addressed to C.L. (email: c.n.lewis@exeter.ac.uk)

while the free-ion concentration of other metals including cadmium (Cd) may decrease or be unaffected^{22,24–26}. Increased metal accumulation under near-future OA conditions has been demonstrated for two bivalves species so far²⁷ and for marine organisms exposed to the same nominal concentrations of any metal, greater metal toxicity effects would be predicted under near-future OA where reduced seawater pH increases free ion availability.

Whilst many transition metals including copper are essential for biological functions, elevated levels can overwhelm an organism's antioxidant defences and induce oxidative damage of cellular components such as lipids, proteins and DNA via the production of reactive oxygen species²⁸. In addition copper is known to exert a number of physiological impacts that are similar to those observed in response to exposures to OA conditions, such as growth reduction, disturbance of acid-base and osmotic regulation, and enzyme inhibition^{29,30}. Marine animals acutely subjected to seawater with elevated $p\text{CO}_2$ experience a corresponding extracellular acidosis^{9,31}. Many fish and crustaceans are able to regulate these acid-base perturbations by the elevation of extracellular bicarbonate ions (HCO_3^-) whilst other invertebrates, such as mussels and some urchin species, are generally considered to be less able to acid-base regulate^{32,33}. Furthermore, copper has been demonstrated to inhibit carbonic anhydrase, a vital enzyme for acid-base regulation and identified as an enzyme of interest for OA physiological studies³⁴. These overlapping physiological and toxicity effects of OA and copper pollution suggest the potential for additional interactions in the responses of an organism to both stressors when exposed in combination on top of those driven purely by the metal speciation changes. Understanding these potential interactions is vital for understanding the impact of OA on coastal metal contamination and its impacts on both commercially and ecologically important biota.

We examined a suite of physiological and toxicity responses to combined OA and copper (nominal $0.1 \mu\text{M}$) exposures in two ecologically important marine invertebrates with known sensitivities to OA as a single stressor. The common mussel *Mytilus edulis* is both an economically important shellfish species and provides a key ecosystem service, forming important substratum for many epibionts and influencing ecosystem functioning via their role in nutrient and mineral cycling³⁵. The sea urchin *Paracentrotus lividus* is an ecologically important herbivore in coastal benthic habitats with additional economic importance as a food source. We used a simple factorial design to test the hypothesis that OA increases the toxicity response to copper of these two key benthic invertebrates.

Results and Discussion

Acid-base responses differed between mussels and urchins. The ability to compensate for OA-induced changes in extracellular pH is believed to be a key determinant of an organisms' ability to tolerate near-future OA^{10,35}. Interestingly, we found very different acid-base responses between mussels and urchins to both the OA conditions and the copper exposures. In mussels, haemolymph $p\text{CO}_2$ levels increased slightly but non-significantly with exposure to OA alone, increased further with exposure to nominal $0.1 \mu\text{M}$ copper alone and showed the greatest increase in $p\text{CO}_2$ in the combined OA-copper exposures (Fig. 1a; two-way GLM model for OA $F_{1,39} = 6.60$, $P = 0.014$; for copper $F_{1,39} = 26.73$, $P < 0.001$; interaction term $F_{1,39} = 1.32$, $P = 0.258$). Haemolymph bicarbonate levels (HCO_3^-) in mussels showed a similar pattern of change in response to this elevated $p\text{CO}_2$ (Fig. 1b), with a small but significant increase of $\sim 0.2 \text{ mM}$ under OA conditions and about double this increase when exposed to copper alone (Fig. 1b; two-way GLM model for OA $F_{1,39} = 9.75$, $P = 0.004$; for copper $F_{1,39} = 26.57$, $P < 0.001$; interaction term $F_{1,39} = 1.95$, $P = 0.171$). Combined exposure to OA and nominal $0.1 \mu\text{M}$ copper caused a further increase in haemolymph bicarbonate level of $\sim 1 \text{ mM}$. These $p\text{CO}_2$ changes combined with limited bicarbonate responses drove a slight but non-significant acidosis of the haemolymph in mussels exposed to OA conditions, whilst exposure to nominal $0.1 \mu\text{M}$ copper induced a stronger and significant acidosis of haemolymph from 7.56 to 7.43 (Fig. 1c). The combined OA and copper exposure caused an even greater acidosis reducing haemolymph pH to 7.33, however statistical analysis revealed this was not an interactive effect (Fig. 1c; two-way GLM model for OA $F_{1,39} = 3.74$, $P = 0.061$; for copper $F_{1,39} = 15.58$, $P < 0.001$; interaction term $F_{1,39} = 0.34$, $P = 0.562$).

The increased haemolymph $p\text{CO}_2$ in the combined OA and copper treatment compared to the OA (no copper) treatment might be explained by changes in mussel ventilation rate as a behavioural mechanism to reduce acidosis, since they appear unable to accumulate substantial bicarbonate as a buffering mechanism. Whilst ventilation rate was not measured here, increased mussel gaping (i.e. greater amount of time ventilating the gills) under reduced seawater pH conditions has been reported elsewhere³⁶. Hyperventilation in response to copper may be less likely as it could be counter-productive by increasing exposure of gills to the waterborne metal. Mussels have been shown to reduce the amount of time spent with their shells open in response to exposures to metals such as copper³⁷. These results support the general consensus that mussels are not good acid-base regulators with a limited ability to buffer their haemolymph using bicarbonate³³, instead responding to periods of hypercapnia with metabolic suppression or changes in ventilation rate.

Conversely, we found that urchins were able to employ physiological mechanisms to regulate coelomic fluid pH against the CO_2 -induced drop in external seawater pH. In sea urchins gas exchange, i.e. uptake of O_2 and elimination of CO_2 , relies solely on a favourable diffusion gradient due to the lack of any active ventilatory mechanism, so is generally considered to be diffusion-limited³⁸. Despite this inability to regulate internal $p\text{CO}_2$ levels, urchins appear to have varying abilities to acid-base regulate depending on species^{32,39,40}. In the present experiments with *P. lividus*, we observed a rise in coelomic fluid $p\text{CO}_2$ in both the OA (alone) and OA with copper treatments, but a slight yet significant reduction in $p\text{CO}_2$ caused by copper alone (Fig. 1d; two-way GLM model for OA $F_{1,32} = 35.14$, $P < 0.001$; for copper $F_{1,32} = 6.33$, $P = 0.018$; interaction term $F_{1,32} = 0.78$, $P = 0.385$). Coelomic fluid bicarbonate levels were significantly elevated by $> 2 \text{ mM}$ in response to this elevated $p\text{CO}_2$ in both OA treatments, whilst exposure to copper alone did not affect bicarbonate levels (Fig. 1e; two-way GLM model for OA $F_{1,32} = 29.25$, $P < 0.001$; for copper $F_{1,32} = 0.07$, $P = 0.793$; interaction term $F_{1,32} = 0.06$, $P = 0.809$).




Figure 1. Acid-base parameters in the haemolymph of *Mytilus edulis* (a,c,e) and coelomic fluid of *Paracentrotus lividus* (b,d,f) following 14 day exposures to elevated pCO₂ with and without the presence of nominal 0.1 μM copper; (a,b) haemolymph/coelomic fluid pH, (c,d) haemolymph/coelomic fluid bicarbonate concentrations, and (e,f) haemolymph/coelomic fluid pCO₂. [N.B. *represent significant differences].

All *P. lividus* in the present study maintained a coelomic fluid pH between 7.60 and 7.76, independent of exposure to OA conditions or elevated copper (Fig. 1f; two-way GLM model for OA $F_{1,32} = 0.56$, $P = 0.461$; for copper $F_{1,32} = 9.31$, $P = 0.005$; interaction term $F_{1,32} = 1.83$, $P = 0.186$), most likely attributed to the 48% and 55% increase, respectively, in the coelomic fluid bicarbonate levels in the two OA treatments buffering against the effect of the increased pCO₂. Increased protein concentrations of coelomic fluid have also been suggested to play




Figure 2. Davenport diagram illustrating the relationship between pH, bicarbonate and pCO_2 in the haemolymph and coelomic fluid of (a) *Mytilus edulis* and (b) *Paracentrotus lividus* respectively. Lines represent isopleths of equal pCO_2 (mmHg). Position calculated from means \pm SEM for haemolymph/coelomic fluid pH and $[HCO_3^-]$ according to pK_1 values calculated from⁵⁷.

a role in this buffering capacity⁴¹ however this was not evident in our data (see below). This agrees with previous studies on *P. lividus* which have shown full compensation of coelomic fluid pH at seawater pCO_2 of 1293 μatm (pH of 7.7) and a partial ability to buffer against an external pCO_2 rise over a wider seawater pCO_2 range (pCO_2 583–2364 μatm ; pH 8.0–7.4)^{32,42}. In contrast, exposing urchins to nominal 0.1 μM copper combined with OA caused a slight but significant alkalosis of their coelomic fluid. This corresponded to slightly lower coelomic fluid pCO_2 in the two copper treatments (alone and combined with OA) compared to their corresponding treatments without added copper (control and OA).

These differences in acid-base responses between mussels and urchins to OA and copper exposures can be illustrated using Davenport diagrams (Fig. 2a,b). In mussels the decrease in haemolymph pH due to OA exposure is increased by the additional presence of copper with very little compensation from the elevation of bicarbonate ions. The pattern observed reflects a primarily respiratory acidosis of varying severity caused by the treatments. In urchins under OA conditions (with or without copper) the coelomic fluid pH is buffered by additional bicarbonate (i.e. a fully compensated respiratory acidosis) whilst copper caused a slight mixed respiratory/metabolic alkalosis. Our contrasting findings for mussels and urchins, and the contrasting findings of other studies across a number of aquatic species to similar copper concentrations^{43–45}, suggest highly species-specific and concentration-dependent acid-base responses to copper.

Oxidative stress responses also differed between mussels and urchins. Superoxide dismutase (SOD) is an important cytosolic anti-oxidant enzyme. SOD activity has been shown to be induced in a number of marine invertebrate species in response to a range of environmental metals including copper^{46,47}, with inhibition being reported for exposures using higher concentrations^{48,49}. After 14 days exposure to our four treatments, SOD activity in mussel haemolymph showed no significant changes in response to any of these treatments, (Fig. 3a, two-way GLM model for copper: $F_{1,39} = 2.57$, $P = 0.118$; for OA: $F_{1,39} = 3.15$, $P = 0.084$; interaction term $F_{1,39} = 1.73$, $P = 0.198$). There was an average ~ 2 fold increase in SOD activity in response to copper alone compared to the treatments with no copper, however this response varied between individuals such that there was no significant difference overall. In urchins, however, SOD activity was significantly increased following exposure to nominal 0.1 μM copper under both ambient and OA conditions (Fig. 3b; two-way GLM model for copper $F_{1,32} = 7.07$, $P = 0.013$). OA conditions had no effect on extracellular fluid SOD activity in urchins (Fig. 3a; two-way GLM model for OA $F_{1,32} = 0.05$, $P = 0.484$; interaction term $F_{1,32} = 0.08$, $P = 0.777$).

Lipid peroxidation was significantly induced by exposure to copper in mussels under both ambient and OA conditions (Fig. 3c, two-way GLM model, for copper: $F_{1,39} = 17.77$, $P < 0.001$). There was no additional




Figure 3. Oxidative stress indicators in the mussel *Mytilus edulis* (a,c,e) and the adult purple urchin *Paracentrotus lividus* (b,d,f) following 14 day exposures to elevated pCO₂ with and without the presence of nominal 0.1 μM copper; (a,b) Activity of the anti-oxidant enzyme superoxide dismutase (SOD) activity; (c,d) lipid peroxidation measured as malondialdehyde (MDA) levels; (e,f) DNA damage, measured as percentage of single strand breaks in haemocytes/coelomocytes.

increase in lipid peroxidation when OA and copper were combined (two-way GLM model, for OA: $F_{1,39} = 0.22$, $P = 0.640$; interaction term $F_{1,39} = 0.05$, $P = 0.486$). Urchins again showed a different response, with no increase in lipid peroxidation induced by exposure to nominal 0.1 μM copper alone, most likely due to the protective

effect of the increased SOD activity. A significant increase in lipid peroxidation was, however, induced by copper when exposed under OA conditions, despite the higher SOD levels (Fig. 3d; two-way GLM model for copper $F_{1,32} = 0.54$, $P = 0.467$; for OA $F_{1,32} = 11.19$, $P = 0.002$). No significant interaction term was present (interaction OA \times copper $F_{1,32} = 3.59$, $P = 0.068$). This suggests a much greater toxicity effect of copper on lipids under the combined treatments that overwhelms the anti-oxidant defence capabilities in urchins in the combined OA and copper treatment.

Protein levels, measured as part of both the SOD and TBARS assays using the Bradford assay, were found to be ~20% lower in haemolymph/ coelomic fluid in response to copper exposure in both mussels and urchins independent of OA treatment. Mussels in the ambient pH and OA treatments were found to have haemolymph protein levels of 3.17 and 3.44 mg protein ml⁻¹. Protein levels were significantly lower at 2.41 and 2.73 mg protein ml⁻¹, respectively, in the treatments with the addition of nominal 0.1 μ M copper (two-way GLM model for copper $F_{1,30} = 8.26$, $P = 0.007$). In urchins protein levels were generally slightly lower with 1.84 and 1.97 mg protein ml⁻¹ in the ambient pH and OA treatments. The additional presence of copper lowered these levels to 1.46 and 1.56 mg protein ml⁻¹, respectively, but this was not a significant effect (two-way GLM model for copper: $F_{1,30} = 3.45$, $P = 0.073$).

DNA Damage was increased under OA in both species. Elevated copper induces DNA damage in the form of single strand breaks by the production of reactive oxygen species via the Fenton reaction and by base modifications such as 8-OHdG (a major product of DNA oxidation)²⁸. In mussels copper-induced DNA damage was only observed in the combined OA and copper exposures, with no increase in DNA damage caused by exposure to nominal 0.1 μ M copper alone (Fig. 3e). Combined exposure of adult mussels to both OA and copper in combination, however, resulted in a 1.9-fold increase in DNA damage from 14% DNA damage in the ambient (pH 8.1, no copper) treatment to 27% in the combined exposures. Statistical analysis reveals a significant interaction term between OA conditions and the presence of copper on DNA fragmentation (Two-way GLM model, for OA $F_{1,39} = 12.54$, $P < 0.001$; for copper $F_{1,39} = 8.25$, $P = 0.007$; interaction term $F_{1,39} = 23.17$, $P < 0.001$).

This indicates that mussels are able to cope with this low concentration of copper under the ambient pH/pCO₂ conditions, either via their antioxidant defences preventing damage occurring in the first place or via efficient DNA repair activities. This does not hold true under OA conditions. The strong increase in DNA damage when exposed to nominal 0.1 μ M copper under OA conditions has two likely explanations; a) the lack of anti-oxidant response (SOD) to the copper under OA conditions, and/or b) the increased availability and presumably uptake of the toxic free Cu²⁺ ion under the reduced seawater pH of OA conditions.

In urchins both OA and copper exposure were found to significantly affect DNA damage in coelomocytes. Exposure to copper induced a significant increase in DNA damage in both the pH 8.1 and pH 7.7 treatments, but as with mussels, DNA damage was significantly higher again in the OA treatment (Fig. 3f two-way GLM model for copper $F_{1,32} = 138.12$, $P < 0.001$; for OA $F_{1,32} = 23.81$, $P = 0.038$). No significant interaction term between copper and OA was found ($F_{1,32} = 4.71$, $P = 0.104$). This suggests that whilst increased SOD levels appear to provide some protection against lipid peroxidation it has not prevented copper-induced DNA damage from occurring in the urchin coelomocytes. This toxicity effect of copper on DNA is then significantly greater under OA conditions.

The urchins' acid-base regulation strategy of increasing extracellular bicarbonate levels may actually act 'protectively' against copper toxicity and explain the overall reduced toxicity driven by the difference in the DNA damage response to combined copper and OA that we observed between mussels and urchins. It has been widely shown that copper is less toxic to freshwater species in hard water (i.e. higher alkalinity) than in soft water⁵⁰. A protective effect of hypercapnia on copper toxicity has also been suggested by the work of Larsen *et al.*⁵¹ in the cod *Gadus morhua*. Free copper (II) ions (Cu²⁺) will form a range of complexes with bicarbonate ions such that the amount of free Cu²⁺ is only a small fraction of the total copper present in a high bicarbonate solution, with the relative concentrations of copper species varying with pH⁵². Elevated bicarbonate levels should therefore reduce the proportion of the toxic free copper ions and so reduce the amount that is bioavailable to cause damage. Whilst the damage to urchin's lipids did not follow the same pattern as for DNA, this might be explained by their different positions within the cell, with the TBARS assay measuring mostly damage to lipids on the outside of cell membranes whilst DNA is within the cell nucleus, perhaps making lipids more susceptible to damage by the remaining free Cu²⁺ present. More detailed biochemical studies of the precise mechanisms of this damage would be required to fully explain these differences. A hypothesis resulting from this work requiring further testing could be that species which regulate their extracellular pH in response to OA by the elevation of internal bicarbonate levels will show reduced overall copper toxicity responses compared to those which are unable to acid-base regulate using bicarbonate.

This increased copper toxicity as a result of OA conditions in both test species is consistent with two other studies looking at metal-induced DNA damage under near-future OA. Roberts *et al.*⁵³ found that DNA damage in the sediment dwelling amphipod *Corophium volutator* exposed to naturally contaminated sediments, which contained a range of metals and polycyclic aromatic hydrocarbons (PAHs), was higher under OA conditions than extant pCO₂ conditions⁵³. More recently an increase in copper-induced DNA damage in the sperm of the polychaete *Arenicola marina* was reported when worms were exposed to copper under elevated pCO₂⁵⁴. Synergistic toxicities have been reported between OA and copper in adults of the copepod *Amphiascoides atopus*²², where the LC₅₀ for copper was reduced from 0.65 mg l⁻¹ at pH 8.1 to 0.32 mg l⁻¹ at a CO₂-driven reduction of pH of 6.5. Increased copper toxicity under OA has also been reported for larvae of the polychaete worms *Pomatoceros lamarckii*⁵⁵ and *Arenicola marina*⁵⁴. Larvae are often considered to be the most sensitive life history stages to environmental stressors, particularly in free spawning marine invertebrates with bi-phasic life histories. We have now demonstrated that this increased toxicity under experimental OA can also be found in adult marine invertebrates at environmentally relevant concentrations.

Treatment	Temperature (°C)	pH _{NBS}	Salinity	Copper (μM)	TA (μmol/kg) [†]	TCO ₂ (μmol/kg)	pCO ₂ (μatm) [†]	HCO ₃ ⁻ (μmol/kg) [†]	CO ₃ ²⁻ (μmol/kg) [†]	ΩCa [†]	ΩAr [†]
8.1	15.0 (±0.1)	8.14 (±0.01)	35.68 (±0.04)	0.016 (±0.006)	2345.8 (±68.6)	2132.4 (±60.4)	436.4 (±10.9)	1960.2 (±53.3)	156.0 (±7.7)	3.7 (±0.2)	2.4 (±0.1)
7.7	15.0 (±0.1)	7.68 (±0.01)	35.96 (±0.06)	0.011 (±0.002)	2303.1 (±63.6)	2264.3 (±63.3)	1373.0 (±69.2)	2152.5 (±60.2)	60.6 (±2.7)	1.4 (±0.1)	0.9 (±0.4)
8.1 + Cu	14.6 (±0.1)	8.09 (±0.01)	35.60 (±0.04)	0.047 (±0.007)	2635.3 (±55.0)	2423.1 (±38.4)	536.2 (±7.6)	2238.5 (±39.7)	164.1 (±7.1)	3.9 (±0.2)	2.5 (±0.1)
7.7 + Cu	14.5 (±0.1)	7.68 (±0.01)	35.74 (±0.07)	0.052 (±0.007)	2662.8 (±56.4)	2568.7 (±41.7)	1481.4 (±16.1)	2486.6 (±49.7)	73.0 (±3.1)	1.7 (±0.1)	1.1 (±0.1)

Table 1. Seawater carbonate chemistry and copper levels for the four experimental treatment solutions for the *Mytilus edulis* exposures.

Conclusions

Our data has clearly shown that near-future OA significantly increases the sub-lethal toxicity responses of two key coastal marine invertebrates to coastally relevant concentrations of the common metal pollutant copper. We found that for both mussels (*Mytilus edulis*) and urchins (*Paracentrotus lividus*) copper-induced damage to DNA was significantly greater when animals were exposed to nominal 0.1 μM copper under OA (high pCO₂/ low pH) conditions compared with animals exposed under extant pCO₂ levels. This relative increase in copper-induced DNA damage under OA was four-times greater in mussels than in urchins (despite the measured seawater copper concentrations actually being lower in the mussel exposures than for the urchins). In mussels OA seemed to suppress the response of the anti-oxidant enzyme SOD to copper exposures, whilst in urchins lipid peroxidation was also increased when exposed to copper under OA. So whilst copper-induced toxicity was clearly increased in both species under near-future OA, there were slight differences in the biochemical details of how the two species responded.

Demonstrating an increase in the toxicity of copper in two ecologically and economically important coastal invertebrates under near future OA is a significant cause for concern given the widespread nature of coastal metal contamination. This is particularly the case given the relatively low concentrations of this coastal contaminant used in the present study, which are relevant to measurements of copper contamination for UK coastal waters generally and which are often exceeded in contaminated locations globally^{20,21}. Furthermore, the changes in seawater copper speciation expected with a reduction in seawater pH will be compounded by an increase in the total copper released from sediments under OA conditions¹⁵. Subsequently the OA-induced increase in bioavailable copper is expected to be two-fold. We cannot fully determine whether these interactions between OA and copper are additive (the sum of the individual stressor effects) or synergistic interactions (greater than the sum of the individual stressor effects) due to the limited number of copper concentrations or seawater pH levels used here. However, the strong interaction terms in the analysis for some of the end points measured here are suggestive that synergies are likely to be present. Recent meta-analysis studies have shown that synergistic interactions between multiple abiotic stressors in the marine environment are more common than additive interactions^{11,14}. This highlights a real need for the potential for interactions between climate driven stressors and coastal pollution to be studied in much greater detail rather than relying on predictive modelling approaches for environmental risk assessments.

Our results have clear implications for ecotoxicological assessments, highlighting a need to better understand how OA will alter the behaviour and toxicity of not just copper or metals generally but a wider range of environmental contaminants. Such knowledge is vital for those charged with protecting our marine habitats in order to provide justifiable predictions of OA impacts in coastal regions. In demonstrating a contrasting toxicity response in two ecologically important marine invertebrate species, related to acid-base regulation, we show that organismal responses may be far more diverse than either single stressor or single species studies have previously indicated. Furthermore, our results imply that not only may we be under-estimating OA impacts for coastal invertebrates exposed to chronic metal pollution, but also that OA may impact a much wider range of species, not traditionally considered as OA sensitive, due to this indirect effect on metal toxicity. This suggests that further investigations on OA-pollutant interactions in a wider range of organisms will be important to better understand the near future impact of toxic coastal contaminants for marine organisms globally.

Methods

Adult *Paracentrotus lividus* specimens (4–6 cm diameter) were purchased from Dunmannus Sea Farm Ltd. in Cork, Ireland. Adult *Mytilus edulis* specimens were collected by hand from the intertidal range of the River Exe estuary, Exmouth, Devon, UK. Individuals were left for 7 days in 30 litre holding tanks at 15 °C in ambient artificial seawater (pH_{NBS} 8.1, 470 μatm pCO₂, salinity = 35) to acclimatise prior to the exposures. Ten individuals per treatment were exposed to one of the following four treatments for 14 days at 15 °C; (1) ambient conditions (pH_{NBS} 8.1) with no added copper; (2) ambient conditions (pH_{NBS} 8.1) with nominal 0.1 μM copper sulphate added; (3) OA conditions (pH_{NBS} 7.7) with no added copper; (4) OA conditions (pH_{NBS} 7.7) with nominal 0.1 μM copper sulphate added.

Seawater pH_{NBS} values of 7.7 were targeted to represent near-future OA treatments as projected according to scenario RCP 8.5 and the 2013 IPCC WGI AR5^{4,56}; full seawater chemistry is provided in Tables 1 and 2. Seawater pH in the OA conditions was nominally maintained at pH_{NBS} 7.7 (to a resolution of 0.05 units) using pH computers (Aqua Medic, Bissendorf, Germany) which continually controlled the release of CO₂ gas directly into the header tanks to maintain stable conditions throughout the experimental exposures. Partial water changes (50%) were carried out every 48 hours using temperature equilibrated water of the correct pH and CO₂ level and copper concentrations were re-dosed appropriately. Seawater pH_{NBS} (Metrohm 827 pH lab), temperature, and salinity

Treatment	Temperature (°C)	pH _{NBS}	Salinity	Copper (μM)	TA (μmol/kg) [†]	TCO ₂ (μmol/kg)	pCO ₂ (μatm) [†]	HCO ₃ ⁻ (μmol/kg) [†]	CO ₃ ²⁻ (μmol/kg) [†]	ΩCa [†]	ΩAr [†]
8.1	15.6 (±0.1)	8.14 (±0.01)	34.06 (±0.02)	0.031 (±0.006)	2599.7 (±35.2)	2364.9 (±25.5)	470.7 (±15.8)	2168.9 (±19.2)	178.7 (±8.0)	4.3 (±0.2)	2.8 (±0.1)
7.7	15.6 (±0.1)	7.72 (±0.01)	34.11 (±0.02)	0.028 (±0.004)	2538.5 (±36.4)	2467.9 (±28.6)	1297.5 (±85.0)	2337.0 (±28.3)	83.3 (±12.4)	2.0 (±0.3)	1.3 (±0.2)
8.1 + Cu	15.3 (±0.1)	8.13 (±0.01)	34.25 (±0.03)	0.084 (±0.002)	2673.0 (±30.6)	2452.6 (±21.4)	528.4 (±17.9)	2262.0 (±15.8)	170.9 (±7.1)	4.1 (±0.2)	2.6 (±0.1)
7.7 + Cu	15.23 (±0.1)	7.71 (±0.01)	34.50 (±0.05)	0.088 (±0.005)	2655.1 (±22.9)	2608.8 (±21.4)	1483.1 (±31.9)	2481.3 (±20.2)	72.2 (±1.8)	1.7 (±0.1)	1.1 (±0.1)

Table 2. Seawater carbonate chemistry and copper levels for the four experimental treatment solutions for the *Paracentrotus lividus* exposure.

(Mettler Toledo SG7), were measured daily in header tanks and experimental chambers. Water samples were taken every third day for measurements of dissolved inorganic carbon (DIC) and metals. Seawater DIC analysis was carried out as described in detail in Lewis *et al.*⁵⁵ using a custom built system and using certified reference materials from Andrew Dickson's Laboratory (Scripps Institution of Oceanography batch 114). Copper concentrations in seawater were determined using ICP-MS.

Following the 14 day exposure samples of extracellular fluid (haemolymph for mussels and coelomic fluid for urchins) were collected from each individual. Extracellular fluid for TCO₂ analysis was stored in 100 μl hematocrit (micro capillary) tubes sealed with paraffin oil and the CritosealTM capillary tube sealant (Fisher) then analysed using a Corning 965 CO₂ analyser (Corning Ltd., UK). Extracellular fluid for oxidative stress assays was snap frozen in liquid nitrogen and stored at −80 °C until analysis. Immediately following extraction extracellular fluid pH was measured at 15 °C using a pH meter (Hannah Instruments HI 8314) and microprobe (Cole Parmer, Accumet) calibrated prior to usage using pH_{NBS} 7.00 and 10.00 specific buffers (calibrated to 7.04 and 10.11 at 15 °C). Acid-base parameters were then calculated using a modified version of the Henderson-Hasselbalch equation using the first dissociation constant (pK) for carbonic acid and solubility constant (αCO₂) for carbon dioxide derived from Truchot⁵⁷.

Superoxide dismutase (SOD) activity was measured using the nitroblue tetrazolium (NBT) microplate assay⁵⁸. Lipid peroxidation was determined using the thiobarbituric acid reactive substances (TBARS) assay⁵⁹ which quantifies malondialdehyde (MDA), a secondary product of lipid peroxidation, via its reaction with thiobarbituric acid. DNA damage in haemocytes/ coelomocytes was measured as single strand DNA breaks using the comet assay. Two hundred microlitres of haemolymph/ coelomic fluid from each individual was used immediately after sampling for the comet assay according to the methods described by Lewis & Galloway⁶⁰, using alkaline conditions at 5 °C.

Data were analysed using 2-way analysis of variance (ANOVA) general linear models with the fixed factors; 'pH' and 'copper concentration'.

References

- Halpern, B. S. *et al.* A global map of human impact on marine ecosystems. *Science* **319**, 948–952, doi: 10.1126/science.1149345 (2008).
- Gattuso, J. P. *et al.* Contrasting futures for ocean and society from different anthropogenic CO₂ emissions scenarios. *Science* **349**, 45–55 (2015).
- Bopp, L. *et al.* Multiple stressors of ocean ecosystems in the 21st Century: projections with CMIP5 models. *Biogeosci.* **10**, 6225–6245 (2013).
- IPCC. Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the IPCC 5th Assessment Report—Changes to the Underlying Scientific/Technical Assessment. (2013).
- Fabry, V. J., Seibel, B. A., Feely, R. A. & Orr, J. C. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci.* **65**, 414–432, doi: 10.1093/icesjms/fsn048 (2008).
- Kroeker, K. J. *et al.* Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob. Chan. Biol.* **19**, 1884–1896, doi: 10.1111/gcb.12179 (2013).
- Orr, J. C. *et al.* Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* **437**, 681–686, doi: 10.1038/nature04095 (2005).
- Fabricius, K. E., De'ath, G., Noonan, S. & Uthicke, S. Ecological effects of ocean acidification and habitat complexity on reef-associated macroinvertebrate communities. *Proc. R. Soc. B-Biol. Sci.* **281**, doi: 10.1098/rspb.2013.2479 (2014).
- Pörtner, H.-O. Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Mar. Ecol. Prog. Ser.* **373**, 203–217, doi: 10.3354/meps07768 (2008).
- Wittmann, A. C. & Pörtner, H.-O. Sensitivities of extant animal taxa to ocean acidification. *Nat. Clim. Chan.* **3**, 995–1001, doi: 10.1038/nclimate1982 (2013).
- Crain, C. M., Kroeker, K. & Halpern, B. S. Interactive and cumulative effects of multiple human stressors in marine systems. *Ecol. Lett.* **11**, 1304–1315, doi: 10.1111/j.1461-0248.2008.01253.x (2008).
- Byrne, M. & Przeslawski, R. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integr. Comp. Biol.* **53**, 582–596, doi: 10.1093/icb/ict049 (2013).
- Schram, J. B., Schoenrock, K. M., McClintock, J. B., Amsler, C. D. & Angus, R. A. Multiple stressor effects of near-future elevated seawater temperature and decreased pH on righting and escape behaviors of two common Antarctic gastropods. *Journal of Exp. Mar. Biol. Ecol.* **457**, 90–96, doi: 10.1016/j.jembe.2014.04.005 (2014).
- Przeslawski, R., Byrne, M. & Mellin, C. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. *Glob. Chan. Biol.* **21**, 2122–2140, doi: 10.1111/gcb.12833 (2015).
- Millero, F. J., Woosley, R., Dittolio, B. & Waters, J. Effect of ocean acidification on the speciation of metals in seawater. *Oceanogr.* **22**, 72–85 (2009).
- Hoffmann, L. J., Breitbarth, E., Boyd, P. W. & Hunter, K. A. Influence of ocean warming and acidification on trace metal biogeochemistry. *Mar. Ecol. Prog. Ser.* **470**, 191–205, doi: 10.3354/meps10082 (2012).
- Ivanina, A. V. *et al.* Short-term acute hypercapnia affects cellular responses to trace metals in the hard clams *Mercenaria mercenaria*. *Aquat. Tox.* **140**, 123–133, doi: 10.1016/j.aquatox.2013.05.019 (2013).

18. Bryan, G. W. & Langston, W. J. Bioavailability, accumulation and effects of heavy-metals in sediments with special reference to United-Kingdom estuaries-A review. *Environ. Poll.* **76**, 89–131, doi: 10.1016/0269-7491(92)90099-v (1992).
19. Schiff, K., Brown, J., Diehl, D. & Greenstein, D. Extent and magnitude of copper contamination in marinas of the San Diego region, California, USA. *Mar. Poll. Bull.* **54**, 322–328, doi: 10.1016/j.marpolbul.2006.10.013 (2007).
20. Jones, B. & Bolam, T. Copper speciation survey from UK marinas, harbours and estuaries. *Mar. Poll. Bull.* **54**, 1127–1138, doi: 10.1016/j.marpolbul.2007.04.021 (2007).
21. Bryan, G. W. & Gibbs, P. E. Heavy metals in the Fal estuary, Cornwall: A study of long term contamination by mining waste and its effects on estuarine organisms. *Occ. Pub. Mar. Biol. Assoc.* **2**, 112 (1983).
22. Pascal, P.-Y., Fleeger, J. W., Galvez, F. & Carman, K. R. The toxicological interaction between ocean acidity and metals in coastal meiobenthic copepods. *Mar. Poll. Bull.* **60**, 2201–2208, doi: 10.1016/j.marpolbul.2010.08.018 (2010).
23. Richards, R., Chaloupka, M., Sano, M. & Tomlinson, R. Modelling the effects of 'coastal' acidification on copper speciation. *Ecol. Mod.* **222**, 3559–3567, doi: 10.1016/j.ecolmodel.2011.08.017 (2011).
24. Lacoue-Labarthe, T. *et al.* Temperature and pCO₂ effect on the bioaccumulation of radionuclides and trace elements in the eggs of the common cuttlefish, *Sepia officinalis*. *J. Exp. Mar. Biol. Ecol.* **413**, 45–49, doi: 10.1016/j.jembe.2011.11.025 (2012).
25. Lacoue-Labarthe, T. *et al.* Effects of increased pCO₂ and temperature on trace element (Ag, Cd and Zn) bioaccumulation in the eggs of the common cuttlefish, *Sepia officinalis*. *Biogeosci.* **6**, 2561–2573 (2009).
26. Lacoue-Labarthe, T. *et al.* Effects of ocean acidification on trace element accumulation in the early-life stages of squid *Loligo vulgaris*. *Aquat. Tox.* **105**, 166–176, doi: 10.1016/j.aquatox.2011.05.021 (2011).
27. Goetze, S., Matoo, O. B., Beniash, E., Saborowski, R. & Sokolova, I. M. Interactive effects of CO₂ and trace metals on the proteasome activity and cellular stress response of marine bivalves *Crassostrea virginica* and *Mercenaria mercenaria*. *Aquat. Tox.* **149**, 65–82, doi: 10.1016/j.aquatox.2014.01.027 (2014).
28. Stohs, S. J. & Bagchi, D. Oxidative mechanisms in the toxicity of metal-ions. *Free Rad. Biol. Med.* **18**, 321–336, doi: 10.1016/0891-5849(94)00159-h (1995).
29. Parveen, A. & Javed, M. Effect of Water-borne copper on the growth performance of fish *Catla catla*. *Int. J. Agricult. Biol.* **12**, 950–952 (2010).
30. Wang, T. *et al.* Copper exposure impairs intra- and extracellular acid-base regulation during hypercapnia in the fresh water rainbow trout (*Oncorhynchus mykiss*). *J. Comp. Phys. B-Biochem. Syst. Environ. Physiol.* **168**, 591–599, doi: 10.1007/s003600050181 (1998).
31. Widdicombe, S. & Spicer, J. I. Predicting the impact of ocean acidification on benthic biodiversity: What can animal physiology tell us? *J. Exp. Mar. Biol. Ecol.* **366**, 187–197, doi: 10.1016/j.jembe.2008.07.024 (2008).
32. Collard, M. *et al.* Buffer capacity of the coelomic fluid in echinoderms. *Comp. Biochem. Physiol. Part A, Molec. Integ. Physiol.* **166**, 199–206, doi: 10.1016/j.cbpa.2013.06.002 (2013).
33. Gazeau, F. *et al.* Impacts of ocean acidification on marine shelled molluscs. *Mar. Biol.* **160**, 2207–2245, doi: 10.1007/s00227-013-2219-3 (2013).
34. Hofmann, G. E., O'Donnell, M. J. & Todgham, A. E. Using functional genomics to explore the effects of ocean acidification on calcifying marine organisms. *Mar. Ecol. Prog. Ser.* **373**, 219–225 (2008).
35. Melzner, F. *et al.* Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosci.* **6**, 2313–2331 (2009).
36. Bamber, R. N. The effects of acidic seawater on three species of lamellibranch mollusc. *J. Exp. Mar. Biol. Ecol.* **143**, 181–191, doi: 10.1016/0022-0981(90)90069-o (1990).
37. Fdíl, M. A. *et al.* Valve movement response of the mussel *Mytilus galloprovincialis* to metals (Cu, Hg, Cd and Zn) and phosphate industry effluents from Moroccan Atlantic coast. *Ecotox.* **15**, 477–486, doi: 10.1007/s10646-006-0083-3 (2006).
38. Farmanfarmaian, A. In *Physiology of Echinodermata* (ed R. A. Boolootian) (Interscience Publishers, 1966).
39. Spicer, J. I., Widdicombe, S., Needham, H. R. & Berge, J. A. Impact of CO₂-acidified seawater on the extracellular acid-base balance of the northern sea urchin *Strongylocentrotus droebachiensis*. *J. Exp. Mar. Biol. Ecol.* **407**, 19–25, doi: 10.1016/j.jembe.2011.07.003 (2011).
40. Spicer, J. I. Oxygen and acid-base status of the sea urchin *Psammechinus miliaris* during environmental hypoxia. *Mar. Biol.* **124**, 71–76, doi:10.1007/bf00349148 (1995).
41. Bookbinder, L. H. & Shick, J. M. Anaerobic and aerobic energy-metabolism in ovaries of the sea-urchin *Strongylocentrotus droebachiensis*. *Mar. Biol.* **93**, 103–110, doi: 10.1007/bf00428659 (1986).
42. Catarino, A. I., Bauwens, M. & Dubois, P. Acid-base balance and metabolic response of the sea urchin *Paracentrotus lividus* to different seawater pH and temperatures. *Environ. Sci. Poll. Res.* **19**, 2344–2353, doi: 10.1007/s11356-012-0743-1 (2012).
43. Biellmyer, G. K., Brix, K. V., Capo, T. R. & Grosell, M. The effects of metals on embryo-larval and adult life stages of the sea urchin, *Diadema antillarum*. *Aquat. Tox.* **74**, 254–263, doi: 10.1016/j.aquatox.2005.05.016 (2005).
44. Boitel, F. & Truchot, J. P. Comparative-study of the effects of copper on hemolymph ion concentrations and acid-base-balance in shore crabs *Carcinus maenas* acclimated to full-strength or dilute seawater. *Comp. Biochem. Phys. C-Pharmacol. Tox. Endocrin.* **95**, 307–312, doi: 10.1016/0742-8413(90)90123-q (1990).
45. Spicer, J. I. & Weber, R. E. Respiratory impairment by water-borne copper and zinc in the edible crab *Cancer-pagurus* (l) (Crustacea, Decapoda) during hypoxic exposure. *Mar. Biol.* **112**, 429–435, doi: 10.1007/bf00356288 (1992).
46. Rhee, J.-S. *et al.* Expression of superoxide dismutase (SOD) genes from the copper-exposed polychaete, *Neanthes succinea*. *Mar. Poll. Bull.* **63**, 277–286, doi: 10.1016/j.marpolbul.2011.04.023 (2011).
47. Geracitano, L., Monserrat, J. M. & Bianchini, A. Physiological and antioxidant enzyme responses to acute and chronic exposure of *Laonereis acuta* (Polychaeta, Nereididae) to copper. *J. Exp. Mar. Biol. Ecol.* **277**, 145–156, doi: 10.1016/s0022-0981(02)00306-4 (2002).
48. Jing, G., Li, Y., Xie, L. P. & Zhang, R. Q. Metal accumulation and enzyme activities in gills and digestive gland of pearl oyster (*Pinctada fucata*) exposed to copper. *Comp. Biochem. Phys. C-Tox. Pharmacol.* **144**, 184–190, doi: 10.1016/j.cbpc.2006.08.005 (2006).
49. Maria, V. L. & Bebianno, M. J. Antioxidant and lipid peroxidation responses in *Mytilus galloprovincialis* exposed to mixtures of benzo(a)pyrene and copper. *Comp. Biochem. Phys. C-Tox. Pharmacol.* **154**, 56–63, doi: 10.1016/j.cbpc.2011.02.004 (2011).
50. Flemming, C. A. & Trevors, J. T. Copper toxicity and chemistry in the environment-a review. *Wat. Air Soil Poll.* **44**, 143–158, doi: 10.1007/bf00228784 (1989).
51. Larsen, B. K., Portner, H. O. & Jensen, F. B. Extra- and intracellular acid-base balance and ionic regulation in cod (*Gadus morhua*) during combined and isolated exposures to hypercapnia and copper. *Mar. Biol.* **128**, 337–346, doi: 10.1007/s002270050099 (1997).
52. Stiff, M. J. Copper/bicarbonate equilibria in solutions of bicarbonate ion at concentrations similar to those found in natural water. *Water Res.* **5**, 171–&, doi: 10.1016/0043-1354(71)90049-2 (1971).
53. Roberts, D. A. *et al.* Ocean acidification increases the toxicity of contaminated sediments. *Glob. Chan. Biol.* **19**, 340–351, doi: 10.1111/gcb.12048 (2013).
54. Campbell, A. L., Mangan, S., Ellis, R. P. & Lewis, C. Ocean acidification increases copper toxicity to the early life history stages of the polychaete *Arenicola marina* in artificial seawater. *Environ. Sci. Technol.* (2014).
55. Lewis, C., Clemow, K. & Holt, W. V. Metal contamination increases the sensitivity of larvae but not gametes to ocean acidification in the polychaete *Pomatoscerus lamarckii* (Quatrefages). *Mar. Biol.* **160**, 2089–2101 (2013).
56. Meinshausen, M. *et al.* The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Clim. Change* **109**, 213–241, doi: 10.1007/s10584-011-0156-z (2011).

57. Truchot, J. P. Carbon dioxide combining properties of blood of shore crab *Carcinus maenas*-I-carbon dioxide solubility coefficient and carbonic acid dissociation constants. *J. Exp. Biol.* **64**, 45–57 (1976).
58. Parry, H. E. & Pipe, R. K. Interactive effects of temperature and copper on immunocompetence and disease susceptibility in mussels (*Mytilus edulis*). *Aquat. Tox.* **69**, 311–325, doi: 10.1016/j.aquatox.2004.06.003 (2004).
59. Camejo, G., Wallin, B. & Enojärvi, M. In *Free radical and antioxidant protocols. Methods in molecular biology* Vol. 108 (ed Armstrong D.) 377–387 (Humana, 1999).
60. Lewis, C. & Galloway, T. Genotoxic damage in polychaetes: A study of species and cell-type sensitivities. *Mutat. Res.-Gen. Tox. Environ. Mutagen.* **654**, 69–75, doi: 10.1016/j.mrgentox.2008.05.008 (2008).

Acknowledgements

CL was supported by a Natural Environment Research Council (NERC) UK Fellowship: NE/G014728/1. CL, RE and RW were supported by a UK-OARP NERC consortium grant NE/H017496/1. SN was supported by a Cefas-Exeter funded studentship. Thanks to Jan Shears, Darren Rowe and John Dowdle for their excellent technical support. The determination of total copper in the seawater media was undertaken by Dr. A. Fisher of the Analytical Research Facility, SoGEES, Plymouth University under ISO 9001:2008 certification. The authors would like to thank John Spicer for his insightful comments on the manuscript.

Author Contributions

C.L. designed and led the study, C.L., R.E., K.E., E.V. and S.N. all contributed to the running of the experiments and the collection and analysis of the data; C.L. wrote the manuscript; R.W.W. advised on all aspects of design and interpretation and contributed to the manuscript. All authors read and approved the final version of the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Lewis, C. *et al.* Ocean acidification increases copper toxicity differentially in two key marine invertebrates with distinct acid-base responses. *Sci. Rep.* **6**, 21554; doi: 10.1038/srep21554 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>