

# SCIENTIFIC REPORTS



OPEN

## Ocean acidification reduces transfer of essential biomolecules in a natural plankton community

J. Rafael Bermúdez<sup>1,2</sup>, Ulf Riebesell<sup>1</sup>, Aud Larsen<sup>3</sup> & Monika Winder<sup>1,4</sup>

Received: 06 January 2016

Accepted: 24 May 2016

Published: 21 June 2016

Ocean acidification (OA), a process of increasing seawater acidity caused by the uptake of anthropogenic carbon dioxide (CO<sub>2</sub>) by the ocean, is expected to change surface ocean pH to levels unprecedented for millions of years, affecting marine food web structures and trophic interactions. Using an *in situ* mesocosm approach we investigated effects of OA on community composition and trophic transfer of essential fatty acids (FA) in a natural plankton assemblage. Elevated *p*CO<sub>2</sub> favored the smallest phytoplankton size class in terms of biomass, primarily picoeukaryotes, at the expense of chlorophyta and haptophyta in the nano-plankton size range. This shift in community composition and size structure was accompanied by a decline in the proportion of polyunsaturated FA (PUFA) to total FA content in the nano- and picophytoplankton size fractions. This decline was mirrored in a continuing reduction in the relative PUFA content of the dominant copepod, *Calanus finmarchicus*, which primarily fed on the nano-size class. Our results demonstrate that a shift in phytoplankton community composition and biochemical composition in response to rising CO<sub>2</sub> can affect the transfer of essential compounds to higher trophic levels, which rely on their prey as a source for essential macromolecules.


Biomolecules synthesized by primary producers such as fatty acids (FA), and in particular polyunsaturated FA (PUFA), are considered essential metabolites for zooplankton and fish larvae<sup>1</sup>. As most consumers cannot synthesize these compounds *de novo*, they have to acquire them through their diet<sup>1</sup>. Laboratory experiments on the PUFA composition of microalgal cultures under different CO<sub>2</sub> scenarios revealed a range of responses, ranging from increased<sup>2</sup>, to stable<sup>3</sup>, to reduced PUFA fractions<sup>4–6</sup>. Considering that food quality, including the FA composition, can be as important as food quantity for the transfer efficiency up the food web and the productivity of higher trophic levels<sup>7</sup>, a better understanding of OA effects on the biochemical composition of primary producers and the trophic transfer of this signal seems essential.

At community level, the food quality to a large extent is determined by the phytoplankton species composition, as different algal taxonomic groups have distinct biochemical signatures<sup>8</sup>. Experiments with natural plankton assemblages indicate that changing *p*CO<sub>2</sub> can have pronounced impacts on phytoplankton community structure<sup>9,10</sup>, particularly small-sized algae like picoeukaryotes and cyanobacteria<sup>11,12</sup>. Thus, OA can affect the quality and quantity of food available for upper trophic levels by changing both the taxonomic composition and the biochemical signature of primary producers.

To investigate the effects of OA on phytoplankton species composition and consequences of biochemical transfer to upper trophic levels, we conducted an *in situ* mesocosm study enclosing a natural North Sea plankton community. The carbonate chemistry in the mesocosms was manipulated via the addition of CO<sub>2</sub>-saturated seawater to yield *p*CO<sub>2</sub> ranging from current to projected levels by end of this century<sup>13</sup>, plus two extreme treatments beyond projected levels. We monitored the plankton community composition, FA content of three size fractions (micro: 100–10 μm, nano: 10–2.7 μm, pico: 2.7–0.3 μm), as well as zooplankton FA in the dominant copepod *Calanus finmarchicus* over the course of the experiment, and classified FA as PUFA, MUFA (monounsaturated fatty acids) and SFA (saturated fatty acids).

<sup>1</sup>GEOMAR | Helmholtz Centre for Ocean Research Kiel, Germany. <sup>2</sup>Facultad de Ingeniería Marítima, Ciencias Biológicas, Oceánicas y Recursos Naturales. Escuela Superior Politécnica del Litoral, ESPOL, Guayaquil, Ecuador.


<sup>3</sup>The Hjort Centre for Marine Ecosystem Dynamics, Uni Research Environment, 5008 Bergen, Norway. <sup>4</sup>Department of Ecology, Environment and Plant Sciences, Stockholm University, Sweden. Correspondence and requests for materials should be addressed to J.R.B. (email: jrbermud@espol.edu.ec)




**Figure 1.** Average calculated biomass of the phytoplankton between sampling days 3 and 25, by size fraction: (a) microplankton ( $>10\mu\text{m}$ ), (b) nanoplankton ( $10\text{--}2.7\mu\text{m}$ ) and (c) picoplankton ( $<2.7\mu\text{m}$ ) in the  $\text{CO}_2$  gradient treatments during the experiment. A Mixed Effect Model analysis showed that all size fractions had significant differences in biomass between the  $p\text{CO}_2$  treatments. The x-axes display the mean  $\text{CO}_2$  levels ( $\mu\text{atm}$ ) during the analyzed time period in each mesocosm, bars show standard error.

## Results and Discussion

The size structure of the phytoplankton community shifted during the experiment in the different  $\text{CO}_2$  treatments. A Mixed Effect Model (MEM) analysis from sampling day 3 to sampling day 25 performed by size fraction revealed that plankton biomass in the micro- (Fig. 1a, Fig. S1) and nano-size (Fig. 1b, Fig. S1) fractions changed significantly through time at different  $p\text{CO}_2$  treatments (MEM micro:  $p\text{CO}_2$   $F = 12.95$ ,  $p < 0.0001$ ,  $df = 85$ ; time  $F = 24.32$ ,  $p < 0.0001$ ,  $df = 85$ ;  $p\text{CO}_2$ : time interaction  $F = 12.09$ ,  $p < 0.0001$ ,  $df = 85$ . MEM Nano:  $p\text{CO}_2$   $F = 28.73$ ,  $p < 0.0001$ ,  $df = 81$ ; time  $F = 5.17$ ,  $p = 0.025$ ,  $df = 81$ ;  $p\text{CO}_2$ : time interaction  $F = 42.37$ ,  $p < 0.0001$ ,  $df = 81$ ), while the pico-size fraction biomass (picocyanobacteria and picoeukaryotes) showed a fast change with  $p\text{CO}_2$ , reaching a significantly higher abundance at elevated treatments in a short time (pico:  $p\text{CO}_2$   $F = 74.47$ ,  $p < 0.0001$ ,  $df = 85$ ; time  $p = 0.8496$ ;  $p\text{CO}_2$ : time interaction  $F = 3.6873$ ,  $p = 0.0582$ ) (Fig. 1c; Fig. S1). These observations are consistent with some studies showing that small-sized phytoplankton benefit from high  $p\text{CO}_2$  levels in terms of growth rate and biomass accumulation<sup>11,12</sup>. A Mann-Kendall trend analyses over the duration of the experiment revealed that this change was due to differential responses of major taxonomic groups to elevated  $p\text{CO}_2$  (Fig. 2a). Chlorophyta (mostly Prasinophyceae in the nano-size range), the most abundant group, gradually increased significantly ( $p < 0.05$ ) in low  $p\text{CO}_2$  treatments and did not change at intermediate and high  $p\text{CO}_2$  conditions over the experimental duration (Fig. 2a). Haptophyta, dominated by the coccolithophore *Emiliania huxleyi* in the nano-size class, increased significantly at low  $p\text{CO}_2$  levels, were unaffected at intermediate values and significantly declined at high  $\text{CO}_2$  treatments (Fig. 2a). The negative  $\text{CO}_2$  effect on the coccolithophores observed in the present experiment is consistent with other mesocosm studies that showed a diminished abundance of calcifying algae at high  $p\text{CO}_2$  levels attributed to reduced growth rates at increased  $\text{CO}_2$  conditions<sup>14</sup>. The same pattern was




**Figure 2.** (a) Mann Kendall trend statistics of major phytoplankton taxonomic group biomass, between sampling days 3 and 25, across mesocosm treatments. Trends are shown as percent change per day (Trend %day $^{-1}$ ); blue dots indicate significant trends ( $p < 0.05$ ), grey dots no significant change (\* $p$ -value = 0.056). The x-axes display the mean  $pCO_2$  levels ( $\mu atm$ ) during the analyzed time period in each mesocosm. (b) Mean of picoeukaryote biomass between day 3 and 25 during the experiment. A Mixed Effect Model analysis revealed a significant increase at elevated  $pCO_2$  levels in the biomass between the different treatments ( $pCO_2$ :  $F = 83.2834$ ,  $p < 0.0001$ ,  $df = 84$ ; time:  $p = 0.360$ ;  $pCO_2$ : time interaction,  $p = 0.102$ ). Mann Kendall trend test could not be applied to this group, as biomass increased non-monotonic during the experiment. Bars show standard error.



**Figure 3.** Relative Polyunsaturated Fatty Acids (PUFAs) content in (a) microplankton ( $>10\mu\text{m}$ ; no significant difference), (b) nanoplankton ( $10\text{--}2.7\mu\text{m}$ ; MEM,  $F = 14.70$ ,  $p < 0.001$ ,  $df = 44$ ) and (c) picoplankton ( $<2.7\mu\text{m}$ ; MEM,  $F = 6.58$ ,  $p = 0.013$ ,  $df = 42$ ) size fractions over the  $p\text{CO}_2$  gradient treatments between sampling days 5 and 29. The x-axes display the mean  $p\text{CO}_2$  levels ( $\mu\text{atm}$ ) during the analyzed time period in each mesocosm. Bars show standard error.

observed for the sum of miscellaneous phytoplankton taxa (Euglenophyta, Heterokontophyta and Dinophyta in the micro-size range) (Fig. 2a, Others), which comprised only a small fraction of algal biomass in all mesocosms. Cryptophyta in the nano-size fraction increased significantly over time with no consistent pattern between  $p\text{CO}_2$  treatments (Fig. 2a). Picocyanobacteria (*Synechococcus*) increased significantly over the course of the experiment in all mesocosms with a stronger increase at high  $p\text{CO}_2$  but represented a small fraction of the total biomass (Fig. 2a, Fig. S1). Picoeukaryote biomass, as indicated by a mixed model effect, increased with the treatment gradient and became the most dominant taxa at elevated  $p\text{CO}_2$  towards the end of the experiment (Fig. 2b, Fig. S1). The change of the community composition in terms of species biomass over time and  $\text{CO}_2$  was confirmed by a Non-metrical Multidimensional Scaling analysis, which showed a strong gradual shift (Fig. S2) confirmed by an Analysis of Similarity ( $R = 0.40$ ;  $p < 0.001$ ), while the change associated with  $\text{CO}_2$  was comparatively smaller although significant ( $R = 0.18$ ;  $p < 0.001$ ).

The change in the plankton community composition was accompanied with a change in the FA composition of the different plankton size fractions. The relative content of the ecologically important PUFA group revealed marked differences between size fractions, with  $\sim 7\%$  ( $3 \pm 2\text{ ng L}^{-1}$ ) of total FA in the micro-,  $\sim 34\%$  ( $39 \pm 14\text{ ng L}^{-1}$ ) in the nano- and  $\sim 20\%$  ( $21 \pm 15\text{ ng L}^{-1}$ ) in the pico-size fraction. The PUFA content of the micro-size plankton fraction did not change with  $p\text{CO}_2$  (Fig. 3a), while there was a significant negative  $p\text{CO}_2$  effect at the nano- and pico-size fraction (Fig. 3b,c) at low  $p\text{CO}_2$  levels. In the pico and nano-size fractions the most important PUFA were Eicosapentaenoic (20:5n3, EPA), Eicosatrienoic (20:3n3, ETE) and Docosahexaenoic (22:6n3, DHA) acids (Fig. S3a,b), representing  $\sim 60\%$  of the total PUFA. The  $\text{CO}_2$ -related PUFA decrease in the nano- and pico-size fractions was associated with an increase of both SFA and mostly MUFA (Fig. S4).



**Figure 4.** (a) Relative Polyunsaturated Fatty Acids (PUFAs) content in the copepod *Calanus finmarchicus* across the different CO<sub>2</sub> treatments (MEM,  $F = 80.74$ ,  $p = 0.001$ ,  $df = 31$ ). The x-axes display the mean  $p\text{CO}_2$  levels ( $\mu\text{atm}$ ) between days 5–33 in each mesocosm. Bars show standard error. (b) Relative PUFAs content of *C. finmarchicus* between days 1–33 across CO<sub>2</sub> treatments. High  $p\text{CO}_2$  treatments (left) showed a significant decrease through time (Linear regression, 1120:  $R^2 = 0.73$ ,  $p = 0.017$ ; 1400:  $R^2 = 0.97$ ,  $p < 0.001$ ; 2000:  $R^2 = 0.86$ ,  $p < 0.01$ ; 3000:  $R^2 = 0.93$ ,  $p = 0.001$ ).

	SFA:PUFA	sd	SFA:MUFA	sd	MUFA:PUFA	sd
<i>Calanus</i>	<b>1.67</b>	<b>0.14</b>	<b>3.48</b>	<b>1.03</b>	<b>0.48</b>	<b>0.1</b>
Micro-fraction	12.13	4.32	5.74	2.1	2.3	0.96
Nano-fraction	<b>1.41</b>	<b>0.33</b>	<b>3.19</b>	<b>1.37</b>	<b>0.52</b>	<b>0.22</b>
Pico-fraction	3.2	1.24	3.1	1.36	1.12	0.54

**Table 1.** Ratios of FA in the copepod *Calanus finmarchicus* to different phytoplankton size fractions (micro: 10–100  $\mu\text{m}$ , nano: 2.7–10  $\mu\text{m}$ , pico: 0.3–2.7  $\mu\text{m}$ ). Ratios are shown for different FA classes: Saturated (SFA), Monounsaturated (MUFA) and Polyunsaturated (PUFA). Bold values show the similarity between *C. finmarchicus* and the nano-size phytoplankton fraction.

The reduced PUFA content in the nano-size fraction is likely a result of the change in plankton community composition, especially due to the reduced Haptophyta biomass at high  $p\text{CO}_2$  level, which are a very good sources of PUFA<sup>8</sup>. The reduced PUFA content in the pico-size fraction can be due to an increase of picocyanobacteria and picoeukaryotes abundance. The first are known for being a poor PUFA source<sup>15</sup>, while the high abundance of the later at elevated  $p\text{CO}_2$  suggests a PUFA loss in this group. Although we have not identified the species composition of the picoeukaryote group, significant increases of pico- chlorophyta (Prasinophyceae) have been observed in plankton assemblages at elevated CO<sub>2</sub> conditions<sup>11</sup>. Chlorophyta have a relatively low PUFA content<sup>8</sup>, which may contribute to the reduced PUFA content at high  $p\text{CO}_2$ . Given that our experiment did not allow the distinction between an indirect effect on PUFA concentration through shifts in species composition and a direct CO<sub>2</sub> response on cell physiology, the latter cannot be excluded. It has been shown for both, Chlorophyta and the Haptophyte *Emiliania huxleyi* that culturing at high CO<sub>2</sub> reduced their PUFA content<sup>4,16</sup>. This suggests that the relative PUFA decrease in both size fractions is most likely a combination of a change in community composition and a direct CO<sub>2</sub> effect on cell physiology.

Our results showing low PUFA levels at high  $p\text{CO}_2$  are contrasting the response of an Arctic plankton community, in which relative PUFA content increased at elevated  $p\text{CO}_2$  levels<sup>17</sup>. The divergent results can be attributed to the different plankton community compositions<sup>8</sup>. However, this can also be due to the size-fractionation of the samples for FA analysis in the present experiment, which was not done in in the Arctic study. A change in the FA of a single species within the community, especially if it has a large biomass, may mask changes in other size classes or taxonomic groups. The observed PUFA increment in the Arctic was associated to an elevated abundance of large-sized dinoflagellates<sup>17</sup>, which are typically high in PUFA content<sup>8</sup> and therefore may not reflect a net positive CO<sub>2</sub> effect on all primary producers, but rather in this specific group. Thus, biochemical changes of primary producers belonging to different size fractions may affect higher trophic levels differently as aquatic predators are size-selective<sup>18</sup>.

The decrease of PUFA in the nano-size fraction observed in this study affected the PUFA profile of the dominant copepod species. The PUFA content in *C. finmarchicus* represented  $\sim 32\%$  ( $120 \pm 40 \text{ ng ind.}^{-1}$ ) of total FA, and the relative PUFA content decreased significantly with higher  $p\text{CO}_2$  levels (Fig. 4a). While staying constant at low  $p\text{CO}_2$  levels, it gradually and significantly decreased at  $p\text{CO}_2$  treatments of 1120  $\mu\text{atm}$  and higher, with an average loss of  $\sim 4\%$ , and strongly declined ( $\sim 10\%$ ) at the highest  $p\text{CO}_2$  treatment over the course of the experiment (Fig. 4b). In the copepod, like in the plankton size fractions, the Eicosapentaenoic (20:5n3, EPA), Eicosatrienoic (20:3n3, ETE) and Docosahexaenoic (22:6n3, DHA) acids where the most abundant (Fig. S3c) and represented around  $\sim 90\%$  of the total PUFA. The relative PUFA decline was associated with an increase of MUFA and SFA (Fig. S5). The ratio of the different FA classes (PUFA, MUFA, and SFA) in copepods closely resembled the corresponding proportions in the nano-size phytoplankton fraction (Table 1). This close resemblance was

maintained throughout the entire experiment (Table S1), indicating that *C. finmarchicus* strongly relied on this size fraction as a PUFA resource. The copepod's prey-dependent FA profile is coherent with previous studies showing that copepods strongly rely on their diet for essential PUFA and that their FA composition mirrors the algae they graze on<sup>19</sup>, and with reports showing that this species capitalizes efficiently on small-sized algal prey when present in high amounts, like in the present study<sup>20,21</sup>. Even though we were not able to evaluate the impact of the PUFA reduction on *C. finmarchicus* life cycle (~1 year long)<sup>21</sup> as the time frame of the experiment (~6 weeks) did not allow the observation of significant changes in the population structure; it has been observed in previous studies that even small changes in the food source of FA have important consequences in copepods and fish<sup>19,22</sup>.

Although a direct CO<sub>2</sub> effects on copepod FA synthesis cannot be excluded, it seems unlikely. Previous experiments showed that *C. finmarchicus* is rather insensitive to elevated CO<sub>2</sub> exposure<sup>23</sup>. Another factor that may have influenced FA transfer to the copepods in the present study is the shift in size of the plankton community, with the pico-size fraction becoming more dominant at high pCO<sub>2</sub> levels. A reduction of mean prey size towards small-sized cells can cause a decline in feeding efficiency of large copepods, such as *C. finmarchicus*, irrespective of their feeding mechanism<sup>18</sup>. Therefore, the observed decrease in the *C. finmarchicus* PUFA content at high pCO<sub>2</sub> conditions may be a combined consequence of a CO<sub>2</sub>-driven deterioration of the food quality and prey availability.

Our study provides the first evidence that a CO<sub>2</sub>-driven shift in community composition and associated change in food quality in terms of FA at the base of the food web can be transferred to primary consumers in a natural marine assemblage. A diatom grown at 750 µatm pCO<sub>2</sub> showed a PUFA decrease of ~20% that, when used to feed a copepod, produced a ~29% PUFA reduction in them with a concomitant decline in both somatic growth and egg production of ~85%<sup>19</sup>. For fish it has been shown that a decrease of just 0.6% in the amount of n-3 FA in the food source can reduce their egg viability by ~10% and larval survival by ~15%<sup>22</sup>. Given that fish is a critical natural resource<sup>24</sup>, acidification-driven FA quality deterioration may reduce the availability of essential PUFA in human diets<sup>25</sup>. Nonetheless, the plankton community response to OA will strongly depend on how changes in CO<sub>2</sub> affect species composition of primary producers, whereby genetic and physiological diversity in natural assemblages may buffer adverse CO<sub>2</sub> effects<sup>17,26</sup>.

## Methods

An *in situ* mesocosm CO<sub>2</sub> perturbation experiment in Raunefjord, southern Norway was performed during late spring 2011 using nine enclosures with a length of 25 m containing ~75 m<sup>3</sup> of natural seawater. The mesocosms were set up and manipulated as described in detail by Riebesell *et al.*<sup>27</sup>. The carbonate system was manipulated through the addition of CO<sub>2</sub>-saturated seawater to seven of the nine mesocosms in four steps between days 0 to 3 to achieve target values of 380, 560, 840, 1120, 1400, 2000 and 3000 µatm plus one control unit with a natural concentration of 280 µatm. Samples for the calculation of the pCO<sub>2</sub> in each mesocosm were taken every sampling day, and was calculated based on measurements of total alkalinity, pH (in total scale), salinity and temperature, using the computer program CO2SYS<sup>28</sup>. Samples for total alkalinity measurements were filtered on a Whatman GF/F filter and measured by open-cell acidimetric titration as describes by Dickson (2010)<sup>29</sup>. pH was measured spectrophotometrically with a VARIAN Cary 100 in 10 cm cuvette at 25°C and then recalculated to *in-situ* temperature as described in Dickson (2010)<sup>29</sup>. The precision on the pH measurements was typically over 0.001 units at high and 0.002 units at low pH. Vertical profiles of temperature and conductivity were taken daily in each mesocosm with a CTD60M (Sun and Sea Technologies). The measurements of pCO<sub>2</sub>, pH and inorganic nutrients for all nine mesocosm are provided in Table S2.

On day 14, nitrate and phosphate were added to all mesocosms to a final concentration of 5 and 0.2 µmol l<sup>-1</sup>, respectively. Samples for phytoplankton counts and flow cytometry were taken every second day and for fatty acids every fourth day using a depth-integrated water sampler (Hydrobios, Kiel, Germany) covering the upper 20 m of the water column from days 1 to 25; integrated zooplankton net tows were taken every 7th day starting the day before the CO<sub>2</sub> manipulation (day-1) until day 33. Phytoplankton cell counts were carried out from 50 ml sample water, fixed with alkaline Lugol's iodine (1% final concentration) using an Utermöhl chamber with an inverted microscope (ZEISS Axiovert 100) and identified up to species level when possible (Table S3). Picoplankton cells were enumerated using a bench-top Becton Dickinson FACSCalibur flow cytometer (FCM) equipped with a 488 nm argon laser. Biovolume was calculated according to geometric shapes and converted to cellular organic carbon using taxon-specific conversion equations; for micro- and nanophytoplankton according to Menden-Deuer<sup>30</sup>, and for picophytoplankton according to Worden *et al.*<sup>31</sup>. For algal FA, 1 L of seawater was filtered in three size fractions: 100 to 10 µm (micro), 10 to 2.7 µm (nano) and 2.7 to 0.3 µm (pico) by using, non-combusted Millipore NY1002500 (10 µm ø pore), Whatman GF/D (2.7 µm ø pore) pre-combusted (450°C, 6 h) and Advantec Grade GF 75 (0.3 µm ø pore) filters, respectively. Individuals of the copepod *Calanus finmarchicus* (copepodite stage V) were sorted for FA measurements. All samples were immediately stored at -80°C until analysis. FA were measured by gas chromatography as fatty acid methyl esters (FAME) following Klein Breteler *et al.*<sup>32</sup>. FAME were analyzed by a Thermo GC Ultra gas chromatograph equipped with a non-polar column (RX11-SIL-MS 0.32 µm, 30 m, company Restek) and Flame ionization detector and were clustered according to their degree of saturation: saturated (SFA), monounsaturated (MUFA) and polyunsaturated (PUFA).

To identify differences in the relative content of FAs between the treatments a nested Mixed Effects Model (MEM) was used to determine the differences in relative fatty acid content (%) between the CO<sub>2</sub> treatments (µatm) through time, with the treatment level as nested random variable (random distribution of CO<sub>2</sub> treatments among the mesocosm). Mann-Kendall trend test was applied to analyze for monotonic trends in the data to determine the temporal change for plankton taxonomic groups in terms of calculated biomass throughout the

experiment. Slopes were expressed as % day<sup>-1</sup> by dividing the experimental mean of the variable; in the case of the picoeukaryotes, where the biomass distribution was non-monotonic, MEM analysis as described above was performed. The similarity in the structure of the plankton community between the treatments in terms of calculated species biomass in the mesocosm was analyzed by Non Metrical Multidimensional Scaling (NMDS) with Bray distance, auto-transformation and 4 dimensions (k = 4). An analysis of similarity (ANOSIM) using a Bray-Curtis distance matrix and 5000 permutations was used to corroborate the NMDS results. All statistical analyses were done using the R software environment 3.0.1<sup>33</sup>.

## References

- Arts, M. & Kohler, C. In *Lipids in aquatic ecosystems* 1st edn (eds Arts, M. et al.) Ch. 9, 237–256 (Springer, 2009).
- Hoshida, H. et al. Accumulation of eicosapentaenoic acid in *Nannochloropsis* sp. in response to elevated CO<sub>2</sub> concentrations. *J. Appl. Phycol.* **17**, 29–34 (2005).
- Tsuzuki, M. et al. Effects of CO<sub>2</sub> concentration during growth on fatty acid composition in microalgae. *Plant Physiol.* **93**, 851–6 (1990).
- Fiorini, S., Gattuso, J.-P., van Rijswijk, P. & Middelburg, J. Coccolithophores lipid and carbon isotope composition and their variability related to changes in seawater carbonate chemistry. *J. Exp. Mar. Biol. Ecol.* **394**, 74–85 (2010).
- Torstensson, A. et al. Synergism between elevated pCO<sub>2</sub> and temperature on the Antarctic sea ice diatom *Nitzschia lecontei*. *Biogeosciences* **10**, 6391–6401 (2013).
- Bermúdez, R. et al. Long-term conditioning to elevated pCO<sub>2</sub> and warming influences the fatty and amino acid composition of the diatom *Cylindrotheca fusiformis*. *PLoS One* **10**, e0123945 (2015).
- Müller-Navarra, D. C. et al. Unsaturated fatty acid content in seston and tropho-dynamic coupling in lakes. *Nature* **427**, 69–72 (2004).
- Galloway, A. W. E. & Winder, M. Partitioning the relative importance of phylogeny and environmental conditions on phytoplankton fatty acids. *PLoS One* **10**(6), e0130053, doi: 10.1371/journal.pone.0130053 (2015).
- Feng, Y. et al. Effects of increased pCO<sub>2</sub> and temperature on the North Atlantic spring bloom. I. The phytoplankton community and biogeochemical response. *Mar. Ecol. Prog. Ser.* **388**, 13–25 (2009).
- Tatters, A. et al. Short- and long-term conditioning of a temperate marine diatom community to acidification and warming. *Phil. Trans. R. Soc. B.* **368**, 20120437 (2013).
- Newbold, L. K. et al. The response of marine picoplankton to ocean acidification. *Environ. Microbiol.* **14**, 2293–307 (2012).
- Brussaard, C. P. D. et al. Arctic microbial community dynamics influenced by elevated CO<sub>2</sub> levels. *Biogeosciences* **10**, 719–731 (2013).
- Pörtner, H. O. et al. in *Climate change 2014: Impacts, adaptation and vulnerability* (eds Drinkwater, K. et al.) Ch. 6. 1–138 (Cambridge University Press, 2014).
- Engel, A. et al. Testing the direct effect of CO<sub>2</sub> concentration on a bloom of the coccolithophorid *Emiliania huxleyi* in mesocosm experiments. *Limnol. Oceanogr.* **50**, 493–507 (2005).
- Patil, V. et al. Fatty acid composition of 12 microalgae for possible use in aquaculture feed. *Aquac. Int.* **15**, 1–9 (2006).
- Sato, N., Tsuzuki, M. & Kawaguchi, A. Glycerolipid synthesis in *Chlorella kessleri* 11 h II. Effect of the CO<sub>2</sub> concentration during growth. *Biochim. Biophys. Acta* **1633**, 35–42 (2003).
- Leu, E. et al. Effect of ocean acidification on the fatty acid composition of a natural plankton community. *Biogeosciences* **10**, 1143–1153 (2013).
- Kjørboe, T. How zooplankton feed: mechanisms, traits and trade-offs. *Biol. Rev.* **86**, 311–39 (2011).
- Rossoll, D. et al. Ocean acidification-induced food quality deterioration constrains trophic transfer. *PLoS One* **7**(4), e34737, doi: 10.1371/journal.pone.0034737 (2012).
- Nejstgaard, J., Båmstedt, U. & Bagoien, E. Feeding and reproduction by *Calanus finmarchicus*, and microzooplankton grazing during mesocosm blooms of diatoms and the coccolithophore *Emiliania huxleyi*. *Mar. Ecol. Prog. Ser.* **147**, 197–217 (1997).
- Harris, R. et al. Feeding, growth, and reproduction in the genus *Calanus*. *ICES J. Mar. Sci.* **57**, 1708–1726 (2000).
- Izquierdo, M., Fernández-Palacios, H. & Tacon, A. G. Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture* **197**, 25–42 (2001).
- Mayor, D. et al. CO<sub>2</sub>-induced acidification affects hatching success in *Calanus finmarchicus*. *Mar. Ecol. Prog. Ser.* **350**, 91–97 (2007).
- Food and Agriculture Organization of the United Nations. in *The State of World Fisheries and Aquaculture 2014* 1st edn. Ch. 1, 62–68 (FAO, 2014).
- Kang, J. X. Omega-3: a link between global climate change and human health. *Biotechnol. Adv.* **29**, 388–90 (2011).
- Rossoll, D., Sommer, U. & Winder, M. Community interactions dampen acidification effects in a coastal plankton system. *Mar. Ecol. Prog. Ser.* **486**, 37–46 (2013).
- Riebesell, U. et al. Technical Note: A mobile sea-going mesocosm system – new opportunities for ocean change research. *Biogeosciences* **10**, 1835–1847 (2013).
- Lewis, E. & Wallace, D. in *Program developed for CO<sub>2</sub> system calculations ORNL/CDIAC-105*. 54 pp. (U.S. Department of Energy, 1998).
- Dickson, A. G. in *Guide to Best Practices in Ocean Acidification Research and Data Reporting* 1st edn. (eds Riebesell, U. et al.) Ch. 1, 17–40 (Publications Office of the European Union, 2010).
- Menden-Deuer, S. & Lessard, E. J. Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. *Limnol. Oceanogr.* **45**, 569–579 (2000).
- Worden, A. Z., Drive, G., Jolla, L. & Nolan, J. K. Assessing the dynamics and ecology of marine picophytoplankton: The importance of the eukaryotic component. *Limnol. Oceanogr.* **49**, 168–179 (2004).
- Breteler, W., Schogt, N. & Baas, M. Trophic upgrading of food quality by protozoans enhancing copepod growth: role of essential lipids. *Mar. Biol.* **191**–198 (1999).
- R Development Core Team. *R: A language and environment for statistical computing*. (2013) Available at: <http://www.r-project.org/> (Accessed 10th September 2014).

## Acknowledgements

We thank Daniela Freese, Hendrik Schultz, Andrea Ludwig and Thomas Hansen for laboratory and technical assistance. This work is a contribution to the BMBF coordinated projects “Surface Ocean Processes in the Anthropocene” (SOPRAN), and “Biological Impacts of Ocean Acidification” (BIOACID) funded by the German Federal Ministry of Education and Research (BMBF) under contract numbers 03F0611C and 03F0608B, and to the Marie Curie IRG grant No. 276917.

### Author Contributions

U.R. designed the experiments; J.R.B., M.W. and U.R. designed the sampling method; J.R.B., M.W. and A.L. collected the samples and analyzed the data; A.L. contributed materials/analysis tools; J.R.B. and M.W. co-wrote the paper, with input from the other authors.

### Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Bermúdez, J. R. *et al.* Ocean acidification reduces transfer of essential biomolecules in a natural plankton community. *Sci. Rep.* **6**, 27749; doi: 10.1038/srep27749 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>