

ENVIRONMENTAL STUDIES

Adaptability as the key to success for the ubiquitous marine nitrite oxidizer *Nitrococcus*

Jessika Füssel,^{1,2,*†} Sebastian Lücker,^{3,4,*†} Pelin Yilmaz,¹ Boris Nowka,⁵ Maartje A. H. J. van Kessel,⁴ Patric Bourceau,^{1,4} Philipp F. Hach,¹ Sten Littmann,¹ Jasmine Berg,¹ Eva Spieck,⁵ Holger Daims,³ Marcel M. M. Kuypers,¹ Phyllis Lam^{1,2}

Nitrite-oxidizing bacteria (NOB) have conventionally been regarded as a highly specialized functional group responsible for the production of nitrate in the environment. However, recent culture-based studies suggest that they have the capacity to lead alternative lifestyles, but direct environmental evidence for the contribution of marine nitrite oxidizers to other processes has been lacking to date. We report on the alternative biogeochemical functions, worldwide distribution, and sometimes high abundance of the marine NOB *Nitrococcus*. These largely overlooked bacteria are capable of not only oxidizing nitrite but also reducing nitrate and producing nitrous oxide, an ozone-depleting agent and greenhouse gas. Furthermore, *Nitrococcus* can aerobically oxidize sulfide, thereby also engaging in the sulfur cycle. In the currently fast-changing global oceans, these findings highlight the potential functional switches these ubiquitous bacteria can perform in various biogeochemical cycles, each with distinct or even contrasting consequences.

INTRODUCTION

Nitrite oxidation is the major pathway that generates nitrate, the dominant form of biologically available nitrogen and often a limiting nutrient to biological production in surface oceans. The prevalence of nitrate below surface, however, attests to the substantial occurrence of nitrite oxidation in the ocean interior, which represents a key conduit to resupply nitrogen to the surface ocean and is thus essential to the global nitrogen cycle (1). Nevertheless, current knowledge of the physiology of marine nitrite-oxidizing bacteria (NOB) remains poor, with only a few species described to date that belong to the genera *Nitrobacter*, *Nitrospira*, *Nitrospina*, *Candidatus Nitromaritima*, or *Nitrococcus* (2, 3). Although several studies have challenged the conventional understanding of nitrate production as the only relevant contribution of NOB to global biogeochemical cycling (4–7), no direct environmental evidence of marine NOB participating in elemental cycles other than nitrogen and carbon has been reported to date.

Most recent research on marine nitrite oxidizers has been heavily biased toward *Nitrospina*, which can be found in a wide range of habitats, including open ocean (8), sediments (9, 10), and suboxic water columns (11, 12). Experimental and genome analyses of the two cultured *Nitrospina* strains, *Nitrospina gracilis* and *Nitrospina watsonii*, strongly suggest that these organisms are chemolithoautotrophs dependent on nitrite oxidation for energy conservation (13–15). In comparison, *Nitrococcus*, with only one species known to date, *Nitrococcus mobilis*, has had its presence reported only twice in the 1980s in the Pacific (16, 17) and once in a high-salinity pond (18) since its first isolation (13); its presence has not been reported again until its recently recorded abundance in the Namibian oxygen minimum zone (OMZ) where they

even outnumbered *Nitrospina* (11). In that study, *Nitrococcus* and *Nitrospina* were the only NOB genera detectable via catalyzed reporter deposition fluorescence in situ hybridization (CARD-FISH) and together comprised almost 10% of the total microbial community. Although *Nitrospina* use a putatively high-affinity *cbb₃*-type terminal oxidase, which should allow them to catalyze nitrite oxidation even under hypoxic conditions (14), the adaptation strategy of *Nitrococcus* to oxygen-deficient conditions remains unclear. *Nitrospina* preferred residing in the oxycline and the upper OMZ, whereas *Nitrococcus*, in contrast, became more abundant in the lower OMZ (4.9% of total microbial abundance) where oxygen levels were minimal (11). Here, a similar trend was again observed for October 2011, with *Nitrospina* contributing up to 1.3% of total microbial community in the upper OMZ (65 m, 95 $\mu\text{M O}_2$), whereas *Nitrococcus* displayed maximal abundance (1.3%) at 119 m (16 $\mu\text{M O}_2$) (Fig. 1A). Overall, *Nitrococcus* showed a moderate yet significant negative correlation with oxygen (Spearman correlation = 0.44, $P < 0.01$) and was undetectable at oxygen concentrations $>20 \mu\text{M}$ (fig. S1A).

To investigate whether *Nitrococcus* is restricted to certain environmental settings, we searched for *Nitrococcus*-like sequences in publicly available databases. Three novel *Nitrococcus*-like phylotypes, in addition to *N. mobilis*, were identified in the SILVA 16S ribosomal RNA (rRNA) database (19, 20), sharing 97.8 to 99.2%, 96.5 to 97.8%, and 96.6 to 97.3% sequence identities with *N. mobilis* (fig. S2 and table S1). Subsequent searches for all phylotypes' 16S rRNA genes in marine amplicon and metagenomic data sets from the MetaGenome Rapid Annotation using Subsystems Technology (MG-RAST) server (21), Ocean Sampling Day 2014 (OSD) (22), and Tara Oceans (23) revealed the presence of *N. mobilis* and *Nitrococcus*-like phylotypes in 69% of all data sets examined. Among the data sets that carry geographical information, *N. mobilis* and *Nitrococcus*-like phylotypes were found to be globally distributed from coastal upwelling regions to oligotrophic open oceans, from tropical to temperate and polar latitudes, and from epipelagic (surface sunlit layer, ~0 to 100 m) to mesopelagic (twilight zone, ~100 to 1000 m) ocean waters (Fig. 1B and tables S2 and S3). Considering only metagenomic data sets (table S2) and excluding the more bias-prone amplicon sequencing data (table S3), higher contributions of *N. mobilis* and *Nitrococcus*-like phylotypes to total microbial community ($>5\%$, up to 17%) were nearly exclusively associated with highly productive environments, such as the

¹Max Planck Institute for Marine Microbiology, 28359 Bremen, Germany. ²Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Southampton SO14 3ZH, UK. ³Department of Microbiology and Ecosystem Science, University of Vienna, 1090 Vienna, Austria. ⁴Department of Microbiology, Institute for Water and Wetland Research, Radboud University, 6525 AJ Nijmegen, Netherlands. ⁵Section Microbiology, Biocenter Klein Flottbek, University of Hamburg, 22609 Hamburg, Germany.

*Corresponding author. Email: jessika.fuessel@soton.ac.uk (J.F.); s.luecker@science.ru.nl (S.L.)

†These authors contributed equally to this work.




Fig. 1. Occurrence and distribution of *Nitrococcus* in a marine OMZ and the global oceans. (A) Water column chemistry and nitrogen cycling rates in the Namibian OMZ. Left: Vertical distribution of oxygen, NO_2^- , NH_4^+ , and NO_3^- at station 2. Middle: Nitrite oxidation rates, N_2O production rates, and cellular abundance of *Nitrococcus* and *Nitrospina*. Right: Rates of ammonia oxidation, nitrate reduction, and anammox measured with ^{15}N -labeling experiments. Error bars show SDs derived from linear regression analyses. (B) Worldwide distribution of *Nitrococcus mobilis* and *Nitrococcus*-like phylotypes in the oceans. Combined abundance of all four phylotypes in different metagenomic data sets is denoted in orange (MG-RAST), turquoise (OSD), and blue (Tara Oceans). Circle size indicates percentage of 16S rRNA gene reads relative to all bacterial and archaeal 16S rRNA gene reads within each sample.

Peruvian OMZ and Guanabara Bay or with hypersaline lagoons, indicating an ecological advantage for them under these conditions (Fig. 1B and table S2). On the basis of Tara Oceans metagenomic data alone, which is currently the most comprehensive data set with consistent sequencing depths and covering the widest geographic range of global oceans to date, 16S rRNA genes affiliated with *Nitrococcus* were present in all samples and represented 1.2 to 5.1% (mean, 2.3%) of the microbial community (table S2).

We analyzed the genome of *N. mobilis* Nb-231 to identify potential functional roles *Nitrococcus* may play in such diverse environments. The resulting genome-derived hypotheses were tested in incubation

experiments with both the pure culture isolate *N. mobilis* Nb-231 and Namibian OMZ seawater samples. *Nitrococcus* only encodes a putatively low-affinity *aa₃*-type cytochrome c oxidase (Fig. 2 and table S4), thus ruling out a specialization on microaerobic respiration as suggested for *Nitrospina* (14). Instead, its abundance in OMZ waters indicates the utilization of alternative anaerobic metabolisms. In agreement with this hypothesis, *Nitrococcus* was abundant even in waters where no nitrite oxidation was detectable in the Namibian OMZ, and no correlation with nitrite oxidation rates was observed (Spearman correlation = 0.05, $P > 0.05$, $n = 35$) (Fig. 1A and fig. S1B). These findings suggest that NOB other than *Nitrococcus*, such as *Nitrospina*, were responsible for most of the nitrite oxidation detected in the Namibian OMZ (Fig. 1A).

On the other hand, *Nitrococcus* abundance was significantly correlated with nitrate reduction rates (fig. S1C) (Spearman correlation = 0.54, $P < 0.05$), suggesting a contribution to this anaerobic process. The key enzyme for nitrite oxidation, nitrite oxidoreductase (NXR), is reversible, and nitrate reduction activities have previously been observed in *Nitrobacter* (24, 25) and *Nitrospira* (6). When the *N. mobilis* pure culture was subjected to anoxic conditions in the presence of formate and acetate, substrates that have previously been reported to support mixotrophic or heterotrophic growth in NOB (6, 25), NO_3^- was reduced at a rate of 3.66 ± 0.32 fmol per cell per day ($\pm$ SD) within the first 230 hours of incubation (Fig. 3A). Because the *N. mobilis* genome does not encode any other known nitrate-reducing enzyme, the NXR of *Nitrococcus* is most likely the candidate for nitrate reduction (Fig. 2). Furthermore, *N. mobilis* encodes duplicated versions of NXR alpha subunit (NxrA), the substrate-binding catalytic subunit of the enzyme complex (fig. S3) (26, 27). NxrA1 is encoded in one genomic region together with genes for the NXR beta, delta, and gamma subunits. The duplicated subunit NxrA2 (fig. S3), with an amino acid identity of 89% to NxrA1, is not clustered with any other genes known to be involved in nitrite oxidation. This duplication of the catalytic NxrA subunit alone has also been observed in all sequenced *Nitrobacter* genomes and in *Nitrolancea hollandica* (28, 29). It is tempting to speculate that *Nitrococcus* uses one NxrA paralog for nitrite oxidation and the other for the reduction of nitrate, but biochemical verification will be necessary.

The observed switch to NO_3^- reduction with formate and acetate indicates that *Nitrococcus* can use organic matter as energy and carbon source. Although whether this truly translates to growth remains to be empirically verified, chemoorganoheterotrophic and autotrophic potentials of *Nitrococcus* are corroborated by the presence of all genes for the oxidative TCA (tricarboxylic acid) cycle, glycolysis, PPP (pentose phosphate pathway), and the Calvin-Benson-Bassham cycle in the genome (Fig. 2 and table S4). Genome annotation also revealed genes for respiratory complexes I to V (Fig. 2 and table S4). Complexes I to III are not required for energy conservation during nitrite oxidation, unlike during the oxidation of organic substrates, but they do mediate the reverse electron flow from nitrite to NAD^+ during chemolithoautotrophic growth. An ETF (electron-transport flavoprotein) complex and an ETF/quinone oxidoreductase probably couple β -oxidation of fatty acids to the respiratory chain.

Consistent with the pure culture experiments and genome analyses, uptake of organic carbon was also demonstrated by single-cell analyses of uncultured *Nitrococcus* in hypoxic Namibian OMZ seawater samples ($O_2 < 2$ μM, $NO_3^- > 15$ μM). Upon incubation with ^{13}C -labeled algal DOM, fluorescence in situ hybridization–secondary ion mass spectrometry (FISH-SIMS) analyses (30) of individual *Nitrococcus* cells showed substantial uptake of ^{13}C -DOM after 29 hours (0.3 to 48.1 atomic % ^{13}C) (Fig. 3B and table S5). The possibility of ^{13}C enrichment as a result of




Fig. 2. Cell metabolic cartoon based on the annotation of the *N. mobilis* Nb-231 genome. ETF, electron-transport flavoprotein; hmp, NO dioxygenase; HYD, hydrogenase; NirBD, assimilatory nitrite reductase; NirK, copper-containing nitrite reductase (NO-forming); NXR, nitrite oxidoreductase; sNOR, NO reductase; PHB, poly-hydroxybutyrate; poly-P, polyphosphates; SOR, sulfite dehydrogenase; SQR, sulfide/quinone oxidoreductase; PPP, pentose phosphate pathway; TRAP, tripartite ATP-independent periplasmic transporter (TRAP-T); TRK, K⁺ transporter; RND, resistance-nodulation-division; ATPase, adenosine triphosphatase; ABC, ATP-binding cassette. Enzyme complexes of the electron transport chain are labeled by Roman numerals. Red and orange diamonds represent cytochrome c proteins and quinones, respectively. The legend indicates transport protein classifications.

rapid remineralization of ¹³C-DOM by other organisms followed by lithoautotrophic uptake of ¹³C-HCO₃⁻ by *Nitrococcus* can be largely excluded, as no detectable ¹³C enrichment was observed in *Nitrococcus* cells in parallel anoxic incubations with ¹³C-HCO₃⁻ and NO₂⁻ (Fig. 3C). The high ¹³C enrichment in combination with our findings from genome analyses and pure culture experiments strongly indicate a chemo-organoheterotrophic potential of *Nitrococcus* under NO₃⁻ reducing conditions. The broad range of ¹³C enrichment observed in our incubations (table S5) can likely be attributed to the presence of different *Nitrococcus* phylotypes or phenotypic heterogeneity of individual cells within a population (31).

When exposed to elevated nitrite levels (~200 μM), pure cultures of *N. mobilis* continuously produced small but significant amounts of N₂O during anoxic incubations [$3.2 (\pm 0.48) \times 10^{-3}$ fmol per cell per day

during the first 230 hours] (Fig. 3B and fig. S4). No organic substrates were amended in these incubations, and denitrification was likely driven by the oxidation of dissolved organic carbon present in the sterilized seawater used for medium preparation (table S6) and/or the oxidation of intracellular storage compounds such as glycogen (Fig. 2) (13). *N. mobilis* encodes a copper-containing nitrite reductase (NirK) (table S4), likely facilitating nitrite reduction to nitric oxide (NO). None of the enzymes known to be responsible for the reduction of NO to N₂O—cytochrome c or quinone-dependent nitric oxide reductase (cNOR or qNOR, respectively)—were identified in the genome. However, the Nb-231 genome carries the genes for a two-subunit oxidase that belongs to a distinct phylogenetic cluster within the haem-copper oxidase family (table S4 and fig. S5). This enzyme group is designated the sNOR family and likely functions as NO reductases to detoxify NO_x




Fig. 3. Denitrifying potential of *Nitrococcus mobilis* Nb-231. (A) Anaerobic nitrogen dissimilation under (left) organoheterotrophic conditions (500 μM $^{15}\text{NO}_3^-$, 250 μM acetate, and 250 μM formate, anoxic) and (right) without the addition of electron donor (200 μM $^{15}\text{NO}_2^-$, anoxic). See the main text for more details. Error bars indicate SD of three biological replicates. (B) FISH-SIMS analyses of single *Nitrococcus* cells in Namibian OMZ bottom water samples (1.20 m above seafloor), incubated with (top) ^{13}C -labeled dissolved organic matter (DOM) or (bottom) ^{13}C -labeled bicarbonate under hypoxic conditions. Left: *Nitrococcus* cells (green) targeted by CARD-FISH among other microbial cells (blue). Middle: $^{32}\text{S}/^{12}\text{C}$ of the corresponding cells. Right: $^{13}\text{C}/^{12}\text{C}$ ratio. Scale bars, 1 μm .

intermediates produced in the periplasm, as proposed for ammonia-oxidizing bacteria (32). To our knowledge, this is the first time that partial denitrification (NO_3^- or NO_2^- reduction to N_2O) has been shown for an organism apparently containing this type of NOR only, thus giving indirect evidence that this enzyme family functions as NOR and might be used for energy conservation under denitrifying conditions. It should, however, be noted that the *N. mobilis* Nb-231 genome is not closed such that the definite absence of a canonical NOR is yet to be confirmed. Although we can exclude strictly abiotic N_2O production based on our abiotic control incubation (fig. S4), we cannot rule out abiotic-biotic hybrid N_2O formation, in which the NO formed from NirK-mediated NO_2^- reduction may react abiotically with medium components to give N_2O . This has recently been reported to be the main N_2O production pathway by ammonia-oxidizing archaea that also lack a known NOR (33). A nitrous oxide reductase (for reduction of N_2O to N_2) has not been identified in the genome, which is consistent with the lack of detectable N_2 production in our pure-culture experiments.

OMZs are a major global source of oceanic N_2O (25 to 59%) (34), a potent greenhouse gas and ozone-depleting agent that has usually been

attributed to denitrifying and ammonia-oxidizing microorganisms (35, 36). In the Namibian OMZ, N_2O production ($\sim 17 \text{ nM day}^{-1}$) was detected particularly in the bottom waters where *Nitrococcus* abundance was maximal and nitrite oxidation was minimal (Fig. 1A), strongly indicating a contribution from *Nitrococcus*. Considering the worldwide distribution and high abundance of *N. mobilis* and *Nitrococcus*-like phylotypes especially within productive coastal ecosystems, *Nitrococcus* might represent a previously overlooked yet significant source of oceanic N_2O . Because OMZs are expanding and intensifying globally (37), *Nitrococcus* may switch from a nitrogen-preserving (NO_2^- -oxidizing) mode to one that facilitates nitrogen loss ($\text{NO}_3^-/\text{NO}_2^-$ reduction to N_2O).

Apart from their importance in nitrogen cycling, *Nitrococcus* may also affect sulfur cycling. The *N. mobilis* genome harbors several key sulfur-metabolizing enzymes, including a sulfide/quinone oxidoreductase (*sqr*) that might oxidize hydrogen sulfide (H_2S) to elemental sulfur or polysulfide (S^0) in the periplasmic space with concomitant reduction of quinone (Fig. 2 and table S4). In addition, a periplasmic sulfite dehydrogenase (*sor*) may oxidize sulfite to sulfate and transfer the electrons onto cytochrome c. Last, a cytoplasmic type 3b bidirectional hydrogenase (38) is present (Fig. 2 and table S4). This so-called sulfhydrogenase can catalyze the reversible oxidation of H_2 with NAD(P)^+ (39) and has been shown to reduce S^0 to H_2S in the hyperthermophilic archaeon *Pyrococcus furiosus* (40).

The ability of *Nitrococcus* to catalyze sulfide oxidation in the presence of O_2 was confirmed with *N. mobilis* pure cultures, which consumed sulfide at a rate of 43 nmol S per microgram of protein per day, whereas the disappearance of sulfide in the abiotic control was negligible even during 5 hours of incubation (Fig. 4B). To our knowledge, this is the first time that capability for sulfur oxidation has ever been reported in association with nitrifying organisms, adding another remarkable facet to the versatile metabolic repertoire already known in NOB.

Among 20 screened marine environmental metagenomes, 50% were found to contain *sqr* genes with high sequence identities to the *N. mobilis* sequence, indicating the prevalent potential of *Nitrococcus* to oxidize reduced S-compounds in the oceans. More than half of the metagenomes from mesopelagic habitats contain *N. mobilis*-like *sqr*, but none do from the deep chlorophyll maxima and epipelagic ocean (fig. S6 and table S7). Hence, sulfide-oxidizing capacity seems to present an ecological advantage for *Nitrococcus* in subsurface waters: This could mean a direct coupling with growth, or facilitation of sulfide detoxification and/or preservation of biologically available sulfur (41, 42).

The prevalence of this functional trait of *Nitrococcus* in the mesopelagic is consistent with the recently reported widespread occurrence of sulfide-oxidizing enzyme complexes in the mesopelagic ocean and within the Global Ocean Sampling (GOS) metagenomic data sets, despite the apparent lack of detectable sulfide (42–44). On the basis of the 138 Tara Oceans data sets that distinguish mesopelagic from surface and deep chlorophyll maximum (DCM) samples, *N. mobilis* and *Nitrococcus*-like phylotypes appeared to be particularly abundant in the mesopelagic ocean (mean, 3.3%, versus 2 and 2.1% for surface and DCM, respectively) (table S2). Particulate organic matter, both sinking and suspended, serves as a major carbon source for heterotrophic communities in the mesopelagic ocean. Although *Nitrococcus* does not necessarily rely on these particles for their carbon demand, active remineralization by other microbes may create oxygen-deficient microniches and facilitate anaerobic processes within marine snow (45–47). Moreover, remineralization processes release inorganic and organic nutrients, including reduced sulfur compounds, proteins, or osmolytes such as




Fig. 4. Sulfide oxidation potential of *N. mobilis* Nb-231. (A) Oxygen consumption in the presence (diamonds; mean of three replicates) and absence (triangles; single measurement) of Na₂S. Only one measurement every 2 min is shown. The arrow indicates the time point of substrate addition. (B) Sulfide consumption by *N. mobilis* Nb-231 (diamonds) and abiotic control (triangles). The curves represent the mean of three biological replicates, and error bars indicate the SD of three technical replicates.

dimethylsulfoniopropionate that may further be metabolized to release sulfide (42, 43). *Nitrococcus*, sometimes found enriched on particles (11), could then use these reduced sulfur compounds to support sulfide oxidation. The actual occurrence and ecological significance, however, remain to be determined in situ.

In coastal OMZs associated with highly productive upwelling areas, such as the Namibian and Peruvian OMZs, sulfide sometimes accumulates to levels that are detrimental to marine life and fisheries. Previous studies have shown that sulfide-oxidizing bacteria affiliated with Gammaproteobacteria and Epsilonproteobacteria participate in detoxifying the harmful sulfide in these oxygen-deficient zones (48, 49). The relatively high abundance of 16S rRNA genes affiliated with *N. mobilis* and *Nitrococcus*-like phylotypes (up to 9.2%; table S2) in Peruvian OMZ metagenomes during a sulfidic event (49) indicates that they are well adapted to such episodic sulfidic conditions and may hint toward a contribution to sulfide detoxification in these waters (48, 49).

NOB are important preservers of biologically available nitrogen in global oceans. Their significance in OMZs, specifically in counteracting nitrogen loss through anammox and denitrification, which both reduce NO₂[−] to N₂, has been increasingly recognized in recent years through direct activity measurements, molecular surveys, and natural stable isotope analyses (11, 12, 50). Thus far, nitrite oxidation is postulated to have prevented oxygen-deficient water bodies from turning into active nitrogen-loss zones (51). This nitrogen conservation role is most certainly becoming more crucial as global ocean deoxygenation continues (37).

Nonetheless, our findings of relatively high abundance of *Nitrococcus* in these waters and of the diverse alternative biogeochemical functions *Nitrococcus* may perform suggest that they may be on the verge of reversing the NOB community's usual role to instead reduce nitrate and produce N₂O, thus promoting nitrogen loss and exacerbating the global greenhouse effect.

In a wider context, the remarkably versatile and sometimes contrasting functional capacities of *Nitrococcus* unveiled in our study—nitrate reduction versus conventional nitrite oxidation, heterotrophy (degradation of organic carbon releasing CO₂) versus autotrophy (CO₂ fixation into organic carbon), along with N₂O production and sulfide oxidation—have conferred on these organisms appreciable adaptability and robustness toward dynamic, changing environmental conditions. Their success is shown by their worldwide distribution across diverse habitats, with particular prominence in dynamic settings such as coastal upwelling, or the vast mesopelagic ocean where resources likely come in pulses. Hence, potential occurrence of any functional switches of *Nitrococcus* could extend beyond oxygen-deficient waters to the wider ocean and may substantially affect the cycling and budgets of nitrogen, carbon, oxygen, and sulfur in global oceans as a result. In the face of rapid global environmental changes, the key regulatory controls and the tipping points of contrasting biogeochemical functions associated with the ubiquitous *Nitrococcus* thus urgently need to be determined.

MATERIALS AND METHODS

Sample collection

Water sampling was conducted on board the R/V Maria S. Merian in October 2011 (MSM19/1c) over the Namibian shelf between 21°59.9'S/13°40.92'E and 28°30'S/15°45.01'E. Salinity, temperature, dissolved oxygen, and chlorophyll *a* fluorescence were measured with a conductivity-temperature-depth (CTD) system, equipped with an oxygen sensor and a fluorometer (Sea-Bird Electronics). Water samples were collected with 12-liter Niskin bottles attached to the CTD system. Oxygen data were calibrated against Winkler titration. The benthic boundary layer (BBL) was sampled at three depths from 30 cm to 2 m above seafloor using a bottom water sampler (52). NO₂[−] and NH₄⁺ were measured on board spectrophotometrically (53) and fluorometrically (54). Water samples were frozen for later NO₃[−] and PO₄^{3−} analyses with an autoanalyzer in a shore-based laboratory (TrAACs 800, Bran & Luebbe). Detection limits for NH₄⁺, NO₂[−], NO₃[−], and PO₄^{3−} were 0.01, 0.01, 0.1, and 0.1 μM, respectively.

Incubation experiments with environmental samples

Incubation experiments were conducted with water collected from two stations in the Namibian OMZ, located at 22°S/13°4'E and 22°6'S/16°6'E. Water samples (250 ml) were collected from five to six depths with the CTD system or the bottom water sampler for samples from the BBL. Different combinations of ¹⁵N and ¹⁴N substrates were added for rate determinations of ammonia and nitrite oxidation as well as nitrate reduction (table S8). Samples were purged with helium for 15 min to eliminate any O₂ contamination from the CTD system (55). For NH₃ and NO₂[−] oxidation, a second set of incubation experiments without helium purging was conducted. Each ¹⁵N(¹⁴N)-amended sample was immediately transferred into five 12-ml Exetainer vials (Labco). These samples were incubated for up to 48 hours in the dark at in situ temperatures. At each time interval (approximately 0, 6, 12, 24, and 48 hours), incubation in one Exetainer was terminated by adding 100 ml of saturated mercuric chloride solution to stop biological activities. Samples were stored upside down in the dark at room temperature until further

processing in a shore-based laboratory [for a more detailed description, see the study of Holtappels *et al.* (56)].

Stable isotopic analyses of N₂

Nitrogen-stable isotopic ratios of N₂ were determined by gas chromatography–isotope ratio mass spectrometry (GC-IRMS; VG Optima). Ammonia oxidation rates and nitrate reduction rates were determined as the ¹⁵NO₂[−] production over time from incubations amended with ¹⁵NH₄⁺/¹⁴NO₂[−] or ¹⁵NO₃[−]/¹⁴NO₂[−] via the conversion of NO₂[−] by sulfamic acid (57). Nitrite oxidation rates were determined in incubations amended with ¹⁵NO₂[−]/¹⁴NO₃[−]. After the removal of residual NO₂[−], NO₃[−] was reduced to NO₂[−] with cadmium and subsequently to N₂ via sulfamic acid, as previously described (11). Process rates were calculated from the slopes of linear regressions with ¹⁵N₂ production as a function of time, and only when the production was instantaneous. All rates were calculated as net rates from five samples obtained over the course of incubation (0 to 48 hours) (*P* < 0.05) and have been corrected for the ¹⁵N-labeling percentages of initial substrate pools.

CARD-FISH

Water samples were fixed with 2% (final concentration) paraformaldehyde (PFA) in phosphate-buffered saline solution for 8 to 12 hours at 4°C, followed by filtration onto polycarbonate membrane filters (GTTP, 0.22 μm pore size, 47 mm diameter; Millipore). CARD-FISH was performed according to the protocol by Pernthaler *et al.* (58). First, cells were immobilized on the GTTP filters by embedding in 0.2% agarose, followed by permeabilization in lysozyme (10 mg/ml) in 50 mM EDTA and 100 mM tris-HCl. Labeled oligonucleotide probes (Biomers) were added and allowed to hybridize for 2 hours at 46°C at varying formamide concentrations, depending on the oligonucleotide applied (table S9). This was followed by tyramide signal amplification for 15 min at 46°C and subsequent staining of cells with DAPI (4',6'-diamidino-2-phenylindole). DAPI-stained cells and positive hybridization signals were enumerated with epifluorescence microscopy (Axioplan 2, Zeiss).

Pure culture experiments

N. mobilis strain Nb-231 was grown in aerobic batch cultures that were moderately stirred for 32 days in 3.5 liters of marine NOB medium (table S6). NO₂[−] was added to a final concentration of 3 mM and resupplied when consumed. The cultures had reached stationary phase when the various incubation experiments were conducted.

For anoxic incubation experiments, cultures were harvested by centrifugation and transferred into 2 liters of anaerobic marine NOB medium, including the various amendments for experiments 1 to 8 as listed in table S10. Subsequently, each experimental step was conducted in an anaerobic chamber (90% N₂/10% CO₂ atmosphere) to avoid oxygen contamination. One hundred fifty milliliters of culture per incubation was transferred into 250-ml serum bottles. All experiments were conducted in triplicate. Control incubations with autoclaved culture were included to test for abiotic reactions (fig. S5B). Through the duration of 28 days, a total of 10 subsamples consisting of 3-ml culture and 2-ml headspace were obtained every 2 to 3 days. The sampling volume was replaced with 90% N₂/10% CO₂ from the anaerobic chamber to avoid underpressure in the incubation vessels. Sulfide-dependent oxygen consumption was measured by using a respiration cell RC-350 (Warner Instruments), equipped with a Clark-type electrode (model 1302) connected to a picoammeter PA2000 (Unisense). Dissolved oxygen concentration was recorded in SensorTrace Basic version 3.0.2 (Unisense), with one measurement every 2 s. Measurements were performed by using

approximately 25× concentrated *N. mobilis* Nb-231 biomass from actively nitrite-oxidizing cultures. Biomass was harvested by centrifugation (4000g for 15 min), washed twice, and resuspended in marine NOB medium. For measurements, the cell chamber was filled with 2 ml of biomass suspension and closed with the electrode inserted into the general electrode holder EH-100 (Warner Instruments) without enclosing air bubbles. Measurements were performed at 28°C. Na₂S was added to an end concentration of 5 mM with a 250-μl glass syringe (Hamilton Gastight, #1725). Biomass without substrate was used as a negative control to measure the electrode drift.

To determine sulfide oxidation in aerobic batch incubation assays, 200 ml of actively nitrite-oxidizing *N. mobilis* Nb-231 culture was harvested and washed, as described above, and resuspended in 100 ml of marine NOB medium. Ten milliliters of nitrite-free culture per incubation was transferred to 30-ml serum bottles, and 250 μM Na₂S was added. Overpressure in the bottles was ensured by adding 5-ml sterile air to each incubation. Incubations were performed in a shaking incubator at 28°C. Cell-free medium was used as a negative control. One milliliter of headspace was sampled every hour. Sulfide concentrations were determined by gas chromatography. One milliliter of approximately 557× diluted headspace samples was injected in triplicate into a gas chromatograph [Agilent 7890B, equipped with a Chrompack ML 848 glass column (2 m × 4 mm) with Carbopack B HT 100 40 to 60 mesh packing, at 80°C; Agilent Technologies] combined with a flame photometric detector (Agilent Technologies).

N₂O and N₂ measurements

The gas samples were injected into sterile water-filled 3-ml Exetainers, and the water was displaced via a compensation needle. The concentration of N₂O in the headspace was measured by a gas chromatograph with a ⁶³Ni electron capture detector (Shimadzu, GC-8A). The linear increase over time was used to calculate N₂O production rates by *Nitrococcus*. Nitrogen-stable isotopic ratios of N₂ (¹⁵N¹⁵N/¹⁴N¹⁴N and ¹⁵N¹⁴N/¹⁴N¹⁴N) were determined by GC-IRMS (VG Optima). Rates were calculated from the slopes of linear regressions with ²⁹N₂/³⁰N₂ and N₂O production as a function of time.

NanoSIMS analyses

During the cruise to the Namibian OMZ, 250-ml water samples were amended with either ¹³C-labeled DOM (¹³C-DOM) or 5 μM ¹⁵NO₂[−] and 200 μM ¹³C-HCO₃[−], alongside incubations for rate measurements. After ~29 hours of incubation, samples were fixed with 1% PFA (final concentration) for 8 to 12 hours at 4°C. Subsequently, 11 ml of the sample was filtered onto gold-palladium precoated polycarbonate filters (GTTP, 0.2 μm pore size, 25 mm diameter; Millipore) and stored at −80°C until further analyses in a shore-based laboratory.

Nitrococcus cells were hybridized on a filter with *Nitrococcus*-specific CARD-FISH probes as described previously (11) and analyzed with a NanoSIMS 50L (Cameca). To obtain a stable ion emission rate and to clean the sample from any contamination, we presputtered the area of interest with a Cs⁺ primary ion beam of 100 pA. The cells were analyzed by rastering a primary Cs⁺ ion beam with a beam current of 0.8 to 1 pA and a beam diameter of <100 nm. Secondary ion images of ¹²C[−], ¹³C[−], ¹⁹F[−], ¹²C¹⁴N[−], and ³²S[−] were recorded in parallel. The analyzed areas ranged from 10 × 10 μm to 20 × 20 μm with an image size of 256 × 256 pixels and a dwell time of 1 ms per pixel. Tuning of the instrument for high mass resolution (~7000 mass resolving power) reduced the interference for ¹³C. Data analyses were performed with the freeware Look@NanoSIMS (59).

Genome annotation of *N. mobilis*

The genome of *N. mobilis* Nb-231 was sequenced by the J. Craig Venter Institute in the framework of the Gordon and Betty Moore Foundation Marine Microbiology Initiative. The sequence was deposited in the GenBank database in 2006 and consists of 43 assembled contigs.

The draft genome of *N. mobilis* Nb-231 (GenBank project PRJNA13475) was integrated into the MicroScope annotation platform (60). After automated prediction and annotation of coding sequences (CDS), the annotation of all CDS in key pathways, including those for nitrite oxidation, respiration, and carbon fixation, was manually refined by using the respective tools of MaGe (60), as described in detail elsewhere (27).

Phylogenetic analyses

16S rRNA sequences classified as *Nitrococcus* were identified within the SILVA small subunit rRNA database release 119 (20). Affiliation with the genus was confirmed by phylogenetic analyses and by selecting for sequence identities $\geq 94.5\%$ (61). The three *N. mobilis* 16S rRNA gene sequences (HM038001, L35510, and AAOF01000010) originate from the *N. mobilis*-type strain and the genome sequence. *Nitrococcus*-like phylotypes were defined on the basis of a 98.7% sequence identity threshold (19, 61). Only sequences ≥ 1300 bases were used for phylogenetic tree reconstruction. Shorter sequences were added to the tree using the parsimony insertion tool in ARB (62). NxrA and heme-copper oxidase (HCO) family member proteins were imported in custom-made databases.

Sequence alignments for all data sets were generated and manually refined using the sequence editor in ARB. Phylogenetic trees were calculated using RAXML version 7.0.4 (63), with 100 (NxrA and HCO) or 1000 (16S rRNA) bootstraps. To test for tree stability, other treeing methods (neighbor joining, maximum parsimony, and Bayesian interference) were used but yielded almost identical branching patterns. For 16S rRNA analyses, the GTRGAMMA substitution model and a 50% conservation filter resulting in 1515 valid alignment positions were used. NxrA and HCO trees were calculated using the PROTMIX rate distribution and the WAG substitution model, with a 10% conservation filter for NxrA and without a conservation filter for HCO, resulting in 1198 and 1367 distinct alignment positions, respectively.

Metagenomic data analyses

In total, 253 metagenomic and 422 amplicon data sets of marine origin were downloaded from MG-RAST (21), 150 metagenomic and 155 amplicon data sets were obtained from OSD 2014, as detailed in <https://github.com/MicroB3-IS/osd-analysis/wiki/Guide-to-OSD-2014-data>, and 138 samples of 16S rDNA miTags Illumina sequence data were obtained from <http://ocean-microbiome.embl.de/companion.html> as fasta files from the Tara Oceans project. Metadata for the data sets are provided in table S1. All sequence reads were processed with the next-generation sequencing analysis pipeline of the SILVA project (available at www.arb-silva.de/ngs). A cutoff of (BLASTN percent query coverage + BLASTN percent alignment identity)/2 > 97% was used to assign *Nitrococcus*-like phylotypes to a read. Of the 1118 data sets, 772 were found to contain at least one *N. mobilis* or *Nitrococcus*-like phylotype (tables S1 and S2). Of these, only metagenomic data sets (Fig. 1B) were used to create a world map showing the global distribution and relative abundance of *N. mobilis* or *Nitrococcus*-like phylotypes.

Selected OSD metagenomes, Tara, and MG-RAST samples were mapped to the *N. mobilis* Nb-231 genome (accession no. [gij211606481](https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/13475)|[ref|NZ_CH672427.1](https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/13475)) using FR-HIT version 0.7 with the following parameters: -e 0.001 -u 0 -p 8 -c 80 -m (dynamically set to 40% of aver-

age read length in a sample). Hits for nitrogen and sulfur cycling genes of interest were considered significant when the e value was $<10^{-5}$, and percent sequence identity was $>80\%$. The results were evaluated on the basis of presence/absence of genes, rather than quantitatively, as the *Nitrococcus* species had rather low relative abundance in most data sets.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/3/11/e1700807/DC1>

Supplementary Text

fig. S1. Abundance of *Nitrococcus*-affiliated cells in the Namibian OMZ based on CARD-FISH counts.

fig. S2. 16S rRNA gene-based phylogenetic tree visualizing the relation of four *Nitrococcus* phylotypes.

fig. S3. Phylogenetic analysis of NxrA.

fig. S4. Incubation experiments with *N. mobilis* Nb-231.

fig. S5. Phylogenetic analysis of the sNOR.

fig. S6. Selection of Tara Oceans, MG-RAST, and OSD samples that were mapped to the *N. mobilis* Nb-231 genome.

fig. S7. Effect of reduced O_2 and enhanced IO_3^- concentrations on nitrite oxidation rates.

table S1. Percent identity of 16S rRNA genes between *N. mobilis* and the newly identified *Nitrococcus*-like phylotype 1.

table S2. List of marine environmental metagenomes that contain at least one of the four *Nitrococcus* phylotypes.

table S3. List of marine amplicon sequencing data sets that contain at least one of the four *Nitrococcus* phylotypes.

table S4. *N. mobilis* strain Nb-231 proteins with predicted functions in key metabolic pathways.

table S5. ^{13}C enrichment of single *Nitrococcus* cells from the Namibian OMZ.

table S6. Composition of marine NOB medium used to grow *N. mobilis*.

table S7. Functional genes associated with *Nitrococcus* spp. in selected Tara Oceans metagenomes.

table S8. Summary of stations, sampling depths, and ^{15}N incubation experiments conducted.

table S9. NOB specific 16S rRNA-targeted oligonucleotide probes.

table S10. Substrate amendments for *N. mobilis* Nb-231 incubation experiments.

References (64–76)

REFERENCES AND NOTES

1. A. Yool, A. P. Martin, C. Fernández, D. R. Clark, The significance of nitrification for oceanic new production. *Nature* **447**, 999–1002 (2007).
2. E. Spieck, E. Bock, The lithoautotrophic nitrite-oxidizing bacteria, in *Bergey's Manual of Systematic Bacteriology*, G. Garrity, D. J. Brenner, N. R. Krieg, J. R. Staley, Eds. (Springer-Verlag, ed. 2, 2005), pp. 149–153.
3. H. Daims, S. Lüscher, M. Wagner, A new perspective on microbes formerly known as nitrite-oxidizing bacteria. *Trends Microbiol.* **24**, 699–712 (2016).
4. H. Koch, A. Galushko, M. Albertsen, A. Schintlmeister, C. Gruber-Döringer, S. Lüscher, E. Pelletier, D. Le Paslier, E. Spieck, A. Richter, P. H. Nielsen, M. Wagner, H. Daims, Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science* **345**, 1052–1054 (2014).
5. H. Daims, E. V. Lebedeva, P. Pjevac, P. Han, C. Herbold, M. Albertsen, N. Jehmlich, M. Palatinszky, J. Vierheilig, A. Bulaev, R. H. Kirkegaard, M. von Bergen, T. Rattei, B. Bendinger, P. H. Nielsen, M. Wagner, Complete nitrification by *Nitrospira* bacteria. *Nature* **528**, 504–509 (2015).
6. H. Koch, S. Lüscher, M. Albertsen, K. Kitzinger, C. Herbold, E. Spieck, P. H. Nielsen, M. Wagner, H. Daims, Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*. *Proc. Natl. Acad. Sci.* **112**, 11371–11376 (2015).
7. M. A. H. J. van Kessel, D. R. Speth, M. Albertsen, P. H. Nielsen, H. J. M. Op den Camp, B. Kartal, M. S. M. Jetten, S. Lüscher, Complete nitrification by a single microorganism. *Nature* **528**, 555–559 (2015).
8. T. J. Mincer, M. J. Church, L. T. Taylor, C. Preston, D. M. Karl, E. F. DeLong, Quantitative distribution of presumptive archaeal and bacterial nitrifiers in Monterey Bay and the North Pacific Subtropical Gyre. *Environ. Microbiol.* **9**, 1162–1175 (2007).
9. R. E. Davis, D. S. Stakes, C. G. Wheat, C. L. Moyer, Bacterial variability within an iron-silica-manganese-rich hydrothermal mound located off-axis at the Cleft Segment, Juan de Fuca Ridge. *Geomicrobiol. J.* **26**, 570–580 (2009).

10. A. M. Durbin, A. Teske, Microbial diversity and stratification of South Pacific abyssal marine sediments. *Environ. Microbiol.* **13**, 3219–3234 (2011).
11. J. Füssel, P. Lam, G. Lavik, M. M. Jensen, M. Holtappels, M. Günter, M. M. M. Kuypers, Nitrite oxidation in the Namibian oxygen minimum zone. *ISME J.* **6**, 1200–1209 (2012).
12. J. M. Beman, J. Leilei Shih, B. N. Popp, Nitrite oxidation in the upper water column and oxygen minimum zone of the eastern tropical North Pacific Ocean. *ISME J.* **7**, 2192–2205 (2013).
13. S. W. Watson, J. B. Waterbury, Characteristics of two marine nitrite oxidizing bacteria, *Nitrospina gracilis* nov. gen. nov. sp. and *Nitrococcus mobilis* nov. gen. nov. sp. *Arch. Microbiol.* **77**, 203–230 (1971).
14. S. Lüscher, B. Nowka, T. Rattei, E. Spieck, H. Daims, The genome of *Nitrospina gracilis* illuminates the metabolism and evolution of the major marine nitrite oxidizer. *Front. Microbiol.* **4**, 27 (2013).
15. E. Spieck, S. Keuter, T. Wenzel, E. Bock, W. Ludwig, Characterization of a new marine nitrite oxidizing bacterium, *Nitrospina watsonii* sp. nov., a member of the newly proposed phylum “*Nitrospinaceae*”. *Syst. Appl. Microbiol.* **37**, 170–176 (2014).
16. B. B. Ward, A. F. Carlucci, Marine ammonia- and nitrite-oxidizing bacteria: Serological diversity determined by immunofluorescence in culture and in the environment. *Appl. Environ. Microbiol.* **50**, 194–201 (1985).
17. B. B. Ward, H. E. Glover, F. Lipschultz, Chemoautotrophic activity and nitrification in the oxygen minimum zone off Peru. *Deep-Sea Res.* **36**, 1031–1051 (1989).
18. R. Ghai, L. Pašić, A. B. Fernández, A.-B. Martín-Cuadrado, C. M. Mizuno, K. D. McMahon, R. T. Papke, R. Stepanauskas, B. Rodríguez-Brito, F. Rohwer, C. Sánchez-Porro, A. Ventosa, F. Rodríguez-Valera, New abundant microbial groups in aquatic hypersaline environments. *Sci. Rep.* **1**, 135 (2011).
19. E. Stackebrandt, J. Ebers, Taxonomic parameters revisited: Tarnished gold standards. *Microbiol. Today* **33**, 152–155 (2006).
20. C. Quast, E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, F. O. Glöckner, The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2013).
21. F. Meyer, D. Paarmann, M. D’Souza, R. Olson, E. M. Glass, M. Kubal, T. Paczian, A. Rodriguez, R. Stevens, A. Wilke, J. Wilkening, R. A. Edwards, The metagenomics RAST server—A public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**, 386 (2008).
22. A. Kopf, B. Bick, R. Kottmann, J. Schnetzer, I. Kostadinov, K. Lehmann, A. Fernandez-Guerra, C. Jeanthon, E. Rahav, M. Ullrich, A. Wichels, G. Gerdt, P. Polymenakou, G. Kotoulas, R. Siam, R. Z. Abdallah, E. C. Sonnenschein, T. Cariou, F. O’Gara, S. Jackson, S. Orlic, M. Steinke, J. Busch, B. Duarte, I. Caçador, J. Canning-Clode, O. Bobrova, V. Marteinsson, E. Reynisson, C. M. Loureiro, G. M. Luna, G. M. Quero, C. R. Löscher, A. Kremp, M. E. DeLorenzo, L. Øvreås, J. Tolman, J. LaRoche, A. Penna, M. Frischer, T. Davis, B. Katherine, C. P. Meyer, S. Ramos, C. Magalhães, F. Jude-Lemellieur, M. L. Aguirre-Macedo, S. Wang, N. Poulton, S. Jones, R. Collin, J. A. Fuhrman, P. Conan, C. Alonso, N. Stambler, K. Goodwin, M. M. Yakimov, F. Baltar, L. Bodrossy, J. Van De Kamp, D. M. Frampton, M. Ostrowski, P. Van Ruth, P. Malthouse, S. Claus, K. Deneudt, J. Mortelmans, S. Pitois, D. Wallom, I. Salter, R. Costa, D. C. Schroeder, M. M. Kandil, V. Amaral, F. Biancalana, R. Santana, M. L. Pedrotti, T. Yoshida, H. Ogata, T. Ingletton, K. Munnik, N. Rodriguez-Ezpeleta, V. Berteaux-Lecellier, P. Wecker, I. Cancio, D. Vault, C. Bienhold, H. Ghazal, B. Chaouni, S. Essayeh, S. Ettamimi, H. Zaid el, N. Boukhatem, A. Bouali, R. Chahboune, S. Barrijal, M. Timouni, F. El Otmani, M. Bennani, M. Mea, N. Todorova, V. Karamfilov, P. Ten Hoopen, G. Cochrane, S. L’Haridon, K. C. Bizsel, A. Vezzi, F. M. Lauro, P. Martin, R. M. Jensen, J. Hinks, S. Gebbels, R. Rosselli, F. De Pascale, R. Schiavon, A. Dos Santos, E. Villar, S. Pesant, B. Cataletto, F. Malfatti, R. Edirisinghe, J. A. Silveira, M. Barbier, V. Turk, T. Tinta, W. J. Fuller, I. Salihoglu, N. Serakini, M. C. Ergoren, E. Bresnan, J. Iriberry, P. A. Nyhus, E. Bente, H. E. Karlsen, P. N. Golyshin, J. M. Gasol, S. Moncheva, N. Dzhenbekova, Z. Johnson, C. D. Sinigalliano, M. L. Gidley, A. Zingone, R. Danovaro, G. Tsiamis, M. S. Clark, A. C. Costa, M. El Bour, A. M. Martins, R. E. Collins, A. L. Ducluzeau, J. Martinez, M. J. Costello, L. A. Amaral-Zettler, J. A. Gilbert, N. Davies, D. Field, F. O. Glöckner, The ocean sampling day consortium. *GigaScience* **4**, 1–5 (2015).
23. S. Sunagawa, L. P. Coelho, S. Chaffron, J. R. Kultima, K. Labadie, G. Salazar, B. Djahanschiri, G. Zeller, D. R. Mende, A. Alberti, F. M. Cornejo-Castillo, P. I. Costea, C. Cruaud, F. d’Ovidio, S. Engelen, I. Ferrera, J. M. Gasol, L. Guidi, F. Hildebrand, F. Kokoszka, C. Lepoivre, G. Lima-Mendez, J. Poulain, B. T. Poulos, M. Royo-Llonch, H. Sarmento, S. Vieira-Silva, C. Dimier, M. Picheral, S. Searson, S. Kandsley-Lewis, Tara Oceans coordinators, C. Bowler, C. de Vargas, G. Gorsky, N. Grimsley, P. Hingamp, D. Iudicone, O. Jaillon, F. Not, H. Ogata, S. Pesant, S. Speich, L. Stemmann, M. B. Sullivan, J. Weissenbach, P. Wincker, E. Karsenti, J. Raes, S. G. Acinas, P. Bork, Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).
24. A. Freitag, M. Rudert, E. Bock, Growth of *Nitrobacter* by dissimilatory nitrate reduction. *FEMS Microbiol. Lett.* **48**, 105–109 (1987).
25. E. Bock, H.-P. Koops, U. C. Möller, M. Rudert, A new facultatively nitrite oxidizing bacterium, *Nitrobacter vulgaris* sp. nov. *Arch. Microbiol.* **153**, 105–110 (1990).
26. H. Sundermeyer-Klinger, W. Meyer, B. Warninghoff, E. Bock, Membrane-bound nitrite oxidoreductase of *Nitrobacter*: Evidence for a nitrate reductase system. *Arch. Microbiol.* **140**, 153–158 (1984).
27. S. Lüscher, M. Wagner, F. Maixner, E. Pelletier, H. Koch, B. Vacherie, T. Rattei, J. S. S. Damsté, E. Spieck, D. Le Paslier, H. Daims, A *Nitrospira* metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing bacteria. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 13479–13484 (2010).
28. S. R. Starkenburg, D. J. Arp, P. J. Bottomley, Expression of a putative nitrite reductase and the reversible inhibition of nitrite-dependent respiration by nitric oxide in *Nitrobacter winogradskyi* Nb-255. *Environ. Microbiol.* **10**, 3036–3042 (2008).
29. D. Y. Sorokin, S. Lüscher, D. Vejmekova, N. A. Kostrikina, R. Kleerebezem, W. I. C. Rijpstra, J. S. S. Damsté, D. Le Paslier, G. Muyzer, M. Wagner, M. C. M. van Loosdrecht, H. Daims, Nitrification expanded: Discovery, physiology and genomics of a nitrite-oxidizing bacterium from the phylum *Chloroflexi*. *ISME J.* **6**, 2245–2256 (2012).
30. N. Musat, H. Halm, B. Winterholler, P. Hoppe, S. Peduzzi, F. Hillion, F. Horreard, R. Amann, B. B. Jørgensen, M. M. M. Kuypers, A single-cell view on the ecophysiology of anaerobic phototrophic bacteria. *Proc. Natl. Acad. Sci.* **105**, 17861–17866 (2008).
31. M. Ackermann, A functional perspective on phenotypic heterogeneity in microorganisms. *Nat. Rev. Microbiol.* **13**, 497–508 (2015).
32. L. Y. Stein, D. J. Arp, P. M. Berube, P. S. G. Chain, L. Hauser, M. S. M. Jetten, M. G. Klotz, F. W. Larimer, J. M. Norton, H. J. M. Op den Camp, M. Shin, X. Wei, Whole-genome analysis of the ammonia-oxidizing bacterium, *Nitrosomonas europaea* C91: Implications for niche adaptation. *Environ. Microbiol.* **9**, 2993–3007 (2007).
33. J. A. Kozlowski, M. Stieglmeier, C. Schleper, M. G. Klotz, L. Y. Stein, Pathways and key intermediates required for obligate aerobic ammonia-dependent chemolithotrophy in bacteria and Thaumarchaeota. *ISME J.* **10**, 1836–1845 (2016).
34. P. Suntharalingam, J. Sarmiento, J. R. Toggweiler, Global significance of nitrous-oxide production and transport from oceanic low-oxygen zones: A modeling study. *Global Biogeochem. Cycles* **14**, 1353–1370 (2000).
35. T. J. Goreau, W. A. Kaplan, S. C. Wofsy, M. B. McElroy, F. W. Valois, S. W. Watson, Production of NO₂[−] and N₂O by nitrifying bacteria at reduced concentrations of oxygen. *Appl. Environ. Microbiol.* **40**, 526–532 (1980).
36. A. R. Ravishankara, J. S. Daniel, R. W. Portmann, Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science* **326**, 123–125 (2009).
37. L. Stramma, G. C. Johnson, J. Sprintall, V. Mohrholz, Expanding oxygen-minimum zones in the tropical oceans. *Science* **320**, 655–658 (2008).
38. P. M. Vignais, B. Billoud, Occurrence, classification, and biological function of hydrogenases: An overview. *Chem. Rev.* **107**, 4206–4272 (2007).
39. P. J. Silva, E. C. D. van den Ban, H. Wassink, H. Haaker, B. de Castro, F. T. Robb, W. R. Hagen, Enzymes of hydrogen metabolism in *Pyrococcus furiosus*. *Eur. J. Biochem.* **267**, 6541–6551 (2000).
40. K. Ma, R. Weiss, M. W. W. Adams, Characterization of hydrogenase II from the hyperthermophilic archaeon *Pyrococcus furiosus* and assessment of its role in sulfur reduction. *J. Bacteriol.* **182**, 1864–1871 (2000).
41. J. L. Luebke, J. Shen, K. E. Bruce, T. E. Kehl-Fie, H. Peng, E. P. Skaar, D. P. Giedroc, The CsoR-like sulfurtransferase repressor (CstR) is a persulfide sensor in *Staphylococcus aureus*. *Mol. Microbiol.* **94**, 1343–1360 (2014).
42. Y. Xia, C. Lü, N. Hou, Y. Xin, J. Liu, H. Liu, L. Xun, Sulfide production and oxidation by heterotrophic bacteria under aerobic conditions. *ISME J.* 10.1038/ismej.2017.125 (2017).
43. B. K. Swan, M. Martínez-García, C. M. Preston, A. Sczyrba, T. Woyke, D. Lamy, T. Reinthaler, N. J. Poulton, E. D. P. Masland, M. L. Gomez, M. E. Sieracki, E. F. DeLong, G. J. Herndl, R. Stepanauskas, Potential for chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean. *Science* **333**, 1296–1300 (2011).
44. A. A. Murillo, S. Ramírez-Flandes, E. F. DeLong, O. Ulloa, Enhanced metabolic versatility of planktonic sulfur-oxidizing γ-proteobacteria in an oxygen-deficient coastal ecosystem. *Front. Mar. Sci.* **1**, 18 (2014).
45. D. M. Karl, G. A. Knauer, J. H. Martin, B. B. Ward, Bacterial chemolithotrophy in the ocean is associated with sinking particles. *Nature* **309**, 54–56 (1984).
46. A. L. Alldredge, Y. Cohen, Can microscale chemical patches persist in the sea? Microelectrode study of marine snow, fecal pellets. *Science* **235**, 689–691 (1987).
47. P. Stief, A. Kamp, B. Thamdrup, R. N. Glud, Anaerobic nitrogen turnover by sinking diatom aggregates at varying ambient oxygen levels. *Front. Microbiol.* **7**, 98 (2016).
48. G. Lavik, T. Stührmann, V. Brüchert, A. Van der Plas, V. Mohrholz, P. Lam, M. Mußmann, B. M. Fuchs, R. Amann, U. Lass, M. M. M. Kuypers, Detoxification of sulphidic African shelf waters by blooming chemolithotrophs. *Nature* **457**, 581–584 (2009).
49. H. Schunck, G. Lavik, D. K. Desai, T. Großkopf, T. Kalvelage, C. R. Löscher, A. Paulmier, S. Contreras, H. Siegel, M. Holtappels, P. Rosenstiel, M. B. Schilhabel, M. Graco, R. A. Schmitz, M. M. M. Kuypers, J. LaRoche, Giant hydrogen sulfide plume in the oxygen minimum zone off Peru supports chemolithoautotrophy. *PLOS ONE* **8**, e68661 (2013).
50. C. Buchwald, A. E. Santoro, R. H. Stanley, K. L. Casciotti, Nitrogen cycling in the secondary nitrite maximum of the eastern tropical North Pacific off Costa Rica. *Global Biogeochem. Cycles* **29**, 2061–2081 (2015).

51. L. A. Bristow, C. M. Callbeck, M. Larsen, M. A. Altabet, J. Dekaezemacker, M. Forth, M. Gauns, R. N. Glud, M. M. M. Kuypers, G. Lavik, J. Milucka, S. W. A. Naqvi, A. Pratihary, N. P. Revsbech, B. Thamdrup, A. H. Treusch, D. E. Canfield, N₂ production rates limited by nitrite availability in the Bay of Bengal oxygen minimum zone. *Nat. Geosci.* **10**, 24–29 (2017).
52. M. Holtappels, M. M. M. Kuypers, M. Schlüter, V. Bruchert, Measurement and interpretation of solute concentration gradients in the benthic boundary layer. *Limnol. Oceanogr. Meth.* **9**, 1–13 (2011).
53. K. Grasshoff, M. Ehrhardt, K. Kremling, *Methods of Seawater Analysis* (Wiley-VCH, Weinheim, 1999).
54. R. M. Holmes, A. Aminot, R. Kérouel, B. A. Hooker, B. J. Peterson, A simple and precise method for measuring ammonium in marine and freshwater ecosystems. *Can. J. Fish. Aquat. Sci.* **56**, 1801–1808 (1999).
55. T. Dalsgaard, D. E. Canfield, J. Petersen, B. Thamdrup, J. Acuña-González, N₂ production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica. *Nature* **422**, 606–608 (2003).
56. M. Holtappels, G. Lavik, M. M. Jensen, M. M. M. Kuypers, ¹⁵N-labelling experiments to dissect the contributions of heterotrophic denitrification and anammox to nitrogen removal in the OMZ waters of the ocean, in *Methods in Enzymology: Research on Nitrification and Related Processes, Part A*, M. Klotz, Ed. (Academic Press, ed. 1, 2011), vol. 486, pp. 223–251.
57. J. Granger, D. M. Sigman, Removal of nitrite with sulfamic acid for nitrate N and O isotope analysis with the denitrifier method. *Rapid Commun. Mass Spectrom.* **23**, 3753–3762 (2009).
58. A. Pernthaler, J. Pernthaler, R. Amann, Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. *Appl. Environ. Microbiol.* **68**, 3094–3101 (2002).
59. L. Polerecky, B. Adam, J. Milucka, N. Musat, T. Vagner, M. M. M. Kuypers, Look@NanoSIMS—A tool for the analysis of nanoSIMS data in environmental microbiology. *Environ. Microbiol.* **14**, 1009–1023 (2012).
60. D. Vallenet, L. Labarre, Z. Rouy, V. Barbe, S. Bocs, S. Cruveiller, A. Lajus, G. Pascal, C. Scarpelli, C. Médigue, MaGe: A microbial genome annotation system supported by synteny results. *Nucleic Acids Res.* **34**, 53–65 (2006).
61. P. Yaza, P. Yilmaz, E. Pruesse, F. O. Glöckner, W. Ludwig, K.-H. Schleifer, W. B. Whitman, J. Euzéby, R. Amann, R. Rosselló-Móra, Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat. Rev. Microbiol.* **12**, 635–645 (2014).
62. W. Ludwig, O. Strunk, R. Westram, L. Richter, H. Meier, Yadhukumar, A. Buchner, T. Lai, S. Steppi, G. Jobb, W. Förster, I. Brettske, S. Gerber, A. W. Ginhart, O. Gross, S. Grumann, S. Hermann, R. Jost, A. König, T. Liss, R. Lüßmann, M. May, B. Nonhoff, B. Reichel, R. Strehlow, A. Stamatakis, N. Stuckmann, A. Vilbig, M. Lenke, T. Ludwig, A. Bode, K.-H. Schleifer, ARB: A software environment for sequence data. *Nucleic Acids Res.* **32**, 1363–1371 (2004).
63. A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
64. B. Thamdrup, H. Fossing, B. B. Jørgensen, Manganese, iron and sulfur cycling in a coastal marine sediment, Aarhus bay, Denmark. *Geochim. Cosmochim. Acta* **58**, 5115–5129 (1994).
65. H. Lees, J. R. Simpson, The biochemistry of the nitrifying organisms. 5. Nitrite oxidation by *Nitrobacter*. *Biochem. J.* **65**, 297–305 (1957).
66. A. M. Farrenkopf, M. E. Dollhopf, S. Ni Chadhain, G. W. Luther III, K. H. Nealson, Reduction of iodate in seawater during Arabian Sea shipboard incubations and in laboratory cultures of the marine bacterium *Shewanella putrefaciens* strain MR-4. *Mar. Chem.* **57**, 347–354 (1997).
67. A. M. Farrenkopf, G. W. Luther III, V. W. Truesdale, C. H. Van der Weijden, Sub-surface iodide maxima: Evidence for biologically catalyzed redox cycling in Arabian Sea OMZ during the SW intermonsoon. *Deep-Sea Res. Pt. II* **44**, 1391–1409 (1997).
68. V. W. Truesdale, G. W. Bailey, Iodine distribution in the Southern Benguela system during an upwelling episode. *Cont. Shelf Res.* **22**, 39–49 (2002).
69. Y.-C. Sung, D. Parsell, P. M. Anderson, J. A. Fuchs, Identification, mapping, and cloning of the gene encoding cyanase in *Escherichia coli* K-12. *J. Bacteriol.* **169**, 2639–2642 (1987).
70. M. Meincke, E. Bock, D. Kastrau, P. M. H. Kroneck, Nitrite oxidoreductase from *Nitrobacter hamburgensis*: Redox centers and their catalytic role. *Arch. Microbiol.* **158**, 127–131 (1992).
71. R. A. Rothery, G. J. Workun, J. H. Weiner, The prokaryotic complex iron–sulfur molybdoenzyme family. *Biochim. Biophys. Acta* **1778**, 1897–1929 (2008).
72. S. Grimaldi, B. Schoep-Cothenet, P. Ceccaldi, B. Guigliarelli, A. Magalon, The prokaryotic Mo/W-bisPGD enzymes family: A catalytic workhorse in bioenergetic. *Biochim. Biophys. Acta* **1827**, 1048–1085 (2013).
73. F. Blasco, J.-P. Dos Santos, A. Magalon, C. Frixon, B. Guigliarelli, C.-L. Santini, G. Giordano, NarJ is a specific chaperone required for molybdenum cofactor assembly in nitrate reductase A of *Escherichia coli*. *Mol. Microbiol.* **28**, 435–447 (1998).
74. F. R. Tabita, T. E. Hanson, H. Li, S. Satagopan, J. Singh, S. Chan, Function, structure, and evolution of the RubisCO-like proteins and their RubisCO homologs. *Microbiol. Mol. Biol. Rev.* **71**, 576–599 (2007).
75. T. O. Yeates, C. A. Kerfeld, S. Heinhorst, G. C. Cannon, J. M. Shively, Protein-based organelles in bacteria: Carboxysomes and related microcompartments. *Nat. Rev. Microbiol.* **6**, 681–691 (2008).
76. G. Bowes, W. L. Ogren, R. H. Hageman, Phosphoglycolate production catalyzed by ribulose diphosphate carboxylase. *Biochem. Biophys. Res. Commun.* **45**, 716–722 (1971).

Acknowledgments: We thank M. Zabel (chief scientist; Marum Bremen), the crew of the R/V Maria S. Merian (MSM19/1c), K. Emeis (cruise leader; University of Hamburg), and the crew of the R/V Meteor (M76/2) for their support at sea. We also thank G. Klockgether, D. Tienken, D. Nini, A. Ellrott, and V. Meyer (Max Planck Institute for Marine Microbiology) for conscientious technical and analytical assistance. LABGeM and the national infrastructure “France Genomique” are acknowledged for providing access to the MaGe annotation system.

Funding: This work was supported by the Max-Planck-Gesellschaft, the Deutsche Forschungsgemeinschaft, Bundesministerium für Bildung und Forschung (for cruises M76/2 and MSM 19/1c), the Vienna Science and Technology Fund (grant LS09-40), the Austrian Science Fund (grant P25231-B21), the Netherlands Organization for Scientific Research (VENI grant 863.14.019), and the U.K. Natural Environment Research Council (SONIC grant NE/N003187/1). **Author contributions:** J.F., S.L., P.Y., B.N., H.D., E.S., M.M.M.K., and P.L. designed the study. J.F. and P.F.H. collected environmental samples. J.F., S.L., B.N., M.A.H.J.v.K., J.B., and P.B. conducted pure culture experiments. J.F. was responsible for sample analyses. S.L. performed NanoSIMS analyses. S.L. annotated and analyzed the *N. mobilis* Nb-231 genome. P.Y. was responsible for analyses of environmental amplicon and metagenome data. J.F., S.L., P.Y., and P.L. wrote the manuscript. All authors revised the manuscript. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Submitted 10 March 2017

Accepted 13 October 2017

Published 1 November 2017

10.1126/sciadv.1700807

Citation: J. Füssel, S. Lucker, P. Yilmaz, B. Nowka, M. A. H. J. van Kessel, P. Bourceau, P. F. Hach, S. Littmann, J. Berg, E. Spieck, H. Daims, M. M. M. Kuypers, P. Lam, Adaptability as the key to success for the ubiquitous marine nitrite oxidizer *Nitrococcus*. *Sci. Adv.* **3**, e1700807 (2017).

Adaptability as the key to success for the ubiquitous marine nitrite oxidizer *Nitrococcus*

Jessika Füssel, Sebastian Lückner, Pelin Yilmaz, Boris Nowka, Maartje A. H. J. van Kessel, Patric Bourceau, Philipp F. Hach, Sten Littmann, Jasmine Berg, Eva Spieck, Holger Daims, Marcel M. M. Kuypers and Phyllis Lam

Sci Adv **3** (11), e1700807.
DOI: 10.1126/sciadv.1700807

ARTICLE TOOLS

<http://advances.sciencemag.org/content/3/11/e1700807>

SUPPLEMENTARY MATERIALS

<http://advances.sciencemag.org/content/suppl/2017/10/30/3.11.e1700807.DC1>

REFERENCES

This article cites 72 articles, 16 of which you can access for free
<http://advances.sciencemag.org/content/3/11/e1700807#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science Advances (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science Advances* is a registered trademark of AAAS.