

The impact of temperature rise on viruses of the picoplankton community in the Adriatic Sea

Ordulj Marin¹, Šolić Mladen², Šantić Danijela², Šestanović Stefanija², Bojanić Natalija², Jozić Slaven² and Vrdoljak Ana²

¹ Department of Marine Studies, University of Split, Ruđera Boškovića 37, 21000 Split, Croatia
E-mail: marin_ordulj@unist.hr

² Institute of Oceanography and Fisheries, Šetalište Ivana Meštrovića 63, 21000 Split, Croatia

Viruses are an important and dynamic component of the marine microbial community. They are shaping microbial populations by lysing the host cells and thus are changing the biogeochemical nutrient cycling. The rise of sea surface temperature caused by the global warming could have a significant effect on viral population and their interaction with the marine microbial community. This could affect the way carbon is exported through the microbial food network. Modified laboratory dilution experiments that allow estimates of viral lysis on marine picoplankton (bacteria, *Prochlorococcus*, *Synechococcus*, picoeukaryotes) under *in situ* and 3°C above *in situ* temperatures were performed at seasonal scale. Viral lysis removed 14-94% day⁻¹ of bacterial standing stock, 3-135% day⁻¹ of *Prochlorococcus* standing stock, 0.9-30% day⁻¹ of *Synechococcus* standing stock, and 0.2-17% day⁻¹ of picoeukaryote standing stock. Rates of viral lysis on picoplankton were more pronounced in the warmer months (July and November), particularly for the autotrophic picoplankton groups (especially *Synechococcus* and picoeukaryotes) that were significant hosts for viral infection. However, the largest part of the total picoplankton biomass loss through the viral lysis was related to bacteria (48-98%). A rise in temperature of 3°C increased the both, picoplankton biomass production and the amount of lysed picoplankton biomass, especially autotrophic hosts. These results suggest that the global warming could increase the virus-mediated flux of picoplankton carbon toward the pool of dissolved organic matter.

Keywords: marine viruses; viral lysis; picoplankton; global warming