



ELSEVIER

Aquaculture 191 (2000) 209–245

Aquaculture

www.elsevier.nl/locate/aqua-online

Osmoregulatory capacity as a tool in monitoring the physiological condition and the effect of stress in crustaceans

J.-H. Lignot, C. Spanings-Pierrot, G. Charmantier *

*Laboratoire d'Ecophysiologie des Invertébrés, Université des Sciences, Montpellier II, Pl. E. Bataillon,
34095 Montpellier cedex 05, France*

Received 1 March 2000; accepted 9 May 2000

Abstract

Exposure to water-borne pollutants, environmental stressors and pathological agents of a crustacean capable of osmoregulation usually results in a decrease of its Na^+ and Cl^- regulation and/or of its osmoregulatory capacity (OC: difference between the osmotic pressures of the hemolymph and of the external medium, at a given salinity). The partial or complete loss of osmoregulatory and ionoregulatory capacity is generally linked to disruptions of the osmotic and ionic regulations. Different causes of these variations include alterations in the structure and ultrastructure of the branchial and excretory organs and changes in Na^+ , K^+ -ATPase activity, ionic fluxes and surface permeability.

OC appears therefore as a nonspecific bioindicator. Hyper-OC and/or hypo-OC can be measured according to the type of osmoregulation of the species, respectively, in low-salinity media and in seawater or high-salinity media. Provided several precautions are taken (control of salinity and temperature, size, nutritional status, developmental stage, molt stage, use of controls), measurement of OC variations is proposed as a convenient and reliable way to monitor the physiological condition and the effect of stressors in crustaceans. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: Crustaceans; Osmoregulation; Osmoregulatory capacity; Pollutants; Stressors

* Corresponding author. Tel.: +33-4-6714-3672; fax: +33-4-6714-3031.

E-mail address: charmantier@crit.univ-montp2.fr (G. Charmantier).

1. Introduction

Rapid methods for the evaluation and understanding of stressors (pollutants, physical disruptions of the environment, pathological agents) on marine organisms are increasingly needed so that effective management decisions can be made and further environmental degradation averted (Malins and Ostrander, 1991; Holdway et al., 1995). Responses to stressors have traditionally been observed at the population, community or ecosystem levels of organization. Effects, however, are also manifested at the organismal level by impairing molecular, cellular and physiological functions (with reviews in Vernberg and Vernberg, 1974; Gilles and Péqueux, 1983; Hebel et al., 1997). Different histological, physiological, biochemical and molecular measures of toxicant-induced stress — referred to as biomarkers — have therefore been suggested and used as sensitive predictors of effects of stressors at the level of response of whole animals (with reviews in Haya and Waiwood, 1983; Giesy et al., 1988; Giesy and Graney, 1989; Mayer et al., 1992; Livingstone, 1993; Cajaraville et al., 1995; Holdway et al., 1995; Walker, 1995; Wedderburn et al., 1998). Osmoregulation maintaining osmotic homeostasis has already been used as a biomarker in fish species (Clarke and Blackburn, 1977; Eddy, 1981; Wendemeyer and McLay, 1981; Wendelaar Bonga and Lock, 1992) and can be considered as another possible marker for crustaceans (Charmantier et al., 1989; Lin et al., 1991, 1992; Young-Lai et al., 1991; Mayer et al., 1992; Bambang et al., 1995a,b; Lignot et al., 1997, 1998a). Osmoregulation, which is one of the most important regulatory functions an aquatic animal has to perform, has been extensively studied in many crustaceans (with reviews in Mantel and Farmer, 1983; Péqueux, 1995). Many crustacean species that effectively osmoregulate, either as hyper-isoregulators or as hyper-hypo-regulators, occur in marine, brackish and freshwater conditions (with reviews in Mantel and Farmer, 1983). Monitoring the physiological condition of crustaceans by using osmoregulation could therefore have a potential use as a biomarker in natural waters, such as inland freshwater lakes and rivers as well as in coastal areas, estuaries, lagoons, i.e., in the most exposed parts of the aquatic environment. The use of osmoregulation might also be considered in aquaculture in the early detection of adverse rearing conditions from different origins, including water quality.

The aim of this manuscript is to review the effects of various stressors on osmoregulation, and to evaluate the potential use of the ability to osmoregulate for monitoring the physiological condition of wild and cultured individuals and species of aquatic crustaceans submitted to different stresses. Temperature, although an important external factor and potential stress, is not taken into account in this review since the interrelations between temperature and osmoregulation have already been reviewed (Charmantier, 1975, 1987; Burton, 1986).

Since Na^+ and Cl^- ions are known to make up 90% or more of the osmotic pressure of hemolymph in most crustaceans (Prossner, 1973; Castille and Lawrence, 1981a,b; Mantel and Farmer, 1983), the changes in ionoregulation following stress will also be reviewed. In aquatic crustaceans and particularly in decapods, the organs of the branchial chambers are the primary site of osmotic and ionic regulation (as well as respiration and excretion) (with reviews in Mantel and Farmer, 1983; Taylor and Taylor, 1992; Péqueux, 1995). The large area for adsorption, the large volume of water passing

over the ion-transporting cells (or ionocytes) and the relatively small biomass compared to their surface area are indications that tissues of the branchial cavities can potentially suffer from structural and functional alterations after exposure to stress. Comparable alterations can also affect the excretory organs, in particular the antennal glands of decapods involved in volume regulation, and also in the regulation of solutes, divalent ions and nutrients (Mantel and Farmer, 1983; Wheatly and Gannon, 1995). The effect of water-borne toxicants and stressors on the structure of crustacean ion-transporting organs will therefore be reviewed. Modifications of the activity of Na^+, K^+ -ATPase, one of the most important enzymes involved in active ion-transport in crustaceans (Towle, 1990; Lucu, 1993), and changes in ionic fluxes and surface permeability will also be considered.

To understand the impact of pollutants on aquatic organisms, the notion of stress that has been extensively reviewed (Bayne, 1975, 1985; Ivanovici and Wiebe, 1981; Broom, 1991) has to be defined. From all those available, Bayne's (1975) definition, derived from mollusc studies, is the best suited to the scope of this review: he defined stress as "a measurable alteration of a physiological (or behavioural, biochemical, or cytological) steady-state that is induced by an environmental change and that renders the individual (or the population, or the community) more vulnerable to further environmental change." An important aspect in this definition is that the modification (i.e., the alteration from a steady-state, or control, value) in the chosen physiological parameter must be "measurable". In addition, the "environmental" qualification of the change will be broadened in this review to changes affecting natural environments and artificial culture conditions.

2. Effect of stress on osmoregulation

2.1. Definition of the osmoregulatory capacity

Osmoregulatory capacity (OC) is defined, for a given osmoregulating species, as the difference between the osmotic pressures of the hemolymph and of the external medium, at a given salinity (Charmantier et al., 1989). Hyper-OC and/or hypo-OC can be measured according to the type of osmoregulation of the species, respectively, in low-salinity media, and in seawater or high-salinity media (Robertson, 1960).

The measurement of hemolymph osmolality that permits OC evaluation has been successful in numerous crustaceans from small isopod and gammarid species (Charmantier, 1975; Einarson, 1993) to penaeid shrimps (Williams, 1960; Pannikar, 1968; Castille and Lawrence, 1981a,b; Dall, 1981; Ferraris et al., 1987; Lin et al., 1991; Bambang et al., 1995a,b; Lignot et al., 1997, 1998a,b, 1999b), homarid lobsters (Dall, 1970; Thuet et al., 1988; Young-Lai et al., 1991) and crabs (Warburg et al., 1987; Diamond et al., 1989). Measurement of hemolymph osmolality can also be evaluated in embryos and early developmental stages (Charmantier et al., 1988; Morritt and Spicer, 1995; reviewed in Charmantier, 1998). The technique implies several precautions. Experimental salinities and temperatures must be chosen in such a range that mortality is kept to a minimum and OC values are kept at maximum, in order to better reflect the changes induced by stress. When the evaluation of OC requires a change in salinity, animals

Table 1

Effects of different stress factors on OC, osmoregulation and ionic regulation of different species of crustaceans. When available, size, weight, sex, developmental stage, dose or value of exposure to stress, and experimental salinity(ies), are indicated. OC: osmoregulatory capacity (difference between osmotic pressures of hemolymph and external medium, at indicated salinity); OR: osmoregulation (osmotic pressure of hemolymph at the indicated salinity); Na⁺R, Cl⁻R: sodium, chloride regulation (ionic concentration in hemolymph at the indicated salinity); CTL: cephalothorax length; CTW cephalothorax width; TL: total length; FW: freshwater; SW: seawater; DSW: diluted seawater; ~ : about; ∼ : significant decrease of; ∗ : significant increase of (*p* < 0.05)

Pollutant or stress	Species, stage, size and weight	Dose	Effect on osmoregulation and/or ionic regulation	References
<i>Oil</i>				
Fuel oil	<i>Penaeus aztecus</i> (5 g)	20% solution of water soluble fraction	no effect on OR at 10‰ and 30‰	1
Crude oil	<i>Homarus americanus</i> (~ 500 g)	40–500 µg l ⁻¹ (total oil concentration)	↘ OR by 2% in SW but no effect on Na ⁺ R and Cl ⁻ R	2
<i>Pesticides-PCBs</i>				
<i>Alachlor</i>				
	<i>Rhithropanopeus harrisii</i>	25, 50 mg l ⁻¹	no effect on OR during adaptation from 20‰ to 1‰	3
Aroclor 1254 (PCB)	<i>P. aztecus</i> (adult, 11.5–13.7 cm TL)	3 µg l ⁻¹	no significant effect on OR ↘ Na ⁺ R by 16% at 10‰ ↘ Cl ⁻ R by 19% at 10‰	4
	<i>Palaemonetes pugio</i> (adult and juvenile)	14–60 µg l ⁻¹	no effect on OR from 32‰ to 2‰	1
	<i>Pa. pugio</i> (adult and juvenile) (juvenile)	7, 21.5 µg l ⁻¹ ; 1.8 µg l ⁻¹ ;	no effect on Cl ⁻ R from 35‰ to 10‰	5
DDT, Mirex	<i>Callinectes sapidus</i> (35–50 mm CTW)	0.53 and 3.2 µg/week (DDT) and 0.03, 0.06, 0.14 µg/week (Mirex) in food	↘ Cl ⁻ R by 10% at 7‰ no effect on OR, Na ⁺ R, Cl ⁻ R from 0.7‰ to 48‰	6
DDT	<i>Carcinus maenas</i> (40 g)	0.2 µg l ⁻¹	↘ Na ⁺ R by 18% at 1.7‰	7
Endosulfan	<i>Ozotelphusa senex senex</i> (32 ± 1 g)	6.2, 18.6 mg l ⁻¹	↘ Na ⁺ R by 7% and 13% in FW ↘ Cl ⁻ R by 10% and 16% in FW	8
Fenitrothion	<i>P. japonicus</i> (early adult at stages C and D ₀ , 43 mm CTL) <i>P. japonicus</i> (juvenile at stage C, 13 ± 1 g and 20 ± 1 g)	0.25, 0.5, 1 µg l ⁻¹ 0.5, 0.75, 1 µg l ⁻¹	↘ OR by 58% in SW (37‰) time- and dose-dependent ↘ OC, up to 62% at 37‰ and up to 18% at 19‰	9 10

	<i>P. stylirostris</i> (juvenile at stage C, 13.5 ± 1.4 g)	6, 8, 10, 12 $\mu\text{g l}^{-1}$	time- and dose-dependent $\searrow$ OC, up to 22% at 36‰	11
	<i>P. vannamei</i> (juvenile at stage C, 14.2 ± 2.0 g)	5, 10, 15, 20 $\mu\text{g l}^{-1}$, 25 $\mu\text{g l}^{-1}$	no effect from 5 to 20 $\mu\text{g l}^{-1}$ in SW (36‰) $\searrow$ OC by 21% at 25 $\mu\text{g l}^{-1}$ in SW (36‰)	11
Lindane	<i>Gammarus duebeni</i> (60–90 mg)	3.2–10 $\mu\text{g l}^{-1}$	$\searrow$ Na ⁺ R by 7–12% at 0.7‰	12
Methoxychlor	<i>Cancer magister</i> (adult, 80–100 mm CTW)	0.01 mg l ⁻¹	no effect on OR from 6‰ to 31‰ no effect on Na ⁺ R	13
	<i>Hemigrapsus nudus</i> (15–30 mm CTW)	0.01 mg l ⁻¹	no effect on OR from 9‰ to 34‰	13
Metals				
Aluminium	<i>Astacus astacus</i> (adult at stage C ₄)	0.25, 0.5, 1 mg l ⁻¹	$\searrow$ Na ⁺ R by 10–15% in FW (at pH 5.0)	14, 15
	<i>A. astacus</i> (7–10.5 cm TL)	180 $\mu\text{g l}^{-1}$	$\searrow$ Cl ⁻ R by 35% in FW (at pH 6.3)	16
	<i>A. astacus</i> (23–40 g)	25 $\mu\text{mol l}^{-1}$	$\searrow$ OR by 38% in FW (at pH 4) $\searrow$ Na ⁺ R by 44% in FW (at pH 4) $\searrow$ Cl ⁻ R by 17% in FW (at pH 4)	17
Arsenite	<i>Pacifastacus leniusculus</i> (adult at stage C ₄)	0.25, 0.5, 1 mg l ⁻¹	$\searrow$ Na ⁺ R by 10% in FW (at pH 5.0)	14
	<i>C. maenas</i> (55–72 mm CL)	0.01, 0.1, 1 mg l ⁻¹	no effect on OR during adaptation from 30‰ to 5‰	18
Cadmium	<i>Jaera albifrons</i>	10, 20 mg l ⁻¹ ; 10 mg l ⁻¹	no effect on OR at 3.4‰ and 34‰ $\nearrow$ OR by $\sim$ 40% at 17‰	19
	<i>Idotea baltica</i>	10 mg l ⁻¹	no effect on OR at 27‰	19
	<i>I. emarginata</i>	10 mg l ⁻¹	no effect on OR at 27‰	19
	<i>I. neglecta</i>	10 mg l ⁻¹	$\searrow$ OR at 27‰	19
	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 13 ± 1 g)	2, 4 mg l ⁻¹	time- and dose-dependent $\searrow$ OC, up to 53% at 38‰ and time- and dose-dependent $\searrow$ OC, up to 47% at 16‰ $\nearrow$ OR by 13% in SW (7.5 mg l ⁻¹), no effect on OR in 60‰ SW, dose-dependent $\searrow$ OR up to 33% in 20‰ SW	20
	<i>Palaeomon elegans</i> (juvenile at stage C, 200–300 mg)	1–7.5 mg l ⁻¹	no effect on OR at 25 ± 1 ‰	21
	<i>H. americanus</i> (adult)	3–6 $\mu\text{g l}^{-1}$	dose-dependent $\searrow$ OR up to 25% at 2.5‰	22
	<i>Ca. sapidus</i> (juvenile, 15–20 mm CTL)	50, 100 $\mu\text{g l}^{-1}$	dose-dependent $\nearrow$ OR by 20–50% according to salinity	23
	<i>C. maenas</i> (18–30 g)	0.5–8 mg l ⁻¹		24

(continued on next page)

Table 1 (continued)

Pollutant or stress	Species, stage, size and weight	Dose	Effect on osmoregulation and/or ionic regulation	References
Chromium	<i>C. maenas</i> (25–50 g)	1 mg l ⁻¹	no effect on OR and Na ⁺ R from 13.6‰ to 24‰ ↘ OR	25
	<i>Can. irroratus</i>	25 µg l ⁻¹	↘ Na ⁺ R in 2.5‰ but unaffected in 30‰	26
	<i>Chasmagnathus granulata</i>	1.25, 5 mg l ⁻¹	no effect on Cl ⁻ R in 2.5‰ and 30‰	27
	<i>Uca rapax</i>	1, 5, 25 mg l ⁻¹	Na ⁺ - and Cl ⁻ R: unaffected at 25‰ and 125‰ SW	28
	<i>Praunus flexuosus</i>	16 mg l ⁻¹	↘ OR by 21, 16 and 12% at 4.5‰, 9‰ and 27‰, respectively. no effect on OR at 18‰	29
Copper	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 13 ± 1 g)	0.5–1.5 mg l ⁻¹	time- and dose-dependent ↘ OC, up to 100% at 37‰ and up to 33% at 17‰	30
	<i>Pal. elegans</i> (juvenile at stage C, 200–300 mg)	1–7.5 mg l ⁻¹	dose-dependent ↗ OR up to 38% in SW dose-dependent ↘ OR up to 19% in 60‰ SW and up to 53% in 20‰ SW	21
	<i>A. leptodactylus</i> and <i>Orconectes limosus</i> (42–56 mm)	1 mg l ⁻¹ (max.)	↘ Na ⁺ R by 30% in FW ↘ Cl ⁻ R by 30% in FW	31
	<i>C. maenas</i> (18–30 g)	2.5–40 mg l ⁻¹	dose-dependent ↘ OC, up to 100% at 15‰	24
	<i>C. maenas</i> (adult)	0.25–1 mg l ⁻¹	dose-dependent ↘ OR, Na ⁺ R and Cl ⁻ R up to 30%, 12% and 16%, respectively, at 13.6‰	32
	<i>C. maenas</i> (male, 40–70 g)	1, 2 mg l ⁻¹	no major effect on Na ⁺ R and Cl ⁻ R at 33‰	33
	<i>C. maenas</i> (male, 40–70 g)	0.5 mg l ⁻¹	no major effect on OR at 33.5‰ but ↘ OR, Na ⁺ R and Cl ⁻ R by ~ 14% at 17.5‰	34
	<i>C. maenas</i> (adult, intermolt male)	10 mg l ⁻¹	↘ Na ⁺ R, up to 40% at 10‰ dose-dependent ↘ OR up to 42% (0.5 mg l ⁻¹) after transfer from SW (30‰) to DSW (5‰)	35
	<i>C. maenas</i> (55–72 mm CL)	0.1, 0.5, 1 mg l ⁻¹		18
	<i>C. maenas</i> (adult)	0.75 mg l ⁻¹	↘ Na ⁺ R and Cl ⁻ R by 21% and 11% at 10‰	36
	<i>Can. irroratus</i> (28–50 g)	0.3–5 mg l ⁻¹	↗ Cl ⁻ R by 30% at 30‰ dose-dependent ↘ OC, up to 90% at 15‰	24

Lead	<i>Cherax destructor</i> (adult at stage C, 75–100 mm)	0.5 mg l ⁻¹	no effect on OR, Na ⁺ R and Cl ⁻ R in FW	37
Mercury	<i>J. albifrons</i>	1 mg l ⁻¹	↗ OR by 5% at 33‰	19
	<i>Pal. elegans</i> (juvenile at stage C, 200–300 mg)	0.1 mg l ⁻¹ 20–150 µg l ⁻¹	↗ OR by 14% at 1.7‰ ↗ OR up to 7% in SW (0.15 mg l ⁻¹) dose-dependent ↘ OR up to 10% in 60% SW and up to 23% in 20% SW	21
	<i>O. propinquus</i>	1 µg/Hg g ⁻¹ mercury in water mercury in food	↘ Na ⁺ R by 21% ↘ Na ⁺ R by 17%	38
	<i>H. americanus</i> (adult) <i>C. maenas</i> (adult)	3–6 µg l ⁻¹ 0.5–10 mg l ⁻¹	no effect on OR at 25 ± 1‰ dose-dependent ↘ OR, Na ⁺ R and Cl ⁻ R, up to 20%, 25% and 35%, respectively, at 13.6‰. Effects vary during the year no effect on Cl ⁻ R from 7‰ to 35‰ ↘ Na ⁺ R by 40% in FW	22 39 1, 40 41
Methylmercury	<i>Petrolisthes armatus</i>	50 µg l ⁻¹	↘ Cl ⁻ R by 30% in FW	12
Nickel	<i>Eriocheir sinensis</i>	10 µg l ⁻¹	↘ Na ⁺ R by ~8% at 0.7‰	29
	<i>G. duebeni</i> (60–90 mg) <i>Pra. flexuosus</i>	56–320 µg l ⁻¹ 64 mg l ⁻¹	↘ OR by ~17% at 4.5‰ and 9‰ no effect on OR at 18‰ and 27‰	29
Zinc	<i>Pra. flexuosus</i>	16 mg l ⁻¹	↘ OR by 11%, 20% and 9% at 4.5‰, 9‰ and 27‰, respectively no effect on OR at 18‰ ↘ OR by 12% at 10‰	42
	<i>G. duebeni</i>	0.5 mg l ⁻¹	↘ Na ⁺ R by 13% at 10‰	19
	<i>J. albifrons</i>	20 mg l ⁻¹	↘ OR by 20% at 3.4‰	19
	<i>I. ballica</i>	20 mg l ⁻¹	↘ OR by 15% at 34‰	19
	<i>I. neglecta</i>	10 mg l ⁻¹	no effect on OR at 27‰ and 34‰	21
	<i>Pal. elegans</i> (juvenile at stage C, 200–300 mg)	1–15 mg l ⁻¹	↗ OR up to 9% in SW (15 mg l ⁻¹) dose-dependent ↘ OR up to 9% in 60% SW and up to 17% in 20% SW	43
Tributyltin oxide (TBTO)	<i>P. japonicus</i> (juvenile at stage C)	100, 200 µg l ⁻¹	time- and dose-dependent ↘ OC, up to 79% at 38.5‰ and up to 86% at 19‰	

(continued on next page)

Table 1 (continued)

Pollutant or stress	Species, stage, size and weight	Dose	Effect on osmoregulation and/or ionic regulation	References
Phenols				
Phenol	<i>Mesidotea entomon</i>	50, 80, 120, 180, 270, 400 mg l ⁻¹	dose-dependent ↘ Na ⁺ R, up to 25% at 6‰	44
4-Chlorophenol	<i>M. entomon</i>	45 mg l ⁻¹	↗ Na ⁺ R by 20% at 6‰	44
		100 mg l ⁻¹	↘ Na ⁺ R by 10% at 6‰	45
Potassium	<i>H. americanus</i> (20–55 mm CTL)	0.8 g l ⁻¹	no effect on iso-OR from 31 to 48‰	46, 47
Ammonia	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 8.5–11.5 g)	16, 48 mg l ⁻¹	dose-dependent ↘ OC, up to 100% at 36‰	48
	<i>H. americanus</i> (st. IV, stage C);	32, 48, 64 mg l ⁻¹	dose-dependent ↘ OC, up to 40% at 15‰	
Nitrite	(adult, 65–75 mm CTL, st. C)	32, 64, 96 mg l ⁻¹	dose-dependent ↘ OC, up to 47% at 17‰	49
	<i>P. japonicus</i> (juvenile at stage C, 11.4 ± 1.4 g)	100, 150, 200 mg l ⁻¹	dose-dependent ↘ OC, up to 45% at 17‰	
	<i>Macrobrachium rosenbergii</i> (23.3 ± 3.5 g)	0.01, 0.07, 0.37, 0.72, 1.43 mM	dose-dependent ↗ OC, up to 47% at 30‰	50
	<i>A. astacus</i> (18–45 g)	2, 10 mg l ⁻¹	dose-dependent ↘ OR, up to 5.5% at pH 7.3 and up to 4.1% at pH 7.7	
Low pH	<i>A. astacus</i> (30–60 g)	0.8 mM	↘ Na ⁺ R by 5% in FW	51
		0.5, 1 mM	↘ Cl ⁻ R by 23% in FW	52
	<i>Pac. leniusculus</i> (33 ± 1 g)	1 mM	dose-dependent ↘ Na ⁺ R up to 17% in FW	53
			dose-dependent ↘ Cl ⁻ R up to 38% in FW	
			no significant effect on OR in FW	54
			↘ Na ⁺ R by 12% in FW	
			↘ Cl ⁻ R by 25% in FW	54
	<i>Artemia franciscana</i> (nauplii)	pH 5.7–8.3	↗ Na ⁺ R by 79% (pH 5.8) and by 32% (pH 6.0) at 35‰	
			No significant variation from pH 6.5 to 8.3	54
			pH-dependent ↘ Na ⁺ R from pH 7.0 to 5.7, up to 39% at 7‰	
			pH-dependent ↘ Cl ⁻ R from pH 7.0 to 5.7, up to 64% at 7‰	
	<i>Ar. franciscana</i> (adult)	pH 4.5–8.3	↗ Na ⁺ R and Cl ⁻ R by ~ 160% (pH 4.5) at 35‰	
			No significant variation from pH 5.0 to 8.3	
			pH-dependent ↘ Na ⁺ R from pH 7.0 to 5.25, up to 42% at 7‰	

Table 1 (continued)

Pollutant or stress	Species, stage, size and weight	Dose	Effect on osmoregulation and/or ionic regulation	References
Starvation	<i>C. maenas</i> (adult)	0 mm Hg	no effect on OR in SW	66
	<i>C. maenas</i> (50–150 g)	20 Torr	no effect on Na ⁺ R at 13.5‰ ➤ Cl ⁻ R by 10% at 13.5‰ ➤ OR	67
	<i>Apus (L. productus)</i> <i>Branchipus</i> sp.		➤ OR	68
	<i>Corophium volutator</i>	72–192 h	➤ OR by 19% at 10‰	68
	<i>D. magna</i>		➤ OR by 30% in FW (?)	69
Fungal infection	<i>P. stylirostris</i> (juvenile at stage C, 9.7 ± 1.3 g)	24 h	➤ OR by 6% in SW	70
	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 13.4 ± 2.8 g)			71
		Injection of <i>F. oxysporum</i> in juveniles	➤ hypo-OC by 53% in SW 14 days after infection ➤ hyper-OC by 26% in DSW (20‰) 16 days after infection	72
	<i>A. leptodactylus</i> and <i>Au. pallipes</i> (adult at stage C)	36 h of exposure to <i>Fusarium oxysporum</i> (conidia)	➤ OR by 14%, 23% and 11% after, respectively, 1, 2 and 3 weeks in FW after infection.	73
	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 9.7 ± 1.3 g)	35 ± 15 NTU	➤ hypo-OC by 69% at 42‰ ➤ hyper-OC by 35% at 15‰	74

References: (1) Anderson et al., 1974; (2) Payne et al., 1983; (3) Diamond et al., 1989; (4) Nimmo and Bahner, 1974; (5) Roesijadi et al., 1976; (6) Leffler, 1975; (7) Jowett et al., 1981; (8) Rajeswari et al., 1988; (9) Langy, 1991; (10) Lignot et al., 1997; (11) Lignot et al., 1998b; (12) Inman and Lockwood, 1977; (13) Caldwell, 1974; (14) Appelberg, 1985; (15) Appelberg, 1987; (16) Fjeld et al., 1988; (17) Jensen and Malte, 1990; (18) Bamber and Depledge, 1997; (19) Jones, 1975; (20) Bambang et al., 1995a; (21) Chen and Slinn, 1980; (22) Thurberg et al., 1977; (23) Guerin and Stickle, 1995; (24) Thurberg et al., 1973; (25) Bjerregaard, 1991; (26) Dawson; (in) Thurberg et al., 1977; (27) Vitale et al., 1998; (28) Rojas and Zanders, 1998; (29) McLusky and Hagerman, 1987; (30) Bambang et al., 1995b; (31) Chaisemartin, 1973; (32) Bjerregaard and Vislie, 1986; (33) Boitel and Truchot, 1989; (34) Boitel and Truchot, 1990; (35) Hansen et al., 1992; (36) Weeks et al., 1993; (37) Ahern and Morris, 1998; (38) Wright and Welbourn, 1993; (39) Bjerregaard and Vislie, 1985; (40) Roesijadi et al., 1974; (41) Bouqueneau and Hibaude, cited in Gilles and Péquereux, 1983; (42) Johnson and Jones, 1990; (43) Lignot et al., 1998a; (44) Oksama and Kristofferson, 1980; (45) Charmanier et al., 1985; (46) Lin et al., 1991; (47) Lin et al., 1993; (48) Young-Lai et al., 1991; (49) Chen and Cheng, 1996; (50) Chen and Lee, 1997; (51) Jensen, 1990; (52) Jeberg and Jensen, 1994; (53) Harris and Coley, 1991; (54) McMahon and Doyle, 1997; (55) Havas and Hutchinson, 1983; (56) Allan and Maguire, 1992; (57) Ellis and Morris, 1995; (58) Wood and Rogano, 1986; (59) Zanotto and Wheatley, 1993; (60) Morgan and McMahon, 1982; (61) McMahon and Stuart, 1989; (62) Kelley and Burbanck, 1972; (63) Charmanier et al., 1994; (64) Hagerman and Uglov, 1981; (65) Hagerman and Uglov, 1982; (66) Spaargaren, 1977; (67) Johnson and Uglov, 1987; (68) Krogh, 1939; (69) McLusky, 1970; (70) Fritsche, 1916; (71) Lignot et al., 1999b; (72) Souheil et al., 1999; (73) Maestracci and Vey, 1989; (74) Lin et al., 1992.

must be kept in the new medium until full osmotic equilibration (Charmantier-Daures et al., 1988). Since the OC value also changes with size, nutritional status, developmental stage and molt stage, only starved animals at the same size and developmental stage, and at intermolt stage C must be used (Lignot et al., 1999b). Controls, not subjected to the studied stress but maintained under the same experimental conditions, must also be used for comparisons of OC values.

2.2. Effect of stress on ionic regulation and OC

The effects of different stress factors on OC, osmoregulation and ionic regulation of crustaceans are summarized in Table 1. They have been reported mostly in decapods, for a lesser part in isopods and amphipods and very seldom in other groups. Stress was mainly applied on animals in hyper-regulating situations, i.e., on marine or brackish water species in dilute media or on freshwater species in their medium. In the vast majority of experimental cases, application of a stress alters osmoregulation and/or ionic regulation and reduces the OC. The reduction is mostly observed for crustaceans living in (or acclimated to) seawater and diluted media, i.e., for hyper-hypo- and hyper-iso-regulators. In isosmotic crustaceans (osmoconformers), alterations in OC and/or ionic regulation are not usually significant since OC is close to zero.

In freshwater and diluted media, significant decreases of hyper-OC range from a few percent to usually 20–50% and up to 90–100% in some cases (Thurberg et al., 1973). Hyper-OC decreases are particularly apparent on exposure to ammonia, low pH, TBTO, fungal infections of the gills, some pesticides and most of the studied metals, particularly aluminium, cadmium and copper (see Table 1). In some cases, exposures to oil, pesticides, PCBs and metals do not affect osmoregulation (Anderson et al., 1974; Caldwell, 1974; Leffler, 1975; Diamond et al., 1989; Bjerregaard, 1991; Bamber and Depledge, 1997; Ahern and Morris, 1998). These observations might result from the application of too low doses, over too short exposure times, from specific differences of sensitivity to pollutants and/or from the absence of interference of these pollutants with osmoregulatory processes. In some rare occurrences, exposure to pollutants increases OC or sodium regulation, as in the case of cadmium and mercury in *J. albifrons* (Jones, 1975) and 4-chlorophenol in *M. entomon* (Oksama and Kristoffersson, 1980). In the latter species, however, a higher dose of 4-chlorophenol lowers sodium regulation. Although little is known about the threshold of pollutant detection, an activation of ionic uptake following pollutant exposure could be initiated in some species as the first reaction to a change of the ambient medium, in order to maintain the internal hemolymph homeostasis. This compensating mechanism might also explain the lack of effect or even some of the observed OC increases.

In seawater, stress exposure generally induces a decrease in hypo-OC and/or Na^+ and Cl^- regulation. This is particularly apparent on exposure to ammonia, low pH and most of the metals studied, particularly aluminium, cadmium and copper. Hypo-OC can decrease up to 100%, rendering the animals isosmotic with the medium, as in *P. japonicus* exposed to ammonia (Lin et al., 1991) or to copper (Bambang et al., 1995b). As in low-salinity media, osmoregulation in seawater may not be affected by pollutants in crustaceans that are either weak hyporegulators (Thurberg et al., 1977) or isosmotic

Table 2
Effects of different stress and pollutants on the morphology, histological structure and ultrastructure of branchial chambers and excretory organs. When several concentrations or several times of exposure to pollutants were used, the maximum deleterious effect on tissues was described. H: hour; CL: carapace length; TL: total length; FW: freshwater; SW: seawater; S‰: salinity

Pollutant	Species, size and weight	Dose, time and medium of exposure	Effect on morphology (M), histological structure (H) and ultrastructure (U)	References
Branchial organs				
<i>Oil and dispersants</i>				
Crude oil	<i>Charybdis lucifera</i>	0.1–0.8 ml oil l ⁻¹ ; 1 day; S = 35.5‰	H: branchial cell necrosis, rupture and fusion of lamellae, accumulation of hemocytes.	1
	<i>Scylla serrata</i>	1–4 ml oil l ⁻¹ ; 1 day; S = 35.5‰	H: branchial epithelial damage but no lamellar disruption or fusion.	1
Oil emulsions and Dispersants	<i>Pal. serratus</i>	200–5000 mg l ⁻¹	H: swelling of gill lamellae, cell vacuolization, nuclear pycnosis, rupture of pillar cells.	2
<i>Pesticides, fungicides</i>				
Busan 85	<i>Pa. pugio</i> (22–28 mm TL)	5–60 µg l ⁻¹ ; 14 days; S = 15‰	M, H: swelling of gill lamellae, blackened tip of lamellae, accumulation of hemocytes. U: cytolysis of branchial epithelial cells, swollen mitochondria with sparse cristae, dilated rough endoplasmic reticulum without attached ribosomes.	3
Fenitrothion	<i>P. japonicus</i> (13–20 g)	0.5, 0.75, 1 µg l ⁻¹ ; 4 days; SW	H: gill and epipodite necrosis, accumulation of particles between the branchial lamellae, breakage of the	4

			3.1 $\mu\text{g l}^{-1}$, 14 days; 0.097 $\mu\text{g l}^{-1}$, 30 days	cuticular lining, hemocytic congestion (thrombosis). H: necrosis of gill tissue, hemocytic congestion and separation of the cuticle from the epithelium.	5
Gusathion A	<i>Ma. kismensis</i> (50 $\pm$ 10 mm TL, 0.6 $\pm$ 0.1 g)	<i>P. monodon</i>	75, 150 $\mu\text{g l}^{-1}$, 4 days	H: hyperplasia of the gill epithelium, degeneration and necrosis of gill tissue.	6
Metals Aluminium	<i>A. astacus</i> (7–10.5 cm TL)		80 $\mu\text{g l}^{-1}$ pH 6.7; 10 days; FW	M, H: brown colored gill filaments, shrunken with an irregular surface. U: branchial epithelial cells severely degraded, electron dense layer on surface of the epithelium.	7
Cadmium	<i>Ja. nordmanni</i>		10, 20 mg l^{-1} ; 8–24 h; S = 10% SW	H: numerous hemocytes in gill haemolymph spaces. U: gill basal membrane infoldings fragmented, distended microvilli, dilated endoplasmic reticulum, swollen mitochondria.	8
	<i>P. duorarum</i> (43–100 mm TL)		5 mg l^{-1} 4 days; 1 mg l^{-1} 21 days; S = 20%	M, H: partly blackened gills, accumulation of hemocytes, necrosis of epithelium, dissolution of exo- and endocuticular layers.	9
	<i>P. duorarum</i>		763 $\mu\text{g l}^{-1}$ 15 days; S = 20%	M, H: partly blackened gills, disorganization of the epithelial structure, necrosis.	10

(continued on next page)

Table 2 (continued)

Pollutant	Species, size and weight	Dose, time and medium of exposure	Effect on morphology (M), histological structure (H) and ultrastructure (U)	References
	<i>P. japonicus</i> (12 ± 1 g)	2000, 4000 µg l ⁻¹ 4 days; SW	U: gill cell vacuolization, cellular membrane degeneration, nuclear pycnosis. M, H: blackened and necrosed gills and epipodites, thickening of the epithelial layer, swollen epipodites, separation of the cuticle from the epithelium. U: increased number of nephrocytes in the gills, autolysis and loss of cytoplasmic organelles, microvilli and basal lamina, swollen mitochondriae, vacuolization and nuclear pycnosis. U: branchial mitochondria degeneration, dilatation of intercellular spaces and of the rough endoplasmic reticular system.	11
	<i>Paratyta tasmaniensis</i>	30, 50 µg l ⁻¹ 4 days; FW		12
	<i>Pal. serratus</i>	5, 25, 50 mg l ⁻¹ ; 44 h; S = 30‰	U: mitochondria in the gills are the most affected organelles (disruption of the bounding membranes, loss of cristae, swelling), impairment of the pinocytotic process and of the membrane system involved in osmoregulation, fewer infoldings of the apical and basal plasma membrane.	13

	<i>Pa. vulgaris</i>	75 $\mu\text{g l}^{-1}$; 10 days; S = 20‰	H: melanized and necrotic gill lamellae distended by numerous hemocytes.	9
	<i>Cr. crangon</i>	5, 25, 50 mg l^{-1} ; 20 h; S = 30‰	U: mitochondria in the gills are the most affected organelles, swollen endoplasmic reticulum, fewer invaginations of the apical plasma membrane and more cytoplasmic vesicles than in controls.	14
	<i>Pr. clarkii</i> (20.5–30.4 g)	1 mg l^{-1} ; 4 days; FW	H: extensive disorganization of the gill filaments, the medial septum cannot be observed.	15
	<i>Pa. pugio</i> (22–28 mm TL)	0.5, 1, 2, 4 mg l^{-1} ; 30 days; S = 10‰	H: relatively few degenerative changes in the gills, lamellar irregularity. U: decrease in epithelial cytoplasmic density in the gill filaments, swollen mitochondria with fragmented cristae, diminution of plasmalemmal infoldings.	16
Chromium			U: numerous hemocytes, irregularly outlined microvilli, large intracellular vacuoles, disrupted infolded basal membrane, swollen mitochondria with a reduction in cristae, dilated and fragmented endoplasmic reticulum.	8
	<i>Ja. nordmanni</i>	1, 10 mg l^{-1} ; 8–24 h; S = 10‰ SW; 8–96 h; S = 50‰ SW		
Copper				
	<i>P. japonicus</i> (12–15 g)	500, 1000 $\mu\text{g l}^{-1}$; 4 days; SW	M, H: dose-dependent blackening of gills and	17

(continued on next page)

Table 2 (continued)

Pollutant	Species, size and weight	Dose, time and medium of exposure	Effect on morphology (M), histological structure (H) and ultrastructure (U)	References
	<i>Ma. kistnensis</i>	50, 100 µg l ⁻¹ ; 6–8 days; FW	epipodites, necrosis, swollen epipodites. U: increased number of gill nephrocytes, swollen epithelial cells, vacuolization, degraded microvilli and basal infoldings and disorganized cytoplasmic structures.	18
	<i>Caridina sp.</i>	50, 100 µg l ⁻¹ ; 6–8 days; FW	M, H: no blackened gills, distension of gill lamellae, vacuolization and necrosis of gill tissue. H: partly blackened gills, distension of gill lamellae, vacuolization and necrosis of gill tissue.	18
	<i>C. maenas</i> (50–80 g)	0.5, 2 mg l ⁻¹ ; 5–6 days; S = 30–33‰	U: gill necrosis, cellular hyperplasia, vacuolization, thickening of the gill epithelium and reduction of haemolymph spaces.	19
	<i>C. maenas</i> (15–20 g; 40–45 mm CL)	50 µg l ⁻¹ ; 10 days; S = 10, 35‰	U: decrease in the number of plasma membrane infoldings and associated mitochondria in the gill epithelium, vacuolization, change in ribosomal distribution, disruption of the microtubular network.	20
Lead	<i>Pr. clarkii</i> (21.5–28.4 g)	200 mg l ⁻¹ ; 4 days; FW	M, H: gill filaments grossly blackened, pronounced	21

			disorganization, medial septum not visible. M, H: gross blackening of almost all gill filaments observed, irregular profiles of the filaments, structural disorganization.	14
	<i>Pr. clarkii</i> (20.5–30.4 g)	100 mg l ⁻¹ ; 4 days; FW	H: branchial epithelial layer separated from the cuticle, reduction of cellular organites, nuclear degeneration, numerous vesicles containing electron dense material.	22
	<i>Pr. clarkii</i>	100 mg l ⁻¹ ; 4 days; FW	H: numerous hemocytes, cellular break down and large sub-cuticular spaces.	8
	<i>Ja. nordmanni</i>	0.1, 1 mg l ⁻¹ ; 8–24 h; S = 10% SW	U: irregularly outlined microvilli, large intracellular vacuoles, disrupted infolded basal membrane, swollen mitochondria with a reduction in cristae, dilated and fragmented endoplasmic reticulum.	23
	<i>Cr. crangon</i> (4.5–6.6 cm TL)	17 µg l ⁻¹ ; 14 days; S = 20‰	H: accumulation of mercury on the gills surface and within the lamellar epithelial cells, the septum and the hemocytes.	24
	<i>Ma. hendersonianum</i> (1.1 g)	3 mg l ⁻¹ 15 days; 6–15 mg l ⁻¹ 4 days; FW	U: mercury deposits in lysosome-like bodies and mainly in nephrocytes. H: gill lamellae disrupted, accumulation of hemocytes, necrosis of epithelium, nuclear pycnosis.	

(continued on next page)

Table 2 (continued)

Pollutant	Species, size and weight	Dose, time and medium of exposure	Effect on morphology (M), histological structure (H) and ultrastructure (U)	References
TBTO	<i>P. japonicus</i> (9–12 g)	200, 400 $\mu\text{g l}^{-1}$; 4 days; SW	H: multiple gill and epipodire necrosis; hemocytic congestion, severe nephrocyte hyperplasia.	25
Fungal infection	<i>A. leptodactylus</i> , <i>Au. pallipes</i> (adults)	36 h of exposure to <i>Fusarium oxysporum</i> (conidia), up to 3 weeks; FW	H: black spots on the gills with melanotic deposit in contact with the hyphae; hemocytic reaction with encapsulation processes. U: cellular alteration of the gill epithelium, vacuolization, dilatation and alteration of mitochondria, microvilli acquiring a loose structure.	26
Antennal glands <i>Fungicides</i> Busan 85 Aquareat (DNM 30)	<i>Pa. pugio</i> (22–28 mm TL)	5–60 $\mu\text{g l}^{-1}$; 14 days; SW 40–120 $\mu\text{g l}^{-1}$; 3–35 days; S = 10‰	U: labyrinth: channel virtually occluded, microvilli distended and ruptured, dilated mitochondria, numerous vesicles; coelomosac: deterioration of podocyte organization, nuclear pycnosis.	27

Metals				
Chromium	<i>Pa. pugio</i> (22–28 mm TL)	0.5, 1, 2, 4 mg l ⁻¹ ; 30 days; S = 10‰	U: labyrinth: cytoplasmic vacuolization, swollen mitochondria with vesiculation of cristae, exfoliation of microvilli, decrease in number and extent of the basal membrane infoldings, exfoliation of cells; coelomosac: epithelium generally normal.	16
Copper	<i>O. rusticus</i> (12–15 g)	0.5 mg l ⁻¹ ; 2 days; FW	H: labyrinth: cells vacuolized then completely necrosed; nephridial canal: less affected.	28
Mercury	<i>Cr. crangon</i> (4.5–6.6 cm TL)	17 µg l ⁻¹ ; 14 days; S = 20‰	H: coelomosac: blackened cells. U: labyrinth: mercury deposits within lysosomes and along the basal lamina; coelomosac: large black inclusions in vacuoles and cytoplasm of podocytes and along the basal membrane.	23
	<i>Pac. leniusculus</i>	2.5 mg l ⁻¹ ; 2 days; FW	H: labyrinth, coelomosac: nuclear pycnosis, cellular disruption.	29

References: (1) Chandty and Kolwalkar, 1984; (2) Papineau and Cheze, 1984; (3) Pawar and Katdare, 1982; (4) Lignot et al., 1997; (5) Doughtie and Rao, 1983a; (6) Baticados and Tendencia, 1991; (7) Fjeld et al., 1988; (8) Bubel, 1976; (9) Nimmo et al., 1977; (10) Couch, 1977; (11) Soegianto et al., 1999a; (12) Lake and Thorp, 1974; (13) Papathanassiou and King, 1983; (14) Papathanassiou, 1985; (15) Torreblanca et al., 1989; (16) Doughtie and Rao, 1984; (17) Soegianto et al., 1999b; (18) Ghate and Mulherkar, 1979; (19) Nonnotte et al., 1993; (20) Lawson et al., 1995; (21) Torreblanca et al., 1987; (22) Rubio et al., 1991; (23) Andersen and Bastrup, 1988; (24) Patil and Kaliwal, 1989; (25) Lignot et al., 1998a; (26) Maestracci and Vey, 1989; (27) Doughtie and Rao, 1983b; (28) Hubschman, 1965; (29) Ellis and Fuller, 1979.

with the external seawater (Roesijadi et al., 1974; Jones, 1975; Thurberg et al., 1977; Boitel and Truchot, 1989, 1990; Bjerregaard, 1991). Osmoregulation can even be increased by pollutants (Jones, 1975; Chen and Slinn, 1980; McMahon and Doyle, 1997).

Most authors do not indicate whether or not they used individuals at selected molt stages (stage C usually). Water permeability, water content and the concentration of inorganic ions in the hemolymph are known to change during the molt cycle (Robertson, 1960; Bursey and Lane, 1971; Lockwood, 1977; Exbrayat and Bourguet, 1982; Mantel and Farmer, 1983; Wheatly and Ignaszewski, 1990), resulting in variations of the hemolymph osmotic pressure and therefore of OC (Mantel and Farmer, 1983; Ferraris et al., 1987; Charmantier et al., 1994). If molt stage selection is not carried out prior to measuring OC or ion concentrations, results can be variable enough to either decrease the difference between control and exposed groups, or even abolish any significant difference. This might account for several “no effect” results in Table 1.

2.3. Effect of toxicant-induced stress on the structure and ultrastructure of the branchial chambers and excretory organs

Ion-transporting cells or ionocytes, located in the gills and epipodites, in extra-branchial epithelia (e.g., pleurae, branchiostegites) and in antennal glands are characterized by the presence of apical microvilli and basolateral membrane infoldings associated with numerous mitochondria (Berridge and Oschman, 1972; Cioffi, 1984; Taylor and Taylor, 1992; Bouaricha et al., 1994; Haond et al., 1998). The ionocytes show extensive structural changes within the same species according to external salinity (Compère et al., 1989; Shires et al., 1994; Freire and McNamara, 1995; Haond et al., 1998).

Ion-transporting branchial organs are severely affected by toxicant-induced stress (Table 2). Gills and epipodites of crustaceans exposed to pollutants and fungal infections generally blacken and are the site of necrosis and hemocytic congestions. At the ultrastructural level, damages to branchial organs have been reported in numerous studies carried out on crustaceans exposed mainly to metals (Table 2). Common features of cellular alterations are fewer and swollen mitochondria, nuclear pycnosis, intracellular vacuolization, decrease or fragmentation of the basolateral membrane infoldings and occurrence of pseudomyelinic structures. In the antennal glands of decapods, the effects of pollutants generally result in alterations in the structure and ultrastructure of the epithelial cells of the coelomosac and labyrinth, with necrosis, nuclear pycnosis, swollen mitochondria and vacuolization.

The blackening and concomitant disorganization of the ion-transporting epithelial cells (tissue necrosis, thrombosis, vacuolization, disruption of the basolateral infoldings) are usually time- and dose-dependent and appear within a few days of exposure. The progressive destruction of the transporting cells after exposure to stressors can be related to the ambient salinity (Lawson et al., 1995) and can partially explain the decrease of OC. It is generally accepted that the influx of Na^+ ions across the ionocytes involves a two-step process implicating a Na^+, K^+ -ATPase pump and a Na^+/H^+ antiporter (Siebers et al., 1985, 1987; Péqueux et al., 1988; Lucu, 1990; Towle, 1990; Pierrot et al.,

1995; Péqueux, 1995), respectively located along the basolateral infoldings and the apical microvilli (Towle and Kays, 1986; Péqueux et al., 1988; Towle, 1990; Taylor and Taylor, 1992; Lignot et al., 1999a). The observed alterations of these plasma membranes may therefore decrease the surface area available for ion-exchange, disrupt the function of the ion-transferring sites, particularly the sodium pumps, and thus impair ionic exchanges and osmoregulation mechanisms. The decrease in the number of mitochondria and the disruption of their internal membranes (swelling) where ATP is synthesized, can also result in the reduction of available energy necessary for the ion-exchanging pumps. Vacuolization may also indicate a failure to regulate water content. The rate of entry of water into the cell and then, into the hemolymph, can furthermore be decreased by the subcuticular spaces and vacuoles creating a longer pathway for water movement (Bubel, 1976; Lawson et al., 1995).

2.4. Stress effects on the activity of the Na^+, K^+ -ATPases in the gills

The effects of stressors on the gill Na^+, K^+ -ATPase activity are summarized in Table 3. Na^+, K^+ -ATPase activity generally decreases after exposure to stress despite varying effects according to the pollutants. A discrepancy between in vivo and in vitro response of Na^+, K^+ -ATPase activity to stress also exists. In *C. maenas* exposed to DDT, a surprising opposite response in vivo and in vitro has been observed (Jowett and Rhead, 1978; Jowett et al., 1981). In *Ch. destructor* exposed to lead and in *Cha. granulata* exposed to cadmium, gill Na^+, K^+ -ATPase also appeared more sensitive to the pollutant in vitro than in vivo (Ahern and Morris, 1998; Rodriguez-Moreno et al., 1998). This could result from the binding of the pollutant to intracellular proteins or organelles (Andersen and Baatrup, 1988; Rodriguez-Moreno et al., 1998).

Since Na^+, K^+ -ATPase is directly involved in active ion transport, its functional disruption can be related to the ionic and osmotic alterations observed in Table 1. This applies to *C. maenas* exposed to DDT (Jowett et al., 1981), to *A. leptodactylus*, *O. limosus* and *C. maenas* exposed to copper concentrations (Chaisemartin, 1973; Hansen et al., 1992) and to *Ch. destructor* exposed to chronic sublethal lead concentrations (Ahern and Morris, 1998). The observed decrease of the Na^+, K^+ -ATPase activity might be due to the inhibition of the phosphorylated enzyme intermediate necessary for ion translocation (Ahern and Morris, 1998) and/or to the disruption of the basolateral infolding membranes within the ionocytes, as mentioned above. A decrease in Na^+, K^+ -ATPase activity has also been observed in *Can. magister* after sublethal acute exposure to methoxychlor (Caldwell, 1974) and in *Ch. destructor* after long-term exposure to lead (Ahern and Morris, 1998), despite the absence of any effect on ionic regulation. These alterations have been explained either by the fact that the degree of inhibition of the enzyme may be insufficient to significantly affect osmotic and ionic regulation, or by increased ion uptake and reabsorption from the urine in the freshwater species in order to regulate the ionic homeostasis.

In a few cases, the Na^+, K^+ -ATPase activity increases in stress-exposed animals (Jowett and Rhead, 1978; Lin et al., 1992, 1993). A short-term increase in Na^+, K^+ -ATPase activity can result from hormonal stimulation by biogenic amines like dopamine

Table 3

Effects of different stress and pollutants on the activity of the gill enzyme Na^+, K^+ -ATPase. Abbreviations as in Tables 1 and 2; in vitro: pollutant added to the incubation medium of gill homogenates or membranes

Pollutant	Species (length and weight)	Dose, time and medium of exposure	Effect on Na^+, K^+ -ATPase activity	References
<i>Oil and dispersants</i>				
Dispersants	<i>Pal. serratus</i>	2000–15,000 mg l^{-1} ; 24 h; SW	↘ 17–30%	1
Emulsions	<i>Pal. serratus</i>	200–1000 mg l^{-1} ; 24 h; SW	↘ 22–57%	1
Oil	<i>Pal. serratus</i>	5000 mg l^{-1} ; 24–96 h; SW	↘ 46–49%	1
<i>Pesticides</i>				
DDT	<i>C. maenas</i>	0.03–3500 $\mu\text{g l}^{-1}$ (in vitro)	↗ 217–140%	2
	<i>C. maenas</i> (~ 40 g)	0.2 $\mu\text{g l}^{-1}$; 25 days; 15% SW	↘ 43.5%	3
	<i>Can. irroratus</i> (170–250 g)	0.5–4 mg l^{-1} (in vitro)	↘ 30–38%	4
		0.1 mg kg^{-1} weight (injection)	↘ 4 h: 26% 30 h: 41% 50 h: 42%	
Endosulfan 35 EC	<i>Barytelphusa guerini</i> (male, 25.0 ± 0.5 mm)	6 $\mu\text{g l}^{-1}$; 4 days; FW	↘ 15% and 22% after 72 and 96 h of exposure	5
Methoxychlor	<i>Can. magister</i> (80–100 mm CTW)	0.01 mg l^{-1} ; 14 days; S = 30–33‰	↘ 50%	6

Metals							
Cadmium	<i>Pr. clarkii</i> (~ 30.5 g)	1 mg l ⁻¹ ; 4 days; FW	no effect				7
	<i>H. americanus</i>	6 µg l ⁻¹ ; 30 days; S = 24–26‰	no effect				8
		control: 4 µg Cd g ⁻¹ dry gills contaminated: 70 µg Cd g ⁻¹ dry gills	no effect				9
	<i>Can. irroratus</i>	1, 10 mg l ⁻¹ (in vitro)	1 mg l ⁻¹ : no effect 10 mg l ⁻¹ : ↘ 37%				10
	<i>Sc. serrata</i> (~ 25 g, 4.5–5.5 cm TL)	0.3–24 mg l ⁻¹ ; 4–30 days; S = 30‰ 4 days 10 days 20 days 30 days	0.3 mg l ⁻¹	0.6 mg l ⁻¹	1.5 mg l ⁻¹	24 mg l ⁻¹ ↘ 62%	11
		↗ 11%	↘ 47%	↘ 58%			
		↘ 26%	↘ 63%	↘ 71%			
		↘ 26%	↘ 30%	↘ 45%			
	<i>Cha. granulata</i> (28.0 ± 0.7 mm CTW)	0.1, 0.25 mg l ⁻¹ ; 18 days; S = 30‰ (in vivo) 0–100 mg l ⁻¹ ; 40 min (in vitro)	no significant effect dose-dependent ↘ up to 97% with 100 mg l ⁻¹ Cd ²⁺				12
Copper	<i>A. leptodactylus</i> and <i>O. limosus</i>	0.03–3 mg l ⁻¹ (in vitro)	0.03 mg l ⁻¹ : no effect 1 mg l ⁻¹ : ↘ 60% 3 mg l ⁻¹ : ↘ 87%				13
	<i>C. maenas</i> (adult)	10 mg l ⁻¹	↘ 50%				14
Lead	<i>Ch. destructor</i> (adult at stage C, 75–100 mm)	0.5 mg l ⁻¹ ; 21 days; FW (in vivo) 120, 240 mg l ⁻¹ ; 21 days; FW (in vitro):	↘ 40% ↘ 50% and 80%, respectively				15

(continued on next page)

Table 3 (continued)

Pollutant	Species (length and weight)	Dose, time and medium of exposure	Effect on Na ⁺ ,K ⁺ -ATPase activity		References
Mercury	<i>Pr. clarkii</i> (~ 30.5 g)	100 mg l ⁻¹ ; 4 days; FW	no effect		7
	<i>Can. irroratus</i>	1, 10 mg l ⁻¹ (in vitro)	no effect		10
	<i>Pr. clarkii</i> (~ 30.5 g)	0.25 mg l ⁻¹ ; 4 days; FW	no effect		7
Zinc	<i>H. americanus</i> (476 ± 47 g)	25 mg l ⁻¹ ; 4 days; SW	↘ 67%		16
Mixture of cadmium, copper and zinc	<i>Nephrops norvegicus</i> (male: 3.63 ± 0.07 cm, female: 3.62 ± 0.10 cm)	Cd: 1, 5, 25 µg l ⁻¹ ; 18 days; SW Cu: 1, 5, 25 µg l ⁻¹ ; 18 days; SW Zn: 8, 40, 200 µg l ⁻¹ ; 18 days; SW	1, 1, 8 µg l ⁻¹ : no effect 5, 5, 40 µg l ⁻¹ : ↘ 43% 25, 25, 200 µg l ⁻¹ : ↘ 30%		17
TBTO	<i>P. japonicus</i> (juvenile at stage C, 9–12 g)	100, 200 µg l ⁻¹ ; 4 days; SW	no effect on female no effect		18
Ammonia	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 8.5–11.5 g)	48 mg N l ⁻¹ ; 2 days; S = 36‰	↗ 20.5%		19
Turbidity	<i>P. japonicus</i> (juvenile at stages C and D ₀ , 9.7 ± 1.3 g)	35 ± 15 NTU; 3 weeks; S = 15‰ and 42‰ 65 ± 15 NTU; 3 weeks; S = 15‰ and 42‰	S = 15‰ ↗ 33.5%	S = 42‰ ↗ 16.5% ↗ 35%	20

References: (1) Papineau and Le Gal, 1983; (2) Jowett and Rhead, 1978; (3) Jowett et al., 1981; (4) Neufeld and Pritchard, 1979; (5) Reddy et al., 1992; (6) Caldwell, 1974; (7) Torreblanca et al., 1989; (8) Tucker, 1979; (9) Haya et al., 1980; (10) Tucker and Matte, 1980; (11) Dhavale et al., 1988; (12) Rodriguez-Moreno et al., 1998; (13) Chaisemartin, 1973; (14) Hansen et al., 1992; (15) Ahern and Morris, 1998; (16) Haya et al., 1983; (17) Canli and Stagg, 1996; (18) Lignot et al., 1998a; (19) Lin et al., 1993; (20) Lin et al., 1992.

known to stimulate Na^+, K^+ -ATPase activity in *C. maenas* (Sommer and Mantel, 1988), while long-term activation can be due to a higher concentration of gill Na^+, K^+ -ATPase (Wheatly and Gannon, 1995). This increased activity could also result from a compensating mechanism for the impairment of ionic regulation, e.g., from an increased surface permeability (Lin et al., 1993).

2.5. Stress effects on water and ionic fluxes and surface permeability

The effects of pollutants and stress on ionic fluxes and surface permeability are presented in Table 4. In general, the permeability and net ion outflux increase whereas the water outflux and net ion influx decrease (Chaisemartin, 1973; Inman and Lockwood, 1977; Spaargaren, 1990; Harris and Coley, 1991; Wright and Welbourn, 1991; Rasmussen et al., 1995). These exchanges thus result in a decrease in hemolymph ionic concentrations and OC (Chaisemartin, 1973; Inman and Lockwood, 1977; Harris and Coley, 1991). In stress-exposed crustaceans, hemolymph circulation and movements of water around the branchial ion-transporting organs are also decreased through reduced ventilation and cardiac activity (Depledge, 1984; Nonnotte et al., 1993), therefore lowering gill perfusion and ionic influx. It has also recently been demonstrated that the cadmium inhibitions of Na^+, K^+ -ATPase and active ion influxes in *C. maenas* are directly interdependent (Postel et al., 1998). Structural lesions of the ion-transporting epithelia also likely contribute to the reduction in ion influx and to the increase in permeability. The decrease in water outflux might be related to the appearance of vacuoles and of subcuticular spaces in ionocytes. Permeability and active transport are also closely interrelated (Spaargaren, 1990). An increase in permeability also usually results in a higher passive ion outflux that would therefore require a compensating increase in Na^+, K^+ -ATPase activity, as observed in *P. japonicus* exposed to low ammonia concentrations (Lin et al., 1993). The reduction of Na^+ influx can thus be correlated to the functional disruption of Na^+, K^+ -ATPase also linked to the impairment of the ion-transporting epithelia.

3. Conclusion

It appears from the available literature that exposure to stress most often results in a disruption of ionic regulation (mostly of Na^+ and Cl^- regulation) and therefore of osmotic regulation in osmoregulating crustaceans. From the data gathered in Table 1, the ability to ionoregulate or to osmoregulate following exposure to stress was affected in 79% of the species, with a decrease of this ability in 90% of them. In cultured species, osmoregulation was disrupted in 93% of penaeid shrimp and in 100% of crayfish exposed to various stresses. Several thorough studies have shown that the effect of stress upon osmotic and ionic metabolism was time- and dose-dependent. When data are available, they most often demonstrate that the changes in iono- or osmoregulation are induced by sublethal doses of stressors.

Table 4

Effects of different stress and pollutants on water and ionic fluxes and surface permeability. Fluxes were measured in total organisms except for Chaisemartin (1973, perfused gills). Abbreviations as in Tables 1 and 2

Pollutant or stress	Species, size and weight	Dose, time, medium of exposure	Effect on water and ionic fluxes and surface permeability	References
<i>Pesticides</i>				
Lindane	<i>G. duebeni</i> (50–80 mg)	1.8 $\mu\text{g l}^{-1}$; 2 days; S = SW and 2% SW	After transfer in 2% SW: Na ⁺ influx: $\searrow$ 28% Na ⁺ outflux: $\searrow$ 17%	1
Aroclor 1254 (PCBs)	<i>Pa. pugio</i>	31–51 $\mu\text{g l}^{-1}$; 4 days; S = 2, 17 and 32‰	Cl [−] outflux: no effect at S = 2‰, 17‰ and 32‰ Water outflux: no effect at S = 2, 32‰ but $\searrow$ 66% at S = 17‰	2
<i>Metals</i>				
Cadmium	<i>C. maenas</i> (31–41 mm CTW)	4, 8 mg l^{-1} ; 5 days; S = 14‰	Water outflux: $\searrow$ 68% (4 mg l^{-1}) and 71% (8 mg l^{-1})	3
Copper	<i>A. leptodactylus</i> , <i>O. limosus</i>	0.3 mg l^{-1} in incubation medium of perfused gills	Na ⁺ influx: $\searrow$ Na ⁺ outflux: $\nearrow$	4
Lead	<i>C. maenas</i> (31–41 mm CTW)	1 mg l^{-1} ; 1 h; S = 15‰	Water outflux: $\searrow$ 62%	3
	<i>Ch. destructor</i> (75–100 mm)	0.5 mg l^{-1} ; 21 days; FW	Na ⁺ influx: $\searrow$ 50% Na ⁺ outflux: no effect	5

Methylmercury	<i>G. duebeni</i> (50–80 mg)	100 $\mu\text{g l}^{-1}$; 2 days; S = SW and 2% SW	After transfer in 2% SW: Na ⁺ influx: $\searrow$ 20% Na ⁺ outflux: $\searrow$ 14%	1
	<i>As. aquaticus</i>	8–1000 $\mu\text{g l}^{-1}$; 2 h; FW	Na ⁺ influx: $\searrow$ 56–79% Ca ²⁺ influx: $\searrow$ 67–72%	6
Mercury	<i>As. aquaticus</i>	8–1000 $\mu\text{g l}^{-1}$; 2 h; FW	Na ⁺ influx: $\searrow$ 50–56% Ca ²⁺ influx: $\searrow$ 43–78%	6
	<i>C. maenas</i> (31–41 mm CTW)	1 mg l^{-1} ; 1 h; S = 15‰	Water outflux: $\searrow$ 71%	3
Zinc	<i>Cr. crangon</i> (36–47 mm TL)	500 $\mu\text{g l}^{-1}$; 1, 3 days; S = 15‰	Water outflux: $\searrow$ $\sim$ 82% (1 day) and $\sim$ 76% (3 days)	3
	<i>Pac. leniusculus</i> (33.0 $\pm$ 1.1 g)	46 mg l^{-1}	Cl [−] influx: $\searrow$ 41% Cl [−] outflux: $\searrow$ 20%	7
Ammonia	<i>A. pallipes</i> (10 g)	0.2–1 mg l^{-1} ; FW	Na ⁺ influx: $\searrow$ 20–80%	8
	<i>C. maenas</i> (15–60 g)	3.6–36 mg l^{-1} ; 2–10 min; S = 24‰	ionic surface permeability: $\nearrow$ net salt outflux: $\nearrow$	9
Low pH	<i>A. pallipes</i> (10 g)	pH 4.0	Na ⁺ influx: $\searrow$ 70–80%	8

References: (1) Inman and Lockwood, 1977; (2) Roesijadi et al., 1976; (3) Rasmussen et al., 1995; (4) Chaisemartin, 1973; (5) Ahern and Morris, 1998; (6) Wright and Welbourn, 1991; (7) Harris and Coley, 1991; (8) Shaw, 1960; (9) Spaargaren, 1990.

The disruption of osmoregulation due to stressors originates from physiological and cellular alterations. The activity of Na^+, K^+ -ATPase was — most often negatively — affected in 74% of species exposed to stressors. Water and ion fluxes were also altered in all reported cases. Stressors also induced different types of damage to the transporting epithelia located in the branchial chambers and/or the excretory organs, with different levels of cellular alterations affecting the ionocytes of these organs.

It therefore appears that an integrated series of events occurs in most crustaceans following exposure to a supra-limnal, but sublethal, level of stressor. According to the physico-chemical nature of the stressor, it can induce deleterious effects on the cellular structure of ion-transporting epithelia and/or variations in the activity of the enzyme Na^+, K^+ -ATPase. These events lead to alterations in water and ionic fluxes and surface permeability, finally resulting in disruption of ionic regulation and thus of osmotic regulation.

As underlined above, the effect of stressors on osmoregulation is usually detectable at sublethal levels, often before any mortality is noticeable among exposed animals. The variations in osmotic and ionic regulation, and thus in OC and/or Na^+ or Cl^- regulation, can therefore be considered as an early warning of sublethal stress in crustaceans.

We thus confirm that one of these parameters, in particular OC, can be used as a reliable biomarker to monitor the physiological condition and the effect of stressors in osmoregulating crustaceans (Charmantier et al., 1989; Young-Lai et al., 1991; Lin et al., 1992, 1993; Bambang et al., 1995a,b; Charmantier et al., 1994; Lignot et al., 1997, 1998a,b). From experimental studies conducted under laboratory conditions, the use of OC as a stress biomarker has been successfully applied in penaeid shrimps under pilot medium-scale culturing conditions (Lignot et al., 1999b) and has begun to be used in aquaculture farms (Cochard et al., 1997).

Future research might concentrate in several directions. Further stressors should be tested on the osmoregulatory ability of diverse species of crustaceans, leading to an improved knowledge and understanding of the effect of different stress on the physiology of ionic and osmotic regulation. The cellular and molecular effects of low levels of stressors should be studied through electron and confocal microscopy, immunocytochemistry, etc..., and through molecular approaches, to detect their effect on the specific structures of ionocytes (apical microvilli, basolateral infoldings, mitochondria), and the location, activity and regulation of Na^+, K^+ -ATPase and of other enzymes involved in osmoregulation such as carbonic anhydrase and V-ATPases. The determination of OC variations can also be used to assay endocrine and neuroendocrine factors susceptible to control osmotic and ionic regulation (McNamara et al., 1991; Freire and McNamara, 1992; Charmantier-Daures et al., 1994). Another direction of research should be to continue studies in cultured species, mostly in penaeid shrimp, and also in crayfish and *Macrobrachium* species. Their objective would concentrate on the practical use of OC as a biomarker. Some types of stress, in particular pathological stress, have been seldom studied as yet (Maestracci and Vey, 1989; Souheil et al., 1999). The effects on osmoregulation of viral, bacterial and fungal infections affecting cultured species should be studied, with the aim of also using OC as an early warning of these fast-spreading diseases.

References

- Ahern, M., Morris, S., 1998. Accumulation of lead and its effects on Na balance in the freshwater crayfish *Cherax destructor*. J. Exp. Zool. 281, 270–279.
- Allan, G.L., Maguire, G.B., 1992. Effects of pH and salinity on survival, growth and osmoregulation in *Penaeus monodon* Fabricius. Aquaculture 107, 33–47.
- Andersen, J.T., Baatrup, E., 1988. Ultrastructural localization of mercury accumulations in the gills, hepatopancreas, midgut, and antennal glands of the brown shrimp, *Crangon crangon*. Aquat. Toxicol. 13, 309–324.
- Anderson, J.W., Neff, J.M., Petrocelli, S.R., 1974. Sublethal effects of oil, heavy metals and PCBs on marine organisms. In: Khan, M.A.Q., Bederka, J.P. Jr. (Eds.), Survival in Toxic Environments. Academic Press, New York, pp. 83–121.
- Appelberg, M., 1985. Changes in haemolymph ion concentrations of *Astacus astacus* L. and *Pacifastacus leniusculus* (Dana) after exposure to low pH and aluminium. Hydrobiology 121, 19–25.
- Appelberg, M., 1987. Some factors regulating the crayfish *Astacus astacus* L. in acid and neutralized waters. Ecophysiology of Acid Stress in Aquatic Organisms. In: Witters, H., Vanderborgh, O. (Eds.), Ann. Soc. R. Zool. Belg. 117 pp. 167–179, Suppl. 1.
- Bambang, Y., Charmantier, G., Thuét, P., Trilles, J.-P., 1995a. Effects of cadmium on survival and osmoregulation of various developmental stages of the shrimp *Penaeus japonicus* Bate (Crustacea: Decapoda). Mar. Biol. 123, 443–450.
- Bambang, Y., Thuét, P., Charmantier-Daures, M., Trilles, J.-P., Charmantier, G., 1995b. Effects of copper on survival and osmoregulation of various developmental stages of the shrimp *Penaeus japonicus* Bate (Crustacea, Decapoda). Aquat. Toxicol. 33, 125–139.
- Bamber, S.D., Depledge, M.H., 1997. Responses of shore crabs to physiological challenges following exposure to selected environmental contaminants. Aquat. Toxicol. 40, 79–92.
- Baticados, M.C.L., Tendencia, E.A., 1991. Effects of Gusathion A on the survival and shell quality of juvenile *Penaeus monodon*. Aquaculture 93, 9–19.
- Bayne, B.L., 1975. Aspects of physiological condition in *Mytilus edulis* L., with respect of the effects of oxygen tension and salinity. Proc. Ninth Eur. Mar. Biol., 213–238.
- Bayne, B.L., 1985. Responses to environmental stress: tolerance, resistance and adaptation. In: Gray, J.S., Christiansen, M.E. (Eds.), Marine Biology of Polar Regions and Effects of Stress on Marine Organisms. Wiley, New York, pp. 331–347.
- Berridge, M.J., Oschman, J.O., 1972. Transporting Epithelia. Academic Press, New York, 91 pp.
- Bjerregaard, P., 1991. Relationship between physiological condition and cadmium accumulation in *Carcinus maenas* (L.). Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol. 99, 75–83.
- Bjerregaard, P., Vislie, T., 1985. Effects of mercury on ion and osmoregulation in the shore crab *Carcinus maenas* (L.). Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol. 82, 227–230.
- Bjerregaard, P., Vislie, T., 1986. Effect of copper on ion- and osmoregulation in the shore crab *Carcinus maenas*. Mar. Biol. 91, 69–76.
- Boitel, F., Truchot, J.-P., 1989. Effects of sublethal and lethal copper levels on hemolymph ion concentrations in the shore crab *Carcinus maenas* kept in undiluted seawater. Mar. Biol. 103, 495–501.
- Boitel, F., Truchot, J.-P., 1990. Comparative study of the effects of copper on hemolymph ion concentrations and acid–base balance in the shore crabs *Carcinus maenas* acclimated to full-strength or dilute seawater. Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol. 95, 307–312.
- Bouaricha, N., Charmantier-Daures, M., Thuét, P., Trilles, J.-P., Charmantier, G., 1994. Ontogeny of osmoregulatory structures in the shrimp *Penaeus japonicus* (Crustacea, Decapoda). Biol. Bull. 186, 29–40.
- Broom, D.M., 1991. Assessing welfare and suffering. Behav. Proc. 25, 117–123.
- Bubel, A., 1976. Histological and electron microscopical observations on the effects of different salinities and heavy metal ions, on the gills of *Jaera nordmanni* (Rathke) (Crustacea, Isopoda). Cell Tissue Res. 167, 65–95.
- Bursey, C.R., Lane, C.E., 1971. Ionic and protein concentration changes during the molt cycle of *Penaeus duorarum*. Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol. 40, 155–162.
- Burton, R.F., 1986. Ionic regulation in crustacea: the influence of temperature on apparent set points. Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol. 84, 135–139.

- Cajaraville, M.P., Abascal, I., Etxeberria, M., Marigomez, I., 1995. Lysosomes as cellular markers of environmental pollution: time- and dose-dependent responses of the digestive lysosomal system of mussels after petroleum hydrocarbon exposure. *Environ. Toxicol. Water. Qual.* 10, 1–8.
- Caldwell, R.S., 1974. Osmotic and ionic regulation in Decapod Crustacea exposed to methoxychlor. In: Vernberg, F.J., Vernberg, W.B. (Eds.), *Pollution and Physiology of Marine Organisms*. Academic Press, New York, pp. 197–223.
- Canli, M., Stagg, R.M., 1996. The effects of in vivo exposure to cadmium, copper and zinc on the activities of gill ATPases in the Norway lobster, *Nephrops norvegicus*. *Arch. Environ. Contam. Toxicol.* 31, 494–501.
- Castille, F.L. Jr., Lawrence, A.L., 1981a. The effect of salinity on the osmotic, sodium and chloride concentrations in the hemolymph of euryhaline shrimp of the genus *Penaeus*. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 68, 75–80.
- Castille, F.L. Jr., Lawrence, A.L., 1981b. A comparison of the capabilities of juvenile and adult *Penaeus setiferus* and *Penaeus stylirostris* to regulate the osmotic, sodium, and chloride concentrations in the hemolymph. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 68, 677–680.
- Chaisemartin, C., 1973. Analyse comparée de la toxicité du Cu^{2+} chez *Astacus leptodactylus* et *Orconectes limosus*: dépression des fonctions ionorégulatrices et de l'activité de l'ATPase Na–K dépendante du système branchial. *C. R. Soc. Biol.* 167, 324–328.
- Chandy, J.P., Kolwalkar, D.G., 1984. Histological changes in the gill and hepatopancreas of the marine crabs *Charybdis lucifera* (Fabricius) and *Scylla serrata* (Forsk.) exposed to crude oil emulsion. *Indian J. Mar. Sci.* 13, 10–13.
- Charmantier, G., 1975. Variations saisonnières des capacités ionorégulatrices de *Sphaeroma serratum* (Fabricius, 1787) (Crustacea, Isopoda, Flabellifera). *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 50, 339–345.
- Charmantier, G., 1987. L'osmorégulation chez les crevettes Penaeidae. *Oceanis* 13, 179–196.
- Charmantier, G., 1998. Ontogeny of osmoregulation in crustaceans; a review. *Invertebr. Reprod. Dev.* 33, 177–190.
- Charmantier, G., Charmantier-Daures, M., Young-Lai, W.W., 1985. Lethal and sublethal effects of potash brine on different stages of the lobster, *Homarus americanus*. *Can. Tech. Rep. Fish. Aquat. Sci.* 1344, 1–13.
- Charmantier, G., Charmantier-Daures, M., Bouaricha, N., Thuet, P., Aiken, D.E., Trilles, J.-P., 1988. Ontogeny of osmoregulation and salinity tolerance in two decapod crustaceans: *Homarus americanus* and *Penaeus japonicus*. *Biol. Bull.* 175, 102–110.
- Charmantier, G., Bouaricha, N., Charmantier-Daures, M., Thuet, P., Trilles, J.-P., 1989. Salinity tolerance and osmoregulatory capacity as indicators of the physiological state of penaeid shrimps. *Eur. Aquat. Soc. Spec. Publ.* 10, 65–66.
- Charmantier, G., Soyeux, C., Aquacop, 1994. Effect of molt stage and hypoxia on osmoregulatory capacity in the penaeid shrimp *Penaeus vannamei*. *J. Exp. Mar. Biol. Ecol.* 178, 233–246.
- Charmantier-Daures, M., Thuet, P., Charmantier, G., Trilles, J.-P., 1988. Tolérance à la salinité et osmorégulation chez les post-larves de *Penaeus japonicus* et *P. chinensis*, Effet de la température. *Aquat. Living Resour.* 1, 267–276.
- Charmantier-Daures, M., Charmantier, G., Janssen, K.P.C., Aiken, D.E., Van Herp, F., 1994. Involvement of eyestalk factors in the neuroendocrine control of osmoregulation in adult American lobster *Homarus americanus*. *Gen. Comp. Endocrinol.* 94, 281–293.
- Chen, H.C., Slinn, J.D., 1980. Osmoregulation of the prawn *Palaemon elegans* exposed to some heavy metals. *J. Fish. Soc. Taiwan* 7, 1–13.
- Chen, J.C., Cheng, S.Y., 1996. Hemolymph osmolality, acid–base balance, and ammonia excretion of *Penaeus japonicus* Bate exposed to ambient nitrite. *Arch. Environ. Contam. Toxicol.* 30, 151–155.
- Chen, J.C., Lee, Y., 1997. Effects of nitrite exposure on acid–base balance, respiratory protein, and ion concentrations of giant freshwater prawn *Macrobrachium rosenbergii* at low pH. *Arch. Environ. Contam. Toxicol.* 33, 290–297.
- Cioffi, M., 1984. Comparative ultrastructure of arthropod transporting epithelia. *Am. Zool.* 24, 139–156.
- Clarke, W.C., Blackburn, J., 1977. A seawater challenge test to measure smolting of juvenile salmon. *Fish. Mar. Serv. Res. Dev. Tech. Rep.* 705, 1–11.
- Cochard, J.-C., Lemonier, H., Bernard, E., Boglio, E., Goarant, C., 1997. Expériences 97 sur l'influence des

- sédiments du bassin d'élevage sur la physiologie de *Penaeus stylirostris*. IFREMER Tech. Rep., (DRV/AQ/TAH 97.42), 19 pp.
- Compère, P., Wanson, S., Péqueux, A., Gilles, R., Goffinet, G., 1989. Ultrastructural changes in the gill epithelium of the green crab *Carcinus maenas* in relation to the external salinity. *Tissue Cell* 21, 299–318.
- Couch, J.A., 1977. Ultrastructural study of lesions in gills of a marine shrimp exposed to cadmium. *J. Invertebr. Pathol.* 29, 267–288.
- Dall, W., 1970. Osmoregulation in the lobster *Homarus americanus*. *J. Fish. Res. Board Can.* 27, 1123–1130.
- Dall, W., 1981. Osmoregulatory ability and juvenile habitat preference in some penaeid prawns. *J. Exp. Mar. Biol. Ecol.* 54, 55–64.
- Depledge, M.H., 1984. Disruption of circulatory and respiratory activity in the shore crabs (*Carcinus maenas* (L.)) exposed to heavy metal pollution. *Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol.* 78, 445–459.
- Dhavalé, D.M., Masurekar, V.B., Giridhar, B.A., 1988. Cadmium induced inhibition of $\text{Na}^+ - \text{K}^+$ ATPase activity in tissues of crab *Scylla serrata* (Forsk.). *Bull. Environ. Contam. Toxicol.* 40, 759–763.
- Diamond, D.W., Scott, L.K., Forward, R.B. Jr., Kirby-Smith, W., 1989. Respiration and osmoregulation of the estuarine crab, *Rhithropanopeus harrisii* (Gould); effects of the herbicide, Alachlor. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 93, 313–318.
- Doughtie, D.G., Rao, K.R., 1983a. Ultrastructural and histological study of degenerative changes leading to black gills in grass shrimp exposed to a dithiocarbonate biocide. *J. Invertebr. Pathol.* 41, 33–50.
- Doughtie, D.G., Rao, K.R., 1983b. Ultrastructural and histological study of degenerative changes in the antennal glands, hepatopancreas and midgut of grass shrimp exposed to two dithiocarbonate biocides. *J. Invertebr. Pathol.* 41, 281–300.
- Doughtie, D.G., Rao, K.R., 1984. Histopathological and ultrastructural changes in the antennal gland, midgut, hepatopancreas and gill of grass shrimp following exposure to hexavalent chromium. *J. Invertebr. Pathol.* 43, 89–108.
- Eddy, F.B., 1981. Effects of stress on osmotic and ionic regulation in fish. In: Pickering, A.D. (Ed.), *Stress and Fish*. Academic Press, London, pp. 77–102.
- Einarson, S., 1993. Effects of temperature, seawater osmolality and season on oxygen consumption and osmoregulation of the amphipod *Gammarus oceanicus*. *Mar. Biol.* 117, 599–606.
- Ellis, B.A., Morris, S., 1995. Effects of extreme pH on the physiology of the Australian “yabby” *Cherax destructor*: acute and chronic changes in haemolymph carbon dioxide, acid–base and ionic status. *J. Exp. Biol.* 198, 397–407.
- Ellis, R.W., Fuller, E.G., 1979. Mercury interaction in crayfish antennal gland and hepatopancreas. *J. Invertebr. Pathol.* 34, 308–309.
- Exbrayat, J.M., Bourguet, J.P., 1982. Variations du milieu intérieur de *Penaeus japonicus* en fonction de certaines conditions naturelles (cycle de mue et âge) et de certaines conditions expérimentales (épédonculation et ablation des organes Y). *Bull. Soc. Zool. Fr.* 107, 33–51.
- Ferraris, R.P., Parado-Estépa, E.G., De Jesus, E.G., Ladja, J.M., 1987. Osmotic and chloride regulation in the hemolymph of the tiger prawn *Penaeus monodon* during molting in various salinity. *Mar. Biol.* 95, 377–385.
- Fjeld, E., Hessen, D.O., Roos, N., Taugbol, T., 1988. Changes in gill ultrastructure and haemolymph chloride concentrations in the crayfish, *Astacus astacus*, exposed to de-acidified aluminium-rich water. *Aquaculture* 72, 139–150.
- Freire, C.A., McNamara, J.C., 1992. Involvement of the central nervous system in neuroendocrine mediation of osmotic and ionic regulation in the freshwater shrimp *Macrobrachium olfersii* (Crustacea, Decapoda). *Gen. Comp. Endocrinol.* 88, 316–327.
- Freire, C.A., McNamara, J.C., 1995. Fine structure of the gills of the fresh-water shrimp *Macrobrachium olfersii* (Crustacea): effect of acclimation to high salinity medium and evidence for involvement of the intralamellar septum in ion uptake. *J. Crustacean Biol.* 15, 103–116.
- Fritzsche, H., 1916. Studien über die Schwankungen des osmotischen Druckes der Körperflüssigkeiten bei *Daphnia magna*. *Int. Rev. Hydrobiol.* 8, 22–80 and 125–203.
- Ghate, H.V., Mulherkar, L., 1979. Histological changes in the gills of two freshwater prawn species exposed to copper sulfate. *Indian J. Exp. Biol.* 17, 838–840.

- Giesy, J.P., Graney, R.L., 1989. Recent developments in and comparisons of acute and chronic bioassays and indicators. *Hydrobiology* 188/189, 21–60.
- Giesy, J.P., Versteeg, D.J., Graney, R.L., 1988. A review of selected clinical indicators of stress-induced changes in aquatic organisms. In: Evans, M.S. (Ed.), *Toxic Contaminants and Ecosystem Health: A Great Lakes Focus*. Wiley, New York, pp. 169–200.
- Gilles, R., Péqueux, A., 1983. Interactions of chemical and osmotic regulation with the environment. The Biology of Crustacea. In: Bliss, D.E., Vernberg, F.J., Vernberg, W.B. (Eds.), *Environmental Adaptations* vol. 8 Academic Press, New York, pp. 109–177.
- Guerin, J.L., Stickle, B., 1995. Effects of cadmium on survival, osmoregulatory ability and bioenergetics of juvenile blue crab *Callinectes sapidus* at different salinities. *Mar. Environ. Res.* 40, 227–246.
- Hagerman, L., Uglow, R.F., 1981. Ventilatory behaviour and chloride regulation in relation to oxygen tension in the shrimp *Palaemon adspersus* Rathke maintained in hypotonic medium. *Ophelia* 20, 193–200.
- Hagerman, L., Uglow, R.F., 1982. Effects of hypoxia on osmotic and ionic regulation in the brown shrimp *Crangon crangon* (L.) from brackish water. *J. Exp. Mar. Biol. Ecol.* 63, 93–104.
- Hansen, J.I., Mustafa, T., Depledge, M., 1992. Mechanisms of copper toxicity in the shore crab, *Carcinus maenas*: II. Effects on key enzymes, metabolites and energy charge potential. *Mar. Biol.* 114, 259–264.
- Haond, C., Flik, G., Charmantier, G., 1998. Confocal laser scanning and electron microscopical studies on osmoregulatory epithelia in the branchial cavity of the lobster *Homarus gammarus*. *J. Exp. Biol.* 201, 1817–1833.
- Harris, R.R., Coley, S., 1991. The effects of nitrite on chloride regulation in the crayfish *Pacifastacus leniusculus* Dana (Crustacea: Decapoda). *J. Comp. Physiol., B* 161, 199–206.
- Havas, M., Hutchinson, T., 1983. Effect of low pH on the chemical composition of aquatic invertebrates from tundra ponds at the Smoking hills, N.W.T., Canada. *Can. J. Zool.* 61, 241–249.
- Haya, K., Waiwood, B.A., 1983. Adenylate energy charge and ATPase activity: potential biochemical indicators of sublethal effects caused by pollutants in aquatic animals. In: Nriagu, J.O. (Ed.), *Aquatic Toxicology*. Wiley, New York, pp. 307–333.
- Haya, K., Johnston, C.E., Waiwood, B.A., 1980. Adenylate energy charge and ATPase activity in American lobster (*Homarus americanus*) from Belledune Harbour. Cadmium Pollution of Belledune Harbour, New Brunswick, Canada. *Can. Tech. Rep. Fish. Aquat. Sci.* 963, 85–91.
- Haya, K., Waiwood, B.A., Johnston, D.W., 1983. Adenylate energy charge and ATPase activity of lobster (*Homarus americanus*) during sublethal exposure to zinc. *Aquat. Toxicol.* 3, 115–126.
- Hebel, D.K., Jones, M.B., Depledge, M.H., 1997. Responses of crustaceans to contaminant exposure: a holistic approach. *Estuarine, Coastal Shelf Sci.* 44, 177–184.
- Holdway, D.A., Brennan, S.E., Ahokas, J.T., 1995. Short review of selected fish biomarkers of xenobiotic exposure with an example using fish hepatic mixed-function oxidase. *Aust. J. Ecol.* 20, 34–44.
- Hubschman, J.H., 1965. Effects of copper on the crayfish *Orconectes rusticus* (Girard): II. Mode of toxic action. *Crustaceana* 12, 141–150.
- Inman, C.B.E., Lockwood, A.P.M., 1977. Some effects of methylmercury and lindane on sodium regulation in the Amphipod *Gammarus duebeni* during changes in the salinity of its medium. *Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol.* 58, 67–75.
- Ivanovici, A.M., Wiebe, W.J., 1981. Towards a working “definition” of “stress”: a review and critique. In: Barrett, G.W., Rosenberg, R. (Eds.), *Stress Effects on Natural Ecosystems*. Wiley, New York, pp. 13–27.
- Jeberg, M.V., Jensen, F.B., 1994. Extracellular and intracellular ionic changes in crayfish *Astacus astacus* exposed to nitrite at two acclimation temperatures. *Aquat. Toxicol.* 29, 65–72.
- Jensen, F.B., 1990. Sublethal physiological changes in freshwater crayfish, *Astacus astacus*, exposed to nitrite: haemolymph and muscle tissue electrolyte status, and haemolymph acid–base balance and gas transport. *Aquat. Toxicol.* 18, 51–60.
- Jensen, F.B., Malte, H., 1990. Acid–base and electrolyte regulation, and haemolymph gas transport in crayfish, *Astacus astacus*, exposed to soft, acid water with and without aluminium. *J. Comp. Physiol., B* 160, 483–490.
- Johnson, I.T., Jones, M.B., 1990. Effect of zinc on osmoregulation of *Gammarus duebeni* (Crustacea, Amphipoda) from the estuary and the sewage treatment works at Love, Cornwall. *Ophelia* 31, 187–196.
- Johnson, I., Uglow, R.F., 1987. The effect of hypoxia on ion regulation and acid–base balance in *Carcinus maenas*. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 86, 261–268.

- Jones, M.B., 1975. Synergistic effects of salinity, temperature and heavy metals on mortality and osmoregulation in marine and estuarine isopods (Crustacea). *Mar. Biol.* 30, 13–20.
- Jowett, P.E., Rhead, M.M., 1978. In vitro changes in the activity of ATPases in the gills of *Carcinus maenas* exposed to various concentrations of *p,p'*-DDT. *Environ. Pollut.* 17, 1–6.
- Jowett, P.E., Rhead, M.M., Bayne, B.L., 1981. In vivo changes in the activity of gill ATPases and haemolymph ions of *Carcinus maenas* exposed to *p,p'*-DDT and reduced salinities. *Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol.* 69, 399–402.
- Kelley, B.J. Jr., Burbanck, W.D., 1972. Osmoregulation in juvenile and adult *Cyathura polita* subjected to salinity changes and ionizing γ irradiation. *Chesapeake Sci.* 13, 201–205.
- Krogh, A., 1939. *Osmotic Regulation in Aquatic Animals*. Cambridge Univ. Press, reprinted (1965) by Dover, New York, 242 pp.
- Lake, P.S., Thorp, V.J., 1974. The gill lamellae of the shrimp *Paratya tasmaniensis* (Atyidae: Crustacea). Normal ultrastructure and changes with low levels of cadmium. *Proc. Eight Int. Congr. Electron Microsc., Canberra* 2, 448–449.
- Langy, S., 1991. Effets d'un insecticide organophosphoré, le Fénitrothion, sur la physiologie de la crevette *Penaeus japonicus* (Crustacé, Décapode). DES, Université Montpellier 2, 70 pp.
- Lawson, S.L., Jones, M.B., Moate, R.M., 1995. Effect of copper on the ultrastructure of the gill epithelium of *Carcinus maenas* (Decapoda: Brachyura). *Mar. Pollut. Bull.* 31, 63–72.
- Leffler, C.W., 1975. Effects of ingested Mirex and DDT on juvenile *Callinectes sapidus* Rathbun. *Environ. Pollut.* 8, 283–299.
- Lignot, J.-H., Trilles, J.-P., Charmantier, G., 1997. Effects of an organophosphorus insecticide, fenitrothion, on survival and osmoregulation of various developmental stages of the shrimp *Penaeus japonicus* (Crustacea: Decapoda). *Mar. Biol.* 128, 307–316.
- Lignot, J.-H., Pannier, F., Trilles, J.-P., Charmantier, G., 1998a. Effects of tributyltin oxide on survival and osmoregulation of various developmental stages of the shrimp *Penaeus japonicus* (Crustacea: Decapoda). *Aquat. Toxicol.* 41, 277–299.
- Lignot, J.-H., Cochard, J.-C., Charmantier, G., 1998b. Effects of fenitrothion, on survival, osmoregulation and acetylcholinesterase activity of *Penaeus stylirostris* and *P. vannamei* (Crustacea, Decapoda). *J. Shellfish Res.*, in press.
- Lignot, J.-H., Charmantier-Daures, M., Charmantier, G., 1999a. Localization of the Na^+ , K^+ -ATPase in the branchial cavity of juvenile lobster, *Homarus gammarus* (Crustacea, Decapoda). *Cell Tissue Res.*, in press.
- Lignot, J.-H., Cochard, J.-C., Soyez, C., Lemaire, P., Charmantier, G., 1999b. Haemolymph osmolality according to nutritional status, molting stage and body weight of *Penaeus stylirostris*. *Aquaculture* 170, 79–92.
- Lin, H.P., Charmantier, G., Trilles, J.-P., 1991. Toxicité de l'ammonium au cours du développement postembryonnaire de *Penaeus japonicus* (Crustacea, Decapoda). Effets sur l'osmorégulation. *C. R. Acad. Sci. Paris* 312, 99–105.
- Lin, H.P., Charmantier, G., Thuet, P., Trilles, J.-P., 1992. Effects of turbidity on survival, osmoregulation and gill Na^+ – K^+ ATPase in the shrimp *Penaeus japonicus*. *Mar. Ecol.: Prog. Ser.* 90, 31–37.
- Lin, H.P., Thuet, P., Trilles, J.-P., Mounet-Guillaume, R., Charmantier, G., 1993. Effects of ammonia on survival and osmoregulation at various developmental stages of the shrimp *Penaeus japonicus*. *Mar. Biol.* 117, 591–598.
- Livingstone, D.R., 1993. Biotechnology and pollution monitoring: use of molecular biomarkers in the aquatic environment. *J. Chem. Tech. Biotechnol.* 57, 195–211.
- Lockwood, A.P.M., 1977. Transport and osmoregulation in Crustacea. In: Gupta, B.L., Moreton, R.B., Oschman, J.L., Wall, B.J. (Eds.), *Transport of Ions and Water in Animals*. Academic Press, New York, pp. 670–707.
- Lucu, C., 1990. Ionic regulatory mechanisms in crustacean gill epithelia. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 97, 297–306.
- Lucu, C., 1993. Ion transport in the gill epithelium of aquatic crustacea. *J. Exp. Zool.* 265, 378–386.
- Malins, D.C., Ostrander, G.K., 1991. Perspective in aquatic toxicology. *Annu. Rev. Pharmacol. Toxicol.* 31, 371–399.
- Mantel, L.H., Farmer, L.L., 1983. Osmotic and ionic regulation. The Biology of Crustacea. In: Bliss, D.E.,

- Mantel, L.H. (Eds.), Internal Anatomy and Physiological Regulation vol. 5 Academic Press, New York, pp. 53–161.
- Mayer, F.L., Versteeg, D.J., McKee, M.J., Folmar, L.C., McCume, D.C., Rattner, B.A., 1992. Physiological and Nonspecific Biomarkers. Biomarkers. Biochemical, Physiological, and Histological Markers of Anthropogenic Stress. In: Huggett, R.J., Kimerle, R.A., Merhle, P.M. Jr., Bergam, H.L. (Eds.), SETAC Spec. Publ. Ser. Lewis Publishers, Boca Raton, FL, pp. 5–85.
- McLusky, D., 1970. Osmoregulation in *Corophium volutator*. The effect of starvation. Comp. Biochem. Physiol. 35, 303–306.
- McLusky, D., Hagerman, L., 1987. The toxicity of chromium, nickel and zinc: effects of salinity and temperature, and the osmoregulatory consequences in the mysid *Praunus flexuosus*. Aquat. Toxicol. 10, 225–238.
- McMahon, B.R., Doyle, J.E., 1997. Acid exposure in euryhaline environments: ion regulation and acid tolerance in larval and adult *Artemia franciscana*. Hydrobiologia 352, 1–7.
- McMahon, B.R., Stuart, S.A., 1989. The physiological problems of crayfish in acid waters. In: Morris, R., Taylor, E.W., Brown, D.J.A., Brown, J.A. (Eds.), Acid Toxicity and Aquatic Animals. Cambridge Univ. Press, Cambridge, pp. 171–199.
- McNamara, J.C., Salomo, L.C., Ribeiro, E.A., 1991. Neuroendocrine regulation of osmotic and ionic concentrations in the hemolymph of the freshwater shrimp *Macrobrachium olfersii* (Wiegmann) (Crustacea, Decapoda). Gen. Comp. Endocrinol. 84, 16–26.
- Maestracci, V., Vey, A., 1989. Fungal infection of gills in crayfish: histological, cytological and physiopathological aspects of the disease. Freshwater Crayfish 7, 187–194.
- Morgan, D.O., McMahon, B.R., 1982. Acid tolerance and effects of sublethal acid exposure on iono-regulation and acid–base status in two crayfish *Procambarus clarkii* and *Orconectes rusticus*. J. Exp. Biol. 97, 241–252.
- Morritt, D., Spicer, J.I., 1995. Changes in pattern of osmoregulation in the brackish water amphipod *Gammarus duebeni* Lilljeborg (Crustacea) during embryonic development. J. Exp. Zool. 273, 271–281.
- Neufeld, G.J., Pritchard, J.B., 1979. Osmoregulation and gill Na,K-ATPase in the rock crab, *Cancer irroratus*: response to DDT. Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol. 62, 165–172.
- Nimmo, D.R., Bahner, L.H., 1974. Some physiological consequences of polychlorinated biphenyl- and salinity-stress in penaeid shrimp. In: Vernberg, F.J., Vernberg, W.B. (Eds.), Pollution and Physiology of Marine Organisms. Academic Press, New York, pp. 427–443.
- Nimmo, D.R., Lightner, D.V., Bahner, L.H., 1977. Effect of cadmium on the shrimp *Penaeus duorarum*, *Palaemonetes pugio* and *Palaemonetes vulgaris*. In: Vernberg, F.J., Calabrese, A., Thurberg, F.P., Vernberg, W.B. (Eds.), Physiological Responses of Marine Biota to Pollutants. Academic Press, New York, pp. 131–183.
- Nonnotte, L., Boitel, F., Truchot, J.-P., 1993. Waterborne copper causes gill damage and hemolymph hypoxia in the shore crab *Carcinus maenas*. Can. J. Zool. 71, 1569–1576.
- Oksama, M., Kristoffersson, R., 1980. Effects of phenol and 4-chlorophenol on ionic regulation in *Mesidotea entomon* (Crustacea) in brackish water. Ann. Zool. Fenn. 17, 243–247.
- Pannikar, N.K., 1968. Osmotic behaviour of shrimps and prawns in relation to their biology and culture. FAO Fish. Rep. 57, 527–538.
- Papathanassiou, E., 1985. Effects of cadmium ions on the ultrastructure of the gill cells of the brown shrimp *Crangon crangon* (L.) (Decapoda, Caridea). Crustaceana 48, 6–17.
- Papathanassiou, E., King, P.E., 1983. Ultrastructural studies on the gills of *Palaemon serratus* (Pennant) in relation to cadmium accumulation. Aquat. Toxicol. 3, 273–284.
- Papineau, C., Cheze, G., 1984. Modifications histopathologiques des branchies de la crevette (*Palaemon serratus*) soumise à l'effet des dispersants et émulsions pétrolières. Cah. Biol. Mar. 25, 75–81.
- Papineau, C., Le Gal, Y., 1983. Effet subléthal des dispersants et des émulsions pétrolières sur l'ATPase des branchies de *Palaemon serratus*. Rev. Int. Oceanogr. Med. 70–71, 39–47.
- Patil, H.S., Kaliwal, M.B., 1989. Histopathological effects of zinc on the gills of the prawn *Macrobrachium kendersodanum*. Z. Angew. Zool. 76, 505–509.
- Pawar, K.R., Katdare, M., 1982. Gill lesions in fresh water prawn *Macrobrachium kistnensis* (Tiwari) after intoxication with organo-phosphate insecticide Sumithion. Pol. Res. 1, 59–62.

- Payne, J.F., Kiceniuk, J., Misra, R., Fletcher, G., Thompson, R., 1983. Sublethal effects of petroleum hydrocarbons on adult American lobsters (*Homarus americanus*). Can. J. Fish. Aquat. Sci. 40, 705–717.
- Péqueux, A., 1995. Osmotic regulation in crustaceans. J. Crustacean Biol. 15, 1–60.
- Péqueux, A., Gilles, R., Marshall, W.S., 1988. NaCl transport in gills and related structures. In: Greger, R. (Ed.), Advances in Comparative and Environmental Physiology. Springer, Berlin, pp. 1–73.
- Pierrot, C., Péqueux, A., Thuet, P., 1995. Effects of ions substitutions and of inhibitors on transepithelial potential difference and sodium fluxes in perfused gills of the crab *Pachygrapsus marmoratus*. Arch. Physiol. Biochem. 103, 466–475.
- Postel, U., Petrausch, G., Riestenpatt, S., Weihrach, D., 1998. Inhibition of Na^+/K^+ -ATPase and of ion-transport functions in the gills of the shore crab *Carcinus maenas* induced by cadmium. Mar. Biol. 130, 407–416.
- Prossner, C.L., 1973. Inorganic ions. In: Prossner, C.L. (Ed.), Comparative Animal Physiology. Saunders, Philadelphia, PA, pp. 79–110.
- Rajeswari, K., Kalarani, V., Reddy, D.C., Ramamurthi, R., 1988. Acute toxicity of endosulfan to crabs: effect on hydromineral balance. Bull. Environ. Contam. Toxicol. 40, 212–218.
- Rasmussen, A.D., Krag, A., Bjerregaard, P., Weeks, J.M., Depledge, M.H., 1995. The effects of trace metals on the apparent water permeability of the shore crab *Carcinus maenas* (L.) and the brown shrimp *Crangon crangon* (L.). Mar. Pollut. Bull. 31, 1–3.
- Reddy, A.N., Venugopal, N.B.R.K., Reddy, S.L.N., 1992. Effect of endosulfan 35 EC on ATPases in the tissues of a freshwater field crab *Barytelphusa guerini*. Bull. Environ. Contam. Toxicol. 48, 216–222.
- Robertson, J.D., 1960. Osmotic and ionic regulation. In: Waterman, T. (Ed.), The Physiology of Crustacea 1. Academic Press, New York, pp. 317–339.
- Rodriguez-Moreno, P.A., Schwarzbaum, P.J., Rodriguez, E.M., 1998. Effects of cadmium on gill Na,K -ATPase of the estuarine crab *Chasmagnathus granulata* (Decapoda, Grapsidae) during postmolt: in vivo and in vitro studies. Bull. Environ. Contam. Toxicol. 61, 629–636.
- Roesijadi, G., Petrocelli, S.R., Anderson, J.W., Presley, B.J., Sims, R., 1974. Survival and chloride ion regulation of the porcelain crab *Petrolisthes armatus* exposed to mercury. Mar. Biol. 2, 213–217.
- Roesijadi, G., Anderson, J.W., Petrocelli, S.R., Giam, C.S., 1976. Osmoregulation of the grass shrimp *Palaemonetes pugio* exposed to polychlorinated biphenyls (PCBs): I. Effect on chloride and osmotic concentrations and water-exchange kinetics. Mar. Biol. 38, 343–355.
- Rojas, W.E., Zanders, I.P., 1998. Cd-related impairment of hemolymph Ca and Mg regulation in fiddler crab (*Uca rapax*), acclimated to dilute or hypersaline seawater. Proc. 4th Int. Crustacean Congr., 20–24 July 1998, Amsterdam. p. 179, Abstract no. 450.
- Rubio, R., Tineo, P., Torreblanca, A., Del Ramo, J., Diaz-Mayans, J., 1991. Histological and electron microscopical observations on the effects of lead on gills and midgut gland of *Procambarus clarkii*. Toxicol. Environ. Chem. 31–32, 347–352.
- Shaw, J., 1960. The absorption of sodium ions by the crayfish *Astacus pallipes* Lereboullet: III. The effect of other cations in the external solution. J. Exp. Biol. 37, 548–556.
- Shires, R., Lane, N.J., Inman, C.B.E., Lockwood, A.P.M., 1994. Structural changes in the gill cells of *Gammarus duebeni* (Crustacea, Amphipoda) under osmotic stress; with notes on the microtubules in association with the septate junctions. Tissue Cell 26, 767–778.
- Siebers, D., Winkler, A., Lucu, C., Thedens, G., Weichart, D., 1985. $\text{Na}-\text{K}$ -ATPase generates an active transport potential in the gills of the hyperregulating shore crab *Carcinus maenas*. Mar. Biol. 87, 185–192.
- Siebers, D., Lucu, C., Winkler, A., Grammerstorf, V., Wille, M., 1987. Effects of amiloride on sodium chloride transport across isolated perfused gills of shore crab *Carcinus maenas* acclimated to brackish water. Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol. 87, 333–340.
- Soegianto, A., Charmantier-Daures, M., Trilles, J.-P., Charmantier, G., 1999a. Impact of cadmium on the structure of gills and epipodites of the shrimp *Penaeus japonicus* (Crustacea: Decapoda). Aquat. Living Resour. 12, 57–70.
- Soegianto, A., Bambang, Y., Charmantier-Daures, M., Charmantier, G., 1999b. Impact of copper on the structure of gills and epipodites of the shrimp *Penaeus japonicus* (Crustacea: Decapoda). J. Crustacean Biol., in press.
- Sommer, M.J., Mantel, L.H., 1988. Effect of dopamine, cyclic AMP, and pericardial organs on sodium uptake

- and Na/K-ATPase activity in the gills of the green crab *Carcinus maenas* (L.). J. Exp. Zool. 248, 272–277.
- Souheil, H., Vey, A., Thuet, P., Trilles, J.-P., 1999. Pathogenic and toxic effects of *Fusarium oxysporum* (Schlecht) on survival and osmoregulatory capacity of *Penaeus japonicus* (Bate). Aquaculture 178, 209–224.
- Spaargaren, D.H., 1977. On the metabolic adaptation of *Carcinus maenas* to reduced oxygen tension in the environment. Neth. J. Sea Res. 11, 325–333.
- Spaargaren, D.H., 1990. The effect of environmental ammonia concentrations on the ion-exchange of shore crabs, *Carcinus maenas* (L.). Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol. 97, 87–91.
- Taylor, H.H., Taylor, E.W., 1992. Gill and lungs: the exchange of gases and ions. Microscopic Anatomy of Invertebrates. In: Harrison, F.W., Humes, A.G. (Eds.), Decapoda, Crustacea vol. 10, Wiley, New York, pp. 203–293.
- Thuet, P., Charmantier-Daures, M., Charmantier, G., 1988. Relation entre osmorégulation et activités d'ATPase $\text{Na}^+ - \text{K}^+$ et d'anhydrase carbonique chez les larves et postlarves de *Homarus gammarus* (L.) (Crustacea: Decapoda). J. Exp. Mar. Biol. Ecol. 115, 249–261.
- Thurberg, F.P., Dawson, M.A., Collier, R.S., 1973. Effects of copper and cadmium on osmoregulation and oxygen consumption in two species of estuarine crabs. Mar. Biol. 23, 171–175.
- Thurberg, F.P., Calabrese, A., Gould, E., Greig, R.A., Dawson, M.A., Tucker, R.K., 1977. Response of the lobster, *Homarus americanus*, to sublethal levels of cadmium and mercury. In: Vernberg, F.J., Calabrese, A., Thurberg, F.P., Vernberg, W.B. (Eds.), Physiological Responses of Marine Biota to Pollutants. Academic Press, New York, pp. 185–197.
- Torreblanca, A., Diaz-Mayans, J., Del Ramo, J., Nunez, A., 1987. Oxygen uptake and gill morphological alterations in *Procambarus clarkii* (Girard) after sublethal exposure to lead. Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol. 86, 219–224.
- Torreblanca, A., Del Ramo, J., Arnau, J.A., Diaz-Mayans, J., 1989. Cadmium, mercury and lead effects on gill tissue of freshwater crayfish *Procambarus clarkii* (Girard). Biol. Trace Elem. Res. 21, 343–347.
- Towle, D.W., 1990. Sodium transport in gills. Comparative Physiology. In: Kinne, R.K.H. (Ed.), Comparative Aspects of Sodium Cotransport Systems vol. VII Basel, Karger, pp. 241–263.
- Towle, D.W., Kays, T.W., 1986. Basolateral localization of $\text{Na}^+ + \text{K}^+$ -ATPase in gill epithelium of two osmoregulating crabs, *Callinectes sapidus* and *Carcinus maenas*. J. Exp. Zool. 239, 311–318.
- Tucker, R.K., 1979. Effects of in vivo exposure of cadmium on ATPases in gill of the lobster, *Homarus americanus*. Bull. Environ. Contam. Toxicol. 23, 33–35.
- Tucker, R.K., Matte, A., 1980. In vitro effects of cadmium and lead on ATPases in the gill of the rock crab, *Cancer irroratus*. Bull. Environ. Contam. Toxicol. 24, 847–852.
- Vernberg, F.J., Vernberg, W.B., 1974. Pollution and Physiology of Marine Organisms. Academic Press, New York, 492 pp.
- Vitale, A.M., Monserrat, J.M., Rodriguez, E.M., 1998. Toxic effects of cadmium on *Chasmagnathus granulata* (Decapoda, Grapsidae) acclimated to different salinities. Proc. 4th Int. Crustacean Congr., 20–24 July 1998, Amsterdam. p. 176, Abstract no. 441.
- Walker, C.H., 1995. Biochemical biomarkers in ecotoxicology — some recent developments. Sci. Total Environ. 171, 189–195.
- Warburg, M.R., Goldenberg, S., Tudiver, B., 1987. Osmotic and ionic regulation in two *Pachygrapsus* crab, under varying salinities and dehydration. Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol. 86, 761–765.
- Wedderburn, J., Cheung, V., Bamber, S., Bloxham, M., Depledge, M.H., 1998. Biomarkers of biochemical and cellular stress in *Carcinus maenas*: an in situ field study. Mar. Environ. Res. 46, 1–5.
- Weeks, J.M., Jensen, F.B., Depledge, M.H., 1993. Acid-base status, haemolymph composition and tissue copper accumulation in the shore crab *Carcinus maenas* exposed to combined copper and salinity stress. Mar. Ecol.: Prog. Ser. 97, 91–98.
- Wendelaar Bonga, S.E., Lock, R.C.A., 1992. Toxicants and osmoregulation in fish. Neth. J. Zool. 42, 478–493.
- Wendemeyer, G.A., McLay, D.J., 1981. Methods for determining the tolerance of fishes to environmental stressors. In: Pickering, A.D. (Ed.), Stress and Fish. Academic Press, London, pp. 247–275.

- Wheatly, M.G., Gannon, A.T., 1995. Ion regulation in crayfish: freshwater adaptations and the problem of moulting. *Am. Zool.* 35, 49–59.
- Wheatly, M.G., Ignaszewski, L.A., 1990. Electrolyte and gas exchange during the moulting cycle of a freshwater crayfish. *J. Exp. Biol.* 151, 469–483.
- Williams, A.B., 1960. The influence of temperature on osmotic regulation in two species of estuarine shrimps (*Penaeus*). *Biol. Bull. (Woods Hole, Mass.)* 119, 560–571.
- Wood, C.M., Rogano, M.S., 1986. Physiological responses to acid stress in crayfish (*Orconectes*): haemolymph ions, acid–base status, and exchanges with the environment. *Can. J. Fish. Aquat. Sci.* 43, 1017–1026.
- Wright, D.A., Welbourn, P.M., 1991. Effect of mercury on unidirectional sodium and calcium influx in *Asellus aquaticus*. *Arch. Environ. Contam. Toxicol.* 21, 567–570.
- Wright, D.A., Welbourn, P.M., 1993. Effects of mercury exposure on ionic regulation in the crayfish *Orconectes propinquus*. *Environ. Pollut.* 82, 139–142.
- Young-Lai, W.W., Charmantier-Daures, M., Charmantier, G., 1991. Effect of ammonia on survival and osmoregulation in different life stages of the lobster *Homarus americanus*. *Mar. Biol.* 110, 293–300.
- Zanotto, F.P., Wheatly, M.G., 1993. The effect of ambient pH on electrolyte regulation during the postmoult period in freshwater crayfish *Procambarus clarkii*. *J. Exp. Biol.* 178, 1–19.