

Erratum

Erratum to “Effect of rapid changes in temperature
and salinity on availability of the rotifers
Brachionus rotundiformis and
Brachionus plicatilis”
[Aquaculture 189 (2000) 85–99][☆]

D.S. Fielder ^{a,*}, G.J. Purser ^b, S.C. Battaglene ^c

^a NSW Fisheries, Port Stephens Research Centre, Taylors Beach Road, Taylors Beach 2315, Australia

^b University of Tasmania, School of Aquaculture, P.O. Box 1214, Launceston 7250, Australia

^c Tasmanian Aquaculture and Fisheries Institute, Nubeena Crescent, Taroona, Tasmania 7053, Australia

Accepted 1 August 2000

The Publisher regrets that there was an error in the heading of column 5 in Table 1 of the above article. It should read “after 6 h”. The table is reproduced correctly, below.

Table 1

Mean number of available rotifers, for control treatments (23°C, 35‰) in experiments 1–4 giving the number potentially available and those sampled 1 and 6 h after stocking into non-turbulent beakers

Experiment number	Rotifer (<i>Brachionus</i> species)	Potentially available number of rotifers (ml ⁻¹) ^a	Mean rotifers ml ⁻¹ after 1 h (%) ^b	Mean rotifers ml ⁻¹ after 6 h (%) ^b
1	<i>B. rotundiformis</i>	25.1 ± 0.6	61.7 ± 4.9	89.5 ± 9.6
2	<i>B. plicatilis</i>	26.2 ± 0.6	57.0 ± 2.3	63.9 ± 3.5
3	<i>B. rotundiformis</i>	27.6 ± 0.5	36.4 ± 4.9	61.7 ± 9.5
4	<i>B. plicatilis</i>	26.1 ± 0.6	45.3 ± 1.9	87.3 ± 11.1

^a Values are means ± standard error of three replicate beakers.

^b Values are percent availability of potentially available number. Data for Experiments 1 and 2 are means ± S.E. of eight replicate beakers. Data for Experiments 3 and 4 are means ± se of four replicate beakers.

[☆] PII of original article S0044-8486(00)00369-0

* Corresponding author. Tel.: +61-2-4982-1232; fax: +61-2-4982-1107.

E-mail address: fielders@fisheries.nsw.gov.au (D.S. Fielder).