

Development of Molecular Identification Method for Genus *Alexandrium* (Dinophyceae) Using Whole-Cell FISH

Choong-Jae Kim,¹ Chang-Hoon Kim,² Yoshihiko Sako³

¹Environmental Biotechnology Laboratory, Korea Research Institute of Bioscience & Biotechnology, Yuseong-gu, 305-333, Daejeon, Korea

²Department of Aquaculture, Pukyong National University, 599-1 Daeyeon-3-dong, Nam Gu, Busan 608-737, South Korea

³Department of Applied Biosciences, Graduate School of Agriculture, Kyoto University, Kyoto, 606-8502, Japan

Received: 18 June 2003 / Accepted: 23 July 2004 / Online publication: 17 May 2005

Abstract

We have developed a method to identify species in the genus *Alexandrium* using whole-cell fluorescent in situ hybridization with FITC-labeled oligonucleotide probes that target large subunit ribosomal rRNA molecules. The probes were designed based on the sequence of the rDNA D1-D2 region of *Alexandrium* species. DNA probes specific for toxic *A. tamarens* and *A. catenella* and nontoxic *A. affine*, *A. fraterculus*, *A. insuetum*, and *A. pseudogonyaulax*, respectively, were applied to vegetative cells of all above *Alexandrium* species to test the sensitivity of the probes. Each DNA probe hybridized specifically with vegetative cells of the corresponding *Alexandrium* species and showed no cross-reactivity to noncorresponding *Alexandrium* species. In addition, no cross-reactivity of the probes was observed in experiments using concentrated natural seawater samples. The TAMAD2 probe, which is highly specific to *A. tamarens*, a common toxic species in Korean coastal waters, provides a simple and reliable molecular tool for identification of toxic *Alexandrium* species.

Key words: DNA probe — *Alexandrium* — whole-cell FISH

Introduction

Harmful algal blooms (HABs) have become a worldwide problem, causing immense economic losses in aquaculture and much human suffering as a result of accidental exposure. These blooms are not a temporary natural phenomenon, but rather an

anthropologic problem caused by the disposal of untreated domestic and industrial waste, and by improvements in aquaculture (Lam and Ho, 1989; Okaichi, 1989; Hallegraeff, 1993).

Certain toxic dinoflagellate species of the genus *Alexandrium* (Halim) are lethal to humans, owing to their production of paralytic shellfish poisoning (PSP) toxins, which block sodium channels in animal nerve cells (Schantz et al., 1966; Baden and Trainer, 1993; Kao, 1993;). The *Alexandrium* species are planktonic-sized marine thecate and photosynthetic dinoflagellates that belong to the phylum Dinophyta, class Dinophyceae, order Gonyaulacales, and family Gonyaulacaceae. The species *A. tamarens* (Lebour) Balech, *A. catenella* (Whedon and Kofoid) Balech, *A. acatenella* (Whedon and Kofoid) Balech, *A. cohorticula* (Balech) Balech, *A. fundyense* Balech, *A. ostenfeldii* (Paulsen) Balech and Tangen, *A. minutum* Halim, *A. tamiyavanichii* Balech, and some related species (Balech, 1995) have caused many PSP-related events worldwide. Human ingestion of aquaculture products that contain bivalves contaminated with these species can result in poisoning episodes, paralytic diseases, and even death. Rapid detection of the presence of toxic dinoflagellates, identification of the species, and characterization of their abundance and distribution in algal blooms would aid in the protection of sensitive aquaculture areas and the prevention of human illness.

In addition to the production of PSP toxin by these species, another long-term problem with the *Alexandrium* genus is confusion in the species nomenclature. There are 28 species in this genus, with 3 equivocal *Alexandrium* spp. in addition to *A. depressum* Gaarder (Balech, 1995). Morphologic identification of *Alexandrium* species is usually based on the microtabulation in the fine thecal

Correspondence to: Choong-Jae Kim; E-mail: chung-jae@han-mail.net

plates, chain formation, cell size, and overall shape. Strong morphologic similarities between *Alexandrium* species have led to much taxonomic confusion. The worldwide dispersion of the genus (Shumway, 1990; Hallegraeff, 1995) also highlights the need for the development of exact species identification and population analysis methods.

In order to eliminate the confusion in the *Alexandrium* taxonomy, molecular phylogenetic analysis has been used to establish more concrete criteria and to provide basal genetic information (Destombe et al., 1992; Scholin and Anderson, 1994; Scholin et al., 1994; Zardoya et al., 1995; Adachi et al., 1996a; Usup et al., 2002). We conducted phylogenetic analysis based on comparison of the newly determined sequences of the large subunit (LSU) ribosomal DNA D1-D2 and small subunit (SSU) rRNA regions from *A. affine*, *A. catenella*, *A. fraterculus*, *A. insuetum*, *A. pseudogonyaulax*, *A. tamiyavanichii*, and *A. tamarensis* with sequences from functionally expressed rRNA and those retrieved from the database. These analyses divided the genus *Alexandrium* into 14 groups: (1) *A. tamarensis*, (2) *A. excavatum*, (3) *A. catenella*, (4) Tasmanian *A. tamarensis*, (5) *A. affine* (and/or *A. concavum*), Thai *A. tamarensis*, (7) *A. tamiyavanichii*, (8) *A. fraterculus*, (9) *A. margalefii*, (10) *A. andersonii*, (11) *A. ostensfeldii*, (12) *A. minutum* (or *A. lusitanicum*), (13) *A. insuetum*, and (14) *A. pseudogonyaulax*. Ribosomal DNA and RNA sequence information can be used to design molecular probes for the detection, identification, and enumeration of these groups (Giovannoni et al., 1988; Amann et al., 1990, 1991, 1995; DeLong and Shah, 1990; Distel et al., 1991; Worden et al., 2000). The use of DNA probes has been applied to only a limited number of HAB species (Adachi et al., 1996b; Scholin et al., 1996, 1997, 1999; Miller and Scholin, 1996; Parson et al., 1999). It is a matter of great interest to develop methods to identify *Alexandrium* species and other PSP producers exactly and rapidly during field monitoring. However, near real-time monitoring of *Alexandrium* species has not been achieved because of difficulties resulting from the highly similar morphologies of the species and a lack of the extensive genetic information required for molecular approaches. In particular, difficulties arise from the so-called tamarensis species complex; this refers to the overlap of the morphologic characteristics of some species with those of others. For example, *A. excavatum* and *A. tamarensis*, as well as *A. tamiyavanichii*, share characteristics with *A. affine* and *A. fraterculus* that are related to chain formation and occurrence in natural seawater columns; this overlap confounds morphologic identification.

Adachi et al. (1996b) used molecular techniques to identify *Alexandrium* species, targeting the genomic rRNA gene (rDNA) to identify *A. tamarensis* and *A. catenella*; however, the hybridization used in this technique is time-consuming and produces a weak labeling signal.

To develop a molecular identification technique for the toxic *A. tamiyavanichii*, we designed a cellular-rRNA-targeted DNA probe for use in whole-cell fluorescent in situ hybridization (FISH) (Kim, 2003). The *A. tamiyavanichii*-specific DNA probe TAMID2, with a design based on the LSU rDNA regions D1-D2, successfully hybridized with the targeted vegetative cells and showed no cross-reactivity either with *Alexandrium* congeners such as *A. tamarensis*, *A. catenella*, *A. affine*, *A. fraterculus*, *A. insuetum*, and *A. pseudogonyaulax* or with concentrated natural seawater samples. Simple and rapid procedures were also established for the use of FISH with whole cells of these species.

In the present study we aimed to design species-specific DNA probes for 6 *Alexandrium* species, *A. tamarensis*, *A. catenella*, *A. affine*, *A. fraterculus*, *A. insuetum*, and *A. pseudogonyaulax*, for use in molecular identification with the whole-cell FISH technique previously developed by Kim (2003). The resulting probes were tested for cross-reactivity to *Alexandrium* congeners and concentrated natural seawater samples and used in the detection and identification of toxic *A. tamarensis* species in Korean coastal waters.

Materials and Methods

Cultivation and Identification of *Alexandrium* Species for Testing with Whole-Cell FISH. Clonal cultures of *Alexandrium* were established from cultures collected in Korea and Japan (Table 1). *A. tamarensis* was incubated at 15°C, and the other *Alexandrium* species were maintained at 20°C in SWIIm medium (Sako et al., 1990) under a 14 hours–10 hours light-dark regimen with 100 $\mu\text{E m}^{-2}\text{s}^{-1}$ provided by cool white bulbs. All *Alexandrium* species were identified ribotypically through determination of the sequences of the LSU rDNA D1-D2 and SSU rDNA regions (Table 1).

Probe Design. Sequences corresponding to the LSU rDNA D1-D2 regions of *Alexandrium* species obtained from GenBank (Table 1) were aligned using CLUSTAL X (Thomson et al., 1997). A species-specific region was identified for use in the design of oligonucleotide DNA probes for 6 *Alexandrium* species: *A. tamarensis*, *A. catenella*, *A. affine*, *A. fraterculus*, *A. insuetum*, and *A. pseudogonyaulax*.

Table 1. *Alexandrium* Species Tested with Whole-Cell FISH

Species	Strain	Origin
<i>A. affine</i>	AFF37	Harima Nada, Japan
<i>A. catenella</i>	DPC7	Pusan, Korea
	ACY12	Harima Nada, Japan
<i>A. fraterculus</i>	DPW9709	Pusan, Korea
<i>A. insuetum</i>	AI104	Utiumi Bay, Japan
<i>A. tamarense</i>	KJC9711	Kojedo, Korea
	HAT4	Hiroshima Bay, Japan
<i>A. pseudogonyaulax</i>	AP391	Harima Nada, Japan

The sequences complementary to the rDNA coding strand, and the sites and melting temperatures of the probes, are shown in Table 2. Portions of the 5' end of each probe were labeled with fluorescein isothiocyanate (FITC). Each DNA probe was synthesized commercially and cartridge-purified (Amersham Pharmacia Biotech). Working stocks of the DNA probes were prepared by resuspending the powder containing the DNA oligonucleotides in 10 mM Tris HCl (pH 7.4) to a final concentration of 100 pmol μl^{-1} of probe.

Reactivity of DNA Probes with *Alexandrium* Cells. Cells of each of the 6 cultured *Alexandrium* species were harvested by centrifugation, and the supernatant was removed. The pelleted cells were dehydrated, and autofluorescence originating from chlorophyll was removed using an ethanol series of 50% for 1 minute and 100% for 30 minutes at room temperature. The dehydrated cells were air-dried completely and then subjected to hybridization for 5 minutes at 35°C, in a water bath, in 200 μl of a hybridization buffer containing 40% formamide (Sigma), 5× SSC (83.3 mM NaCl, 83.3 mM trisodium citrate, pH 7.0), and DNA probe (final concentration, 0.5 pmol). Hybridized cells were then washed twice in 5× SSC at 45°C for 5 minutes to remove unhybridized DNA probe. Afterward, the pellets, which contained 10 to 20 μl of cells, were gently resuspended by pipetting and observed at 400× magnification under an epifluorescence microscope (Nikon ECLIPSE E800) equipped with a fluorescein band pass filter set for FITC and counterstain (TRITC. PI)

(excitation, 450–490 nm; DM, 505 nm; BA, 520 nm). Fluorescence and light images of the cells were taken using a color-chilled 3-CCD camera (Hamamatsu C5810) with a fixed exposure time of 0.4 seconds.

Reactivity of DNA Probes with Nontargeted *Alexandrium* Congeners and Natural Seawater Samples. Each DNA probe was tested for cross-reaction to cultured vegetative cells of the *Alexandrium* congeners. Whole-cell FISH was performed as described above. Natural seawater samples were collected in netting with a pore size of 20 μm on August 29, 2002, at Maizuru, Kyoto, Japan. Concentrated samples were prepared in polypropylene bottles and not treated for fixation; they were maintained in an icebox at 5° to 10°C during transportation to the laboratory, and then subjected to whole-cell FISH with each DNA probe, within 5 hours after sampling. In the laboratory, field samples were concentrated using a sieve with 20- μm pores and transferred to 50-ml polypropylene disposable centrifuge tubes at room temperature. One milliliter of cultured *Alexandrium* cells (approx. 5000–10,000 cells/ml) was added to 1 ml of the wild samples to test for cross-reaction. The hybridization and washing temperatures were set at the relatively high 40° and 45°C, respectively, to improve the specificity of the probes. After the treatments the cells were resuspended and observed under an epifluorescence microscope (Nikon ECLIPSE E800) under the same conditions as those described above.

Identification of *Alexandrium* Species Isolated from Korea and Japan, Using DNA Probes. Forty-three *Alexandrium* strains from Korea and Japan were identified using 6 *Alexandrium*-species-specific DNA probes. Whole-cell FISH was performed, as described above, and the strains tested are listed in Table 1. The results of the molecular identification were reconfirmed by comparison with ribotypically determined *Alexandrium* strains.

In order to detect and identify the *Alexandrium* species in natural seawater, field sampling was conducted at Sujongri in Chinhae Bay, Korea, in April 2003, during monitoring for *A. tamarense* and

Table 2. *Alexandrium*-Species-Specific DNA Probes with Designs Based on Sequences of LSU D1-D2 Region of rDNA

Species	Probe	Sequences ^a (site)	T_m (°C)
<i>A. affine</i>	AFFD2	TAGGTTTGTATTGTGGGTGT (445–464)	52.0
<i>A. catenella</i>	CATED2	TGGGTTTGGCTGCAAGTGC (444–463)	70.4
<i>A. fraterculus</i>	FRAD2	TCGATGTGAATTGCAAATGC (447–466)	64.6
<i>A. insuetum</i>	INSUD2	TATATGTTGGATGTGGGTGC (447–466)	60.6
<i>A. tamarense</i>	TAMAD2	GAGCTTTGGGCTGTGGGTGT (444–463)	68.6
<i>A. pseudogonyaulax</i>	PSEDOD2	GTGKAATTGTATGAGCYTT (444–463)	51.5

^aSequences are listed 5' to 3' and include primer regions.

A. catenella. Two-liter water samples were collected from the surface layer, and the samples were transported to the laboratory under cool conditions without applying fixation methods. Ten-milliliter samples were placed into a 2-cm-diameter filter apparatus that contained a 20- μ m-pore-size nylon sieve. Whole-cell FISH was performed as described above, except that the dehydrated samples were divided into 2 identical volumes and subjected to hybridization with the DNA probes TAMAD2 and CATED2, which are specific to *A. tamarens* and *A. catenella*, respectively. After 2 washes in 5 $\times$ SSC buffer, the filtrates were mounted onto a glass slide for observation under an epifluorescence microscope, and images were taken as described above.

Results

Labeling of Six *Alexandrium* Species with Specific DNA Probes. Figure 1 shows the results of whole-cell FISH using 6 oligonucleotide DNA probes specific for each targeted *Alexandrium* species. All DNA probes reacted with the targeted cultured *Alexandrium* vegetative cells, as indicated by a fluorescent green color (bright green). Specifically hybridized *Alexandrium* cells were easily identified owing to the retention of their original cell shapes.

Specificity of DNA Probes to *Alexandrium* and Wild Samples. To confirm the species specificity of the DNA probes, each probe was tested for cross-reaction to cultured cells of *A. tamarens*, *A. catenella*, *A. fraterculus*, *A. affine*, *A. insuetum*, *A. tamiyavanichii*, and *A. pseudogonyaulax*. All of the conditions and whole-cell FISH procedures were as described above, except that the hybridization and washing temperatures used were relatively low at 35°C and 40°C, respectively, in order to detect very weak cross-reactivity.

None of the 6 DNA probes reacted with non-targeted *Alexandrium* species, indicating that all of the DNA probes are highly specific (Table 3). None of the probes showed cross-reactivity to field samples composed of various diatoms, dinoflagellates, and zooplanktons, and all of the targeted *Alexandrium* species were detectable with each probe, although the fluorescence intensity was variable (Figure 2). This detection can be attributed to the poor condition of some *Alexandrium* cells in the field samples. Occasionally, detritus-like amorphous clusters were labeled by the DNA probes, a phenomenon that could probably be avoided by size fractionation of the samples using a filter apparatus.




Fig. 1. Epifluorescence (A to F) and light (A' to F') micrographs of the 6 *Alexandrium* species *A. affine* (A and A'), *A. fraterculus* (B and B'), *A. insuetum* (C and C'), *A. pseudogonyaulax* (D and D'), *A. catenella* (E and E'), and *A. tamarens* (F and F'), hybridized with the following DNA probes: AFFD2 specific for *A. affine* (A and A'), FRAD2 specific for *A. fraterculus* (B and B'), INSUD2 specific for *A. insuetum* (C and C'), PSEUDOD2 specific for *A. pseudogonyaulax* (D and D'), CATED2 specific for *A. catenella* (E and E'), and TAMAD2 specific for *A. tamarens* (F and F'). Scale bar = 30 μ m.

Identification of *Alexandrium* Species Using DNA Probes. Analysis of specimens of 7 *Alexandrium* species and 18 strains using the 6 DNA probes resulted in the same speciation pattern that had been determined using the rDNA D1-D2 sequences summarized in Table 2. The Japanese strains ACY12 (*A. catenella*) and HAT4 (*A. tamarens*) reacted with the *A. catenella*- and *A. tamarens*-specific DNA probes, as did the Korean strains. The ribotypically undetermined *Alexandrium* species SMG0205 was identified as *A. affine*; SOC2, SOC6, and SOC9, as *A. catenella*; and SOC3, YOC2, and YOC3 as *A. tamarens*.

Considerable quantities of *Alexandrium* were observed in the natural seawater samples collected from Chinhae Bay, Korea. In whole-cell FISH, all of the *Alexandrium* cells collected reacted only with

Table 3. Summary of Molecular Identification of *Alexandrium* Species Using Six DNA Probes

Strain (ribotypic species)	Origin	Acc. no. ^a	DNA probe ^b					
			AFFD2	CATED2	FRAD2	INSUD2	PSEUDOD2	TAMAD2
AFF37 (<i>A. affine</i>)	Harima Nada, Japan	AB088227	+	-	-	-	-	-
SMG0205	Saemangeum (Yellow Sea), Korea	Unidentified	+	-	-	-	-	-
DPC7 (<i>A. catenella</i>)	Busan Dadaepo, Korea	AB088234	-	+	-	-	-	-
DPC8 (<i>A. catenella</i>)	Busan Dadaepo, Korea	AB088239	-	+	-	-	-	-
CMC2 (<i>A. catenella</i>)	Thongyong, Korea	AB088229	-	+	-	-	-	-
CMC3 (<i>A. catenella</i>)	Thongyong, Korea	AB088233	-	+	-	-	-	-
YSC9811 (<i>A. catenella</i>)	Yosu, Korea	AB088273	-	+	-	-	-	-
SOC2	Yosu open sea, Korea	Unidentified	-	+	-	-	-	-
SOC6	Yosu open sea, Korea	Unidentified	-	+	-	-	-	-
SOC9	Kojedo open sea, Korea	Unidentified	-	+	-	-	-	-
ACY12 (<i>A. catenella</i>)	Harima Nada, Japan	AB088225	-	+	-	-	-	-
DPW9709 (<i>A. fraterculus</i>)	Busan Dadaepo, Korea	AB088244	-	-	+	-	-	-
SPW9709 (<i>A. fraterculus</i>)	Chinhae Bay, Korea	AB088262	-	-	+	-	-	-
AI104 (<i>A. insuetum</i>)	Utsunomiya Bay, Japan	AB088248	-	-	-	+	-	-
AP391 (<i>A. pseudogonyaulax</i>)	Harima Nada, Japan	AB088250	-	-	-	-	+	-
KJC971 (<i>A. tamarensis</i>)	Kojedo, Korea	AB088279	-	-	-	-	-	-
SW97043 (<i>A. tamarensis</i>)	Chinhae Bae, Korea	AB088258	-	-	-	-	-	+
SW9516 (<i>A. tamarensis</i>)	Chinhae Bae, Korea	AB088255	-	-	-	-	-	+
ULW9903 (<i>A. tamarensis</i>)	Ulsan, Korea	AB088270	-	-	-	-	-	+
SOC3	Busan open sea, Korea	Unidentified	-	-	-	-	-	+
YOC2	Saemangeum (Yellow Sea), Korea	Unidentified	-	-	-	-	-	+
YOC3	Saemangeum (Yellow Sea), Korea	Unidentified	-	-	-	-	-	+
TAMI2201 (<i>A. tamiyavanichii</i>)	Harima Nada, Japan	AB088264	-	-	-	-	-	-

^aSequences of rDNA D1-D2 region.^bAFFD2, CATED2, FRAD2, INSUD2, PSEUDOD2, and TAMAD2 are specific to *A. affine*, *A. catenella*, *A. fraterculus*, *A. insuetum*, *A. pseudogonyaulax*, and *A. tamarensis*, respectively.

+ and - indicate positive and negative reaction to each DNA probe, respectively.




Fig. 2. Epifluorescence (A to F) and light (A' to F') micrographs of natural seawater samples containing the 6 added *Alexandrium* species *A. affine* (A and A'), *A. fraterculus* (B and B'), *A. insuetum* (C and C'), *A. pseudogonyaulax* (D and D'), *A. catenella* (E and E'), and *A. tamarensis* (F and F'), hybridized with the following DNA probes: AFFD2 specific for *A. affine* (A and A'), FRAD2 specific for *A. fraterculus* (B and B'), INSUD2 specific for *A. insuetum* (C and C'), PSEUDOD2 specific for *A. pseudogonyaulax* (D and D'), CATED2 specific for *A. catenella* (E and E'), and TAMAD2 specific for *A. tamarensis* (F and F'). Scale bar = 30 μ m.

the DNA probe TAMAD2, indicating that the cells were *A. tamarensis* (Figure 3). Despite the large size range of 25–50 μ m of the *A. tamarensis* cells in the natural seawater samples, the reactivity of the cells to the DNA probe was strong and consistent.

Discussion

Whole-cell FISH has proven a highly useful molecular tool in the detection and identification of microbes, particularly in the study of human diseases. Although the need for rapid and exact identification of HABs is pressing, to date whole-cell FISH has been used only rarely to monitor HABs.

The results of this study indicate that rapid and exact whole-cell FISH using oligonucleotide DNA




Fig. 3. Epifluorescence (A and B) and light (A' and B') micrographs of natural seawater samples hybridized with the DNA probes TAMAD2 (A and A') and CATED2 (B and B'). Arrows indicate *Alexandrium* cells. Scale bar = 30 μ m.

probes is an extremely effective molecular tool for the identification of species in the genus *Alexandrium*. We designed 6 DNA probes targeting rRNA molecules specific to *A. tamarensis*, *A. tamarensis*, *A. affine*, *A. insuetum*, *A. fraterculus*, and *A. pseudogonyaulax*. The FISH methods used allowed the rapid and exact molecular identification of *Alexandrium* species. Probes exposed to concentrated natural seawater samples containing cultured *Alexandrium* vegetative cells showed high specificity and almost no cross-reactivity. Use of these probes in the field appears promising, as *A. tamarensis* cells were positively detected and identified in Korean coastal waters with the specific DNA probe TAMAD2.

This whole-cell FISH approach did not require additional fixation, which can negatively affect the physical condition of the cells, nor did it require time-consuming treatment with cetyltrimethyl ammonium bromide (CTAB). The method was possible owing to the direct hybridization of a DNA probe to targeted molecules that have a planar structure; this constitutes one of the most important advantages of rapid whole-cell FISH. Another significant obstacle, interference of the fluorescence illumination of the labeling signal by the autofluorescence derived from the cells (Lee et al., 1993; Christensen et al., 1999), was almost completely removed by merely using an ethanol series for fixation.

During field trials we conducted several preliminary experiments to find ways to preserve whole cells for a longer period and to eliminate the auto-fluorescence originating from chlorophyll. Samples were diluted to final concentrations of 40%, 50%, and 80% ethanol and 80% acetone immediately after collection. The results were very similar to fixation with glutaraldehyde, which caused undesirable modifications in the targeted cells. Instead, we used low concentrations of ethanol: 10%, 20%, and 30%. This treatment was satisfactory for whole-cell FISH, but the fluorescence intensity was not as strong as in untreated cells. In addition, we analyzed samples that had been stored for 2 days at room temperature (15–25°C), at 4°C, or at –20°C immediately after collection. The rRNA remained in the best condition in samples that had been stored at room temperature. The samples stored at 4°C remained in relatively good condition, but there was considerable decrease in the fluorescence intensity, which may be attributable to autolysis, followed by cell death. Cells that had been frozen broke easily during the thawing process, although the level of autolysis was low. Many authors have reported decreases in the cellular rRNA content during storage in batch mode (Neidhardt and Magasanik, 1960; DeLong et al., 1989; Kemp et al., 1993; Lim et al., 1993, 1996; Poulsen et al., 1993; Ruimy et al., 1994; Zheng et al., 1996; Worden et al., 2000). We are now in search of even more effective procedures for the handling of whole cells, for use in the field without fixation, from dehydration using the ethanol series to hybridization. Nevertheless, our whole-cell FISH technique is very simple, and once hybridized, the targeted rRNA molecules will be more resistant to autolysis.

In the future whole-cell FISH systems employing multiple probes will provide powerful molecular tools for the identification of HABs caused not only by *Alexandrium*, but also by other organisms. Development of a portable system to perform these tests has strong commercial possibilities.

Acknowledgments

This work was supported by a grant (R02-2000-00226) from the Basic Research Program of the Korea Science and Engineering Foundation, and supported in part by a grant-in-aid for scientific research (13556033) from the Ministry of Education, Culture, Sports, Science and Technology of JAPAN. We declare that the experiments performed comply with the current laws of the country in which the experiments were performed.

References

1. Adachi M, Sako Y, Ishida Y (1996a) Analysis of *Alexandrium* (Dinophyceae) species using sequences of the 58S ribosomal DNA and internal transcribed spacer regions. *J Phycol* 32, 424–432
2. Adachi M, Sako Y, Ishida Y (1996b) Identification of the toxic dinoflagellates *Alexandrium catenella* and *A. tamarense* (Dinophyceae) using DNA probes and whole-cell hybridization. *J Phycol* 32, 1049–1052
3. Amann RI, Krumholz L, Stahl DA (1990) Fluorescent oligonucleotide probing of whole cells for determinative, phylogenetic, and environmental studies in microbiology. *J Bacteriol* 172, 762–770
4. Amann R, Springer N, Ludwig W, Gortz H-D, Schleifer K-H (1991) Identification *in situ* and phylogeny of uncultured bacterial endosymbionts. *Nature* 351, 161–164
5. Amann RI, Ludwig W, Schleifer K-H (1995) Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiol Rev* 59, 143–169
6. Baden DG, Trainer VL (1993) "Mode of action of toxins of seafood poisoning". In: *Algal Toxins in Seafood and Drinking Water*, Falconer IR, ed. (London, UK: Academic Press) pp 49–74
7. Balech E (1995) The genus *Alexandrium* Halim (Dinophyceae). (Sherkin Island, Co Cork, Ireland: Sherkin Island Marine Station)
8. Cristensen H, Hansen M, Sorensen J (1999) Counting and size classification of active soil bacteria by fluorescence *in situ* hybridization with an rRNA oligonucleotide probe. *Appl Environ Microbiol* 65, 1753–1761
9. DeLong EF, Shah J (1990) Fluorescent, ribosomal RNA probes for clinical application: a research review. *Diagn Clin Testing* 28, 41–44
10. DeLong EF, Wickham FS, Pace NR (1989) Phylogenetic stains: ribosomal RNA-based probes for the identification of single cells. *Science* 243, 1360–1363
11. Destombe C, Cembella AD, Murphy CA, Ragan MA (1992) Nucleotide sequence of the 18S ribosomal RNA genes from the marine dinoflagellate *Alexandrium tamarense* (Gonyaulacales, Dinophyta). *Phycologia* 31, 121–4
12. Distel DL, DeLong EF, Waterbury JB (1991) Phylogenetic characterization and *in situ* localization of the bacterial symbiont of shipworms (Teredinidae: Bivalvia) by using 16S RNA sequence analysis and oligodeoxynucleotide probe hybridization. *Appl Environ Microbiol* 57, 2376–2382
13. Giovannoni SJ, DeLong EF, Olsen GJ, Pace NR (1988) Phylogenetic group-specific oligodeoxynucleotide probes for identification of single microbial cells. *J Bacteriol* 170, 720–726
14. Hallegraeff GM (1993) Review of harmful algal blooms and their apparent global increase. *Phycologia* 32, 79–99
15. Hallegraeff GM (1995) "Harmful algal blooms: a global overview". In: *Manual on Harmful Marine Microal-*

- gae, Hallegraeff GM, Anderson DM, Cembella AD, eds. (Paris, France: IOC (UNESCO)) pp 1–18
16. Kao CY (1993) "Paralytic shellfish poisons". In: *Algal Toxins in Seafood and Drinking Water*, Faalconer IR, ed. (London, UK: Academic Press) pp 75–86
 17. Kemp PF, Lee S, LaRoche J (1993) Estimating growth rate of slowly growing marine bacteria from rRNA content. *Appl Environ Microbiol* 59, 2594–2601
 18. Kim CJ (2003) "Toxicological and molecular biological study on the toxic dinoflagellate genus *Alexandrium*". PhD thesis, Kyoto University, pp 52–69
 19. Lam CWY, Ho KC (1989) "Red tides in Tolo Harbour, Hong Kong". In: *Red Tides: Biology, Environmental Science and Toxicology*, Okaichi T, Anderson DM, Nemoto T, eds. (New York, NY: Elsevier Science Publishing Co.) pp 49–52
 20. Lee S, Malone C, Kemp PF (1993) Use of multiple 16S rDNA-targeted fluorescent probes to increase signal strength and measure cellular RNA from natural planktonic bacteria. *Mar Ecol Prog Ser* 101, 193–201
 21. Lim EL, Amaral LA, Caron DA, DeLong EF (1993) Application of RNA-based probes for observing marine nanoplanktonic protists. *Appl Environ Microbiol* 59, 1647–1655
 22. Lim EL, Caron DA, DeLong EF (1996) Development and field application of a quantitative method for examining natural assemblages of protists with oligonucleotide probes. *App Environ Microbiol* 62, 1416–1423
 23. Miller PE, Scholin CA (1996) Identification of cultured *Pseudo-nitzschia* (Bacillariophyceae) using species-specific LSU rRNA-targeted fluorescent probes. *J Phycol* 32, 646–655
 24. Neidhardt FC, Magasanik B (1960) Studies on the role of ribonucleic acid in the growth of bacteria. *Biochim Biophys Acta* 42, 99–116
 25. Okaichi T (1989) "Red tide problems in the Seto Inland Sea, Japan". In: *Red Tides: Biology, Environmental Science and Toxicology*, Okaichi T, Anderson DM, Nemoto T, eds. (New York, NY: Elsevier Science Publishing Co.) pp 137–142
 26. Parson ML, Scholin CA, Miller PE, Doucette GJ, Powell CL, Fryxell GA, Dortch Q, Soniat TM (1999) *Pseudo-nitzschia* species (Bacillariophyceae) in Louisiana coastal waters: molecular probe field trials, genetic variability, and domoic acid analyses. *J Phycol* 35, 1368–1378
 27. Poulsen LK, Ballard G, Stahl DA (1993) Use of rRNA fluorescence *in situ* hybridization for measuring activity of single cells in young and established biofilms. *Appl Environ Microbiol* 59, 1354–1360
 28. Ruimy R, Breittmayer V, Boivin V, Christen R (1994) Assessment of the state of individual bacterial cells by hybridization with a ribosomal rRNA targeted fluorescently labeled oligonucleotide probe. *FEMS Microbiol Ecol* 15, 207–214
 29. Sako Y, Kim CH, Ninomiya H, Adachi M, Ishida Y (1990) "Isozyme and cross analysis of mating populations in the *Alexandrium catenellaltamarensis* species complex". In: *Toxic Marine Phytoplankton*, Graneli E, Sundstrom K, Anderson DM, eds. (New York, NY: Elsevier) p 320–323
 30. Schantz EJ, Lynch JM, Vayvada G, Matsumoto K, Rapoport H (1966) The purification and characterization of the poison produced by *Gonyaulax catenella* in axenic culture. *Biochemistry* 5, 1191–1195
 31. Scholin CA, Anderson DM (1994) Identification of species and strain-specific genetic markers for globally distributed *Alexandrium* (Dinophyceae), I: RFLP analysis of SSU rRNA genes. *J Phycol* 30, 744–754
 32. Scholin CA, Herzog M, Sogin M, Anderson DM (1994) Identification of group- and strain-specific genetic markers for globally distributed *Alexandrium* (Dinophyceae), II: sequence analysis of a fragment of the LSU rRNA gene. *J Phycol* 30, 999–1011
 33. Scholin CA, Buck KR, Britschgi T, Gangelosi G, Chavez FP (1996) Identification of *Pseudo-nitzschia australis* (Bacillariophyceae) using rRNA-targeted probes in whole cell and sandwich hybridization formats. *Phycologia* 35, 190–197
 34. Scholin CA, Miller P, Buck K, Chavez FP, Harris P, Haydock P, Howard J (1997) Detection and quantification of *Pseudo-nitzschia australis* in cultured and natural populations using LSU rRNA-targeted probes. *Limnol Oceanogr* 42, 1265–1272
 35. Scholin CA, Martin R III, Miller PE, Doucette GJ, Powell CL, Haydock P, Howard J, Ray J (1999) DNA probes and a receptor-binding assay for detection of *Pseudo-nitzschia* (Bacillariophyceae) species and domoic acid activity in cultured and natural samples. *J Phycol* 35, 1356–1367
 36. Shumway SE (1990) A review of the effects of algal blooms on shellfish and aquaculture. *J World Aquacult Soc* 21, 65–104
 37. Thomson JD, Gibson TJ, Plewniak R, Jeanmougm F (1997) The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 24, 4876–4882
 38. Usup G, Pin LC, Ahmad A, Teen LP (2002) Phylogenetic relationship of *Alexandrium tamiyavanichii* (Dinophyceae) to other *Alexandrium* species based on ribosomal RNA gene sequences. *Harmful Algae* 1, 59–68
 39. Worden AZ, Chisholm S, Binder B (2000) *In situ* hybridization of *Prochlorococcus* and *Synechococcus* (marine cyanobacteria) spp with rRNA-targeted peptide nucleic acid probes. *Appl Environ Microbiol* 66, 284–289
 40. Zardoya R, Costas E, Lopez-Rodas V, Carrido-Pertierra A, Bautista JM (1995) Revised dinoflagellate phylogeny inferred from molecular analysis of large subunit ribosomal RAN gene sequences. *J Mol Evol* 41, 637–645
 41. Zheng D, Alm EW, Stahl DA, Raskin L (1996) Characterization of universal small-subunit rRNA hybridization probes for quantitative molecular microbial ecology studies. *Appl Environ Microbiol* 62, 4504–4513