

the antennal scale, the form of the endopod of the second thoracic limbs, the number of joints in the tarsus of the posterior thoracic limbs, the form and armature of the telson, and the absence of spines on the inner uropod combine to differentiate this species from all its congeners.

LIST OF REFERENCES.

- CZERNIAVSKY, V. 1882. *Monographia Mysidarum imprimis Imperii Rossici*. Fasc. 1.
 CZERNIAVSKY, V. 1882. *Ibid.* Fasc. 2.
 CZERNIAVSKY, V. 1883. *Ibid.* Fasc. 3.
 HANSEN, H. J. 1910. The Schizopoda of the 'Siboga' Expedition. 'Siboga' Reports, No. xxxvii.
 ILLIG, G. 1906. Bericht über die neuen Schizopoden-gattungen und Arten der Deutschen Tiefsee-expedition, 1898-99. *Zool. Anz.* Bd. xxx. No. 7, pp. 194-211.
 KOSSMANN, R. 1880. *Zoologische Ergebnisse Reise in die Küsten des Rothen Meeres*, Hft. ii. Lief. i. Theil 2: Anomura.
 NORMAN, A. M. 1892. On British Mysidæ, a family of Crustacea Schizopoda. *Ann. Mag. Nat. Hist.* ser. 6, vol. x. pp. 143-166, 242-263.
 PAULSON, O. 1875. Carcinological Notes. *Zap. Obshch. Estestv. Kieff*, tom. iv. pp. 27-32.
 PAULSON, O. 1875. *Crustacea Mari Rubri*. Pars i.
 SÆRS, G. O. 1877. Nye bidrag til Kundskaben om Middelhavets Invertebratfauna. I. Middelhavets Mysider. *Arch. Math. Nat.* Bd. ii. pp. 10-119.
 TATTERSALL, W. M. 1922. Indian Mysidacea. *Rec. Ind. Mus.* vol. xxiv. pt. iv. pp. 445-504.
 ZIMMER, C. 1915 (1). Die Systematik der Tribus Mysini H. J. Hansen. *Zool. Anz.* Bd. xlvi. Nr. 7, pp. 202-216.
 ZIMMER, C. 1915 (2). Schizopoden des Hamburger Naturhistorischen (Zoologischen) Museums. *Mitt. Nat. Mus. Hamburg*, Bd. xxxii. pp. 159-182.

Appendix to the Report on the Crustacea Mysidacea.

By H. MUNRO FOX.

The Mysid Report brings out the following facts as regards penetration into the Canal. Eight Mediterranean species were collected, of which none have penetrated into the Canal beyond Port Said. Three Red Sea species were found, all of which have penetrated into the Canal. In addition, two forms new to science were discovered, one of them in the Canal. The only occasion on which Mysids (*Neomysis longicornis*) were found to be remarkably abundant was on December 15th at P. 3, in 1 fathom of

water on a sandy bottom. The following table gives the distribution of Mysids found in the Canal itself:—

TABLE 21.—Distribution of Mysids within the Canal.

		Red Sea & Indian Ocean species.			New species.
		<i>Siriella serrata.</i>	<i>Pseudanchialina inermis.</i>	<i>Siriella brevicaudata.</i>	<i>Heteromysis digitata.</i>
Ferry Post, Km. 76.		+			
Lake Timsah.	T. 9.		+		
	T. 15.		+		
	T. 8.	+		+	
Great Bitter Lake, K. 3.				+	
Kabret.	K. 1.			+	
	K. 6.	+	+	+	+
	K. 0.		+		
	K. 4.	+			