

Impacts of ocean acidification on marine fauna and ecosystem processes

Victoria J. Fabry, Brad A. Seibel, Richard A. Feely, and James C. Orr

Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. – *ICES Journal of Marine Science*, 65: 414–432.

Oceanic uptake of anthropogenic carbon dioxide (CO_2) is altering the seawater chemistry of the world's oceans with consequences for marine biota. Elevated partial pressure of CO_2 ($p\text{CO}_2$) is causing the calcium carbonate saturation horizon to shoal in many regions, particularly in high latitudes and regions that intersect with pronounced hypoxic zones. The ability of marine animals, most importantly pteropod molluscs, foraminifera, and some benthic invertebrates, to produce calcareous skeletal structures is directly affected by seawater CO_2 chemistry. CO_2 influences the physiology of marine organisms as well through acid-base imbalance and reduced oxygen transport capacity. The few studies at relevant $p\text{CO}_2$ levels impede our ability to predict future impacts on foodweb dynamics and other ecosystem processes. Here we present new observations, review available data, and identify priorities for future research, based on regions, ecosystems, taxa, and physiological processes believed to be most vulnerable to ocean acidification. We conclude that ocean acidification and the synergistic impacts of other anthropogenic stressors provide great potential for widespread changes to marine ecosystems.

Keywords: anthropogenic CO_2 , calcification, ecosystem impacts, hypercapnia, ocean acidification, physiological effects, zooplankton.

Received 11 July 2007; accepted 14 February 2008

V. J. Fabry: Department of Biological Sciences, California State University San Marcos, San Marcos, CA 92096–0001, USA. B. A. Seibel: Department of Biological Sciences, University of Rhode Island, Kingston RI 02881, USA. R. A. Feely: Pacific Marine Environmental Laboratory, NOAA, Seattle, WA 98115–6349, USA. J. C. Orr: Marine Environmental Laboratories, International Atomic Energy Agency, Monaco MC-98000, Monaco. Correspondence to V. J. Fabry: tel: +1 760 7504113; fax: +1 760 7503440; e-mail: fabry@csusm.edu.

Introduction

Rising atmospheric carbon dioxide (CO_2) concentration is causing global warming and ocean acidification (Caldeira and Wickett, 2003, 2005; Feely *et al.*, 2004; Orr *et al.*, 2005), which increasingly are recognized as important drivers of change in biological systems (Lovejoy and Hannah, 2005). For at least 650 000 years prior to the industrial revolution, atmospheric CO_2 concentrations varied between 180 and 300 ppmv (Siegenthaler *et al.*, 2005). As a result of human activity, today's atmospheric CO_2 concentration is 380 ppmv and currently is rising at a rate of $\sim 0.5\%$ year^{−1} (Forster *et al.*, 2007), which is ~ 100 times faster than any change during the past 650 000 years (Royal Society, 2005; Siegenthaler *et al.*, 2005). Approximately one-third of the anthropogenic CO_2 produced in the past 200 years has been taken up by the oceans (Sabine *et al.*, 2004). The global ocean inventory of anthropogenic carbon was 118 ± 19 Pg C in 2004 (Sabine *et al.*, 2004), which can be adjusted upwards to 140 Pg C in 2005 based on Denman *et al.* (2007, Table 7.1). Without this ocean sink, the anthropogenic change in atmospheric CO_2 concentration would be 55% higher than the observed change from 280 to 380 ppmv (Sabine *et al.*, 2004). Although oceanic uptake of anthropogenic CO_2 will lessen the extent of global warming, the direct effect of CO_2 on ocean chemistry may affect marine biota profoundly.

Elevated partial pressure of CO_2 ($p\text{CO}_2$) in seawater (also known as hypercapnia) can impact marine organisms both via decreased calcium carbonate (CaCO_3) saturation, which affects

calcification rates, and via disturbance to acid–base (metabolic) physiology. Recent work indicates that the oceanic uptake of anthropogenic CO_2 and the concomitant changes in seawater chemistry have adverse consequences for many calcifying organisms, and may result in changes to biodiversity, trophic interactions, and other ecosystem processes (Royal Society, 2005; Kleypas *et al.*, 2006). Most research has focused on tropical coral reefs and planktonic coccolithophores. Little information is available for other important taxa, for processes other than calcification, or for potential ecosystem-level consequences emerging from the oceanic $p\text{CO}_2$ levels that are predicted to occur over the next 100 years. Here we discuss the present and projected changes in ocean carbonate chemistry, and assess their impacts on pelagic and benthic marine fauna and ecosystem processes. We exclude corals from this discussion, but note that excellent recent reviews on this topic exist (Langdon and Atkinson, 2005; Guinotte *et al.*, 2006; Kleypas and Langdon, 2006). We highlight many of the gaps in our knowledge and identify critical questions for future research.

The ocean's inorganic carbon system: present and future changes

The CO_2 –carbonate system in seawater

The inorganic carbon system is one of the most important chemical equilibria in the ocean and is largely responsible for controlling the pH of seawater. Dissolved inorganic carbon (DIC) exists in

	Glacial	Pre-Industrial	Present	2XCO ₂	3XCO ₂	Change from pre-industrial to 3XCO ₂
pCO ₂	180	280	380	560	840	200%
CO ₂ (g) + H ₂ O ⇌ H ₂ CO ₃ Gas exchange Carbonic acid	7	9	13	18	25	178%
H ₂ CO ₃ ⇌ H ⁺ + HCO ₃ ⁻ Bicarbonate	1666	1739	1827	1925	2004	15%
HCO ₃ ⁻ ⇌ H ⁺ + CO ₃ ²⁻ Carbonate	279	222	186	146	115	-48%
DIC	1952	1970	2026	2090	2144	8.8%
pH _(sws)	8.32	8.16	8.05	7.91	7.76	-0.4
Ω _{calcite}	6.63	5.32	4.46	3.52	2.77	-48%
Ω _{aragonite}	4.26	3.44	2.90	2.29	1.81	-47%

Figure 1. Concentrations of carbon species (in units of $\mu\text{mol kg}^{-1}$), pH values, and aragonite and calcite saturation states of average surface seawater for $p\text{CO}_2$ concentrations (ppmv) during glacial, preindustrial, present day, two times pre-industrial CO_2 , and three times pre-industrial CO_2 . Changes in the inorganic carbon system were computed by assuming equilibrium with atmospheric CO_2 , assuming $\text{PO}_4 = 0.5 \mu\text{mol l}^{-1}$ and $\text{Si} = 4.8 \mu\text{mol l}^{-1}$, and using the carbonic acid dissociation constants of Mehrbach *et al.* (1973) as refitted by Dickson and Millero (1987). pH is based on the seawater scale. The last column shows the changes from the pre-industrial levels to three times atmospheric CO_2 (modified from Feely *et al.* (2004) and Kleypas *et al.* (2006)).

seawater in three major forms: bicarbonate ion (HCO_3^-), carbonate ion (CO_3^{2-}), and aqueous carbon dioxide ($\text{CO}_{2(\text{aq})}$), which here also includes carbonic acid (H_2CO_3). At a pH of 8.2, $\sim 88\%$ of the carbon is in the form of HCO_3^- , 11% in the form of CO_3^{2-} , and only $\sim 0.5\%$ of the carbon is in the form of dissolved CO_2 . When CO_2 dissolves in seawater, H_2CO_3 is formed (Figure 1). Most of the H_2CO_3 quickly dissociates into a hydrogen ion (H^+) and HCO_3^- . A hydrogen ion can then react with a CO_3^{2-} to form bicarbonate. Therefore, the net effect of adding CO_2 to seawater is to increase the concentrations of H_2CO_3 , HCO_3^- , and H^+ , and decrease the concentration of CO_3^{2-} and lower pH ($\text{pH} = -\log[\text{H}^+]$). These reactions are fully reversible, and the basic thermodynamics of these reactions in seawater are well known (Millero *et al.*, 2002). The atmospheric CO_2 value today is ~ 100 ppmv greater than the pre-industrial value (280 ppmv), and the average surface ocean pH has dropped by 0.1 unit, which is about a 30% increase in $[\text{H}^+]$. Under the IPCC emission scenarios (Houghton *et al.*, 2001), average surface ocean pH could decrease by 0.3–0.4 pH units from the pre-industrial values by the end of this century (Caldeira and Wickett, 2005; Figure 2).

Present and future changes in carbonate saturation

The reaction of CO_2 with seawater reduces the availability of carbonate ions that are necessary for marine calcifying organisms, such as corals, molluscs, echinoderms, and crustaceans, to produce their CaCO_3 shells and skeletons. The extent to which such organisms are affected depends largely upon the CaCO_3 saturation state (Ω), which is the product of the concentrations of Ca^{2+} and CO_3^{2-} , divided by the apparent stoichiometric solubility product (K_{sp}^*) for either aragonite or calcite, two types of CaCO_3





Figure 2. Atmospheric CO_2 concentration projected under the IS92a “business-as-usual” IS92a CO_2 emissions scenario, bounded by the most and least conservative SRES scenarios, B1 and A1F1, respectively, and projected global average surface seawater pH (modified from Meehl *et al.* (2007)).

commonly secreted by marine organisms:

$$\Omega = [\text{Ca}^{+2}][\text{CO}_3^{2-}]/K_{\text{sp}}^*, \quad (1)$$

where the calcium concentration is estimated from the salinity, and $[\text{CO}_3^{2-}]$ is calculated from DIC and total alkalinity (TA) measurements (Feely *et al.*, 2004). Increasing CO_2 concentrations in the atmosphere, and thus in the surface ocean, will continue to decrease the $[\text{CO}_3^{2-}]$ in the upper ocean, thereby lowering CaCO_3 saturation levels by means of the reaction:



In regions where Ω_{arag} or Ω_{cal} is > 1.0 , the formation of shells and skeletons is favoured. For values < 1.0 , seawater is corrosive to CaCO_3 and, in the absence of protective mechanisms (e.g. Corliss and Honjo, 1981; Isaji, 1995), dissolution will begin. Saturation states are generally highest in the tropics and lowest in the high latitudes, because the solubility of CaCO_3 increases with decreasing temperature and increasing pressure. Consequently, there is significant shoaling of the aragonite saturation horizons in the Pacific, north of $\sim 40^\circ\text{N}$, at the equator, and 10°N , especially towards the east, because of the higher DIC concentrations relative to TA at shallower depths. These patterns result from enhanced upwelling that brings deeper waters rich in nutrients and DIC to the upper ocean (Figure 3) and supports high animal biomass. As one moves north, the aragonite saturation depth shoals from ~ 1000 m near 30°S to 300 m at the equator. Moving farther north, it deepens to 550 m near 30°N , then shoals to ~ 100 m north of 50°N (Figure 3). In the North Pacific, the upward migration of the aragonite saturation horizon from anthropogenic CO_2 uptake is currently $\sim 1\text{--}2$ m year^{-1} (Feely *et al.*, 2006).

Orr *et al.* (2005) developed model scenarios of future changes in surface ocean carbonate chemistry as a function of changes in atmospheric CO_2 , using the IPCC IS92a “business-as-usual” CO_2 emission scenario, with the median projection of DIC changes from 13 ocean models that participated in the OCMIP-2 project. Based on their model outputs and global gridded data (Key *et al.*, 2004), we plotted the projected aragonite




Figure 3. Distribution of (a) aragonite saturation; (b) partial pressure of CO_2 seawater ($p\text{CO}_2$); and (c) dissolved oxygen along the March 2006 P16 N transect along 152°W in the North Pacific.

saturation state of the surface oceans for the years 1765, 1994, 2050, and 2100 (Figure 4). The model results indicate that, by the time atmospheric CO_2 reaches 780 ppmv near the end of this century under the IPCC IS92a “business-as-usual” CO_2 emission scenario, portions of the Subarctic North Pacific and all of the Southern Ocean south of $\sim 60^\circ\text{S}$ will become undersaturated with respect to aragonite (Orr *et al.*, 2005). At that point, the global average surface water CO_3^{2-} concentration and aragonite and calcite saturation state will be nearly half of what they are today. The aragonite saturation horizons would also shoal from its present average depth of 730 m to the surface in the Southern Ocean, from 2600 to 115 m in the North Atlantic, and from 140 m to the surface in parts of the North Pacific (Orr *et al.*, 2005). In the cold, high-latitude surface waters typical of polar and subpolar regions of the Southern Ocean, aragonite and calcite undersaturation will occur when seawater $p\text{CO}_2$ values

reach ~ 560 and 900 ppmv, respectively. In the slightly warmer surface waters of the subpolar North Pacific, aragonite and calcite undersaturation will occur later, when $p\text{CO}_2$ reaches 740 and 1040 ppmv, respectively. The cold waters of the Arctic Ocean are also naturally low in CO_3^{2-} concentration. Continuing research is evaluating how the Arctic Ocean’s changes in carbonate chemistry during the 21st century will differ from those in the Southern Ocean (Orr *et al.*, 2006). The warm surface waters of the tropics and subtropics will not become undersaturated with respect to aragonite or calcite over the range of these projected conditions although, in some regions associated with upwelling, shoaling aragonite saturation horizons now impinge on the depth ranges of many pelagic animals (Feely *et al.*, 2004).

Priority areas for ocean acidification research are therefore high-latitude regions, which are projected to experience the greatest changes in carbonate chemistry over decadal to century




Figure 4. Surface water aragonite saturation state (Ω_{arag}) for the pre-industrial ocean (nominal year 1765), and years 1994, 2050, and 2100. Values for years 1765 and 1994 were computed from the global gridded data product GLODAP (Key *et al.*, 2004), whereas the saturation state for years 2050 and 2100 are the median of 13 ocean general circulation models forced under the IPCC's IS92a "business-as-usual" CO_2 emission scenario (Orr *et al.*, 2005).

time-scales. Moreover, coastal regions, which can be greatly impacted by anthropogenic-driven ocean acidification (Doney *et al.*, 2007), as well as eutrophication leading to low oxygen zones, are not well-represented in global ocean-atmosphere coupled models, because of lack of data, complexities of nearshore circulation processes, and too-coarse model resolution. Given the importance of coastal areas to fisheries and other marine resources and services, coastal ecosystems constitute another target region where research is urgently needed.

Effects of elevated $p\text{CO}_2$ on calcification

The secretion of CaCO_3 skeletal structures is widespread across animal phyla, and evolved independently and repeatedly over geologic time since the late Precambrian period (Knoll, 2003). Protection is one probable advantage of possessing a calcareous skeleton. Additionally, various other biotic and abiotic factors have probably contributed to selection for CaCO_3 hard parts in diverse groups of fauna at different times in evolutionary history.

Most calcifying organisms investigated to date demonstrate reduced calcification in response to increased $p\text{CO}_2$ and decreased $[\text{CO}_3^{2-}]$, CaCO_3 saturation state, and pH (e.g. Gattuso *et al.*, 1998; Langdon *et al.*, 2000, 2003; Riebesell *et al.*, 2000). The majority of work has tested warm-water corals and coccolithophorid algae (Royal Society, 2005; Kleypas *et al.*, 2006). Evidence suggests that the calcification rate in corals is controlled by the CaCO_3 saturation state (Gattuso *et al.*, 1998; Langdon *et al.*, 2000, 2003; Marubini *et al.*, 2001, 2003; Leclercq *et al.*, 2002; Ohde and Hossain, 2004; Langdon and Atkinson, 2005; Schneider and Erez, 2006; Silverman *et al.*, 2007), rather than pH or another parameter of the seawater CO_2 system. Because the $[\text{Ca}^{2+}]$ in the ocean today is approximately constant (depending predominantly on salinity), changes in the $[\text{CO}_3^{2-}]$ are reflected directly as changes in the CaCO_3 saturation state. In contrast, high $[\text{Ca}^{2+}]$ in the Cretaceous (Horita *et al.*, 2002) allowed planktonic calcifiers to flourish and large chalk deposits to accumulate (Bown *et al.*, 2004), despite CO_3^{2-} concentrations that were only $\sim 25\%$ of the current value (Tyrrell and Zeebe, 2004; Ridgwell and Zeebe, 2005). This observation supports the idea that the CaCO_3 saturation state [proportional to the product of Ca^{2+} and CO_3^{2-} concentrations as shown in Equation (1)] is the key component of the seawater carbonate system that controls calcification rates.

Holoplankton

The major planktonic CaCO_3 producers are the coccolithophores, foraminifera, and euthecosomatous pteropods. These three groups of calcifiers account for nearly all the export flux of CaCO_3 from the upper ocean to the deep sea. Planktonic foraminifera and coccolithophores secrete tests or shells made of calcite, whereas pteropods form shells made of aragonite, a metastable polymorph of CaCO_3 , which is $\sim 50\%$ more soluble in seawater than calcite (Mucci, 1983). On a global basis, coccolithophores and foraminifera are thought to produce the majority of pelagic CaCO_3 (Schiebel, 2002), while the labile aragonitic shells of pteropods account for a smaller fraction of the total CaCO_3 produced by planktonic organisms. The relative contributions of these three groups of calcifiers can vary substantially on regional and temporal scales, however. There are few concurrent measurements of the abundances of all three groups, and estimates of their contributions to global calcification rates are poorly constrained.

Both planktonic foraminifera and euthecosomatous pteropods are widely distributed in the upper ocean, with highest species diversity in tropical and subtropical regions. Shelled pteropods can reach densities of 1000s to $>10\,000$ of individuals m^{-3} in high-latitude areas (e.g. Bathmann *et al.*, 1991; Pane *et al.*, 2004) and are important components of polar and subpolar ecosystems. Data are limited on the response of calcification in pteropods and foraminifera to elevated $p\text{CO}_2$ and decreased CaCO_3 saturation state. Currently, evidence is available for only two of the ~ 50 species of planktonic foraminifera and only one of the ~ 34

euthecosome species. Although these data suggest that both groups reduce calcification in response to ocean acidification, the small number of species tested precludes the identification of general trends. Species-specific responses are likely, and it is possible that the calcification rates of some species may not be sensitive to elevated $p\text{CO}_2$, as has been found in coccolithophores (Riebesell *et al.*, 2000; Langer *et al.*, 2006).

In laboratory experiments with the symbiont-bearing planktonic foraminifera *Orbulina universa* and *Globigerinoides sacculifer*, shell mass decreased in response to reduced CO_3^{2-} concentration and calcite saturation state, even though the seawater was supersaturated with respect to calcite (Spero *et al.*, 1997; Bijma *et al.*, 1999, 2002). When grown in seawater chemistry equivalent to $p\text{CO}_2$ of 560 and 740 ppmv, shell mass in these species declined by 4–8 and 6–14%, respectively, compared with the pre-industrial $p\text{CO}_2$ value. Evidence from microelectrode and culture experiments suggests that elevated pH and CO_3^{2-} concentration in the micro-environment immediately adjacent to the shell are critical in promoting shell growth (Spero and Lea, 1993; Rink *et al.*, 1998; Köhler-Rink and Kühl, 2005). When *O. universa* is grown under high light, the pH of the micro-environment surrounding the test can be increased up to 8.8 (0.5 units above ambient seawater) as a result of CO_2 removal during symbiont photosynthesis. In contrast, *O. universa* grown in the dark has a near-shell micro-environment pH of 7.9, owing to respiratory CO_2 release (Köhler-Rink and Kühl, 2005). Because of the large effect of symbiont photosynthesis on seawater CO_2 chemistry at the shell surface, it may be that the impact of ocean acidification on adult symbiont-bearing foraminifera will occur primarily during night calcification. It is unknown whether foraminifera that do not possess photosynthetic symbionts are more susceptible to reduced CO_3^{2-} concentration and calcite saturation state than those species with symbionts. Similarly, the post-zygote, prolocular ontogenetic stage of foraminifera, during which calcification is weak or absent and symbiotic algae have not yet been acquired (Brummer *et al.*, 1987), may be particularly vulnerable to elevated $p\text{CO}_2$. Because it is currently impossible to maintain the prolocular stage in the laboratory, however, this hypothesis awaits testing.

A positive correlation between foraminiferal shell mass and ambient $[\text{CO}_3^{2-}]$ is observed in the sedimentary record as a response to known glacial–interglacial changes in atmospheric CO_2 of the past 50 000 years (Barker and Elderfield, 2002). However, Barker and Elderfield (2002) demonstrate an increase in *Globigerinoides bulloides* shell mass from 11 to 19 μg for a small change in $[\text{CO}_3^{2-}]$ from 210 to 250 $\mu\text{mol kg}^{-1}$, whereas Bijma *et al.* (2002) report a much greater increase, from 40 to 60 μg in shell mass for *O. universa* for a change in $[\text{CO}_3^{2-}]$ from 200 to 600 $\mu\text{mol kg}^{-1}$, and $\sim 10 \mu\text{g}$ increase in shell mass for *G. sacculifer* for a change in $[\text{CO}_3^{2-}]$ from 100 to 600 $\mu\text{mol kg}^{-1}$. Although it appears that culture experiments may underestimate foraminiferal response to altered $[\text{CO}_3^{2-}]$ compared with the change observed in paleo-reconstructions over glacial-to-interglacial time-scales, temperature and food supply also strongly affect foraminiferal calcification (cf. Barker and Elderfield, 2002; Bijma *et al.*, 2002). Hence, future increases in sea surface temperatures could lead to higher foraminiferal growth rates.

Owing to their highly soluble aragonitic shells, pteropods may be particularly sensitive to ocean acidification. When live *Clio pyramidata* collected in the Subarctic Pacific were exposed to a level of aragonite undersaturation similar to that projected for Southern Ocean surface waters in year 2100 under the IS92a emissions

scenario, there was marked shell dissolution within 48 h (Feely *et al.*, 2004; Orr *et al.*, 2005). In additional experiments, *C. pyramidata* were placed in 1 l jars, and ^{45}Ca was added to measure shell calcification rates. The jars were then closed and incubated at 10°C for 4–48 h. At the start of this experiment, the seawater Ω_{arag} was ~ 2.4 . Forty-eight hours later, respiratory CO_2 from the actively swimming pteropods had gradually forced the aragonite saturation state to drop below 1. The ^{45}Ca uptake experiments reveal progressively reduced calcification rates in concert with decreased aragonite saturation state that resulted from the accumulation of metabolic CO_2 produced by animals during the experiment (Figure 5). After 36–48 h, most of the ^{45}Ca that had been incorporated into shells had dissolved back into solution. Hence, by 36 h, net dissolution exceeded net calcification, although animals were alive and actively swimming. Scanning electron microscopic photographs of experimental shells confirm that dissolution occurred along the leading edge of the shell by 48 h during incubation in closed jars, but no dissolution was visible in control shells incubated in open jars, where the aragonite saturation remained >1 (Figure 6). Based on pteropod oxygen consumption rates (Smith and Teal, 1973), it is unlikely that animals were oxygen-stressed during these experiments, but future experiments should explore this possibility. Additional experiments are needed to measure calcification directly in euthecosomatous pteropods and foraminifera, as a function of CaCO_3 saturation state. It will also be necessary to study the interactive effects of CaCO_3 saturation state, temperature, and nutrition for multiple species and life stages of these calcareous holoplankton.

As $p\text{CO}_2$ rises and the CaCO_3 saturation state of surface ocean decreases, euthecosomatous pteropods and foraminifera may secrete under-calcified or thinner structures (Bijma *et al.*, 1999). Should the habitat of these organisms approach CaCO_3 undersaturation, first with respect to aragonite, then with respect to calcite, the question of whether net rates of calcification can still exceed dissolution will likely depend on the degree of CaCO_3 undersaturation and the duration that animals are exposed to undersaturated waters. As yet, no decreases in calcification have been documented in field populations of either group. However,




Figure 5. Net calcification ($\mu\text{mol CaCO}_3 \text{ h}^{-1}$) as a function of shell CaCO_3 concentration for the Subarctic Pacific euthecosomatous pteropod *Clio pyramidata*. Different sizes of pteropods were incubated at 10°C in closed 1 l jars for 4–48 h. Net calcification rates decreased with time, as the aragonite saturation state of seawater was progressively reduced owing to respiratory CO_2 in sample containers.




Figure 6. SEM photographs of the shell of the pteropod *Clio pyramidata*. (a) Whole shell of an animal incubated for 48 h in a closed container, wherein animal respiration forced the aragonite saturation state < 1 ; (b) magnified portion of the leading edge of the shell section shown in (a), revealing dissolution of aragonitic rods; (c) magnified leading edge of the shell of a control animal incubated in seawater that remained supersaturated with aragonite for the duration of the experiment.

baseline data on their present-day vertical distributions and calcification rates are insufficient to detect possible changes that may result from ocean acidification.

The capacity of euthecosome and foraminifera species to adapt to progressively acidified ocean waters is not known, but may be related to species' generation time. If unabated CO_2 emissions continue and surface waters of the Southern Ocean and portions of the Subarctic Pacific become undersaturated with respect to aragonite by 2100 as projected (Orr *et al.*, 2005), then shelled pteropods in these regions would have only ~ 50 – 150 generations to adapt to corrosive seawater, given that high-latitude pteropods are thought to have generation times of 0.6–1.5 years (Kobayashi, 1974; Bathmann *et al.*, 1991; Dadon and de Cidre, 1992; Gannefors *et al.*, 2005). Generation times for spinose species of foraminifera are frequently linked with the lunar cycle such that they reproduce every 2–4 weeks; however, non-spinose species probably have longer reproductive cycles (Hemleben *et al.*, 1989). Shorter generation time affords increased opportunities for microevolutionary adaptation.

In addition to euthecosomes and foraminifera, other holoplankton calcify during part or all of their life cycles. These include the heteropods, visual predators that are found in the epipelagic zone of all tropical and subtropical oceans, but which are absent from high latitudes. Two of the three heteropod families possess aragonitic shells as adults, and species in the third family cast off their larval shells at metamorphosis. Gymnosomes, the highly specialized predators of euthecosomatous pteropods, possess a large veliger shell, presumed to be aragonite, which is also cast off at metamorphosis. Similarly, pseudothecosomatous pteropods possess a veliger shell that is discarded at metamorphosis to the gelatinous adult.

Benthic invertebrates

Nine multicellular invertebrate phyla have benthic representatives with CaCO_3 skeletal hard parts (Lowenstam and Weiner, 1989). These taxa secrete CaCO_3 in the form of aragonite, calcite, high-magnesium calcite (>5 mole % MgCO_3), amorphous CaCO_3 , or a mixture of these CaCO_3 phases. Amorphous CaCO_3 is less stable than the crystalline phases of CaCO_3 , and the seawater solubility of high-magnesium calcite is similar to or greater than that of aragonite (Walter and Morse, 1985; Bischoff *et al.*, 1987). Many

benthic calcifying fauna are prominent in nearshore communities and are economically and/or ecologically important. For example, bivalves, such as mussels and oysters, have high commercial value as fisheries and are also important as ecosystem engineers in coastal areas, providing habitat and other services for a rich diversity of organisms (Gutiérrez *et al.*, 2003). Recent work suggests that benthic adult molluscs and echinoderms are sensitive to changes in seawater carbonate chemistry. In response to an elevated $p\text{CO}_2$ level projected to occur under the IS92a emissions scenario (~ 740 ppmv in 2100), calcification rates in the mussel *Mytilus edulis* and the Pacific oyster *Crassostrea gigas* decreased by 25 and 10%, respectively (Gazeau *et al.*, 2007). When grown for more than six months in seawater bubbled with air containing 560 ppmv CO_2 , a decrease in shell growth was observed in the edible snail *Strombus luhuanus*, and a reduction in wet weight was reported in both this snail and two species of sea urchins (Shirayama and Thorton, 2005).

Early calcifying stages of benthic molluscs and echinoids demonstrate a strong response to increased seawater $p\text{CO}_2$ and decreased pH, CO_3^{2-} concentration, and CaCO_3 saturation state. In the sea urchins *Hemicentrotus pulcherrimus* and *Echinodetra mathaei*, fertilization success, developmental rates, and larval size all decreased with increasing CO_2 concentration (Kurihara and Shirayama, 2004). Abnormal skeletogenesis of the highly soluble high-magnesium CaCO_3 spicules in urchin larvae was also observed. Green *et al.* (2004) found that newly settled juveniles of the hard-shell clam *Mercenaria mercenaria* revealed substantial shell dissolution and increased mortality when they were introduced to surface sediments that were undersaturated with respect to aragonite ($\Omega_{\text{arag}} \sim 0.3$), a level that is typical of near-shore, organic-rich surficial sediments. Within two weeks of settlement, the CaCO_3 shells were completely dissolved, leaving only the organic matrix of the shell.

The mineralogy and calcification mechanisms of mollusc and echinoid larval stages may render them particularly sensitive to ocean acidification. Although adult gastropods and bivalves secrete aragonite, calcite, or both phases in a diverse array of structural patterns, Weiss *et al.* (2002) suggest that veliger shells of gastropods and bivalves all contain aragonite in similar crystalline ultrastructures, and hence, the mollusc larval shell is highly conserved in evolution. Moreover, recent work using infrared

spectrometry and Raman imaging spectroscopy reveals that larvae of the clam *M. mercenaria* (adult shell is aragonitic), as well as that of the oyster *C. gigas* (adult shell is nearly entirely calcitic), form amorphous CaCO_3 as a transient precursor to aragonite (Weiss *et al.*, 2002). Similarly, during the embryonic development of two species of sea urchins, an amorphous CaCO_3 precursor transforms to calcite during spicule formation (Beniash *et al.*, 1997; Raz *et al.*, 2003). Because this unstable, transient, amorphous CaCO_3 is more soluble than the crystalline minerals of aragonite or calcite, biomineralization processes that occur during the embryonic and larval development of sea urchins, and in gastropod and bivalve molluscs, may be particularly vulnerable to ocean acidification.

CaCO_3 skeletal elements are also present in species of other benthic invertebrates, such as crustaceans, cnidarians, sponges, bryozoans, annelids, brachiopods, and tunicates. Apart from warm-water corals, nothing is known about the effect of elevated ambient $p\text{CO}_2$ on calcification rates in these taxa. Some of these animals may use shell dissolution to support acid-based regulation at high internal $p\text{CO}_2$, as has been observed in mussels (Michaelidis *et al.*, 2005) and other organisms.

Other non-skeletal, calcified secretions of marine fauna

In addition to using CaCO_3 for strengthening skeletal structures, the use of calcium minerals in gravity sensory organs is widespread among ocean fauna. In the many zooplankton and benthic invertebrates that possess statoliths, statocysts, or statocontia, the mineralogy is indeterminate or is reported to be amorphous Ca-Mg-phosphate or gypsum (Lowenstam and Weiner, 1989). In squid and fish, however, the statoliths and otoliths are composed of aragonite. Gravity sensory organs can have additional functions in some organisms. For example, in planktonic gymnosome snails, statocysts also are actively involved in the motor neural programme that underlies search movements for prey during hunting behaviour (Levi *et al.*, 2004). Whether mineralization of the various types of gravity receptors would be affected by the changing carbonate chemistry of seawater and, if so, how that might impact overall fitness of the organism, are questions that have not been investigated. Presumably, potential impacts would depend on the ability of the organisms to regulate the acid–base balance in the tissues surrounding those structures.

Other carbonate secretions of marine fauna include gastroliths, mineralized structures formed in the lining of the cardiac stomachs of some decapods that serve as storage sites for calcium during moult intervals; gastroliths can be calcite, amorphous CaCO_3 , or calcium phosphate (cf. Lowenstam and Weiner, 1989). Widespread among marine fish is the intestinal secretion of calcium- and magnesium-rich carbonate complexes, which are then excreted via the rectum; this process appears to play a critical role in osmoregulation (Walsh *et al.*, 1991; Grosell, 2006; Taylor and Grosell, 2006).

Consequences of reduced calcification

The effects of chronic exposure to increased CO_2 on calcifiers, as well as the long-term implications of reduced calcification rates within individual species and their ecological communities, are unknown. Calcification probably serves multiple functions in carbonate producers. Decreased calcification would presumably compromise the fitness of these organisms and could shift the competitive advantage towards non-calcifiers. Such a consequence

is supported by recent work with warm-water reef organisms in which a decreased aragonite saturation state induced the transition from a CaCO_3 -dominated system to one dominated by organic algae (Kuffner *et al.*, 2008).

Increased $p\text{CO}_2$ and other physiological processes

In addition to calcification, a number of other physiological indices appear to correlate with the capacity for acid–base tolerance, and new data are emerging that test the survival, growth, development, metabolism, and pH balance of organisms under elevated $p\text{CO}_2$. Prior to 1995, most studies used CO_2 values that were well above what is expected in the future ocean and from convenient marine animal models, in order to reveal the fundamental mechanisms associated with acid–base regulation. Studies conducted more recently have tested the physiological response to very high CO_2 levels as would be associated with purposeful sequestration of CO_2 in the ocean. There is now a critical need to test the physiological consequences of ocean acidification at lower $p\text{CO}_2$ levels, such as those that are projected to occur over the next century.

Mechanisms to deal with hypercapnia

When $p\text{CO}_2$ levels increase in seawater, dissolved CO_2 more readily diffuses across animal surfaces and equilibrates in both intra- and extracellular spaces. Internal levels rise until a new value is reached that is sufficient to restore CO_2 excretion against the elevated environmental level. As in seawater, CO_2 reacts with internal body fluids causing H^+ to increase and, therefore, pH to decrease. Mechanisms available to counteract this acidification are limited and relatively conserved across animal phyla. The mechanisms are the same as those evolved to deal with metabolically produced CO_2 and hydrogen ions. They include (i) passive buffering of intra- and extracellular fluids; (ii) transport and exchange of relevant ions; (iii) transport of CO_2 in the blood in those species that have respiratory pigments; (iv) metabolic suppression to wait out periods of elevated CO_2 (e.g. Somero, 1985; Truchot, 1987; Cameron, 1989; Walsh and Milligan, 1989; Hand, 1991; Heisler, 1993; Guppy and Withers, 1999; Clairborne *et al.*, 2002; Seibel and Walsh, 2003; Pörtner *et al.*, 2004). Those species adapted to environments with steep CO_2 gradients, such as hydrothermal vents or stagnant tide pools, and those species with high capacity for metabolic production of CO_2 have evolved greater capacities for buffering, ion exchange, and CO_2 transport (Seibel and Walsh, 2001, 2003). Whether such elevated capacity translates into greater tolerance of chronic ocean acidification remains to be seen.

Buffering capacity

When concentrations are elevated, CO_2 readily crosses biological membranes and enters the blood and intracellular spaces. Passive buffering is the only mechanism immediately available to limit pH changes within the body. Locomotory muscles of active animals, such as epipelagic fish and cephalopods, have high activities of anaerobic metabolic enzymes and, consequently, have high capacity for buffering pH changes associated with anaerobically fuelled burst locomotion (Castellini and Somero, 1981; Seibel *et al.*, 1997). Organisms with low buffering capacity will experience greater fluctuations in intracellular pH during hypercapnia than others with higher capacity. For example, an increase in seawater $p\text{CO}_2$ sufficient to lower intracellular pH by 0.2 in a sluggish benthic fish may cause only a 0.02 pH unit drop in an active

epipelagic fish such as tuna (Seibel and Walsh, 2003). Buffering of extracellular fluid is, in some cases, also provided by formation of bicarbonate from dissolution of CaCO_3 stores or exoskeletons (shells or tests) as discussed below.

Ion transport

In the longer term (hours to days), compensation of acid–base imbalance relies on the ability to transport acid–base equivalent ions across cell membranes. The CO_2 that is produced in the cells during routine metabolism is typically hydrated to form bicarbonate and H^+ , a reaction catalyzed by the enzyme carbonic anhydrase. These hydrogen ions are then buffered in the intracellular space as discussed above, while the bicarbonate is transported out of the cell in exchange for Cl^- via ion transport proteins. Species with ineffective ion transport capacities are poorly equipped for acid–base regulation (Heisler, 1989; Walsh and Milligan, 1989). Low rates of metabolism typically correlate with lower concentrations of ion transport proteins such as Na^+/K^+ and H^+ -ATPases (Gibbs and Somero, 1990), suggesting reduced capacities of acid–base balance. In many cases, the gelatinous tissues (e.g. mesoglea) found in diverse zooplankton are distinct from the muscle tissues that are most metabolically active (Thuesen *et al.*, 2005a, b). Therefore, the active tissues of gelatinous animals may have greater CO_2 tolerance than would be estimated, based on rates of whole-animal metabolism. Similarly, species not exposed to environmental fluctuations in CO_2 may also be ill-equipped to handle ocean acidification. For example, species from hydrothermal vents have high activities of carbonic anhydrase relative to species from shallower environments. Even lower still are carbonic anhydrase activities of benthic deep-sea species, far removed from venting water, that experience very little fluctuation in environmental CO_2 (Figure 7). Compensation of acidosis via adjustments in ionic composition appears to be a trade-off that is not likely sustainable on longer time-scales, such as that associated with anthropogenic increases in seawater $p\text{CO}_2$. Nevertheless, a survey of species that are more or less able to regulate the pH of their internal fluids may be informative.

Acid–base regulation via bicarbonate accumulation

A common component of pH compensation in animals is the intracellular accumulation of HCO_3^- (Walsh and Milligan, 1989; Pörtner and Reipschläger, 1996) that drives an elevation in pH. A “bicarbonate threshold” is hypothesized (Heisler, 1993; Pörtner and Reipschläger, 1996), above which the capacity for further compensation is limited. Although additional bicarbonate accumulation will always further compensate reduced pH, there may be an upper limit beyond which acid–base regulation begins to compromise ionic balance (Cameron and Iwama, 1987). Those species that can tolerate accumulation of bicarbonate to levels 4–10 times above control conditions appear generally more tolerant of hypercapnia, whereas those with limited bicarbonate accumulation may be more vulnerable.

Miles *et al.* (2007) recently found incomplete compensation of coelomic fluid pH, despite elevated bicarbonate levels under all CO_2 exposures (pH 6.2–7.4), in an intertidal sea urchin. Dissolution of the high magnesium calcite test was inferred from elevations in coelomic Mg^{2+} , and the authors suggested that reductions in surface seawater pH below 7.5 would be severely detrimental to this species, and probably other sea urchins as well. These results are consistent with those of Burnett *et al.* (2002)




Figure 7. Carbonic anhydrase activity ($\Delta\text{pH min}^{-1} \text{g wet mass}^{-1}$) in gas exchange tissue of benthic macrofauna in habitats with large fluctuations in environmental $p\text{CO}_2$. Bivalves (hatched bars), worms (grey bars), and crustaceans (open bars). Each bar represents the mean of several species (note the x-axis is on a log scale). Species are grouped as ionic regulators (euryhaline species including those from hydrothermal vents), shallow living ionic conformers (those from stable shallow-water habitats), and deep-sea conformers. Data for worms and bivalves are from Kochevar and Childress (1996). Data for crustaceans are from Henry (1984) and unpublished observations (J. Company, pers. comm.).

and Spicer (1995), demonstrating that sea urchins are unable to compensate acidosis resulting from short-term emersion and hypoxia, respectively.

Similarly, the mussel *M. edulis* compensated both short- and long-term exposure to 1% CO_2 (~10 000 ppmv) by dissolution of its shell as indicated by increased Ca^{2+} levels (Lindinger *et al.*, 1984; Michaelidis *et al.*, 2005). Not surprisingly, long-term exposure resulted in reduced growth and metabolism. A deep-sea crab measured recently by Pane and Barry (2007) failed to accumulate any bicarbonate or control haemolymph pH values over 24 h exposure to hypercapnia. This poor performance was attributed to low rates of metabolism, stable environmental conditions, and reduced oxygen at depth that may have limited ion exchange capacity. In short-term experiments, the sipunculid worm *Sipunculus nudus* demonstrated hypercapnic tolerance, with only a 50% elevation in extracellular bicarbonate relative to control levels (Pörtner and Reipschläger, 1996). The reduced metabolic rate observed in this species (40% of control values), under natural, short-term elevation of CO_2 , allows survival until well-aerated waters return with high tide. However, over the longer term (3–6 week), such metabolic suppression resulted in 100% mortality (Langenbuch and Pörtner, 2004).

Species that are more tolerant exhibit greater bicarbonate accumulation and, consequently, compensate more completely the acidosis caused by exposure to elevated CO_2 . For example, exposure of the subtidal crab *Necora puber* to $p\text{CO}_2$ ~10 000 ppmv resulted in haemolymph bicarbonate concentrations more than four times the control levels, in part supplied by shell dissolution (Spicer *et al.*, 2007). Similarly, the crab *Cancer magister* fully compensated its haemolymph pH over

24 h by accumulating more than 12 mM bicarbonate (Pane and Barry, 2007). Fish appear most tolerant among marine animals. The Mediterranean fish *Sparus aurata* was able to compensate completely both blood plasma pH and intracellular pH in the face of 10 000 ppm $p\text{CO}_2$ via elevations in bicarbonate to five times the control levels. No mortality occurred after ten days of exposure (Michaelidis *et al.*, 2007). Although feed intake was reduced at a seawater pH of 7.25, compensation of internal acid–base imbalance was accomplished in the sea bass *Dicentrarchus labrax* via a fivefold elevation in plasma bicarbonate (Cecchini *et al.*, 2001).

Mortality

Mortality occurred in three fish species tested, including yellowtail and flounder, only at very high CO_2 levels ($>50\,000$ ppmv) after 24 h exposure, and the authors concluded that fish mortality caused by anthropogenic CO_2 is never expected in marine environments (Hayashi *et al.*, 2004). Although we believe that this statement is premature, marine fish do appear highly tolerant of CO_2 (Kikkawa *et al.*, 2004, 2006). The hatchling stages of some species appeared fairly sensitive to pH decreases on the order of 0.5 or greater, but high CO_2 tolerance developed within a few days of hatching (Ishimatsu *et al.*, 2004). The relative tolerance of fish and others may relate to high capacity for internal ion and acid–base regulation via direct proton excretion (Ishimatsu *et al.*, 2004) and an intracellular respiratory protein that results in a high oxygen-carrying capacity and substantial venous oxygen reserve. When compensation of pH fails, mortality of all marine animals increases with the level of CO_2 and the duration of exposure (Yamada and Ikeda, 1999; Hayashi *et al.*, 2004; Watanabe *et al.*, 2006; Table 1).

Metabolic suppression

If compensation of acid–base imbalance is not achieved, reduced pH and elevated $p\text{CO}_2$ may depress metabolism in some species (Hand, 1991; Pörtner and Reipschläger, 1996; Guppy and Withers, 1999; Figure 8). Metabolic suppression is considered an adaptive strategy for the survival of short-term hypercapnia and hypoxia. During periods of environmental oxygen limitation, many organisms are able to suppress ATP demand, thereby extending the duration of tolerance. In many cases, oxygen limitation is coincident with internal acid–base disturbance. Metabolic suppression is not advantageous, however, under chronic elevations of CO_2 (e.g. *S. nudus*, as cited above; Langenbuch and Pörtner, 2004). Metabolic suppression is typically achieved by shutting down expensive processes. Chief among these is protein synthesis (Hand, 1991). Reduced protein synthesis, by definition, will reduce growth and reproductive potential. Although suppression of metabolism under short-term experimental conditions is a “sublethal” reversible process, reductions in growth and reproductive output will effectively diminish the survival of the species on longer time-scales.

Blood-oxygen binding

Many marine animals rely on specialized respiratory proteins to bind oxygen at respiratory surfaces (e.g. gills) and deliver it to the tissues for cellular metabolism. A high gradient from the environment to the blood promotes oxygen binding at the gills, and the gradient from the blood to the metabolizing tissues promotes its release. However, these gradients alone are often inadequate to facilitate sufficient oxygen saturation and unloading of the respiratory proteins. CO_2 produced by cellular metabolism

interacts with body fluids to produce hydrogen ions that bind to respiratory proteins, altering their affinity for oxygen. That is, CO_2 production causes acidosis that promotes oxygen release at the tissues, whereas CO_2 excretion elevates pH and promotes oxygen binding at the gills. The sensitivity of oxygen binding to pH is expressed as the Bohr coefficient ($\Delta\log P_{50}/\Delta\text{pH}$, where P_{50} is the $p\text{O}_2$ required to achieve 50% oxygen saturation of the respiratory protein). The biotic and abiotic factors that contribute to selective pressure for pH sensitivity are complex and not easily predicted, and the measurement of pH sensitivity and blood oxygen binding depends on various ionic and organic modulators that vary tremendously from one study to another (Lallier and Truchot, 1989; Mangum, 1991). All else being equal, however, greater pH sensitivity (a larger Bohr effect) may allow more complete release of oxygen in support of high oxygen demand, or from high-affinity respiratory proteins, such as those of species adapted to hypoxic environments (Childress and Seibel, 1998; Hourdez and Weber, 2005).

Pörtner and Reipschläger (1996) predicted that species with high metabolic rates would be more severely impacted by ocean acidification because oxygen binding in their blood is more pH sensitive. Epipelagic squid (e.g. Ommastrephidae, Gonatidae, Loliginidae) are hypothesized to be most severely impacted by the interference of CO_2 with oxygen binding at the gills, because their metabolic rates are higher than other aquatic animals (Seibel, 2007; Seibel and Drazen, 2007). Furthermore, oxygen-carrying capacity is constrained in squid relative to active fish (Pörtner, 1994). As a result, it is necessary for active squid to utilize all of the oxygen carried in the blood on each pass through the body, even at rest, leaving no venous oxygen reserve. Unloading the entire oxygen store at the tissues requires extreme pH sensitivity (Bohr coefficient less than -1.0 ; Figure 9; Pörtner, 1994). One downside of this adaptation is that an increase in CO_2 in the environment will inhibit oxygen binding at the gills. Pörtner (1990, 1994) estimates that a reduction in environmental seawater pH by as little as 0.15 unit will reduce the scope for activity in the squid *Illex illecebrosus*. Recent work demonstrates that elevated $p\text{CO}_2$ (~ 1000 ppmv) can create measurable reductions in the oxygen consumption rate and scope for activity of another ommastrephid squid *Dosidicus gigas* (Figure 8; R. Rosa and B. Seibel, unpublished data). However, squid may be exceptional both metabolically and in their sensitivity to low pH. We review the pH sensitivity of oxygen binding in marine animals and find no correlation with metabolic rate or environmental oxygen levels and find no obvious phylogenetic signal (Figure 9). For example, several metabolically active species have pH insensitive respiratory proteins (low Bohr coefficients), while several others have high pH sensitivity despite low oxygen demand.

Predicting population and ecosystem responses

Table 1 lists the responses of a variety of animals to low pH–high $p\text{CO}_2$ conditions. The data indicate that foraminifera, molluscs, and echinoderms demonstrate reduced calcification and sometimes dissolution of CaCO_3 skeletal structures when exposed to elevated $p\text{CO}_2$ and decreasing pH and CO_3^{2-} concentration. Fertilization rates and early development are also negatively impacted by high CO_2 conditions in a number of groups such as sea urchins, molluscs, and copepods. Significantly, the data are limited with regard to the number of species tested at climate-relevant $p\text{CO}_2$ levels. For example, although teleost fish

Table 1. Examples of the response of marine fauna to ocean acidification.

Species	Description	CO ₂ system parameters	Sensitivity	Reference
Planktonic foraminifera				
<i>Orbulina universa</i>	Symbiont-bearing	pCO ₂ 560–780 ppmv	8–14% reduction in shell mass	Spero <i>et al.</i> (1997); Bijma <i>et al.</i> (1999, 2002)
<i>Globigerinoides sacculifer</i>	Symbiont-bearing	pCO ₂ 560–780 ppmv	4–8% reduction in shell mass	Bijma <i>et al.</i> (1999, 2002)
Cnidaria				
Scyphozoa Hydrozoa	Jellyfish	North Sea seawater pH drop from 8.3 to 8.1	Increase in frequency as measured by CPR from 1958 to 2000	Attrill <i>et al.</i> (2007)
Mollusca				
<i>Clio pyramidata</i>	Shelled pteropod	$\Omega_{\text{arag}} < 1$	Shell dissolution	Feely <i>et al.</i> (2004); Orr <i>et al.</i> (2005); this work
<i>Haliotis laevisgata</i>	Greenlip abalone	pH 7.78; pH 7.39	5% and 50% growth reductions	Harris <i>et al.</i> (1999).
<i>Haliotis rubra</i>	Blacklip abalone	pH 7.93; pH 7.37	5% and 50% growth reductions	
<i>Mytilus edulis</i>	Mussel	pH 7.1 / 10 000 ppmv pCO ₂ 740 ppmv	Shell dissolution 25% decrease in calcification rate	Lindinger <i>et al.</i> (1984) Gazeau <i>et al.</i> (2007)
<i>Crassostrea gigas</i>	Oyster	pCO ₂ 740 ppmv	10% decrease in calcification rate	
<i>Mytilus galloprovincialis</i>	Mediterranean mussel	pH 7.3, ~5000 ppmv	Reduced metabolism, growth rate	Michaelidis <i>et al.</i> (2005)
<i>Placopecten magellanicus</i>	Giant scallop	pH < 8.0	Decrease in fertilization and embryo development	Desrosiers <i>et al.</i> (1996)
<i>Tivela stultorum</i>	Pismo clam	pH < 8.5	Decrease in fertilization rates	Alvarado-Alvarez <i>et al.</i> (1996)
<i>Pinctada fucada martensii</i>	Japanese pearl oyster	pH 7.7 pH > 7.4	Shell dissolution, reduced growth Increasing mortality	Reviewed in Knutzen (1981)
<i>Mercenaria mercenaria</i>	Clam	$\Omega_{\text{arag}} = 0.3$	Juvenile shell dissolution leading to increased mortality	Green <i>et al.</i> (2004)
<i>Illex illecebrosus</i>	Epipelagic squid	2000 ppmv	Impaired oxygen transport	Pörtner and Reipschläger (1996)
<i>Dosidicus gigas</i>	Epipelagic squid	0.1% CO ₂ , ~1000 ppmv	Reduced metabolism/scope for activity	Rosa and Seibel (unpublished)
Arthropoda				
<i>Acartia steueri</i>	Copepod	0.2–1%CO ₂	Decrease in egg hatching success;	Kurihara <i>et al.</i> (2004)
<i>Acartia erythraea</i>	Copepod	~2000–10 000 ppmv	increase in nauplius mortality rate	
Copepods	Pacific, deep vs. shallow	860–22 000 ppmv CO ₂	Increasing mortality with increasing CO ₂ concentration and duration of exposure	Watanabe <i>et al.</i> (2006)
<i>Euphausia pacifica</i>	Krill	pH < 7.6	Mortality increased with increasing exposure time and decreasing pH	Yamada and Ikeda (1999)
<i>Paraeuchaeta elongata</i>	Mesopelagic copepod			
<i>Conchoecia</i> sp.	Ostracod			
<i>Cancer pagurus</i>	Crab	1% CO ₂ , ~10 000 ppmv	Reduced thermal tolerance, aerobic scope	Metzger <i>et al.</i> (2007)
Chaetognatha				
<i>Sagitta elegans</i>	Chaetognath	pH < 7.6	Mortality increased with increasing exposure time and decreasing pH	Yamada and Ikeda (1999)
Echinodermata				
<i>Strongylocentrotus purpuratus</i>	Sea urchin	pH ~6.2–7.3	High sensitivity inferred from lack of pH regulation and passive buffering via test dissolution during emersion	cf. Burnett <i>et al.</i> (2002)
<i>Psammechinus miliaris</i>	Sea urchin			Spicer (1995); Miles <i>et al.</i> (2007)
<i>Hemicentrotus pulcherrimus</i>	Sea urchin	~500–10 000 ppmv	Decreased fertilization rates, impacts larval development	Kurihara and Shirayama (2004)
<i>Echinometra mathaei</i>	Sea urchin			
<i>Cystechinus</i> sp.	Deep-sea urchin	pH 7.8	80% mortality under simulated CO ₂ sequestration	Barry <i>et al.</i> (2002)

Continued

Table 1. Continued

Species	Description	CO ₂ system parameters	Sensitivity	Reference
Sipuncula				
<i>Sipunculus nudus</i>	Peanut worm	1% CO ₂ , 10 000 ppmv	Metabolic suppression	Pörtner and Reipschläger (1996)
			Pronounced mortality in 7-week exposure	Langenbuch and Pörtner (2004)
Vertebrata				
<i>Scyliorhinus canicula</i>	Dogfish	pH 7.7 / 0.13%CO ₂ 7% CO ₂ , ~70 000 ppmv	Increased ventilation 100% mortality after 72 h	Reviewed in Truchot (1987) Hayashi et al. (2004)
<i>Sillago japonica</i>	Japanese whiting	7% CO ₂ , ~70 000 ppmv	Rapid mortality in 1-step exposure	Kikkawa et al. (2006)
<i>Paralichthys olivaceus</i>	Japanese flounder	5% CO ₂ , ~50 000 ppmv	100% mortality within 48 h	Hayashi et al. (2004)
<i>Euthynnus affinis</i>	Eastern little tuna	15%CO ₂ , ~150 000 ppmv	100% mortality of eggs after 24 h	Kikkawa et al. (2003)
<i>Pagrus major</i>	Red sea bream	5%CO ₂ , ~50 000 ppmv	>60% larval mortality after 24 h	Ishimatsu et al. (2005)
<i>Seriola quinqueradiata</i>	Yellowtail/ amberjack	5% CO ₂ , 50 000 ppmv	Reduced cardiac output; 100% mortality after 8 h	Ishimatsu et al. (2004)
<i>Sparus aurata</i>	Mediterranean fish	pH 7.3, ~ 5000 ppmv	Reduced metabolic capacity	Michaelidis et al. (2007)
<i>Dicentrarchus labrax</i>	Sea bass	pH 7.25, 24 mg l ⁻¹ CO ₂	Reduced feed intake	Cecchini et al. (2001)




Figure 8. Oxygen consumption rates under elevated CO₂ for marine animals as a percentage of control rates (air saturation). Decreases in routine metabolism, an adaptive strategy to short-term hypercapnia, of the squid *Dosidicus gigas* ~1000 ppmv (0.1% at 20°C), the pteropod mollusc *Limacina helicina antarctica* under 789 ppmv (−1.86°C), the worm *Sipunculus nudus* and an amphipod *Phronima sedentaria* under 10 000 ppm (1.0%), and the bivalve *Mytilus edulis* under ~5000 ppmv (0.5%, pH 7.3, 18°C) carbon dioxide. (R. Rosa, and B. Seibel, unpublished data; Pörtner and Reipschläger (1996); Michaelidis et al. (2005)).

may appear less sensitive to decreased pH-elevated $p\text{CO}_2$ compared with other faunal groups, it must be emphasized that there is no information on the response of fish to the $p\text{CO}_2$ values that are projected to occur over the next century. Moreover, most empirical evidence comes from short-term experiments, and we know little or nothing about the response of

marine biota to continuous, long-term exposure to elevated $p\text{CO}_2$ or the capacity of these organisms to adapt. Nevertheless, the data in Table 1 clearly demonstrate that elevated $p\text{CO}_2$ can adversely impact marine fauna both via decreased carbonate saturation state, which directly affects calcification rates, and via disturbance to acid–base physiology. We use the available evidence to speculate on possible ecological winners and losers during the 21st century, and to identify priorities for critically needed research.

Species distributions

Euthecosomatous pteropods will be first among the major groups of planktonic calcifiers to experience persistent <1 saturation states in surface waters of their current geographical ranges (Figures 3 and 4). If we assume that these animals are restricted to aragonite-saturated waters, then euthecosomatous pteropod habitat would become increasingly limited, first vertically in the water column, then latitudinally, by shoaling of the aragonite saturation horizon (Orr et al., 2005). For example, the pteropod *C. pyramidata* is widely distributed and, in the North Pacific, its range extends to nearly 55°N. In other oceans, this species is typically found at 400–500 m during the day and in surface waters at night (Bé and Gilmer, 1977). If *C. pyramidata* has a similar pattern of vertical migration in the North Pacific, then this species would already be experiencing seawater corrosive to aragonite ($\Omega < 1$) during part of its diel cycle at ~10°N and near 50°N (Figure 3). When *C. pyramidata* is transported from coastal waters via anticyclonic Haida eddies that form along the eastern margin of the Subarctic Pacific and move towards the Alaskan gyre, this pteropod can be a dominant member of the eddy zooplankton community (Mackas and Galbraith, 2002; Tsurumi et al., 2005). Yet, *C. pyramidata* demonstrated only weak diel vertical migration within an eddy, with most individuals remaining in the upper




Figure 9. Bohr coefficients ($\Delta \log P_{50} \Delta \text{pH}^{-1}$), an indication of the sensitivity of oxygen transport to pH, as a function of oxygen consumption rate at approximate environmental temperature for marine animals from both benthic and pelagic environments. 1. *Vampyroteuthis infernalis*; 2. *Glyphocrangon vicara*; 3. *Bythograea thermydron*; 4. *Gnathopausia ingens*; 5. *Octopus vulgaris*; 6. *Coryphanoides armatus*; 7. *Acanthephyra curtirostris*; 8. *Acanthephyra acutifrons*; 9. *Systelaspis debilis*; 10. *Architeuthis* sp.; 11. *Loligo pealei*; 12. *Acanthephyra smithi*; 13. *Loligo vulgaris*; 14. *Oplophorus gracilirostris*; 15. *Illex illecebrosus*; 16. *Sepia officinalis*; 17. *Octopus dofleini*; 18. *Trematomus bernacchii*; 19. *Artedidraco* sp.; 20. *Lolliguncula brevis*; 21. *Megaleledone senoi*. Bohr coefficients from Grigg (1967); Arp and Childress (1985); Noble *et al.* (1986); Sanders and Childress (1990); Eastman (1993); Bridges (1994); Tamburrini *et al.* (1998); Seibel *et al.* (1999); Lowe *et al.* (2000); Zielinski *et al.* (2001). Metabolic rates reviewed in Childress (1995); Seibel and Drazen (2007)). Metabolic rate for *Architeuthis* sp. is estimated from activity of citrate synthase.

75 m during both day and night (Mackas and Galbraith, 2002). Apart from this Haida eddy study, however, data on the vertical distribution of *C. pyramidata* in the Subarctic Pacific are lacking.

The euthecosome *Limacina helicina* is an important high-latitude species in both the northern and southern hemispheres. In the Arctic Ocean, this species is most abundant between 50–100 m during winter and in the upper 50 m during summer (Kobayashi, 1974). South of the Antarctic Polar Front, where aragonite undersaturation of the entire water column is projected to occur within the next 50–100 years, this pteropod species comprises nearly all of the CaCO_3 export to the ocean interior (Collier *et al.*, 2000; Honjo *et al.*, 2000; Accornero *et al.*, 2003).

As aragonite saturation states approach 1 with progressive acidification during the 21st century (Figure 4), we hypothesize that shelled pteropod species, such as *C. pyramidata* and *L. helicina*, either will have to adapt to living continuously in seawater undersaturated with respect to aragonite or restrict their vertical and latitudinal distributions to warmer, more carbonate-rich regions that remain supersaturated with respect to aragonite. This latter possibility may be limited by the extreme adaptations to low temperature that may prevent equatorward movement of polar animals (Stillman, 2003; Somero, 2005; Seibel *et al.*, 2007). Waters

undersaturated with respect to aragonite are currently impinging upon the depth ranges of pteropods in several other regions, such as upwelling areas associated with the Benguela Current, western Arabian Sea, and Peru Current, where pteropod abundances can be high (e.g. Bé and Gilmer, 1977; Fabry, 1990; Boltovskoy *et al.*, 1993; Kalberer *et al.*, 1993; Hitchcock *et al.*, 2002; Mohan *et al.*, 2006).

Although future anthropogenically induced reductions in the saturation state of calcite will not be as severe as those for aragonite, species of the calcitic foraminifera may also change their geographic distributions in response to decreased calcite saturation states. As calcite undersaturation is projected to occur ~50–100 years subsequent to that of aragonite (Orr *et al.*, 2005), foraminifera could be displaced from high latitudes, where they can be abundant. Changes in species composition could also occur, as has been reported in the California Current in response to anthropogenic warming (Field *et al.*, 2006). Currently, there are few high-quality data on the diel vertical distributions of euthecosomes and foraminifera in these areas to test whether shelled pteropod and foraminiferal populations will shift their depth ranges as the carbonate chemistry of seawater changes.

In addition to calcification, other physiological indices indicate general trends that may be useful for predicting species' vulnerability to ocean acidification. Those with low metabolic rates, including perhaps gelatinous zooplankton and those that experience little natural variation in CO_2 , appear to demonstrate lesser CO_2 tolerance. Thus, at first glance, zooplankton inhabiting the open ocean may appear highly susceptible to ocean acidification, given the constancy of the pelagic environment relative to hydrothermal vents or the intertidal zone (Truchot and Duhamel-Jouve, 1980). However, large, vertical gradients in environmental variables, including oxygen, CO_2 , and pH, exist in the upper 1000 m (Figure 3), and most zooplankton species migrate daily from near-surface waters to depths of 200–700 m. In the expansive regions with pronounced oxygen minimum layers (Figure 3c), these diel migrations expose zooplankton to wide variations in $p\text{CO}_2$, values (Figure 3b) greater than those expected for average surface waters as a result of anthropogenic ocean acidification over the next 100 years (Figure 1). Therefore, although the open ocean environment has fluctuated historically only over millennial time-scales (Kennett and Ingram, 1995), giving species ample time to adapt (Childress and Seibel, 1998), the effective environment of many species in the open ocean is not constant.

Many species migrating into oxygen minimum layers may do so by suppressing metabolism and supplementing remaining energy demands with anaerobic metabolic pathways (Childress and Seibel, 1998; Hunt and Seibel, 2000). Anaerobic metabolism itself may exacerbate internal acid–base imbalance (Hochachka and Somero, 2002). Therefore, vertically migrating species, like those living intertidally or near hydrothermal vents, experience oscillating periods of simultaneous hypoxia and high $p\text{CO}_2$ that require specific adaptations for tolerance (Childress and Seibel, 1998). Those adaptations may make zooplankton in hypoxic regions more tolerant of elevated $p\text{CO}_2$, at least on short time-scales, than those in well-oxygenated regions. This hypothesis is supported by the recent work of Watanabe *et al.* (2006), who found greater mortality during short-term exposure to high $p\text{CO}_2$ in shallow-living and subtropical copepods than in deep-living species in the Subarctic Pacific, where $p\text{CO}_2$ is naturally much higher. However, Pane and Barry (2007) suggested that low oxygen exacerbated internal acid–base imbalance in a

deep-sea crab. Currently, there is no evidence that adaptation to variably hypercapnic environments promotes tolerance of chronic ocean acidification such as that expected over the next century. Furthermore, as warming alters ocean stratification, the vertical and horizontal extent of the oxygen minimum layer is expected to change, and ocean acidification will elevate $p\text{CO}_2$ levels at subsurface depths. Warming is also expected to act synergistically to exacerbate oxygen limitation and further hinder CO_2 tolerance (Metzger *et al.*, 2007).

Predictive ability of the response of zooplankton populations to ocean acidification is hampered by a paucity of measurements at climate-relevant CO_2 concentrations (Table 1). In many cases, no information exists for ecologically important taxa such as larvae, salps, amphipods, and euphausiids. Gelatinous zooplankton have not been examined at all for CO_2 tolerance. Their low metabolic rates may make them highly susceptible; however, they may also exhibit species-specific responses to increased $p\text{CO}_2$, similar to the differential response observed among medusae species to hypoxia (Rutherford and Thuesen, 2005; Thuesen *et al.*, 2005a, b). Attrill *et al.* (2007) reported a significant correlative relationship between reduced pH and increased frequency of medusae, as sampled by the Continuous Plankton Recorder for 40 years in the North Sea. The authors suggest that the frequency of medusae in the North Sea will increase over the next century as surface water pH values decrease. As yet, no causative mechanism linking jellyfish abundance with ocean acidification is known.

Fertilization success and early developmental stages of many faunal groups appear to be particularly vulnerable to elevated $p\text{CO}_2$ (Table 1). The tolerance of early life stages may impact recruitment success and, ultimately, species abundances and distributions. If the formation of highly soluble, amorphous CaCO_3 as a transient precursor to crystalline phases in embryonic and larval stages (Beniash *et al.*, 1997; Weiss *et al.*, 2002) is widespread among mollusc and echinoderm species, then these phyla may be particularly at risk from progressive ocean acidification in many oceanic regions. Additional investigation is needed to determine if ocean acidification-induced mortality of these stages could drive a reorganization of benthic and pelagic communities and also adversely impact commercially important fisheries.

Trophic dynamics and other ecosystem processes

The relative rate of change in surface seawater carbonate ion concentration is greatest in high-latitude regions (Orr *et al.*, 2005; Figure 4). In polar and subpolar areas, the progressive shoaling of the aragonite saturation horizon and the decreasing calcite saturation state of the euphotic zone over future decades will impact trophic dynamics and other ecosystem processes, including the cycling of CaCO_3 and organic matter. Euthecosomatous pteropods are functionally important components of high-latitude ecosystems with the potential to influence phytoplankton stocks (Hopkins, 1987), carbon fluxes (Noji *et al.*, 1997; Collier *et al.*, 2000; Honjo *et al.*, 2000), and dimethyl sulphide levels (Levasseur *et al.*, 1994) that, in turn, influence global climate through ocean–atmosphere feedback loops. The possible extirpation of euthecosomatous pteropods from the high-latitude regions would impact the downward organic carbon flux associated with pteropod faecal pellets (Thibault *et al.*, 1999; Collier *et al.*, 2000) and remove a major source of CaCO_3 in such regions (e.g. Bathmann *et al.*, 1991; Gardner *et al.*, 2000; Honjo *et al.*, 2000; Accornero *et al.*, 2003; Tsurumi *et al.*, 2005). Similarly, if

foraminifera densities decrease in some high-latitude regions, where they are currently abundant (e.g. Subarctic Pacific), CaCO_3 export to the ocean interior will be reduced, which in turn would decrease their potential to act as ballast in the transport of organic carbon to the deep sea (Schiebel, 2002).

Most of the carnivorous zooplankton and fish (e.g. cod, pollock, haddock, mackerel) that feed on euthecosomatous pteropods (Ito, 1964; LeBrasseur, 1966; Lalli and Gilmer, 1989) would be able to switch to other prey types, which could result in greater predation pressure on juvenile fish such as salmon (Willette *et al.*, 2001). In contrast, gymnosomes prey exclusively on shelled pteropods (Lalli and Gilmer, 1989) and would likely shift their geographic distribution in concert with their euthecosome prey, assuming both predator and prey are able to overcome possible thermal tolerance limitations (Seibel *et al.*, 2007). Another specialized predator, the myctophid *Centrobrachus brevirostris*, which feeds predominantly on the pteropod *C. pyramidata* in the Kuroshio waters of the western North Pacific (Watanabe *et al.*, 2002), would also be highly impacted if euthecosomes were excluded from its habitat in the future; typically, the trade-off in such highly specialized feeding is a reduced ability to diversify when the preferred prey is absent.

In the North Pacific, the euthecosomes *L. helicina* and, to a lesser extent, *C. pyramidata* can be important prey of juvenile pink salmon (*Oncorhynchus gobuscha*), which comprise a large part of the commercial catch of salmon in the North Pacific (cf. Armstrong *et al.*, 2005). Pink salmon have a short, two-year lifespan, and their recruitment is thought to be affected by diet and availability of prey during their early marine life history. In a three-year study designed to examine the interannual variability of the feeding habits of juvenile pink salmon, Armstrong *et al.* (2005) found that *L. helicina* accounted for $\geq 60\%$ by weight of the juvenile salmon diet in two of three years, but only 15% in the third year. Juvenile pink salmon rapidly increase in size during summer in the Subarctic Pacific, feeding on progressively larger prey items and switching from *L. helicina* to the larger *C. pyramidata* by October (Boldt and Haldorson, 2003; Armstrong *et al.*, 2005). In a model study linking oceanic foodwebs to production and growth rates of pink salmon, Aydin *et al.* (2005) found that decreased energetic foraging costs for zooplanktivorous juvenile salmon and the ontogenetic diet shift from zooplankton to squid were both key factors that strongly influenced the biomass of mature pink salmon. Because euthecosomatous pteropods can reach swarm densities in near-surface waters during daylight, either through concentration by eddies (Tsurumi *et al.*, 2005) or life-history traits (Bathmann *et al.*, 1991; Gannefors *et al.*, 2005), visual predators such as juvenile pink salmon may be able to reduce forage costs by feeding within pteropod patches. Other preliminary model results suggest that a 10% decrease in pteropod production could lead to a 20% drop in mature pink salmon body weight (Aydin, pers. comm.). During the ontogenetic diet shift of juvenile salmon, the gonatid squid *Berrytheuthis anonychus* is an important control on adult salmon biomass; availability of this lipid-rich prey species substantially accelerates the growth of both pink and sockeye salmon in bioenergetics models (cf. Aydin *et al.*, 2005). Similarly, *Gonatus fabricii*, among the most abundant squid species in the North Atlantic, is the most important prey item to a number of marine mammals and may be responsible for their seasonal occurrence in some regions (Bjorke, 2001; Hooker *et al.*, 2001). Although the respiratory physiology of gonatid squid has not been investigated in detail, their metabolic

rates are high (Seibel, 2007), which may lead to high CO₂ sensitivity, as described above for ommastrephids (e.g. Pörtner and Reipschläger, 1996; Pörtner *et al.*, 2004). Ommastrephids are also important components of ecosystems and are important commercial fisheries worldwide (Rodhouse and White, 1995; Clarke, 1996; Nigmatullin *et al.*, 2001).

Ocean acidification could also affect foodwebs and carbon cycling through bottom-up controls involving pH-dependent speciation of nutrients and metals (Huesemann *et al.*, 2002), which, in turn, may alter species composition and rates of primary productivity. The interactive effects and feedback of changing seawater CO₂ chemistry with other stressors, such as warming, eutrophication, introduced species, and overfishing, may act to alter ecosystem responses that would otherwise result from only one of these stressors (Schippers *et al.*, 2004; Hutchins *et al.*, 2007). Quantification of these complex ecosystem processes requires additional empirical data, as well as new modelling efforts, particularly on regional scales.

Research needs and conclusions

Most experimental work on the impacts of ocean acidification on marine biota at climate-relevant pCO₂ values has investigated the calcification response of corals and coccolithophores (cf. Kleypas *et al.*, 2006). There is a critical need for information on the sublethal calcification and energetic responses of a diverse suite of zooplankton and micronekton. We need to move forward on several fronts in parallel.

- In sensitive regions and for critical species, we need to track the abundances and depth distributions of calcareous and non-calcifying fauna, measure calcification and metabolic rates of these groups, and relate these data to changes in the CO₂ chemistry of the water column. This requires commitment to long-term monitoring programmes at appropriate temporal and spatial scales to detect possible shifts, and distinguish between natural variability and anthropogenically induced changes.
- Using pCO₂ levels projected to occur over the next century, manipulative laboratory experiments are needed to investigate the calcification and dissolution responses, identify physiological indices useful in predicting CO₂ tolerance, determine the costs of acid–base regulation, and quantify sensitive energetic processes, such as skeletal and tissue growth, reproduction, and metabolism for critical life stages of key species.
- Mesocosm and field experiments are necessary to quantify ecosystem impacts from ocean acidification that may include forcing from bottom-up controls, changes in foodweb structure, biogeochemical cycling, and feedback mechanisms.
- High-priority areas for research include high-latitude regions, which may become undersaturated with respect to aragonite as early as 2050, and regions with pronounced oxygen minimum layers or coastal hypoxia, which are already characterized by high pCO₂ and may be particularly at risk, owing to the combined effects of low oxygen with elevated pCO₂, warming, and eutrophication.
- Target species for investigation in the above regions include euthecosomatous pteropods, foraminifera, epipelagic squid, and larval stages and adults of commercially and ecologically important benthic invertebrates such as bivalves, sea urchins,

crabs, and lobsters. We also need to test taxa for which there are currently no data available, including medusae, larvaceans, and various crustaceans. Additional experiments should examine the interactive effects of seawater CO₂ chemistry with temperature, dissolved oxygen, food availability, and other variables that may change as a result of human activities.

- New approaches (e.g. functional genomics and DNA barcoding) and advances in existing technologies (e.g. autonomous chemical sensors and optical plankton samplers) are necessary to investigate the *in situ* response of organisms that are difficult to maintain in the laboratory, identify sublethal effects of chronic exposure to elevated pCO₂ on marine fauna, and address questions of long-term impacts and potential for adaptation over decadal to centennial time-scales.
- Models are critical to scale up results from manipulative experiments and field observations to predict ecosystem impacts on regional and global scales.

Although the changes in seawater chemistry that result from the oceanic uptake of anthropogenic CO₂ are well characterized over most of the ocean, the biological impacts of ocean acidification on marine fauna are only beginning to be understood. New technologies and advances, as well as integrated, multidisciplinary efforts among biologists and chemists, experimentalists and modellers will be required to quantify the effects of ocean acidification on marine fauna and changes in ecosystem structure and function. Nevertheless, sufficient information exists to state with certainty that deleterious impacts on some marine species are unavoidable, and that substantial alteration of marine ecosystems is likely over the next century.

Acknowledgements

This work was supported jointly by the National Science Foundation (NSF grants OCE-0551726 and OPP-0538710 to VJE, and OCE-0526493 and OPP-0538479 to BAS) and the National Oceanic and Atmospheric Administration (NOAA). We specifically acknowledge programme managers Don Rice, Phil Taylor, and Roberta Marinelli of the NSF Chemical and Biological Oceanography Programs and Office of Polar Programs, respectively; and Kathy Tedesco and Mike Johnson of the NOAA Climate Change Program for their support. We thank H. Spero, B. Hönisch, and A. Maas for discussions and comments on earlier drafts of this paper. The IAEA is grateful for the support provided to its Marine Environmental Laboratory by the Government of the Principality of Monaco.

References

- Accornero, A., Manno, C., Esposito, F., and Gambi, M. C. 2003. The vertical flux of particulate matter in the polynya of Terra Nova Bay. Part II. Biological components. *Antarctic Science*, 15: 175–188.
- Alvarado-Alvarez, R., Gould, M. C., and Stephano, J. L. 1996. Spawning, *in vitro* maturation, and changes in oocyte electrophysiology induced serotonin in *Tivela stultorum*. *Biological Bulletin*, 190: 322–328.
- Armstrong, J. L., Boldt, J. L., Cross, A. D., Moss, J. H., Davis, N. D., Myers, K. W., Walker, R. V., *et al.* 2005. Distribution, size, and interannual, seasonal and diel food habits of northern Gulf of Alaska juvenile pink salmon, *Oncorhynchus gorbuscha*. *Deep Sea Research II*, 52: 247–265.

- Arp, A. J., and Childress, J. J. 1985. Oxygen binding properties of the blood of the deep-sea shrimp, *Glyphocrangon vicaria*. *Physiological Zoology*, 58: 38–45.
- Attrill, M. J., Wright, J., and Edwards, M. 2007. Climate-related increases in jellyfish frequency suggest a more gelatinous future for the North Sea. *Limnology and Oceanography*, 52: 480–485.
- Aydin, K. Y., McFarlane, G. A., King, J. R., Megrey, B. A., and Myer, K. W. 2005. Linking oceanic foodwebs to coastal production and growth rates of Pacific salmon (*Oncorhynchus* spp.), using models on three scales. *Deep Sea Research II*, 52: 757–780.
- Barker, S., and Elderfield, H. 2002. Foraminiferal calcification response to glacial–interglacial changes in atmospheric CO₂. *Science*, 297: 833–836.
- Barry, J., Seibel, B. A., Drazen, J., Tamburri, M., Lovera, C., and Brewer, P. 2002. Field experiments on direct ocean CO₂ sequestration: the response of deep-sea faunal assemblages to CO₂ injection at 3200 m off central California. *EOS Transactions of the American Geophysical Union*, 83: OS51F-02.
- Bathmann, U. V., Noji, T. T., and von Bodungen, B. 1991. Sedimentation of pteropods in the Norwegian Sea in autumn. *Deep Sea Research*, 38: 1341–1360.
- Bé, A. W. H., and Gilmer, R. W. 1977. A zoogeographic and taxonomic review of euthecosomatous Pteropoda. In *Oceanic Micropaleontology*, Vol. 1, pp. 733–808. Ed. by A. T. S. Ramsey. Academic Press, London.
- Beniash, E., Aizenber, J., Addai, L., and Weiner, S. 1997. Amorphous calcium carbonate transforms into calcite during sea urchin larval spicule growth. *Proceedings of the Royal Society of London, Series B*, 264: 461–465.
- Bijma, J., Honisch, B., and Zeebe, R. E. 2002. Impact of the ocean carbonate chemistry on living foraminiferal shell weight: comment on “Carbonate ion concentration in glacial-age deepwaters of the Caribbean Sea” by W. S. Broecker and E. Clark. *Geochemistry, Geophysics, Geosystems*, 3: 1064. doi:10.1029/2002GC000388.
- Bijma, J., Spero, H. J., and Lea, D. W. 1999. Reassessing foraminiferal stable isotope geochemistry: impact of the oceanic carbonate systems (experimental results). In *Use of Proxies in Paleooceanography: Examples from the South Atlantic*, pp. 489–512. Ed. by G. Fisher, and G. Wefer. Springer-Verlag, New York.
- Bischoff, W. W., Mackenzie, F. T., and Bishop, F. C. 1987. Stabilities of synthetic magnesian calcites in aqueous solution: comparison with biogenic materials. *Geochimica et Cosmochimica Acta*, 51: 1413–1424.
- Bjorke, H. 2001. Predators of the squid *Gonatus fabricii* (Lichtenstein) in the Norwegian Sea. *Fisheries Science*, 52: 113–120.
- Boldt, J., and Haldorson, L. J. 2003. Seasonal and geographic variation in juvenile pink salmon diets in the northern Gulf of Alaska and Prince William Sound. *Transactions of the American Fisheries Society*, 132: 1035–1052.
- Boltovskoy, D., Alder, V. A., and Abelman, A. 1993. Annual flux of Radiolaria and other shelled plankters in the eastern Atlantic at 853 m: seasonal variations and polycystine species-specific responses. *Deep Sea Research*, 40: 1863–1895.
- Bown, P. R., Lees, J. A., and Young, J. R. 2004. Calcareous nanoplankton evolution and diversity through time. In *From Molecular Processes to Global Impact*, pp. 481–508. Ed. by H. R. Thierstein, and J. R. Young. Springer-Verlag, Berlin.
- Bridges, C. R. 1994. Bohr and Root effects in cephalopod haemocyanins—paradox or pressure in *Sepia officinalis*? In *Physiology of Cephalopod Molluscs: Lifestyle and Performance Adaptations*, pp. 121–130. Ed. by H. O. Portner, R. K. O'Dor, and D. L. MacMillan Gordon and Breach, London.
- Brummer, G.-J. A., Hemleben, C., and Spindler, M. 1987. Ontogeny of extant spinose planktonic foraminifera (Globigerinidae): a concept exemplified by *Globigerinoides sacculifer* (Brady) and *G. ruber* (d'Orbigny). *Marine Micropaleontology*, 12: 357–381.
- Burnett, L., Terwilliger, N., Carroll, A., Jorgensen, D., and Scholnick, D. 2002. Respiratory and acid-base physiology of the purple sea urchin, *Strongylocentrotus purpuratus*, during air exposure: presence and function of a facultative lung. *Biological Bulletin*, 203: 42–50.
- Caldeira, K., and Wickett, M. E. 2003. Anthropogenic carbon and ocean pH. *Science*, 425: 365.
- Caldeira, K., and Wickett, M. E. 2005. Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *Journal of Geophysical Research*, 110: C09S04. doi:10.1029/JC002671.
- Cameron, J. N. 1989. *The Respiratory Physiology of Animals*. Oxford University Press, New York. 353 pp.
- Cameron, J. N., and Iwama, G. K. 1987. Compensation of progressive hypercapnia in channel catfish and blue crabs. *Journal of Experimental Biology*, 133: 183–197.
- Castellini, M. A., and Somero, G. N. 1981. Buffering capacity of vertebrate muscle: correlations with potentials for anaerobic function. *Journal of Comparative Physiology*, 143: 191–198.
- Cecchini, S., Saroglia, M., Caricato, G., Terrova, G., and Sileo, L. 2001. Effects of graded environmental hypercapnia on sea bass (*Dicentrarchus labrax* L.) feed intake and acid-base balance. *Aquaculture Research*, 32: 499–502.
- Childress, J. J. 1995. Are there physiological and biochemical adaptations of metabolism in deep-sea animals? *Trends in Ecology and Evolution*, 10: 30–36.
- Childress, J. J., and Seibel, B. A. 1998. Life at stable low oxygen levels: adaptations of animals to oceanic oxygen minimum layers. *Journal of Experimental Biology*, 201: 1223–1232.
- Clairborne, J. B., Edwards, S. L., and Morrison-Shetlar, A. I. 2002. Acid-base regulation in fishes: cellular and molecular mechanisms. *Journal of Experimental Zoology*, 293: 302–319.
- Clarke, M. R. 1996. The role of cephalopods in the world's oceans: general conclusions and the future. *Philosophical Transactions of the Royal Society of London, Series B—Biological Sciences*, 351: 1105–1112.
- Collier, R., Dymond, J., Honjo, S., Manganini, S., Francois, R., and Dunbar, R. 2000. The vertical flux of biogenic and lithogenic material in the Ross Sea: moored sediment trap observations 1996–1998. *Deep Sea Research*, 47: 3491–3520.
- Corliss, B. H., and Honjo, S. 1981. Dissolution of deep-sea benthonic foraminifera. *Micropaleontology*, 27: 356–378.
- Dadon, J. R., and de Cidre, L. L. 1992. The reproductive cycle of the Thecosomatous pteropod *Limacina retroversa* in the western South Atlantic. *Marine Biology*, 114: 439–442.
- Denman, K. L., et al. 2007. Couplings between changes in the climate system and biogeochemistry. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 500–556. Ed. by S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, H. L. Miller, et al. Cambridge University Press, Cambridge.
- Desrosiers, R. R., Désilets, J., and Dubé, F. 1996. Early developmental events following fertilization in the giant scallop *Placopecten magellanicus*. *Canadian Journal of Fisheries and Aquatic Sciences*, 53: 1382–1392.
- Dickson, A. G., and Millero, F. J. 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research*, 34: 1734–1743.
- Doney, S. C., Mahowald, N., Lima, I., Feely, R. A., Mackenzie, F. T., and Lamarque, F. 2007. The impacts of anthropogenic nitrogen and sulfur deposition on ocean acidification and the inorganic carbon system. *Proceedings of the National Academy of Sciences of the USA*, 104: 14 580–14 585.
- Eastman, J. T. 1993. *Antarctic Fish Biology: Evolution in a Unique Environment*. Academic Press, New York. 322 pp.

- Fabry, V. J. 1990. Shell growth rates of pteropod and heteropod mollusks and aragonite production in the open ocean: implications for the marine carbonate system. *Journal of Marine Research*, 48: 209–222.
- Field, D. B., Baumgartner, T. R., Charles, C. D., Ferreira-Bartrina, V., and Ohman, M. 2006. Planktonic foraminifera of the California Current reflect 20th-century warming. *Science*, 311: 63–66.
- Feely, R. A., Sabine, C. L., Byrne, R. H., and Greeley, D. 2006. Direct evidence for ocean acidification of the North Pacific Ocean. *EOS Transactions of the American Geophysical Union*, 87(52), Fall Meeting Supplement, Abstract OS12B-04.
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleyvas, J., Fabry, V. J., and Millero, F. J. 2004. Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans. *Science*, 305: 362–366.
- Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D. W., Haywood, J., *et al.* 2007. Changes in atmospheric constituents and in radiative forcing. In *Climate Change 2007: the Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 129–234. Ed. by S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, *et al.* Cambridge University Press, Cambridge.
- Gannefors, C., Böer, M., Kattner, G., Graeve, M., Eiane, K., Gulliksen, B., Hop, H., *et al.* 2005. The Arctic sea butterfly *Limacina helicina*: lipids and life strategy. *Marine Biology*, 147: 169–177.
- Gardner, W. D., Richardson, M. J., and Smith, W. O. 2000. Seasonal patterns of water column particulate organic carbon and fluxes in the Ross Sea, Antarctica. *Deep Sea Research*, 47: 3423–3449.
- Gattuso, J.-P., Frankignoulle, M., Bourge, I., Romaine, S., and Buddemeier, R. W. 1998. Effect of calcium carbonate saturation of seawater on coral calcification. *Global and Planetary Change*, 18: 37–46.
- Gazeau, F., Quibler, C., Jansen, J. M., Gattuso, J.-P., Middelburg, J. J., and Heip, C. H. R. 2007. Impact of elevated CO₂ on shellfish calcification. *Geophysical Research Letters*, 34: L07603. doi 10.1029/2006GL028554.
- Gibbs, A. H., and Somero, G. N. 1990. Na⁺-K⁺ adenosine triphosphatase activities in gills of marine teleost fishes, changes with depth, size and locomotory activity level. *Marine Biology*, 106: 315–321.
- Green, M. A., Jones, M. E., Boudreau, C. L., Moore, R. L., and Westman, B. A. 2004. Dissolution mortality of juvenile bivalves in coastal marine deposits. *Limnology and Oceanography*, 49: 727–734.
- Grigg, G. C. 1967. Some respiratory properties of the blood of four species of Antarctic fishes. *Comparative Biochemistry and Physiology*, 23: 139–148.
- Grosell, M. 2006. Intestinal anion exchange in marine fish osmoregulation. *Journal of Experimental Biology*, 209: 2813–2827.
- Guinotte, J. M., Orr, J., Cairns, S., Freiwald, A., Morgan, L., and George, R. 2006. Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? *Frontiers in Ecology and the Environment*, 4: 141–146.
- Guppy, M., and Withers, P. 1999. Metabolic depression in animals: physiological perspectives and biochemical generalizations. *Biological Reviews*, 74: 1–40.
- Gutiérrez, J. L., Jones, C. G., Strayer, D. L., and Iribarne, O. O. 2003. Mollusks as ecosystem engineers: the role of shell production in aquatic habitats. *Oikos*, 101: 79–90.
- Hand, S. C. 1991. Metabolic dormancy in aquatic invertebrates. In *Advances in Comparative and Environmental Physiology*, Vol. 8, pp. 1–50. Ed. by R. Gilles. Springer-Verlag, Heidelberg.
- Harris, J. O., Maguire, G. B., Edwards, S. J., and Hindrum, S. M. 1999. Effect of pH on growth rate, oxygen consumption rate, and histopathology of gill and kidney tissue for juvenile greenlip abalone, *Halitosis laevisgata* and blacklip abalone, *Halitosis rubra* leach. *Journal of Shellfish Research*, 18: 611–619.
- Hayashi, M., Kita, J., and Ishimatsu, A. 2004. Acid-base responses to lethal aquatic hypercapnia in three marine fishes. *Marine Biology*, 144: 153–160.
- Heisler, N. 1989. Interactions between gas exchange, metabolism, and ion transport in animals: an overview. *Canadian Journal of Zoology*, 67: 2923–2935.
- Heisler, N. 1993. Acid-base regulation. In *The Physiology of Fishes*, pp. 343–377. Ed. by D. H. Evans. CRC Press, Boca Raton, FL.
- Hemleben, C., Spindler, M., and Anderson, O. R. 1989. *Modern Planktonic Foraminifera*. Springer-Verlag, New York. 363 pp.
- Henry, R. P. 1984. The role of carbonic anhydrase in blood ion and acid-base regulation. *American Zoologist*, 24: 241–251.
- Hitchcock, G. L., Lane, P., Smith, S., Luo, J., and Ortner, P. B. 2002. Zooplankton spatial distributions in coastal waters of the northern Arabian Sea, August, 1995. *Deep Sea Research*, 49: 2403–2423.
- Hochachka, P. W., and Somero, G. N. 2002. *Biochemical Adaptation*. Princeton University Press, Princeton, NJ. 466 pp.
- Honjo, S., Francois, R., Manganini, S., Dymond, J., and Collier, R. 2000. Particle fluxes to the interior of the Southern Ocean in the Western Pacific sector along 170°W. *Deep Sea Research II*, 47: 3521–3548.
- Hooker, S. K., Iverson, S. J., Ostrom, P., and Smith, S. C. 2001. Diet of northern bottlenose whales inferred from fatty-acid and stable isotope analyses of biopsy samples. *Canadian Journal of Zoology*, 79: 1442–1454.
- Hopkins, T. L. 1987. Midwater food web in McMurdo Sound, Ross Sea, Antarctica. *Marine Biology*, 96: 93–106.
- Horita, J., Zimmerman, H., and Holland, H. D. 2002. Chemical evolution of seawater during the Phanerozoic: implications from the recode of marine evaporates. *Geochimica et Cosmochimica Acta*, 66: 3733–3756.
- Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der Linden, P. J., and Xiaosu, D. 2001. *Climate change 2001: the scientific basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge. 944 pp.
- Hourdez, S., and Weber, R. E. 2005. Molecular and functional adaptations in deep-sea hemoglobins. *Journal of Inorganic Biochemistry*, 99: 130–141.
- Huesemann, M. H., Skillman, A. D., and Crecelius, E. A. 2002. The inhibition of marine nitrification by disposal of carbon dioxide. *Marine Pollution Bulletin*, 44: 142–148.
- Hunt, J. C., and Seibel, B. A. 2000. Life history of *Gonatus onyx* (Teuthoidea: Cephalopoda): ontogenetic changes in habitat, behavior and physiology. *Marine Biology*, 136: 543–552.
- Hutchins, D. A., Fu, F.-X., Zhang, Y., Warner, M. E., Feng, Y., Portune, K., Berhardt, P. W., *et al.* 2007. CO₂ control of *Trichodesmium* N₂ fixation, photosynthesis, growth rates, and elemental ratios: implications for past, present, and future ocean biogeochemistry. *Limnology and Oceanography*, 52: 1293–1304.
- Isaji, S. 1995. Defensive strategies against shell dissolution in bivalves inhabiting acidic environments: the case of *Geloina* (Corbiculidae) in mangrove swamps. *The Veliger*, 38: 235–246.
- Ishimatsu, A., Kikkawa, T., Hayashi, M., Lee, K., and Kita, J. 2004. Effects of CO₂ on marine fish: larvae and adults. *Journal of Oceanography*, 60: 731–741.
- Ishimatsu, A., Hayashi, M., and Lee, S. 2005. Physiological effects on fishes in a high-CO₂ world. *Journal of the American Geophysical Union*, 110: C09S09.
- Ito, J. 1964. Food and feeding habit of Pacific salmon (genus *Oncorhynchus*) in their oceanic life. *Bulletin of the Hokkaido Regional Fisheries Research Laboratory*, 29: 85–97.
- Kalberer, M., Fischer, G., Patzold, J., Donner, B., Segl, M., and Wefer, G. 1993. Seasonal sedimentation and stable isotope records of pteropods off Cap Blanc. *Marine Geology*, 113: 305–320.

- Kennett, J. P., and Ingram, B. L. 1995. A 20 000-year record of ocean circulation and climate change from the Santa Barbara basin. *Nature*, 377: 510–514.
- Key, R. M. A., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., *et al.* 2004. A global ocean carbon climatology: results from Global Data Analysis Project (GLODAP). *Global Biogeochemical Cycles*, 18: 4031. doi:10.1029/2004GB002247.
- Kikkawa, T., Ishimatsu, A., and Kita, J. 2003. Acute CO₂ tolerance during the early developmental stages of four marine teleosts. *Environmental Toxicology*, 18: 375–382.
- Kikkawa, T., Sata, T., Kita, J., and Ishimatsu, A. 2006. Acute toxicity of temporally varying seawater CO₂ conditions on juveniles of Japanese sillago (*Sillago japonica*). *Marine Pollution Bulletin*, 52: 621–625.
- Kikkawa, T., Kita, J., and Ishimatsu, A. 2004. Comparison of the lethal effect of CO₂ and acidification on red sea bream (*Pagrus major*) during the early development stages. *Marine Pollution Bulletin*, 48: 108–110.
- Kleypas, J. A., Feely, R. A., Fabry, V. J., Langdon, C., Sabine, C. L., and Robbins, L. L. 2006. Impacts of ocean acidification on coral reefs and other marine calcifiers: a guide for future research. Report of a workshop held 18–20 April 2005, St Petersburg, FL, sponsored by NSF, NOAA, and the US Geological Survey. 88 pp.
- Kleypas, J. A., and Langdon, C. 2006. Coral reefs and changing seawater chemistry. In *Coral Reefs and Climate Change: Science and Management*, pp. 73–110. Ed. by J. T. Phinney, O. Hoegh-Guldberg, J. Kleypas, W. Skirving, and A. Strong, AGU Monograph Series Coastal Estuarine Studies 61. American Geophysical Union, Washington, DC.
- Knoll, A. H. 2003. Biomineralization and evolutionary history. In *Biomineralization*, Vol. 54, pp. 329–350. Ed. by P. Dove, J. J. De Young, and S. Weiner. The Mineralogical Society of America, Washington, DC. 381 pp.
- Knutzen, J. 1981. Effects of decreased pH on marine organisms. *Marine Pollution Bulletin*, 12: 25–29.
- Kobayashi, H. A. 1974. Growth cycle and related vertical distribution of the thecosomatous pteropod *Spiratella (Limacina) helicina* in the central Arctic Ocean. *Marine Biology*, 26: 295–301.
- Kochevar, R. E., and Childress, J. J. 1996. Carbonic anhydrase in deep-sea chemoautotrophic symbioses. *Marine Biology*, 125: 375–383.
- Köhler-Rink, S., and Kühl, M. 2005. The chemical microenvironment of the symbiotic planktonic foraminifer *Orbulina universa*. *Marine Biology Research*, 1: 68–78.
- Kuffner, I. B., Andersson, A. J., Jokiel, P. L., Rodgers, K., and Mackenzie, F. T. 2008. Decreased abundance of crustose coralline algae due to ocean acidification. *Nature Geoscience*, 1: 114.
- Kurihara, H., and Shirayama, Y. 2004. Effects of increased atmospheric CO₂ on sea urchin early development. *Marine Ecological Progress Series*, 274: 161–169.
- Kurihara, H., Shinji, S., and Shirayama, Y. 2004. Effects of raised CO₂ on the egg production rate and early development of two marine copepods (*Acartia steuri* and *Acartia erythraea*). *Marine Pollution Bulletin*, 49: 721–727.
- Lalli, C. M., and Gilmer, R. W. 1989. *Pelagic Snails: The Biology of Holoplanktonic Gastropod Mollusks*. Stanford University Press, Stanford, CA.
- Lallier, F., and Truchot, J. P. 1989. Modulation of haemocyanin oxygen-affinity by L-Lactate and urate in the prawn, *Penaeus japonicus*. *Journal of Experimental Biology*, 147: 133–146.
- Langdon, C., and Atkinson, M. J. 2005. Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *Journal of Geophysical Research, Oceans*, 110: C09S07. doi:10.1029/2004JC002576.
- Langdon, C., Broecker, W. S., Hammond, D. E., Glenn, E., Fitzsimmons, K., Nelson, S. G., Pend, T.-H., *et al.* 2003. Effect of elevated CO₂ on the community metabolism of an experimental coral reef. *Global Biogeochemical Cycles*, 17: 1011. doi:10.1029/2002GB00.
- Langdon, C., Takahashi, T., Sweeney, C., Chipman, D., Goddard, J., Marubini, F., Aceves, H., *et al.* 2000. Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochemical Cycles*, 14: 639–654.
- Langenbuch, M., and Pörtner, H. O. 2004. High sensitivity to chronically elevated CO₂ levels in a eurybathic marine sipunculid. *Aquatic Toxicology*, 70: 55–61.
- Langer, G., Geisen, M., Baumann, K.-H., Kläs, J., Riebesell, U., Thoms, S., and Young, J. R. 2006. Species-specific responses of calcifying algae to changing seawater carbonate chemistry. *Geochemistry, Geophysics, Geosystems*, Volume 7, Q09006, doi:10.1029/2005GC001127.
- LeBrasseur, R. J. 1966. Stomach contents of salmon and steelhead trout in the northeastern Pacific Ocean. *Journal of Fisheries Research Board Canada*, 23: 85–100.
- Leclercq, N., Gattuso, J.-P., and Jaubert, J. 2002. Primary production, respiration, and calcification of a coral reef mesocosm under increased CO₂ partial pressure. *Limnology and Oceanography*, 47: 558–564.
- Levasseur, M., Keller, M. D., Bonneau, E., D'Amours, D., and Bellows, W. K. 1994. Oceanographic basis of a DMS-related Atlantic cod (*Gadus morhua*) fishery problem: blackberry feed. *Canadian Journal of Fisheries and Aquatic Sciences*, 51: 881–889.
- Levi, R., Varona, P., Arshavsky, Y. I., Rabinovich, M. I., and Selverston, A. I. 2004. Dual sensory-motor function for a molluscan statocyst network. *Journal of Neurophysiology*, 91: 336–345.
- Lindinger, M. I., Lauren, D. J., and McDonald, D. G. 1984. Acid-base balance in the sea mussel, *Mytilus edulis*. III. Effects of environmental hypercapnia on intra- and extracellular acid-base balance. *Marine Biology Letters*, 5: 371–381.
- Lovejoy, T. E., and Hannah, L. 2005. *Climate Change and Biodiversity*. Yale University Press, New Haven, CT. 418 pp.
- Lowe, T. E., Brill, R. W., and Cousins, K. L. 2000. Blood oxygen-binding characteristics of bigeye tuna (*Thunnus obesus*), a high-energy-demand teleost that is tolerant of low ambient oxygen. *Marine Biology*, 136: 1087–1098.
- Lowenstam, H. A., and Weiner, S. 1989. *On Biomineralization*. Oxford University Press, Oxford.
- Mackas, D. L., and Galbraith, M. D. 2002. Zooplankton distribution and dynamics in a North Pacific eddy of coastal origin: I. Transport and loss of continental margin species. *Journal of Oceanography*, 58: 725–738.
- Mangum, C. P. 1991. Salt sensitivity of the hemocyanin of eury- and stenohaline squids. *Comparative Biochemistry and Physiology*, 99A, 159–161.
- Marubini, F., Barnett, H., Langdon, C., and Atkinson, M. J. 2001. Dependence of calcification on light and carbonate ion concentration for the hermatypic coral *Porites compressa*. *Marine Ecology Progress Series*, 220: 153–162.
- Marubini, F., Ferrier-Pages, C., and Cuif, J. P. 2003. Suppression of skeletal growth in scleractinian corals by decreasing ambient carbonate-ion concentration: a cross-family comparison. *Proceedings of the Royal Society of London, Series B*, 270: 179–184.
- Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., *et al.* 2007. Global climate projections. In *Climate Change 2007: the Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. pp.747–846. Ed. by S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, *et al.* Cambridge University Press, Cambridge.

- Metzger, R., Sartoris, F. J., Langebuch, M., and Pörtner, H. O. 2007. Influence of elevated CO₂ concentrations on thermal tolerance of the edible crab, *Cancer pagurus*. *Journal of Thermal Biology*, 32: 144–151.
- Michaelidis, B., Ouzounis, C., Paleras, A., and Pörtner, H. O. 2005. Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, 293: 109–118.
- Michaelidis, B., Spring, A., and Pörtner, H. O. 2007. Effects of long-term acclimation to environmental hypercapnia on extracellular acid-base status and metabolic capacity in Mediterranean fish *Sparus aurata*. *Marine Biology*, 150: 1417–1429.
- Miles, H., Widdicombe, S., Spicer, J. I., and Hall-Spencer, J. 2007. Effects of anthropogenic seawater acidification on acid-base balance in the sea urchin *Psammechinus miliaris*. *Marine Pollution Bulletin*, 54: 89–96.
- Millero, F. J., Pierrot, D., Lee, K., Wanninkhof, R., Feely, R. A., Sabine, C. L., Key, R. M., et al. 2002. Dissociation constants for carbonic acid determined from field measurements. *Deep Sea Research I*, 49: 1705–1723.
- Mohan, R., Verma, K., Mergulhao, L. P., Sinha, D. K., Shanvas, S., and Guptha, V. S. 2006. Seasonal variation of pteropods from the Western Arabian Sea sediment trap. *Geo-Marine Letters*, 26: 265–273.
- Mucci, A. 1983. The solubility of calcite and aragonite in seawater at various salinities, temperatures and 1 atmosphere total pressure. *American Journal of Science*, 238: 780–799.
- Nigmatullin, C. M., Nesis, K. N., and Arkhipkin, A. I. 2001. A review of the biology of the jumbo squid *Dosidicus gigas* (Cephalopoda: Ommastrephidae). *Fisheries Research*, 54: 9–19.
- Noble, R. W., Kwiatkowski, L. D., deYoung, A., Davis, B. J., Haedrich, R. L., Tam, L. T., and Riggs, F. A. 1986. Functional properties of hemoglobins from deep-sea fish: correlations with depth distribution and presence of a swimbladder. *Biochimica et Biophysica Acta*, 870: 552–563.
- Noji, T. T., Bathmann, U. V., von Bodungen, B., Voss, M., Antia, A., Krumbholz, M., Klein, B., et al. 1997. Clearance of picoplankton-sized particles and formation of rapidly sinking aggregates by the pteropod, *Limacina retroversa*. *Journal of Plankton Research*, 19: 863–875.
- Ohde, S., and Hossain, M. M. M. 2004. Effect of CaCO₃ (aragonite) saturation state of seawater on calcification of *Porites* coral. *Geochemical Journal*, 38: 613–621.
- Orr, J. C., Anderson, L. G., Bates, N. R., Bopp, L., Fabry, V. J., Jones, E., and Swingedouw, D. 2006. Arctic Ocean acidification. *EOS Transactions of the American Geophysical Union*, 87(36), Ocean Sciences Meeting Supplement, Abstract OS14B-01.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., et al. 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, 437: 681–686.
- Pane, E. F., and Barry, J. P. 2007. Extracellular acid-base regulation during short-term hypercapnia is effective in a shallow-water crab, but ineffective in a deep-sea crab. *Marine Ecology Progress Series*, 334: 1–9.
- Pane, L., Feletti, M., Francomacaro, B., and Mariottini, G. L. 2004. Summer coastal zooplankton biomass and copepod community structure near the Italian Terra Nova Base (Terra Nova Bay, Ross Sea, Antarctica). *Journal of Plankton Research*, 26: 1479–1488.
- Pörtner, H. O. 1990. Determination of intracellular buffer values after metabolic inhibition by fluoride and nitrilotriacetic acid. *Respiration Physiology*, 81: 275–288.
- Pörtner, H. O. 1994. Coordination of metabolism acid-base regulation and haemocyanin function in cephalopods. In *Physiology of Cephalopod Molluscs: Lifestyle and Performance Adaptations*, pp. 131–148. Ed. by H. O. Pörtner, R. K. O'Dor, and D. L. MacMillan. Gordon and Breach, London.
- Pörtner, H. O., and Reipschläger, A. 1996. Ocean disposal of anthropogenic CO₂: physiological effects on tolerant and intolerant animals. In *Ocean Storage of Carbon Dioxide. Workshop 2—Environmental Impact*, pp. 57–81. Ed. by B. Ormerod, and M. V. Angel. IEA Greenhouse Gas R&D Programme, Cheltenham, UK.
- Pörtner, H. O., Langebuch, M., and Reipschläger, A. 2004. Biological impact of elevated ocean CO₂ concentration: lessons from animal physiology and Earth history. *Journal of Oceanography*, 60: 705–718.
- Raz, S., Hamilton, P. C., Wilt, F. H., Weiner, S., and Addadi, L. 2003. The transient phase of amorphous calcium carbonate in sea urchin larval spicules: the involvement of proteins and magnesium ions in its formation and stabilization. *Advanced Functional Materials*, 13: 480–486.
- Ridgwell, A., and Zeebe, R. E. 2005. The role of the global carbonate cycle in the regulation and evolution of the Earth system. *Earth and Planetary Science Letters*, 234: 299–315.
- Riebesell, U., Zondervan, I., Rost, B., Tortell, P. D., Zeebe, R. E., and Morel, F. M. M. 2000. Reduced calcification of marine plankton in response to increased atmospheric CO₂. *Nature*, 407: 364–367.
- Rink, S., Kühl, M., Bijma, J., and Spero, H. J. 1998. Microsensor studies of photosynthesis and respiration in the symbiotic foraminifera *Orbulina universa*. *Marine Biology*, 131: 583–595.
- Rodhouse, P. G., and White, M. G. 1995. Cephalopods occupy the ecological niche of epipelagic fish in the Antarctic Polar Frontal Zone. *Biological Bulletin*, 189: 77–80.
- Royal Society. 2005. Ocean acidification due to increasing atmospheric carbon dioxide. Policy Document 12/05, The Royal Society, London. 60 pp.
- Rutherford, L. D., and Thuesen, E. V. 2005. Metabolic performance and survival of medusae in estuarine hypoxia. *Marine Ecology Progress Series*, 294: 189–200.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., et al. 2004. The oceanic sink for CO₂. *Science*, 305: 367–371.
- Sanders, N. K., and Childress, J. J. 1990. Adaptations to the deep-sea oxygen minimum layer: oxygen binding by the hemocyanin of the bathypelagic mysid, *Gnathophausia ingens* Dohrn. *Biological Bulletin*, 178: 286–294.
- Schiebel, R. 2002. Planktonic foraminiferal sedimentation and the marine calcite budget. *Global Biogeochemical Cycles*, 16: doi: 10.1029/2001GB1459.
- Schippers, P., Lüring, M., and Scheffer, M. 2004. Increase of atmospheric CO₂ promotes phytoplankton productivity. *Ecology Letters*, 7: 446–451.
- Schneider, K., and Erez, J. 2006. The effect of carbonate chemistry on calcification and photosynthesis in the hermatype coral *Acropora eurystoma*. *Limnology and Oceanography*, 51: 1284–1293.
- Seibel, B. A. 2007. On the depth and scale of metabolic rate variation: scaling of oxygen consumption and enzymatic activity in the Class Cephalopoda (Mollusca). *Journal of Experimental Biology*, 210: 1–11.
- Seibel, B. A., Chausson, F., Lallier, F., Childress, J. J., and Zal, F. 1999. Vampire blood: respiratory physiology of the vampire squid (Vampyromorpha: Cephalopoda) in relation to the oxygen minimum layer. *Experimental Biology Online*, 4(1): 1–10. ISSN:1430-3418.
- Seibel, B. A., and Drazen, J. C. 2007. The rates of metabolism in marine animals: environmental constraints, ecological demands and energetic opportunities. *Philosophical Transactions of the Royal Society, Series B – Biological Sciences*, 362: 2061–2071.
- Seibel, B. A., Dymowska, A., and Rosenthal, J. 2007. Metabolic temperature compensation and coevolution of locomotory performance in pteropod molluscs. *Integrative and Comparative Biology*, 46: 880–891.
- Seibel, B. A., and Walsh, P. J. 2001. Potential impacts of CO₂ injection on deep-sea biota. *Science*, 294: 319–320.

- Seibel, B. A., and Walsh, P. J. 2003. Biological impacts of deep-sea carbon dioxide injection inferred from indices of physiological performance. *Journal of Experimental Biology*, 206: 641–650.
- Seibel, B. A., Thuesen, E. V., Childress, J. J., and Gorodezky, L. A. 1997. Decline in pelagic cephalopod metabolism with habitat depth reflects differences in locomotory efficiency. *Biological Bulletin*, 192: 262–278.
- Shirayama, Y., and Thorton, H. 2005. Effect of increased atmospheric CO₂ on shallow water marine benthos. *Journal of Geophysical Research*, 110: C09S08. doi: 10.1029/2004JC002618.
- Siegenthaler, U., Stocker, T. F., Monnin, E., Luethi, D., Schwander, J., Stauffer, B., Raynaud, D., *et al.* 2005. Stable carbon cycle-climate relationship during the late Pleistocene. *Science*, 310: 1313–1317.
- Silverman, J., Lazar, B., and Erez, J. 2007. Effect of aragonite saturation, temperature, and nutrients on the community calcification rate of a coral reef. *Journal of Geophysical Research*, 112: C05004. doi:10.1029/2006JC003770.
- Smith, K. L., and Teal, J. M. 1973. Temperature and pressure effects on respiration of thecosomatous pteropods. *Deep Sea Research*, 20: 853–858.
- Somero, G. N. 1985. Intracellular pH, buffering substances and proteins: imidazole protonation and the conservation of protein structure and function. In *Transport Processes, Iono- and Osmoregulation*, pp. 454–468. Ed. by R. Gilles, and M. Gilles-Baillien. Springer-Verlag, Berlin.
- Somero, G. N. 2005. Linking biogeography to physiology: evolutionary and acclimatory adjustments of thermal limits. *Frontiers in Zoology*, 2: 1. doi:10.1186/1742-9994-2-1.
- Spero, H. J., Bijma, J., Lea, D. W., and Bemis, B. E. 1997. Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes. *Nature*, 390: 497–500.
- Spero, H. J., and Lea, D. W. 1993. Intraspecific stable isotope variability in the planktic foraminifera *Globigerinoides sacculifer*: results from laboratory experiments. *Marine Micropaleontology*, 22: 221–234.
- Spicer, J. I. 1995. Oxygen and acid-base status of the sea urchin *Psammechinus miliaris* during environmental hypoxia. *Marine Biology*, 124: 71–76.
- Spicer, J. I., Raffo, A., and Widdicombe, S. 2007. Influence of CO₂-related seawater acidification on extracellular acid-base balance in the velvet swimming crab, *Necora puber*. *Marine Biology*, 151: 1117–1125.
- Stillman, J. H. 2003. Acclimation capacity underlies susceptibility to climate change. *Science*, 301: 65.
- Tamburrini, M., Romano, M., Carratore, V., Kunzmann, A., Coletta, M., and di Prisco, G. 1998. The hemoglobins of the Antarctic fishes *Artedidraco orianae* and *Pogonophryne scottii*. *Journal of Biological Chemistry*, 273: 32 452–32 459.
- Taylor, J. R., and Grosell, M. 2006. Evolutionary aspects of intestinal bicarbonate secretion in fish. *Comparative Biochemistry and Physiology, Part A*, 143: 423–529.
- Thibault, D., Roy, S., Wong, C. S., and Bishop, J. K. 1999. The downward flux of biogenic material in the NE subarctic Pacific: importance of algal sinking and mesozooplankton herbivory. *Deep Sea Research*, 46: 2669–2697.
- Thuesen, E. V., McCullough, K. D., and Childress, J. J. 2005a. Metabolic enzyme activities in swimming muscle of medusae: is the scaling of glycolytic activity related to oxygen availability? *Journal of the Marine Biological Association of the UK*, 85: 603–611.
- Thuesen, E. V., Rutherford, L. D., Brommer, P. L., Garrison, K., Gutowska, M. A., and Towanda, T. 2005b. Intragel oxygen promotes hypoxia tolerance of scyphomedusae. *Journal of Experimental Biology*, 208: 2475–2482.
- Truchot, J. P. 1987. Comparative aspects of extracellular acid-base balance. Springer-Verlag, Berlin.
- Truchot, J. P., and Duhamel-Jouve, A. 1980. Oxygen and carbon dioxide in the marine intertidal environment: diurnal and tidal changes in rockpools. *Respiration Physiology*, 39: 241–254.
- Tsurumi, M., Mackas, D. L., Whitney, E. A., Dibacco, C., Galbraith, M. D., and Wong, C. S. 2005. Pteropods, eddies, carbon flux and climate variability in the Alaska Gyre. *Deep Sea Research II*, 52: 1037–1053.
- Tyrrill, T., and Zeebe, R. E. 2004. History of carbonate ion concentration over the last 100 million years. *Geochimica et Cosmochimica Acta*, 68: 3521–3530.
- Walsh, P. J., Blackwelder, P., Gill, K. A., Danulat, E., and Mommsen, T. P. 1991. Carbonate deposits in marine fish intestines: a new source of biomineralization. *Limnology and Oceanography*, 36: 1227–1232.
- Walsh, P. J., and Milligan, C. L. 1989. Coordination of metabolism and intracellular acid-base status: ionic regulation and metabolic consequences. *Canadian Journal of Zoology*, 67: 2994–3004.
- Walter, L. M., and Morse, J. W. 1985. Magnesian calcite stabilities: a reevaluation. *Geochimica et Cosmochimica Acta*, 49: 1503–1513.
- Watanabe, H., Kawaguchi, K., and Hayashi, A. 2002. Feeding habits of juvenile surface-migratory myctophid fishes (family Myctophidae) in the Kuroshio region of the western North Pacific. *Marine Ecology Progress Series*, 236: 263–272.
- Watanabe, Y., Yamaguchi, A., Ishida, H., Harimoto, T., Suzuki, S., Sekido, Y., Ikeda, T., *et al.* 2006. Lethality of increasing CO₂ levels on deep-sea copepods in the western North Pacific. *Journal of Oceanography*, 62: 185–196.
- Weiss, I. M., Tuross, N., Addadi, L., and Weiner, S. 2002. Mollusc larval shell formation: amorphous calcium carbonate is the precursor phase for aragonite. *Journal of Experimental Biology*, 293: 478–491.
- Willette, T. M., Cooney, R. T., Patrick, V., Mason, D. M., Thomas, G. L., and Scheel, D. 2001. Fisheries Oceanography. 10: 14–41.
- Yamada, Y., and Ikeda, T. 1999. Acute toxicity of lowered pH to some oceanic zooplankton. *Plankton Biology and Ecology*, 46: 62–67.
- Zielinski, S., Sartoris, F. J., and Pörtner, H. O. 2001. Temperature effects on hemocyanin oxygen binding in an Antarctic cephalopod. *Biological Bulletin*, 200: 67–76.

doi:10.1093/icesjms/fsn048