

Dissolved Organic Matter and Heterotrophic Microneuston in the Surface Microlayers of the North Atlantic

Abstract. Dissolved organic carbon, carbohydrates, and adenosine triphosphate in the size fractions 0.2 to 3 micrometers and 3 to 1000 micrometers are significantly enriched in the upper 150-micrometer surface layer compared to subsurface water, mean enrichment factors being 1.6, 2.0, 2.5, and 3.1, respectively. When calculated as a 0.1-micrometer microlayer of wet surfactants, the mean concentration of organic matter was 2.9 grams per liter, of which carbohydrates accounted for 28 percent. The data for plant pigments and particulate adenosine triphosphate indicated that bacterioneuston was enriched at seven of nine stations while phagotrophic protists were enriched at five stations. Instances of enrichment and inhibition were verified by cultural data for bacteria and amoebas. The observations indicate that the surface microlayers are largely heterotrophic microcosms, which can be as rich as laboratory cultures, and that an appreciable part of the dissolved organic carbon is carbohydrate of phytoplankton origin, released and brought to the surface by migrating and excreting phagotrophic protists.

The sea-air interface has attracted attention because of its importance in the detection and limitation of pollution (1). However, the sea surface is a major site of biological activity, and the organisms that have adapted to this specialized environment (2) have been termed neuston (3). Despite the importance of neuston in the diet of both sea birds and migrating nekton and the growing knowledge of the biology of the neuston (4), the chemical and microbiological nature of the surface film and the physical and biological mechanisms for its formation require further study. Light intensities at the sea surface are phototoxic to most phytoplankters; exceptions are *Trichodesmium*, *Sargassum*, and some pennate diatoms that are lifted onto the surface film in intertidal waters (5). Therefore the microneuston that nurtures the conspicuous blue invertebrate fauna on the open sea must be largely heterotrophic in nature and be based on bacterioneuston (6) growing on nutrients produced by the phytoplankton in the shaded euphotic zone below the sea surface.

It has been assumed that the surfactant nature of the surface layer is due to hydrocarbons, fatty acids and fatty alcohols, and similar dry surfactants (1, 7), which produce a 0.01- μm microlayer above the sea-air interface (8). However, it has been shown that except for heavily traveled waterways polluted by hydrocarbons, the surface microlayer is dominated by a 0.1- μm -thick microlayer of wet surfactants below the sea-air interface that behaves like a polysaccharide-protein complex (9). The study reported here was designed to further characterize the nature and concentration of the dissolved organic carbon (DOC) and microorganisms in the surface microlayers in an attempt to show the relationship of the microbial plankton and neuston to the chemistry of the microlayer. Data are

presented to show that there is a small but consistent and statistically significant enrichment of both DOC and microneuston in the surface film. If the dissolved materials are actually concentrated in a 0.1- μm microlayer (8, 9) while the microorganisms are concentrated in a 1.0- μm microlayer, then the nutrient concentrations and microbial populations approach or equal those in laboratory cultures.

Stations were occupied at the locations shown in Table 1. A Nyltex screen sampler (1) that had been scrubbed in alcohol was used to obtain the approximately 150- μm -thick surface layer. When the sea state allowed, an inflatable boat with low freeboard was used to sample away from the ship. Subsurface samples, taken with 5- and 30-liter Niskin bottles that had been scrubbed with 0.1N hydrochloric acid and rinsed in water in situ for 20 minutes before sampling (10), were filtered with a 1-mm Nyltex screen and processed as rapidly as possible. The chemical and adenosine triphosphate (ATP) analyses were made in triplicate. The samples for DOC and carbohydrates were passed through Gelman glass-fiber filters (type A/E) without vacuum to remove suspended particles. Samples for total DOC (11) were prepared at sea and analyzed on shore. The determination of monosaccharides and total carbohydrates and the estimation of polysaccharides by difference were done aboard ship by using procedures based on precise and reliable chemical reactions (12). Organisms smaller than 1000 μm were divided into fractions $< 3 \mu\text{m}$ and $> 3 \mu\text{m}$ by Nuclepore filtration; the amount of ATP was determined at sea between stations (13, 14), and plant pigments (15) were used to detect accumulations of phytoplankton. Large volumes of water (up to 200 liters) were passed through a continuous reverse flow con-

centrator (16) that reduced the volume containing the particles $> 3 \mu\text{m}$ by a factor of 1000 in 2 hours or less. Fresh concentrates were examined at sea with phase contrast microscopes fitted with head braces (necessary on a rolling ship); portions fixed at sea by the addition of an equal volume of 6 percent glutaraldehyde in 0.1M sodium cacodylate buffer (pH 7.20) with 0.25M sucrose were examined in greater detail on shore by light microscopy and scanning and transmission electron microscopy. Fresh concentrates were also used to prepare serial decimal dilutions for the extinction dilution culture of bacteria and amoebas. For bacteria, the peptone-yeast extract medium contained 1 g of each per liter of seawater, the carbohydrate medium contained 1 g each of glucose, mannitol, and glycolic acid per liter, and the chitin medium contained approximately 2 g of crab chitin suspended in 1 liter of seawater. Either a single rice grain or a 0.1-cm³ 1.5 percent agar plug of the peptone-yeast extract medium was used to sustain low bacterial populations for the amoeba cultures.

The data obtained for the screen and subsurface samples at nine stations from Block Island Sound along the outer continental shelf to the Grand Banks (stations 2, 5, 6, and 7) and across the North Atlantic to east of the Azores (stations 9, 11, 12, 13, and 14) are given in Table 1. The differences for all samples are significant at the 95 percent level except for two values. The concentration of materials in the surface microlayers was calculated by subtracting the subsurface (mean depth, 8 m) value from the screen value, and the remainder was multiplied by a factor based on a screen efficiency of 70 percent (17) and a dilution of the microlayer with the subsurface water (1). Assumed thicknesses of the components used to estimate the concentration of dissolved organic matter and microorganisms in the surface microlayer are shown in Table 1.

In contrast to the DOC concentrations of ≤ 1.3 mg of carbon per liter in the subsurface waters, the DOC in the microlayer has an estimated effective carbon concentration of 471 to 2250 mg/liter (mean value, 1427 mg/liter) or a dry weight of 2.9 g of organic matter per liter. This is the concentration of organic matter used to grow bacteria in seawater media. A mean value for total carbohydrates of 875 mg/liter indicates that carbohydrates account for some 28 percent of the DOC in the microlayer. The particulates $< 3 \mu\text{m}$ are primarily bacteria, although some of the smaller apochlorotic microflagellates, which presumably func-

135271

2

tion like bacteria, may also be present, depending on the sample. The large, chloroplast-containing phytoplankton were apparently not present in these preparations. An idea of the order of magnitude of the biomass of the bacterioneuston can be obtained by using the ATP : C conversion factor of 1 : 250 (14), which indicates concentrations of 0.08 to 5 mg of cellular carbon per liter.

The ATP concentration for the protists in the fraction > 3 μm , when present, was a third that of the bacterioneuston. Except at the shallow-water station 2, there was no appreciable concentration of chlorophyll a. These observations indicate that this > 3- μm fraction was usually due to phagotrophic protists, presumably grazing on the bacterioneuston. This was confirmed by microscopic ob-

servation of concentrates. At two stations the ATP concentration in the fraction < 3 μm was lower in the microlayer than in the subsurface sample; this may have been caused by grazing or by growth inhibition. The data on dilution extinction cultures for bacteria and amoebas at stations 2 and 12 are given in Table 2. The apparent inhibition at station 2 indicated in Table 1 is also con-

Table 1. Concentrations of dissolved organic carbon (DOC) and living particulates in the surface microlayers of the North Atlantic in July and August 1975. In the calculations a thickness commensurate with the dimensions of the material was assumed: a 0.1- μm layer for DOC gives a dilution factor of 1 : 1500; a 1.0- μm layer for the fraction < 3 μm , 1 : 150; and a 10- μm layer for the protists > 3 μm , 1 : 15. Microlayer concentration = (surface concentration - subsurface concentration) (70 percent screen efficiency \times dilution with subsurface water). Enrichment is the screen enrichment factor, that is, (screen concentration)/(subsurface concentration). Differences between the means for the surface and subsurface determinations were statistically significant at the 95 percent level with two exceptions noted below. Letters in parentheses under station numbers indicate screen sampling method: Z means sampled from the Zodiac inflatable, and V means sampled from the bow of the research vessel.

| Materials assayed | Station 2 (Z) | Station 5 (V) | Station 6 (V) | Station 7 (V) | Station 9 (V) | Station 11 (Z) | Station 12 (Z) | Station 13 (Z) | Station 14 (V) | Means |
|--|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|-------|
| Location | 41°48'N, 67°05'W | 43°31'N, 59°29'W | 43°19'N, 59°41'W | 43°40'N, 52°40'W | 40°36'N, 39°48'W | 37°28'N, 28°43'W | 37°28'N, 28°44'W | 36°59'N, 21°22'W | 36°47'N, 21°20'W | |
| Depth (m) | 50 | 1794 | 160 | 296 | 4972 | 2000 | 1880 | 3803 | 4517 | |
| Dissolved organic carbon (mg/liter) | | | | | | | | | | |
| Microlayer | 1714 | 471 | 2186 | 921 | 2207 | 1671 | 1029 | 600 | 1221 | 1336 |
| Screen | 1.88 | 1.55* | 2.11 | 1.70 | 2.15 | 1.72 | 1.47 | 1.32 | 1.64 | 1.73 |
| Subsurface | 1.08 | 1.33 | 1.09 | 1.27 | 1.12 | 0.94 | 0.99 | 1.04 | 1.07 | 1.10 |
| Enrichment | 1.7 | 1.2 | 1.9 | 1.3 | 1.9 | 1.8 | 1.5 | 1.3 | 1.5 | 1.6 |
| Carbohydrate carbon (mg/liter) | | | | | | | | | | |
| Monosaccharides | | | | | | | | | | |
| Microlayer | 214 | 111 | 96 | 86 | 334 | 244 | 107 | 103 | 150 | 161 |
| Screen | 0.150 | 0.126 | 0.124 | 0.182 | 0.230 | 0.192 | 0.128 | 0.131 | 0.144 | 0.156 |
| Subsurface | 0.050 | 0.074 | 0.079 | 0.142 | 0.074 | 0.078 | 0.078 | 0.083 | 0.074 | 0.081 |
| Enrichment | 3.0 | 1.7 | 1.6 | 1.3 | 3.1 | 2.5 | 1.6 | 1.6 | 1.9 | 2.0 |
| Polysaccharides | | | | | | | | | | |
| Microlayer | 221 | NE† | 193 | 257 | 491 | 154 | 139 | 103 | 411 | 246 |
| Screen | 0.220 | 0.154 | 0.190 | 0.205 | 0.273 | 0.168 | 0.137 | 0.141 | 0.301 | 0.199 |
| Subsurface | 0.117 | 0.171 | 0.100 | 0.085 | 0.044 | 0.096 | 0.072 | 0.093 | 0.109 | 0.099 |
| Enrichment | 1.9 | 0.9 | 1.9 | 2.4 | 6.2 | 1.8 | 1.9 | 1.5 | 2.8 | 2.3 |
| Total carbohydrates | | | | | | | | | | |
| Microlayer | 435 | 75 | 289 | 343 | 825 | 398 | 246 | 206 | 561 | 375 |
| Screen | 0.370 | 0.280 | 0.314 | 0.387 | 0.503 | 0.360 | 0.265 | 0.272 | 0.445 | 0.355 |
| Subsurface | 0.167 | 0.245 | 0.179 | 0.227 | 0.118 | 0.174 | 0.150 | 0.176 | 0.183 | 0.180 |
| Enrichment | 2.2 | 1.1 | 1.7 | 1.7 | 4.3 | 2.1 | 1.8 | 1.5 | 2.4 | 2.0 |
| Percentage of DOC | | | | | | | | | | |
| Microlayer | 25 | 16 | 13 | 37 | 37 | 24 | 24 | 34 | 46 | 28 |
| Screen | 20 | 18 | 15 | 23 | 23 | 21 | 18 | 21 | 27 | 21 |
| Subsurface | 15 | 18 | 16 | 18 | 10 | 18 | 15 | 17 | 17 | 16 |
| Biomass (10 ⁻⁸ g ATP/liter) | | | | | | | | | | |
| Fraction < 3 μm | | | | | | | | | | |
| Microlayer | NE† | 793 | 28 | NE | 1072 | 165 | 2145 | 362 | 1779 | 906 |
| Screen | 5.19 | 6.96 | 2.73 | 3.45 | 9.32 | 3.51 | 11.67 | 3.37 | 10.14 | 6.26 |
| Subsurface | 6.14 | 3.26 | 2.60 | 5.89 | 4.32 | 2.74 | 1.66 | 1.68 | 1.84 | 3.35 |
| Enrichment | 0.8 | 2.1 | 1.1 | 0.6 | 2.2 | 1.3 | 7.0 | 2.0 | 5.5 | 2.5 |
| Fraction > 3 μm | | | | | | | | | | |
| Microlayer | 664 | NE | NE | NE | 273 | 58 | 204 | NE | 245 | 289 |
| Screen | 44.98 | 3.59 | 5.26* | 3.81 | 21.18 | 4.58 | 11.10 | 4.71 | 12.64 | 12.43 |
| Subsurface | 14.00 | 13.45 | 5.87 | 7.44 | 8.43 | 1.88 | 1.59 | 5.72 | 1.23 | 6.62 |
| Enrichment | 3.2 | 0.3 | 0.9 | 0.5 | 2.5 | 2.4 | 7.0 | 0.8 | 10.3 | 3.1 |
| Pigments ($\mu\text{g/liter}$) | | | | | | | | | | |
| Chlorophyll a | | | | | | | | | | |
| Microlayer | 50 | NE | NE | 1.28 | 0.64 | NE | 0.21 | 0.08 | 1.50 | 8.95 |
| Screen | 3.78 | 1.09 | 0.00 | 0.19 | 0.14 | 0.05 | 0.07 | 0.08 | 0.14 | 0.62 |
| Subsurface | 1.46 | 2.05 | 0.10 | 0.13 | 0.11 | 0.07 | 0.06 | 0.08 | 0.07 | 0.45 |
| Enrichment | 2.6 | 0.5 | | 1.5 | 1.3 | 0.7 | 1.2 | 1.0 | 2.0 | 1.4 |
| Phaeopigments | | | | | | | | | | |
| Microlayer | NE | NE | NE | 0.02 | NE | 0.21 | 1.07 | 0.03 | 0.43 | 0.35 |
| Screen | 0.13 | 1.14 | 0.00 | 0.02 | 0.03 | 0.03 | 0.06 | 0.03 | 0.03 | 0.16 |
| Subsurface | 0.60 | 6.75 | 0.06 | 0.02 | 0.06 | 0.02 | 0.01 | 0.03 | 0.03 | 0.84 |
| Enrichment | 0.2 | 0.2 | | 1.0 | 0.5 | 1.5 | 6.0 | 1.0 | 3.0 | 1.7 |

*Not significant. †NE = not enriched.

firmed by the cultural data. If inhibition rather than grazing occurred, it could have been due to substances (18) concentrated in the microlayer that were released from the appreciable *Nitzschia* bloom at this station indicated by the chlorophyll value. An instance of bacterial inhibition in the surface film has been reported (19). In contrast to the cultural data from station 2, the data from station 12 indicated a marked enrichment. When corrected for the thickness of the surface microlayer, bacterial populations on all three media were in line with the biomass estimates from ATP. Data on the amoebas associated with the microlayer correlated with those for the bacteria: extensive bacterial populations supported extensive amoeba populations.

The blue-pigmented pontellid copepods floating on the surface at several stations occurred in concentrations sufficient to be noticed during screen sampling. At station 12 half the specimens were infested with the suctorian *Ephelota gemmipara*, with as many as 80 individuals per copepod. A group of individuals from juvenile to an exogenously budding mature form are shown in Fig. 1A. The suctorians were present mainly in a band along the waterline of these surface-dwelling copepods. The only other copepod infested with stalked ciliates was *Acartia tonsa* from nearshore station 1, which was heavily infested with *Zoothamnium alternans* (20). The restriction of these epizotic stalked ciliates to nearshore waters and to the surface microlayer and fouled surfaces (21) of open ocean waters indicates that they probably occur only when there is an abundant source of microorganisms in the water. Thin sections of the suctorians (Fig. 1) indicate that bacteria are a substantial part of their diet and must occur in significant concentrations in the surface microlayer.

Carbohydrates accounted for 12 to 46 percent of the DOC (mean, 28 percent). They are produced by the phytoplankton as photosynthate, reserve, extracellular, and cell wall materials (22), and may occur in part as side chains on protein backbones—that is, glycoproteins and mucopolysaccharides. The regular diurnal accumulation of monosaccharides, polysaccharides, and DOC in the subsurface waters at these stations was strongly associated with phagotrophic protist maxima occurring near phototrophic protist maxima (23). Such a continual replenishment of dissolved organic matter would be needed to sustain the microbial activity we observed. The enrichment of wet surfactants in the surface microlayers is probably achieved through a

Table 2. Cultivable populations of bacteria and amoebas in a theoretical 1- μ m surface microlayer at station 2 (inhibitory) and station 12 (enriched).

| Station | Sample | Bacteria per milliliter | | | Amoebas per liter | |
|---------|---------------------------|-------------------------|-----------------------|-----------------|-------------------|--------------------|
| | | Peptone yeast medium | Monosaccharide medium | Chitin medium | Rice grain | Nutrient agar plug |
| 2 | Microlayer (0.1 μ m)* | NE† | NE | NE | NE | NE |
| | Screen (150 μ m) | 5×10^2 | 5×10^1 | 5×10^1 | <0.01 | <0.01 |
| | Subsurface (37 m) | 1×10^3 | 5×10^1 | 1×10^2 | 0.50 | 2.30 |
| 12 | Microlayer (0.1 μ m) | $>2 \times 10^7$ | $>2 \times 10^7$ | 1×10^7 | 1200 | 1350 |
| | Screen (150 μ m) | $>1 \times 10^5$ | $>1 \times 10^5$ | 5×10^4 | 6.5 | 6.5 |
| | Subsurface (30 m) | 5×10^2 | 1×10^1 | 5×10^1 | 0.9 | 0.2 |

* Estimated from the screen minus subsurface concentrations $\times 214$ (which allows for 70 percent screen efficiency and a 1 : 150 dilution with subsurface water). † NE = not enriched.




Fig. 1. Electron micrographs (EM) of the suctorian *Ephelota gemmipara* colonizing the lateral surface of the blue copepod *Pontella scutifer* floating in the surface microlayer of the North Atlantic. (A) Scanning EM of several *E. gemmipara* cells in various stages of development. Three of the cells show the prehensile (outer) and feeding (inner) tentacles used to capture and ingest food particles. One exogenously budding individual (arrow) has five ciliated telotrochs. Scale bar, 100 μ m. (B) Transmission EM of a thin section of part of an *E. gemmipara* cell with nucleus (Nu) and stalk (s). Food vacuoles (v) show that these suctorians have been feeding mainly on bacteria. Scale bar, 5 μ m. (C) Higher magnification of the ingested bacteria (b) in a vacuole. Scale bar, 1 μ m.

combination of subsurface phototrophic production and phagotrophic release, Langmuir circulations (24), extraction and transport through bubble action during water turbulence (25), and the attraction of the hydrophobic groups to the sea-air interface.

JOHN MCN. SIEBURTH

PAULA-JEAN WILLIS

KENNETH M. JOHNSON

CURTIS M. BURNEY

DENNIS M. LAVOIE

KENNETH R. HINGA

DAVID A. CARON

FREDERICK W. FRENCH III

PAUL W. JOHNSON, PAUL G. DAVIS

Graduate School of Oceanography,

University of Rhode Island,

Kingston 02881

References and Notes

1. R. A. Duce, J. G. Quinn, C. E. Olney, S. R. Piotrowicz, B. J. Ray, T. L. Wade, *Science* **176**, 161 (1972).
2. P. M. David, *Endeavour* **24**, 95 (1965).
3. E. Naumann, *Biol. Zentralbl.* **37**, 98 (1917).
4. Yu. P. Zaitsev, *Marine Neustonology*, K. A. Vinogradov, Ed. (Israel Program for Scientific Translations, Jerusalem, 1971), pp. 51-171; L. Cheng, *Oceanogr. Mar. Biol.* **13**, 181 (1975).
5. J. L. Gallagher, *Limnol. Oceanogr.* **20**, 120 (1975).
6. J. McN. Sieburth, in *Ocean Science and Ocean Engineering* (Marine Technology Society and American Society of Limnology and Oceanography, Washington, D.C., 1965), pp. 1064-1068; S. A. Crow, D. G. Ahearn, W. L. Cook, *Limnol. Oceanogr.* **20**, 664 (1975).
7. W. D. Garrett, *Deep-Sea Res.* **14**, 221 (1967); N. L. Jarvis, W. D. Garrett, M. A. Schelman, C. O. Timmons, *Limnol. Oceanogr.* **12**, 88 (1967).
8. F. MacIntyre, *Sci. Am.* **230** (No. 5), 62 (1974).
9. R. E. Baier, *J. Geophys. Res.* **77**, 5062 (1972); ———, D. W. Gouppil, S. Perlmutter, R. King, *J. Rech. Atmos.* **8**, 571 (1974).
10. Yu. I. Sorokin, *Int. Rev. Gesamten Hydrobiol.* **56**, 1 (1971).
11. D. W. Menzel and R. F. Vaccaro, *Limnol. Oceanogr.* **9**, 138 (1964).
12. K. M. Johnson and J. McN. Sieburth, *Mar. Chem.*, in press; C. M. Burney and J. McN. Sieburth, *ibid.*, in press.
13. P. D. Allen III, *Dev. Ind. Microbiol.* **14**, 67 (1972).
14. R. D. Hamilton and O. Holm-Hansen, *Limnol. Oceanogr.* **12**, 319 (1967).
15. O. Holm-Hansen, C. J. Lorenzen, R. W. Holmes, J. D. H. Strickland, *J. Cons. Cons. Perm. Int. Explor. Mer* **30**, 3 (1965).
16. K. R. Hinga and J. McN. Sieburth, in preparation.
17. W. D. Garrett, *Limnol. Oceanogr.* **10**, 602 (1965).
18. M. Aubert, J. Aubert and M. Gauthier, *Rev. Int. Oceanogr. Med.* **10**, 137 (1968).
19. J. McN. Sieburth, *Mar. Biol.* **11**, 98 (1971).
20. S. S. Herman and J. A. Mihursky, *Science* **146**, 543 (1964).
21. J. McN. Sieburth, *Microbial Seascapes* (University Park Press, Baltimore, 1975).
22. J. A. Hellebust, *Limnol. Oceanogr.* **10**, 192 (1965); S. Mykkestad, *J. Exp. Mar. Biol. Ecol.* **15**, 261 (1974); ——— and A. Haug, *ibid.* **9**, 125 (1972); ———, B. Larsen, *ibid.*, p. 137; B. Smestad, A. Haug, S. Mykkestad, *Acta Chem. Scand. Ser. B* **28**, 662 (1974); *ibid.* **29**, 337 (1975).
23. K. M. Johnson, C. M. Burney, J. McN. Sieburth, in preparation.
24. A. J. Faller, *Ann. Rev. Ecol. Syst.* **2**, 201 (1971).
25. W. H. Sutcliffe, Jr., E. R. Baylor, D. W. Menzel, *Deep-Sea Res.* **10**, 233 (1963); F. MacIntyre, *J. Rech. Atmos.* **8**, 515 (1974).
26. Supported in part by the Biological Oceanography Program of the National Science Foundation under grants DES74-01537-A01 and OCE74-01537-A02. The technical assistance of J. Hannon, B. McBurney, and S. Sieburth, as well as the help of the crew and officers of R.V. *Trident* during TR-170, is gratefully acknowledged.

7 July 1976; revised 15 September 1976

Reprinted from SCIENCE

24 December 1976, Volume 194, pp. 1415-1418

Copyright© 1976 by the
American Association for the Advancement of Science