



## ICES meets marine historical ecology: placing the history of fish and fisheries in current policy context

Georg H. Engelhard<sup>1,2\*</sup>, Ruth H. Thurstan<sup>3</sup>, Brian R. MacKenzie<sup>4</sup>, Heidi K. Alleway<sup>5</sup>,  
R. Colin A. Bannister<sup>1</sup>, Massimiliano Cardinale<sup>6</sup>, Maurice W. Clarke<sup>7</sup>, Jock C. Currie<sup>8</sup>,  
Tomaso Fortibuoni<sup>9</sup>, Poul Holm<sup>10</sup>, Sidney J. Holt<sup>11</sup>, Carlotta Mazzoldi<sup>12</sup>, John K. Pinnegar<sup>1,2</sup>,  
Saša Raicevich<sup>9,13</sup>, Filip A. M. Volckaert<sup>14</sup>, Emily S. Klein<sup>15</sup>, and Ann-Katrien Lescauwae<sup>16</sup>

<sup>1</sup>Centre for Environment, Fisheries and Aquaculture Science (Cefas), Pakefield Road, Lowestoft NR33 0HT, UK

<sup>2</sup>School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK

<sup>3</sup>University of Queensland, St Lucia, QLD, Australia

<sup>4</sup>Technical University of Denmark (DTU-Aqua), Charlottenlund, Denmark

<sup>5</sup>University of Adelaide, Adelaide, SA 5005, Australia

<sup>6</sup>Swedish University of Agricultural Sciences, Institute of Marine Research, Turistgatan 5, Lysekil, Sweden

<sup>7</sup>Marine Institute, Rinville, Oranmore, Galway, Ireland

<sup>8</sup>South African Environmental Observation Network and Marine Research Institute, Biological Sciences Department, University of Cape Town, Cape Town, South Africa

<sup>9</sup>Italian National Institute for Environmental Protection and Research (ISPRA), Loc. Brondolo, 30015 Chioggia (VE), Italy

<sup>10</sup>Trinity College Dublin, School of Histories and Humanities, Dublin, Ireland

<sup>11</sup>Voc. Palazzetta 68, 06060 Paciano (PG), Italy

<sup>12</sup>Biology Department, University of Padova, Via U. Bassi, 58/B 35121 Padova, Italy

<sup>13</sup>National Research Council (CNR), Institute of Marine Sciences (ISMAR), Largo Fiera della Pesca, 60125 Ancona, Italy

<sup>14</sup>University of Leuven, Ch. Deberiotstraat 32, B-3000 Leuven, Belgium

<sup>15</sup>Princeton University, Ecology and Evolutionary Biology, Guyot Hall, Princeton, NJ 08544, USA

<sup>16</sup>Flanders Marine Institute (VLIZ), Wandelaarkaai 7, 8400 Ostend, Belgium

\*Corresponding author: tel: + 44 1502 527747; fax: + 44 1502 513865; e-mail: [georg.engelhard@cefas.co.uk](mailto:georg.engelhard@cefas.co.uk)

Engelhard, G. H., Thurstan, R. H., MacKenzie, B. R., Alleway, H. K., Bannister, R. C. A., Cardinale, M., Clarke, M. W., Currie, J. C., Fortibuoni, T., Holm, P., Holt, S. J., Mazzoldi, C., Pinnegar, J. K., Raicevich, S., Volckaert, F. A. M., Klein, E. S., and Lescauwae, A-K. ICES meets marine historical ecology: placing the history of fish and fisheries in current policy context. – ICES Journal of Marine Science, doi: 10.1093/icesjms/fsv219.

Received 17 June 2015; revised 26 October 2015; accepted 31 October 2015.

As a discipline, marine historical ecology (MHE) has contributed significantly to our understanding of the past state of the marine environment when levels of human impact were often very different from those today. What is less widely known is that insights from MHE have made headway into being applied within the context of present-day and long-term management and policy. This study draws attention to the applied value of MHE. We demonstrate that a broad knowledge base exists with potential for management application and advice, including the development of baselines and reference levels. Using a number of case studies from around the world, we showcase the value of historical ecology in understanding change and emphasize how it either has already informed management or has the potential to do so soon. We discuss these case studies in a context of the science–policy interface around six themes that are frequently targeted by current marine and maritime policies: climate change, biodiversity conservation, ecosystem structure, habitat integrity, food security, and human governance. We encourage science–policy bodies to actively engage with contributions from MHE, as well-informed policy decisions need to be framed within the context of historical reference points and past resource or ecosystem changes.

**Keywords:** history, management, marine conservation, marine historical ecology, science–policy interface, shifting baseline, sustainable exploitation.

## Introduction

The last two decades have seen the emergence of marine historical ecology (MHE), defined as a distinct discipline within the marine sciences that brings a systematic, long-term perspective to the study of interactions between human society and the seas and oceans (Jackson *et al.*, 2001; Holm *et al.*, 2010; McClenachan *et al.*, 2012; Thurstan, 2013; Schwerdtner Máñez *et al.*, 2014). To an extent inspired by Pauly's (1995) seminal study on the "shifting baselines syndrome", and later the Jackson *et al.* (2001) study on historical changes in marine populations and ecosystems, the field has developed through the collaborative efforts of researchers across disciplines (Figure 1). Although MHE could build on a body of earlier work on long-term changes in marine ecosystems, particularly on exploited fish stocks (e.g. Beverton and Holt, 1957; Cushing, 1980; Pope and Macer, 1996), the field has especially developed through the integration of fisheries science, ecology, history, archaeology, sociology, and economics (Lotze and Milewski, 2002; Holm *et al.*, 2010; Lotze and McClenachan, 2013). As a result, knowledge of the past state of marine animal populations and environments has increased substantially (e.g. Lotze *et al.*, 2006; Bolster, 2008; McClenachan, 2009a; Jackson *et al.*, 2011; Cardinale *et al.*, 2015). This information is crucial to providing baselines for defining sustainable levels of exploitation of marine living resources, understanding the functioning and natural variability of marine environments (Jackson *et al.*, 2001), and planning for resilient human communities that depend on these resources (Lotze *et al.*, 2006; Foley, 2011; FAO, 2014).

Our increased knowledge base on the past marine environment and past stock sizes is widely recognized in mainstream scientific journals (e.g. Jackson *et al.*, 2001; Rosenberg *et al.*, 2005; Eero *et al.*, 2011; Cardinale *et al.*, 2015) and popular books (e.g. Kurlansky, 1997; Butcher, 2004; Roberts, 2007; Bolster, 2008). This is also reflected in applied scientific books (Jackson *et al.*, 2011; Kittinger *et al.*, 2015) and university textbooks (Lotze and McClenachan, 2013; Schwerdtner Máñez and Poulsen, 2016). MHE has contributed to marine science through raised awareness of, often, significant resource depletions and habitat degradation related to human pressures. Such results highlight the global nature of the shifting baseline phenomenon (Pinnegar and

Engelhard, 2008). Despite this worldwide reach, MHE has often been conducted in relative isolation from contemporary ecological studies. Some authors argue that due to the large scale of changes within marine ecosystems, current conservation objectives should not necessarily be underpinned by historical baselines (e.g. Hobday, 2011), and that we should not necessarily expect recovering populations to achieve historical levels. However, the rapidly growing field of restoration ecology counters this view, as do the growing calls among policy-makers for greater knowledge of past states to set more appropriate baselines for sustainable management (Piha and Zampoukas, 2011; ICES, 2014a). For the future of fishing communities, information of the past is deemed to be of critical importance. Science Europe (2013) singled out MHE as one of "12 compelling cases for policy makers" of the relevance of research.


In this study, we argue that, in addition to its contribution to scientific knowledge, MHE has the potential to contribute positively to current and future marine policy and management in a tangible, operational way. In several cases, MHE has already made such a difference (Kittinger *et al.*, 2011; Jennings *et al.*, 2014). To demonstrate the value of MHE to marine policy and management, we outline a number of case studies across four continents—North America, Europe, Africa, and Australia—and cover latitudes from the polar oceans to the tropics, including developed countries with highly managed fisheries and countries lacking in fisheries regulation. For each of the case studies, we outline the ecological problem, show how MHE has contributed knowledge or baselines of past change, and illustrate how it has informed management or has the potential to do so in future.

We discuss each case study within a science–policy framework that involves six themes systematically addressed by marine and maritime policies in many countries. They are also closely aligned with the ecosystem-based approach to fisheries management (Pikitch *et al.*, 2004; Laugen *et al.*, 2014). These themes are: (1) climate change and climate-change adaptation; (2) biodiversity and biodiversity loss; (3) ecosystem structure and function; (4) habitat and seabed integrity; (5) food security including human consumption patterns and exploitation; and (6) human dimensions and governance.

## Policy context and themes

Globally, marine and maritime policies are being developed in different domains and along different lines of action. In general, these policies aim at managing, planning, and developing multiple uses of the marine environment—e.g. fisheries, shipping, tourism, mining, energy production, and conservation—in ways that address the multiple needs and desires of societies and take into account sustainability for both current and future generations (Jennings *et al.*, 2014).

In Europe, the *Integrated Maritime Policy* (IMP, 2007/575/EU; EC, 2007) is a cornerstone of these policies and is based on the principles of ecosystem-based management (see Box 1 for further details about maritime policies developed by the European Commission). In Australia, the *National Strategy for Ecologically Sustainable Development* (ESD) also follows an ecosystem-based approach to fisheries (Fletcher *et al.*, 2002). *Canada's Oceans Strategy: Our Oceans, Our Future* is a policy framework for *Integrated Management* (IM) of estuarine, coastal, and marine environments (Fisheries and Oceans Canada, 2002). In the United States, the *Interagency Ocean Policy Task Force*, established by President Obama in 2009, named ecosystem-based management a top priority for its *National Ocean Policy Implementation Plan* (Interagency Ocean Policy Task



Force, 2010), and the *Magnuson–Stevens Act* provides authority for its implementation through setting goals in regional and state management (US Department of Commerce, 2007). In South Africa, the *Marine Living Resources Act (MLRA)* has governed fisheries since 1998, providing “for the conservation of the marine ecosystem and long-term sustainable utilisation of marine living resources”. Recently, however, preparation of an additional act is under way, with the publication of a *White Paper on National Environmental Management of the Oceans (NEMO; DEA, 2014)*. This white paper includes an ocean governance guiding principle that reads “sustainable use and management of ocean resources and ecosystem services in order to benefit present and future generations” and lists as a strategic priority, the intention to advance sustainable ecosystem-based management in waters under South African jurisdiction. These recent policy developments in countries worldwide require consideration of multiple themes beyond traditional sectoral and single-species management.

**Box 1.** Description of marine and maritime policies developed by the European Commission (EC). Elsewhere, policies have been or are being developed, often along similar lines.

The EC’s cornerstone marine strategy is the *Integrated Maritime Policy* [IMP, COM (2007) 575: EC, 2007], as it integrates a number of transversal policy instruments: in the fields of Blue Growth [COM (2012) 494: EC, 2012] and Blue Innovation [COM (2014) 254: EC, 2014], Marine Knowledge [COM (2010) 461: EC, 2010a], Marine and Maritime Research [COM (2008) 534: EC, 2008b], Maritime Spatial Planning [COM (2013) 133: EC, 2013a], Integrated Maritime Surveillance [COM (2009) 538: EC, 2009a], and Sea Region Strategies in EU waters (Baltic Sea: Helsinki Convention; Mediterranean Sea: Barcelona Convention; Black Sea: Bucharest Convention; Northeast Atlantic: OSPAR Convention). The EU *Marine Strategy Framework Directive* [MSFD, COM (2008) 56: EC, 2008a] addresses the environmental aspects of the IMP, and provides a common framework and goals for EU member states to achieve GES (also abbreviated as GEnS) in national marine waters by 2020. In this framework, GES represents the status where human pressures are considered sustainable since they do not irreversibly hamper biodiversity and the recoverability potential for ecosystem components (Borja *et al.*, 2012). GES needs to be assessed according to a set of indicators and criteria established under the Commission Decision [COM (2010) 477: EC, 2010b] for a range of descriptors including, among others, “Commercial fish and shellfish” (Descriptor 3), “Biodiversity” (Descriptor 1), and “Seabed integrity” (Descriptor 6).


*Marine Knowledge 2020* [COM (2010) 461: EC, 2010a] is an essential component within the IMP as it focuses on centralizing marine data from different sources with the aim to reduce operational costs, provide wider access to quality-checked data, and reduce uncertainty in knowledge of the oceans and the seas as a sound basis for managing future changes. A number of policy instruments also require indicators, targets, time-series, and historical trends to underpin implementation, e.g. the *Common Fisheries Policy* [CFP, Regulation COM (2013) 1380: EC, 2013b], the *Water Framework Directive* [WFD, COM (2000) 60: EC, 2000], the *Habitats Directive* [COM (1992) 43: EEC, 1992], and the *Birds Directive* [COM (2009) 147: EC, 2009b] in marine areas and coastal zones.

The section below provides brief descriptions of the six policy themes we use here to describe the application of our marine historical case studies in supporting and informing management. Note that each of these themes has worldwide relevance, although some may feature more prominently in some parts of the world than in others. Because fisheries management is a relevant topic common to all cases, it is not explicitly included as a policy theme. However, aspects of fisheries resource management that explicitly address one or more of the topics below are highlighted.

- (1) *Climate change.* Global temperatures are rising, glaciers are melting, rainfall patterns are shifting, extreme weather events are becoming more common, and there is concern that ocean acidification will increasingly affect marine life. These trends are likely to continue and potentially worsen (IPCC, 2014). Marine research needs to focus beyond the impacts of climate change on commercially valuable stocks to include the broader consequences for marine ecosystem structure and functioning. Policies are also increasingly addressing climate-change adaptation and planning for resilient human communities.
- (2) *Biodiversity conservation.* Worldwide, biodiversity loss is happening at an unprecedented rate, on land as well as in the marine environment (Dirzo and Raven, 2003; McCauley *et al.*, 2015). In the past 10 years, both policies addressing biodiversity loss and indicators assessing progress towards sustaining diversity have been developed. In 1992, the United Nation’s Convention on Biological Diversity (CBD) marked the international community’s commitment to addressing biodiversity loss. In response, regional-scale and national policies have been put in place in line with the CBD Strategic Action Plan 2011–2020 and the Aichi Biodiversity Targets (CBD, 2011).
- (3) *Ecosystem structure and function.* Researchers and decision-makers are progressively recognizing the need to consider ecosystem structure and functioning, beyond the focus on targeted species, to achieve sustainability of marine resource use. Marine foodwebs play a key role in regulating ecosystem services beyond fisheries, as trophic interactions control energy flow between ecosystem components and contribute to the diversity and structure of marine ecosystems and their responses to change (Worm *et al.*, 2006). Knowledge and effective governance of ecosystem structure and functioning are essential to maintain societal services for future generations (Pikitch *et al.*, 2004; EC, 2008a).
- (4) *Habitat and seabed integrity.* Marine habitats are affected by many anthropogenic pressures. A range of human activities including seabed mining, trawl fisheries, offshore energy platforms, moorings, and dredging for shipping has caused damage to the seabed over large areas (Benn *et al.*, 2010). Nearshore habitats, as nursery and spawning areas often critical for the self-renewal of marine populations, are impacted by land-based sediments and pollutants flowing from rivers and by coastal development. Knowledge of the levels and effects of these impacts is needed to preserve critical seabed structures and healthy habitats for many forms of marine life at multiple life history stages.
- (5) *Food security including human consumption patterns and exploitation.* With a global population projected to reach 9

billion by 2050, consequent human needs will necessitate a doubling of food production and drastic reductions in losses of food and waste, in addition to sustainable management of our natural resources (Foley, 2011; FAO, 2014). These food security demands are a strong incentive for effective policies aimed at sustainable aquaculture and fisheries, while maintaining biodiversity and the provision of other ecosystem services.

- (6) *Human dimensions and governance.* For a substantial part of the global population, livelihoods are partly or wholly dependent on the marine environment, and fish and shellfish are a crucial source of protein (FAO, 2014). In turn, human impacts are likely some of the most important acting on marine ecosystems. Consequently, inclusion of human dimensions and governance is an essential “pillar” within ecosystem-based marine management. The importance of “bottom-up” governance to complement the traditional “top-down” policies is being increasingly recognized (Levin et al., 2009). Along with the reliance of people on marine ecosystems, this means that effective governance involves the “empowerment” of people through knowledge exchange, i.e. effective ways of sharing knowledge to enhance environmental awareness and sustainable policies (Fazey et al., 2012, 2014; e.g. *Marine Knowledge 2020* in EU context, see Box 1), and increased awareness, known as “ocean literacy”.



**Figure 2.** South Australian oyster fishers with their gear (top; State Library of SA, B54098) and a snippet from a late 19th century newspaper article airing concerns about the overexploitation of oyster reefs (Adelaide Observer, 14 January 1899).

## MHE case studies demonstrating applied value in policy context

To illustrate the applied value of MHE, we have made use of a selection of case studies in the context of the ICES Working Group on the History of Fish and Fisheries [ICES WGHIST: see, for example, ICES (2011, 2013)]. The cases are representative of the global approach taken in WGHIST, although the preponderance of examples from the Northeast Atlantic reflects ICES’ strong tradition of work here. In total, 13 case studies were examined, and these are outlined in Table 1; six of these are described below, and the remaining seven are described in Supplementary Material. For each case study, we highlight the ecological problem, show how MHE contributed, and the policy relevance. Where applicable, we point to references to similar case studies elsewhere (Table 1).

### Case 1: Australia—Recovering *Ostrea angasi* oyster reefs lost from coastal ecosystems and society’s memory

#### *The problem*

Oyster reefs sustain important marine ecosystems and economies, but widespread overexploitation has occurred worldwide contributing to changes in ecosystem function, including the loss of water filtering capacity, complex hard substratum, and reef-associated species. In southern Australia, native *O. angasi* reefs have become functionally extinct (Beck et al., 2011), but little attention has been given to this loss in relation to declines in water quality and productivity. Considerable investment is now made in the treatment of poor quality coastal water along these same coastlines, a problem which may have been compounded by the decline of oyster reefs.

#### *Contribution of MHE*

Historical records were used to evaluate the past presence, distribution, and abundance of oyster reefs formed by *O. angasi* in South Australia and their loss due to overexploitation (Alleway and Connell, 2015). Using diaries and correspondence of fisheries inspectors, government reports, legislation, photographs, and maps, a profile of their status during the 1800s and 1900s was created, and a “collective amnesia” regarding their past ecological and economic value identified. Commercial oyster fishing began in the early days of the colony (ca. 1836), and continued for more than a century (Figure 2). Since this time, the interim 180 years and a previous lack of data have contributed to the loss of an important ecological baseline. This resulted in a disparity between the amount of attention historically paid to this habitat and what it is afforded today. Research regained this baseline and clarified the current lack of awareness.

#### *Policy relevance*

Recovering the historical ecological baseline provided the impetus to begin affecting the actual recovery of oyster reefs (*Theme 2 Biodiversity; Theme 4 Habitat and seabed integrity*). Stakeholders, led by the University of Adelaide and South Australian Government, are collaboratively implementing a holistic restoration programme from policy changes through physical restoration (*Theme 6 Human dimensions and governance*). This “living reefs programme” works to improve the lost ecological values of water filtration and habitat complexity by rebuilding oyster reefs as a habitat significantly impacted by human activity.

**Table 1.** Overview of 13 MHE case studies from across the globe, with a brief description of the ecological problem, the contribution of MHE, and the policy relevance. For each case study, we indicate to which policy theme(s) it contributes, and references are provided to similar case studies elsewhere. Notation of policy themes: 1—climate change; 2—biodiversity conservation; 3—ecosystem structure and function; 4—habitat and seabed integrity; 5—food security; 6—human dimensions and governance. See text for further detail on case studies 1–6, and see Supplementary Material for detail on case studies S1–S7.

Case study	Problem	Contribution of historical data	Policy relevance	Policy themes	Similar case studies elsewhere
1. South Australia oysters ( <i>Ostrea angasi</i> )	Loss of oyster reefs from coastal system	Evaluated distribution and abundance of oyster reefs in the past (Alleway <i>et al.</i> , 2014)	Data incorporated into restoration effort <i>Living Reefs Programme</i>	2, 4, 6	Zu Ermgassen <i>et al.</i> (2012) and Grabowski <i>et al.</i> (2012): evaluation of importance lost oyster beds, Chesapeake Bay, United States
2. South Africa linefish	Long-term population declines	Illustrated substantial changes in catch composition and declines in catch rates over time (Penney <i>et al.</i> , 1999; Griffiths, 2000)	Historical evidence revealed the magnitude of catch rate declines and supported strong policy intervention	2, 4, 5, 6	Worm <i>et al.</i> (2006): haddock collapse, and recovery through marine reserve and fishery closure, Georges Bank
3. North Sea herring ( <i>Clupea harengus</i> )	Loss of spatial diversity and stock resilience	Extended time-series on key stock parameters to inform changes in spawning components and spatial diversity (Lescrauwaet, 2013)	Data used to set local management targets, enhancing the potential for recovery of spatial resilience	4, 5, 6	Harma <i>et al.</i> (2012): changes in herring population structure, Celtic Sea; Hutchinson <i>et al.</i> (2003): loss of cod substock, western North Sea
4. North Sea cod ( <i>Gadus morhua</i> )	Slow recovery from overfishing, and unknown drivers of change	Extended fishing and environmental time-series to provide evidence for drivers of cod biomass dynamics (Engelhard <i>et al.</i> , 2014a; ICES, 2014b)	Provides evidence for main drivers of change and information to support management decisions	1, 3, 5	Frank <i>et al.</i> (2005, 2011): similar patterns in climate, plankton, herring, and cod dynamics, Scotian Shelf
5. Baltic Sea fisheries	Unknown drivers of change	Extended biomass time-series demonstrating differing drivers of change over time (Eero <i>et al.</i> , 2007; Eero, 2012)	Contributes to defining ecological and stock reference points, supported <i>Baltic Sea Action Plan</i>	1, 3, 6	Pope and Macer (1996): extended biomass time-series, North Sea gadoids; Toresen and Østvedt (2000): herring, Norwegian Sea
6. Adriatic Sea elasmobranchs	Unknown baseline abundance	Evidenced declines in elasmobranch abundance and diversity (Ferretti <i>et al.</i> , 2013; Raicevich and Fortibuoni, 2013; Barausse <i>et al.</i> , 2014)	Data supported the setting of management and conservation measures	2, 6	Ellis <i>et al.</i> (2005): elasmobranchs, UK waters; Ward-Paige <i>et al.</i> (2010): loss of sharks, Caribbean
S1. Queensland snapper ( <i>Pagrus auratus</i> )	Unknown baseline abundance	Provided catch and effort data (popular media) before official landings records; showed early impact recreational fisheries (Thurstan <i>et al.</i> , 2016)	Potential for data to validate and be included in stock assessment models	3, 6	McClenachan (2009a, b): declines in large trophy fish including groupers, Florida
S2. South Australia fisheries	Use of mean trophic level (MTL) index	Linked declines in MTL to establishment of fisheries for low TL species, not overfishing (Alleway <i>et al.</i> , 2014)	Reconsideration of MTL from fisheries catches as an indicator of marine ecosystem health	3, 5	Pinnegar <i>et al.</i> (2002): Celtic Sea fisheries; Essington <i>et al.</i> (2006): showing "fishing through the foodweb" for global fisheries
S3. Bay of Fundy ecosystem	Loss of resilience and ecosystem structure	Revealed system resilience was driven by local dynamics and highlighted importance of forage fish, notably anadromous species (Klein, 2013)	Provides information to support and prioritize management decisions, including conservation anadromous forage fish	3, 5	Wolff (2000): loss of forage fish, southern North Sea; Lotze and Milewski (2002): declines, Bay of Fundy; Ames and Lichter (2013): cod and alewife, Gulf of Maine
S4. Scotland oysters ( <i>Ostrea edulis</i> )	Loss of oysters from coastal system	Demonstrated fundamental changes in the Firth of Forth benthic ecosystem over the last 200 years (Thurstan <i>et al.</i> , 2013)	Potential to inform EU Habitats Directive and Marine Strategy on GES baselines	3, 4, 6	Riesen and Reise (1982), Lotze (2005): benthos, Wadden Sea; Edgar and Samson (2004): shellfish, Tasmania; Lotze (2010): loss of oysters, Chesapeake Bay
S5. Ireland herring ( <i>C. harengus</i> )	Uncertainty in rebuilding targets	Demonstrated long-term dynamics of the stock and reasons for previous stock collapse (Clarke <i>et al.</i> , 2011)	Demonstrated that rebuilding targets and other reference points are appropriate	1, 5	Toresen and Jakobsson (2002): collapse, management response, and rebuilding in herring, Norwegian Sea

Continued

Table 1. Continued

Case study	Problem	Contribution of historical data	Policy relevance	Policy themes	Similar case studies elsewhere
S6. Kattegat – Skagerrak fisheries	Unknown stock structure	Showed past stock structure and subpopulation richness, revealed different dynamics local substocks (Cardinale et al., 2009, 2011, 2012)	Results applied to redefinition of management units for plaice, turbot, and pollack by ICES	3, 5	Rosenberg et al. (2005): reconstruction of cod stocks on Scotian Shelf; Payne (2010): different dynamics herring substocks
S7. Northeast Atlantic	Evidence of genetic impacts of fishing	Historical data and archived otoliths support evidence of links between fishing and genetic change (fisheries-induced evolution; Diopere, 2014; Laugen et al., 2014; Pinsky and Palumbi, 2014)	Improves understanding and evidence base for management decisions	2, 5, 6	Olsen et al. (2004): fisheries-induced evolution in Newfoundland cod; Engelhard and Heino (2004): herring, Norway; Mollet et al. (2007): North Sea sole

### Case 2: South Africa—Linefish depletion: historical evidence contributes towards drastic management interventions (“state of emergency”)

#### The problem

A comprehensive management framework was introduced in South Africa in 1985 in an attempt to regulate commercial and recreational hook and line fishers (Penney et al., 1989), collectively referred to as the “linefishery”. These measures followed the conviction among fishers and fishery scientists that many line-caught fish were over-exploited. However, it is likely that the management of linefish at the time suffered from a shifting baseline bias (Griffiths, 2000), and convincing scientific evidence required to drive strong policy interventions was lacking. The regulations introduced did not effectively curb rising effort in both recreational and commercial sectors, and catch rates remained low, decreased further, or collapsed (Winker et al., 2012).

#### Contribution of MHE

Van der Elst (1989) and Penney et al. (1989) were among initial efforts to investigate historical (early 1900s) catch records from South Africa’s first research vessel, SS “Pieter Faure” (Figure 3). These records, compared with contemporary data, revealed substantial changes in catch composition and showed how some previously dominant species had almost disappeared from catches during the 20th century. Penney et al. (1999) and Griffiths (2000) expanded this work by collating available historical catch and effort records for the KwaZulu-Natal and Cape Provinces, respectively. Set against modern data, these historical records provided quantitative evidence of the drastic extent to which line-caught species (several of them endemic) were depleted.

#### Policy relevance

Both Penney et al. (1999) and Griffiths (2000) emphasized the inadequacy of linefish regulations in slowing or reversing the overexploitation of nearshore, line-caught fish which was threatening biodiversity and eroding valuable fishery resources (*Theme 2 Biodiversity; Theme 5 Food security and exploitation*). Both studies advocated a reduction in fishing effort and recommended the use of well-placed protected areas to safeguard healthy populations of resident fish (*Theme 4 Habitat and seabed integrity*). Based on this and other evidence, the Minister of Fisheries declared a state of emergency in the fishery in 2000, leading to a drastic curtailment of commercial linefishing effort (*Theme 6 Human dimensions and*

*governance*). Following this drastic reduction resulting from the inclusion of historical data, recent analyses of catch rates have shown evidence of recovery in many linefish stocks (Winker et al., 2012).

### Case 3: Western Europe—North Sea herring: a key area for post-spawning aggregations revealed

#### The problem


The North Sea herring (*Clupea harengus*) stock consists of four “spawning components”, which differ in life history traits and migrate to separate spawning areas at different times of year (Payne, 2010). Of these, Belgian fisheries traditionally targeted the “Downs component” after spawning on the Flemish Banks in the southern North Sea between November and March (Gilson, 1931; Gilis, 1962; Figure 4). These “autumn herring” fisheries disappeared when Downs herring collapsed in the 1950s, preceding the North Sea-wide collapse of the 1970s (Cushing, 1992). Later, North Sea herring recovered, but the Downs component recovered far slower than the rest of the stock, by some two decades (Payne, 2010). However, management objectives for conserving Downs herring were based on recent data only and may have missed information on spatial diversity crucial to stock resilience.

#### Contribution of MHE

Time-series on abundance, age distribution, and mortality of Downs herring post-spawning aggregations were reconstructed along with environmental data from hitherto overlooked sources in national archives (scientific surveys, fisheries statistics, and biological studies; Lescrauwaet, 2013). This extended available time-series on key stock parameters back to the 1930s, including World War II (a period for which few fisheries data exist). The data also demonstrate the historical importance of the Flemish Banks (Sandettie sandbank area) for autumn herring fisheries in a socio-economic context.

#### Policy relevance

This newly obtained historical information is now actively being used as additional evidence to set local management targets and programmes of measures, such as criteria for establishing marine protected areas and marine spatial planning (Special Area of Conservation “The Flemish Banks”; *Theme 4 Habitat and seabed integrity*). These measures enhance the protection of the Downs herring component and therefore preservation of North Sea



**Figure 3.** Drawing of SS “Pieter Faure” trawling in Cape waters in 1898. She was the first research vessel (and steam trawler) in the region and conducted trawl surveys in South African waters between 1898 and 1906. Source: [Department of Agriculture, Cape of Good Hope Parliament](#).

herring overall (*Theme 5 Food security and exploitation*). The recovery of the spatial components of North Sea herring requires local measures in key reproductive habitat, which complement management at the stock level (*Theme 6 Human dimensions and governance*).

#### Case 4: Western Europe—North Sea cod: interactions between fishing, fisheries management, and ecosystem change

##### *The problem*


Since 1960, North Sea cod (*Gadus morhua*) has experienced phases of expansion, overfishing, and recovery. Assessments (e.g. [ICES, 2014b](#)) show that cod recruitment was prolific during 1963–1986 (“gadoid outburst”), when spawning-stock biomass (SSB), fishing mortality (F), and landings reached a peak. After 1987, recruitment halved, but F remained high, so SSB and landings were depleted. Following warnings about stock collapse ([Cook et al., 1997](#)), recovery measures through strict fishery regulations started in 2001, despite socio-political difficulties ([Bannister, 2004](#)). F has now reduced to the 1963 level, and SSB is increasing gradually, but since 1998, recruitment has halved again to a historical minimum, resulting in concern whether North Sea cod can recover ([Horwood et al., 2006](#)) to former levels of biomass (Figure 5).



**Figure 4.** Large catch of herring landed at the quayside of Ostend, Belgium during the 1930s (source: [Gilson, 1931](#)). In these years, very large quantities of post-spawning herring were taken from the Flemish Banks, just off the Belgian coast; knowledge of the former importance of these grounds was highlighted by [Lescrauwaet \(2013\)](#).

##### *Contribution of MHE*

Cod data from 1920 ([Pope and Macer, 1996](#)) show that the high level of recruitment in the gadoid outburst was unprecedented (Figure 5).



**Figure 5.** North Sea cod: long-term changes in (a) landings; (b) spawning-stock biomass (SSB, shown at the same scale); (c) recruitment at age 1; and (d) fishing mortality (F; averaged over ages 2–4; no estimates for WWII), compared against (e) the annual mean sea surface temperature (SST) in the North Sea, and (f) the NAO winter index. Redrawn from Engelhard et al. (2014a). Stock estimates for 1920–1962 from Pope and Macer (1996); data after 1963, and all landings data from ICES (2014b). Due to technical differences between these studies, recruitment estimates pre- and post-1963 not directly comparable (hence different symbols used); it is, however, well established that recruitment levels during the “gadoid outburst” (1960s to early 1980s) were unprecedentedly high.

It coincided with cold sea temperatures and a negative phase of the North Atlantic Oscillation (NAO; ICES, 2014b), and with a herring collapse (1965–1976). The latter is important because herring prey on gadoid eggs and juveniles, while cod larval diet overlaps that of adult herring (Cushing, 1980). After 1987, reduced cod recruitment coincided with changes in abundance, size, and species composition of calanoid copepods (Beaugrand et al., 2003), as well as with warmer seas (O’Brien et al., 2000) and changes in ecological stability, all linked to positive phases of the NAO (Beaugrand et al., 2008) and the Arctic Oscillation (NOAA, 2015). Recovering herring biomass

since 1980 (Payne, 2010) could be an additional factor explaining why cod recruitment is not recovering.

#### Policy relevance

MHE shows that the expansion, overfishing, and recovery of North Sea cod are influenced by climate-related changes in water temperature and plankton (*Theme 1 Climate change*), and possibly by the collapse and recovery of North Sea herring (*Theme 3 Ecosystem function and structure*). Continuing minimal cod recruitment means that despite the marked fall in harvest rate, cod recovery is likely

to be constrained; to ensure sustainable exploitation of this resource, it remains imperative to keep fishing mortality low (*Theme 5 Food security and exploitation*).

### Case 5: Northern Europe—The Baltic Sea Action Plan supported by extended cod and sprat time-series

#### The problem

The Baltic Sea ecosystem changed drastically during the 20th century, owing to increased exploitation, eutrophication, reduction of marine mammal abundance (especially seals), and, more recently, climate change (MacKenzie *et al.*, 2002; HELCOM, 2007; Eero *et al.*, 2011). Many of these changes occurred before systematic biomass time-series for key fish species were recorded, notably cod and sprat (*Sprattus sprattus*). Longer time-series on key fish species are needed to resolve impacts of various drivers for fishery and ecosystem management.

#### Contribution of MHE

Cod and sprat biomass time-series were too short (starting in 1966 and 1974, respectively) to cover the period when the Baltic Sea was less eutrophic, colder, contained higher biomasses of seals (documented predators of cod and sprat; MacKenzie *et al.*, 2011), and experienced more frequent inflows of saline, oxygen-rich water from the North Sea (needed for successful cod reproduction). New, longer stock biomass time-series for both species were constructed from historical fishery reports (Eero *et al.*, 2007; Eero, 2012). Analyses of stock development showed that different forcings (exploitation, seal predation, eutrophication, and climate variability/change) were important during different periods, influencing how multiple foodweb and ecosystem processes affect stock development (Eero *et al.*, 2011; MacKenzie *et al.*, 2011, 2012).

#### Policy relevance

The studies by Eero *et al.* (2011) and MacKenzie *et al.* (2011, 2012) identified major historical drivers of cod and sprat biomass and carrying capacity in the Baltic Sea (*Theme 1 Climate change; Theme 3 Ecosystem structure and function*). These indicators can contribute directly to providing new estimates of stock reference points such as  $B_{MSY}$  (biomass at which maximum sustainable yield is theoretically achieved; Köster *et al.*, 2009). The findings provided ecological background for the development of the *Baltic Sea Action Plan* (HELCOM, 2007) adopted in 2007 as an ambitious multinational programme to restore the good ecological status of the Baltic marine environment by 2021, and regularly updated in ministerial meetings (*Theme 6 Human dimensions and governance*).

### Case 6: Southern Europe—New historical baselines on sharks and rays in the Adriatic Sea influence the Marine Strategy Framework Directive

#### The problem

In the Adriatic Sea, fish have been exploited for thousands of years (Lotze *et al.*, 2011). Over the past century, marked changes in marine communities have been documented, with a dramatic decline in large predators (Fortibuoni *et al.*, 2010). Among exploited fish, elasmobranchs (sharks and rays) showed the strongest decline, owing to direct fishing and bycatch (Ferretti *et al.*, 2008, 2013; Fortibuoni *et al.*, 2010; Barausse *et al.*, 2014). This has resulted in biodiversity loss, potentially influencing the entire community and consequently ecosystem functioning (Ferretti *et al.*, 2013).

#### Contribution of MHE


The combination of multiple historical data sources (naturalists' accounts, documents from archives, libraries, natural history museums, and fish markets; Figure 6) allowed the establishment of more accurate baselines for elasmobranch diversity and abundance in the Adriatic Sea. This is particularly relevant since systematic monitoring surveys only began in the 1980s. According to historical documents, 43 elasmobranch species were previously present in the area (Raicevich and Fortibuoni, 2013). Most of these were common until the 1950s, but declined in the following decades (Figure 6; Ferretti *et al.*, 2013; Barausse *et al.*, 2014) and are currently threatened or even locally extinct (Dulvy *et al.*, 2003). This is the case for common skate (*Dipturus batis*), white skate (*Rostroraja alba*), sandy ray (*Leucoraja circularis*), tope shark (*Galeorhinus galeus*), and angel shark (*Squatina squatina*; Ferretti *et al.*, 2013; Raicevich and Fortibuoni, 2013).

#### Policy relevance

The reconstruction of historical levels of elasmobranch diversity, as well as improved understanding of the population trends (*Theme 2 Biodiversity loss*), supported the setting of management measures set for their conservation in the Adriatic. Importantly, this included the regulation of minimum landing sizes for all commercial elasmobranch species as an environmental target to be met by 2020 within Italy's implementation of the EU *Marine Strategy Framework Directive* (*Theme 6 Human dimensions and governance*).

#### Cases S1–S7: supplementary case studies

Seven additional examples where MHE research contributes to policy are provided by Cases S1–S7 (see Table 1 for an overview, and Supplementary Material for full detail). Taken from Australia, North America, and Europe, these are moreover illustrative of the



**Figure 6.** Long-term declines in landings of (a) skates and rays (Rajidae) and (b) catsharks (Scyliorhinidae) at the fish markets of Venice and nearby Chioggia, Italy, between 1900 and 2000 [redrawn from Raicevich and Fortibuoni (2013)]. The declining trends are illustrative of significant population declines of these and several other vulnerable elasmobranch species in the northern Adriatic Sea (Fortibuoni *et al.*, 2010).



**Figure 7.** Historical sketch, “A Kettle of Fish”, by A. Collingridge illustrating Queensland’s recreational snapper fishery already popular during the 1880s. Data extracted from short, popular newspaper articles describing this fishery formed an important source of information in the [Thurstan et al. \(2016\)](#) study on long-term trends in snapper fisheries. See case study S1 (Supplementary Material). Source: [Trove, National Library of Australia](#).

richness of data sources that MHE has drawn upon, in addition to conventional biological and fisheries datasets. In Australia, for example, 19th century narratives from newspapers and popular magazine articles (Figure 7) were used to provide catch and effort data on Queensland snapper (*Pagrus auratus*), pre-dating official catch statistics by some 70 years (Case S1). In Scotland, maritime charts from the 1850s were used to map historical oyster beds that are long gone, but of relevance in the context of marine spatial planning (Figure 8, Case S4). Other case studies have informed about multidecadal change in the mean trophic level of fisheries catches (Case S2, Australia) and about ecosystem resilience (Case S3, Bay of Fundy, North America), or have pushed stock assessments back in time (Case S5, herring off Ireland; and Case S6, gadoid stocks off Sweden and Denmark). Historical datasets, combined with genetic information taken from decades-old otoliths stored in archives, have supported the evidence base on fisheries-induced evolution, i.e. the selective effects of fishing impacting the genetic composition of fish stocks (Case S7; see Supplementary Material).

## Discussion

The case studies presented illustrate the diversity of ways in which MHE can contribute to topics currently high on the global marine policy agenda. Although far from exhaustive, our examples—stemming from four different continents—bring demonstrable evidence of value to specific marine management and policy instruments in the regional or national context. The six themes used for classification represent broad policy areas relevant to science-policy assessment. They were pragmatically chosen as they broadly correspond to clusters of policy domains (environmental, economic, and social) or to key concepts embedded in more recent policy action (ecosystem-based approach, ocean, and/or resource governance). Arguably, there may be a considerable degree of overlap between the themes as applied to the specific cases, and fisheries policies are a common theme throughout all case studies.

## Policy themes revisited

*Theme 1 Climate change* is a priority for at least three case studies (4, 5, and S5; Table 2). Aspects in which MHE contributes to this policy theme include better characterization of species population parameters (recruitment, biomass, and growth) and their ecology (feeding and distribution patterns) under different climate conditions (Table 2). MHE helps to identify past and current cause-effect relationships and to fine-tune predictive modelling power for assessing potential climate-change impact. The long-term time-series of MHE are particularly relevant considering the temporal scale on which climate change operates and was a major feature in the IPCC Fifth Assessment Report [IPCC, 2014; see also [Fromentin and Powers \(2005\)](#); [Rijnsdorp et al. \(2010\)](#); [Engelhard et al. \(2011, 2014a\)](#); [Brander \(2015\)](#) and [Fortibuoni et al. \(2015\)](#)].

At least four case studies (1, 2, 6, and S7) address past conditions in marine biodiversity or even documented cases of extirpation, relevant to *Theme 2 Biodiversity* (Table 2). When these reference conditions are described in relation to varying levels of exploitation in the past, they may contain key information for current management as they help inform levels of sustainable use of resources. As an example, the “Large Fish Indicator”, which was developed for the North Sea within the OSPAR Convention and used in several ICES Working Groups, was also adopted as an indicator for the EU MSFD ([EC, 2008a](#); [Greenstreet et al., 2010](#); [Engelhard et al., 2015](#)). Its target value of 0.3 at the scale of the North Sea was based on the use of historical trawl survey data, carried out between 1920 and 1982 ([Greenstreet et al., 2010](#)).

Often, changes in biological diversity are also relevant to *Theme 3 Ecosystem structure and function* (case studies 4, 5, S1, S2, S3, S4, and S6; Table 2), especially where changes in community composition were documented ([Essington et al., 2006](#); [Raicevich and Fortibuoni, 2013](#)). *Theme 4 Habitat and seabed integrity* receives support from MHE studies on oyster beds or reefs or other biogenic habitat (case studies 1 and S4), and their preservation or restoration, or from studies highlighting key habitat for specific life stages of fish (case studies 2 and 3; Table 2). Combined, the three policy themes on biodiversity, ecosystem structure and function, and seabed integrity are instrumental to marine conservation and sustainable management throughout the world today, and explicit support MHE can provide for the ecosystem-based approach in marine policies is evidenced from the case studies presented here [see also [Lotze et al. \(2006\)](#), [Kittinger et al. \(2011\)](#), and [Ferretti et al. \(2013\)](#)]. In the EU MSFD, these policy themes, together with the concept of *maximum sustainable yield* in fisheries, are considered an integral part of achieving good environmental status (GES, GEnS) in EU marine waters by 2020 ([EC, 2008a](#)).

Similarly, the adoption of the ecosystem-based approach in fisheries management requires taking these concepts on board and taking a broader historical perspective to patterns of human exploitation of marine living resources in support of *Theme 5 Food security and exploitation* (case studies 2, 3, 4, S2, S3, S5, S6, and S7; Table 2). MHE can help highlighting past stock abundance, placing current management targets in context (e.g. [Rosenberg et al., 2005](#); [Clarke et al., 2011](#)). Importantly, analyses of long-term datasets can lead to better understanding of the relative effects of different drivers on stock dynamics, including fishing pressure, climate change, eutrophication, habitat degradation, and predator-prey interactions; time-series of short duration often do not allow disentangling these factors ([Rijnsdorp and van Leeuwen, 1996](#); [Beaugrand et al., 2008](#); [Eero et al., 2011](#); [Engelhard et al., 2014a, b](#); [Brander, 2015](#)).

**Table 2.** Overview of policy themes, ways through which MHE case studies can contribute to each of these, and their potential relevance for policy instruments.

Policy theme	Focus on case studies	MHE contributions	Examples of policy with targets/indicators
Climate change	4 North Sea cod 5 Baltic fisheries S5 Ireland herring	Long time-series, variability in population parameters, and environmental parameters	UN: IPCC EU: MSFD, UK Climate Change Act 2008 United States: National Ocean Policy Implementation Plan
Biodiversity convention	1 Australian oysters 2 South Africa linefish 6 Adriatic elasmobranchs S7 Genetic impacts fishing	Reference conditions, species conservation status	UN: Aichi Biodiversity Targets (CBD) Australia: Marine Biodiversity (Fisheries Management Act) EU: MSFD, Natura 2000 South Africa: NEMO United States: Marine Mammal Act, Endangered Species Act
Ecosystem structure and function	4 North Sea cod 5 Baltic Sea fisheries S1 Queensland snapper S2 South Australia fisheries S3 Bay of Fundy ecosystem S4 Scotland oysters S6 Kattegat – Skagerrak fish	Reference conditions, cause – effect relationships between pressures and ecosystem components	Australia: Fisheries ESD EU: MSFD, Natura 2000 South Africa: NEMO United States: Magnuson – Stevens Act
Habitat and seabed integrity	1 Australian oysters 2 South Africa linefish 3 North Sea herring S4 Scotland oysters	Reference conditions (including spatial extent) under specified pressure levels in the past	EU: MSFD, Natura 2000, Landscape Convention, Underwater Heritage, Trilateral Wadden Sea Government Policy South Africa: NEMO United States: Magnuson – Stevens Act
Food security and exploitation	2 South Africa linefish 3 North Sea herring 4 North Sea cod S2 South Australia fisheries S3 Bay of Fundy ecosystem S5 Ireland herring S6 Kattegat – Skagerrak fish S7 Genetic impacts fishing	Reference conditions, cause – effect relationships between pressures and ecosystem components, patterns of resource use	Australia: Fisheries ESD EU: CFP South Africa: MLRA, NEMO United States: Magnuson – Stevens Act
Human dimensions and governance	1 Australia oysters 2 South Africa linefish 3 North Sea herring 5 Baltic Sea fisheries 6 Adriatic elasmobranchs S1 Queensland snapper S4 Scotland oysters S7 Genetic impacts fishing	Management plans, management structures, best practices for sustainable and resilient communities	EU: Aarhus Convention, Europe 2020 Strategy, Marine Knowledge, Land and Waterscape Conventions South Africa: MLRA, NEMO United States: Magnuson – Stevens Act

Contributing to *Theme 6 Human dimensions and governance*, eight MHE cases reported here (1, 2, 3, 5, 6, S1, S4, and S7; Table 2) reveal the past importance of resources, techniques, exploitation patterns, and food sources that otherwise are threatened to be lost to humanity, or—if well managed—may open new opportunities for coastal communities in the future. A practical field of application includes the Heritage conventions or policies in coastal and waterscape management (e.g. the Trilateral Wadden Sea Government policy: [TWSC, 2010](#)). Examples of best practice in MHE case studies include the sharing of data and information in open environments and outreach in formats that are relevant to different branches of the community (Clodia database, Adriatic Sea: [Mazzoldi et al., 2014](#); HiFiData, North Sea: [Lescrauwaet et al., 2010](#)).

Although EU policies and case studies from European waters are well represented here, they are demonstrative of similar policy evolutions in other regions around the world (see the section “Policy context and themes”). Most policy targets aim for a state where

human pressures can be considered sustainable if they do not irreversibly damage biodiversity and the potential for recovery of ecosystem components ([Borja et al., 2012](#)). The process of reconstructing historical baselines and trends in support of ecosystem-based management therefore refers to ecosystem components as well as to pressures, requiring a multidisciplinary approach ([Bolster, 2008](#)). In the EU MSFD, three approaches were suggested for setting baselines in relation to GES: (i) baseline as a state at which the anthropogenic effects are considered to be negligible (i.e. pristine state); (ii) baseline set in the past; and (iii) current baseline ([WGGES, 2011](#)). MHE contributes mainly towards informing the first two of these approaches. When contrasted with the current status of environmental parameters, historical data inform the setting of reference points, directions, and targets for management ([Borja et al., 2012](#)). The contribution of MHE has proved to be particularly influential and enlightening when reconstructed baselines are able to characterize pristine states or states where anthropogenic



human activities in the past are not necessarily instrumental for predicting the impact of future activities. Still, even in the latter context, MHE can give guidance in defining thresholds of pressure that may trigger structural and functional degradation in ecosystems and in setting reference levels, especially in data-poor situations.

Notwithstanding the challenges presented by historical data and baselines, they are essential to understand past changes and the impacts that human activities have had on the marine environment. Many countries have absorbed marine historical research outputs in their environmental policies. This uptake has been particularly effective when the research has been supported by published output and reference conditions were clearly linked to pressures exerted on the variables under investigation.

### Implications for management

There is an increasing demand for policies to be more evidence-based. This is particularly true for ocean policies, as awareness regarding the role of oceans in global climate regulation and productivity patterns is growing, and many emerging economic opportunities are ocean-based (European Marine Board, 2013).

This is an opportunity and good timing for the recognition of the added value of MHE. The case studies included here were selected to illustrate the diversity of scale, geographical range, and data products that are relevant to policy development and implementation. We have been far from exhaustive, and there are considerable efforts in many other marine and terrestrial systems globally.

However, the cases demonstrate that MHE has developed beyond proof of concept and regularly provides applied value to science and policy in a variety of ways:

- (i) provides data on historical baselines and historical range of variability of biological and environmental parameters (including previous species distributions), broadening the knowledge base on which to build sustainable policies;
- (ii) enhances our understanding of the effects of anthropogenic disturbances on marine ecosystems, their structure and functioning, facilitating the development of appropriate policy instruments, targets, and indicators;
- (iii) provides excellent case studies that can be used to increase public awareness on the role played by humans in shaping the current status of the seas contributing to the growing of ocean literacy and the direct involvement of citizens in their protection;
- (iv) enhances insights into the long-term responses of marine biota to climate change, helping society to assess its impacts and moreover to adapt to future climate change;
- (v) supports the development of fisheries based on historical conditions of the marine environment and the recovery of sustainable practices from the past; and
- (vi) new algorithms and models are being developed by the MHE community to deal with integrating qualitative and quantitative data, and lessons can be drawn from the innovative approaches used to integrate biological data, environmental data, and data on human uses in support of policy development and assessment.

Recently, MHE has become established as a discipline, with scores of new case studies being published each year. A few are chosen


immediately for their explicit policy relevance, but also for their ability to capture the public imagination. Far too often, however, studies stay within the academic domain even if potentially relevant to policy and fail to reach out more widely in the absence of a dedicated translational research practice (Holm *et al.*, 2010; Fazey *et al.*, 2012). The MHE community should be further encouraged to improve knowledge exchange by highlighting their research in view of current policies; essentially, findings will have to be presented clearly, concisely, and in a defensible way (Raymond *et al.*, 2010). For MHE to achieve its full potential, a number of challenges situated in different domains of science, policy, and governance must also be overcome. While researchers should continue to further the accessibility of their work and facilitate communication and integration as much as possible (Bainbridge, 2014), there is an equally important role for fisheries advisers and marine managers worldwide. In balancing the multiple uses of the marine environment and trading off the current use of marine resources with that of future generations, decision-makers must be made aware of the holistic and long-term view that MHE has to offer as crucial support for informed decisions. It is mostly at the *science-policy interface* that targeted efforts will generate the strong and urgent catalytic effect that is required for this transformation. Considerable progress in this respect has been achieved through strategic action (Fazey *et al.*, 2012).

We, therefore, call upon science-policy bodies to actively engage with the contribution of MHE in policy and management from an early phase. In practice, this could be achieved by making explicit reference to and prioritize the uptake of MHE in strategic marine science policy documents as well as in data and information management plans. Furthermore, it could become an integrated component in capacity-building and in education and training programmes for early-career scientists. Continued and increased efforts for further integration of the MHE science community and policy support will benefit sustainable management and conservation of marine resources as well as the communities that depend on these.

### Looking forward and recommendations

With the advent of the internet and increases in computing power, it has never been easier to locate and explore scores of historical information and make this available for the benefit of the wider community. Previously invisible and thought lost, datasets are being rediscovered in archives and basements around the world, digitized and analysed using modern computational methods, and deployed for applied and important purposes. Building on these developments and as an increasingly recognized discipline, MHE can make a difference to long-term marine management. Often, there is the clear *potential* for outputs to be included in policy; in a subset of cases, outputs have *already* been incorporated into policy; and in a further subset, MHE has already had a (demonstrable) *impact* on marine conservation or resource recovery (illustrated in Figure 9 for our case studies). We suggest the following:

- In cases where MHE has clear potential to contribute towards improved policy, historical ecologists are encouraged to (i) engage with policy- and decision-makers about the management implications of the historical evidence; (ii) inform the public of their research; and (iii) consider packaging and promoting their key findings to achieve real impact and uptake.



**Figure 9.** Policy influence of the documented MHE case studies (numbering follows text), distinguishing cases that have potential to be included in policy (white), cases that have already been taken on-board policy (grey), and cases that have already had a (demonstrable) impact on marine conservation and/or sustainable exploitation (black). Notice, however, that the distinction between these categories is somewhat arbitrary and scale-dependent. For example, in the Bay of Fundy (Case S3, see Supplementary Material), MHE has demonstrated fundamental importance of anadromous forage fish impacted by hundreds of dams, but no policy change has yet occurred; however, recent removal of two dams has already resulted in localized recovery of anadromous fish (Klein, 2013).

- In cases where MHE outputs have already been incorporated in policy, historical ecologists are encouraged to (i) follow up to assess whether the policy change indeed has desired effects; (ii) use the uptake to guide their future research efforts; and (iii) highlight successful application as evidence of the value of their work.
- In cases where MHE has already had a (demonstrable) impact, historical ecologists are encouraged to (i) highlight this as a demonstration of the value of MHE; (ii) continue monitoring the situation to assess whether desired effects are reached and/or maintained; and (iii) use the lessons learned to guide their research efforts towards similar cases where an effective difference to improved governance of marine resources could be made, while at the same time feeling empowered to maintain some “blue skies” research.

We envisage a future where MHE scientists will increasingly “think policy”, so that their work might more readily make a difference, and where policy-makers will increasingly “think MHE” in support of long-term marine conservation and sustainable resource use.

### Supplementary data

Supplementary material is available at the ICES/JMS online version of the manuscript.

### Acknowledgements

This is a position paper synthesizing outputs from the ICES Working Group on the History of Fish and Fisheries (ICES WGHIST), in particular the 2012–2014 meetings, but also building on those in 2009–2011. This study was supported by the EU Framework Programme H2020 [COST Action IS1403 “Oceans Past

Platform” (European Cooperation in Science and Technology)] with additional support from Cefas (Project “Trawling Through Time”). We are grateful for feedback received from Carl O’Brien, Henn Ojaveer, and David Righton, and highly constructive reviews by four anonymous referees.

### References

- Allaway, H. K., and Connell, S. D. 2015. Loss of an ecological baseline through the eradication of oyster reefs from coastal ecosystems and human memory. *Conservation Biology*, 29: 795–804.
- Allaway, H. K., Connell, S. D., Ward, T. M., and Gillanders, B. M. 2014. Historical changes in mean trophic level of southern Australian fisheries. *Marine and Freshwater Research*, 65: 884–893.
- Ames, E. P., and Lichter, J. 2013. Gadids and alewives: Structure within complexity in the Gulf of Maine. *Fisheries Research*, 141: 70–78.
- Bainbridge, I. 2014. How can ecologists make conservation policy more evidence based? Ideas and examples from a devolved perspective. *Journal of Applied Ecology*, 51: 1153–1158.
- Bannister, R. C. A. 2004. The rise and fall of cod (*Gadus morhua*, L.) in the North Sea. In *Management of Shared Fish Stocks*, pp. 316–338. Ed. by A. I. L. Payne, C. M. O’Brien, and S. I. Rogers. Blackwell Publishing, Oxford, UK.
- Barusse, A., Correale, V., Curkovic, A., Finotto, L., Riginella, E., Visentin, E., and Mazzoldi, C. 2014. The role of fisheries and the environment in driving the decline of elasmobranchs in the northern Adriatic Sea. *ICES Journal of Marine Science*, 71: 1593–1603.
- Beaugrand, G., Brander, K. M., Lindley, J. A., Souissi, S., and Reid, P. C. 2003. Plankton effect on cod recruitment in the North Sea. *Nature*, 426: 661–664.
- Beaugrand, G., Edwards, M., Brander, K., Luczak, C., and Ibanez, F. 2008. Causes and projections of abrupt climate-driven ecosystem shifts in the North Atlantic. *Ecology Letters*, 11: 1157–1168.
- Beck, M. W., Brumbaugh, R. D., Airolidi, L., Carranza, A., Coen, L. D., Crawford, C., Defeo, O., et al. 2011. Oyster reefs at risk and

- recommendations for conservation, restoration, and management. *BioScience*, 61: 107–116.
- Benn, A. R., Weaver, P. P., Billet, D. S., van den Hove, S., Murdock, A. P., Doneghan, G. B., and le Bas, T. 2010. Human activities on the deep seafloor in the North East Atlantic: An assessment of spatial extent. *PLoS ONE*, 5: e12730.
- Beverton, R. J. H., and Holt, S. J. 1957. On the dynamics of exploited fish populations. *Fisheries Investigations, Series II*, 19: 1–533.
- Bolster, W. J. 2008. Putting the ocean in Atlantic history: Maritime communities and marine ecology in the Northwest Atlantic, 1500–1800. *American Historical Review*, 113: 19–47.
- Borja, A., Dauer, D. M., and Grémare, A. 2012. The importance of setting targets and reference conditions in assessing marine ecosystems quality. *Ecological Indicators*, 12: 1–7.
- Brander, K. M. 2015. Improving the reliability of fishery predictions under climate change. *Current Climate Change Reports*, 1: 40–48.
- Butcher, J. G. 2004. *The Closing of the Frontier: A History of the Marine Fisheries of Southeast Asia, c. 1850–2000*. ISEAS Publications, Singapore.
- Cardinale, M., Bartolino, V., Llope, M., Maiorano, L., Sköld, M., and Hagberg, J. 2011. Historical spatial baselines in conservation and management of marine resources. *Fish and Fisheries*, 12: 289–298.
- Cardinale, M., Bartolino, V., Svedäng, H., Sundelöf, A., Poulsen, R. T., and Casini, M. 2015. A centurial development of the North Sea fish megafauna as reflected by the historical Swedish longlining fisheries. *Fish and Fisheries*, doi:10.1111/faf.12074.
- Cardinale, M., Linder, M., Bartolino, V., Maiorano, L., and Casini, M. 2009. Conservation value of historical data: Reconstructing stock dynamics of turbot during the last century in the Kattegat-Skagerrak. *Marine Ecology Progress Series*, 386: 197–206.
- Cardinale, M., Svedäng, H., Bartolino, V., Maiorano, L., Casini, M., Hjelm, J., and Linderholm, H. 2012. Spatial and temporal depletion of haddock and pollack during the last century in the Kattegat-Skagerrak. *Journal of Applied Ichthyology*, 28: 200–208.
- CBD (Convention on Biological Diversity). 2011. Conference of the Parties Decision X/2. The Strategic Plan for Biodiversity 2011–2020 and the Aichi Biodiversity Targets. [www.cbd.int/decision/cop/?id=12268](http://www.cbd.int/decision/cop/?id=12268).
- Clarke, M. W., Egan, A., Mariani, S., and Miller, D. 2011. Long Term Trends in the Population Dynamics of Northwestern Ireland Herring Revealed by Data Archaeology. *ICES Document CM 2011/D: 04*.
- Cook, R. M., Sinclair, A., and Stéfansson, G. 1997. Potential collapse of North Sea cod stocks. *Nature*, 385: 521–522.
- Cushing, D. H. 1980. The decline of herring stocks in the North Sea and the gadoid outburst. *Journal du Conseil International pour l'Exploration de la Mer*, 39: 70–81.
- Cushing, D. H. 1992. A short history of the Down stock of herring. *ICES Journal of Marine Science*, 49: 437–443.
- DEA (Department of Environmental Affairs). 2014. White Paper on National Environmental Management of the Ocean (NEMO). Government Notice 426 (29 May 2014). *Government Gazette/Staatskoerant*, 37692: 3–27.
- Department of Agriculture, Cape of Good Hope Parliament. Report of the Government Marine Biologist for the Year 1898.
- Diopere, E. 2014. Genetic signatures of selection in sole—A spatio-temporal approach. PhD thesis, University of Leuven, Belgium. 246 pp.
- Dirzo, R., and Raven, P. H. 2003. Global state of biodiversity loss. *Annual Review of Environment and Resources*, 28: 137–167.
- Duarte, C. M., Borja, A., Carstensen, J., Elliott, M., Krause-Jensen, D., and Marbà, N. 2015. Paradigms in the recovery of estuarine and coastal ecosystems. *Estuaries and Coasts*, doi:10.1007/s12237-013-9750-9.
- Dulvy, N. K., Sadovy, Y., and Reynolds, J. D. 2003. Extinction vulnerability in marine populations. *Fish and Fisheries*, 4: 25–64.
- EC (European Commission). 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 Establishing a Framework for Community Action in the Field of Water Policy. *Official Journal of the European Union*, L327: 1–72.
- EC (European Commission). 2007. An Integrated Maritime Policy for the European Union. Brussels, 10.10.2007, COM(2007) 575 final.
- EC (European Commission). 2008a. Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 Establishing a Framework for Community Action in the Field of Marine Environment Policy (Marine Strategy Framework Directive). *Official Journal of the European Union*, L164: 19–40.
- EC (European Commission). 2008b. A European Strategy for Marine and Maritime Research: A Coherent European Research Area Framework in Support of a Sustainable Use of Oceans and Seas. Brussels, 3.9.2008, COM(2008) 534 final.
- EC (European Commission). 2009a. Towards the Integration of Maritime Surveillance: A Common Information Sharing Environment for the EU Maritime Domain. Brussels, 15.10.2009, COM(2009) 538 final.
- EC (European Commission). 2009b. Directive 2009/147/EC of the European Parliament and of the Council of 30 November 2009 on the Conservation of Wild Birds. *Official Journal of the European Union*, L20: 7–25.
- EC (European Commission). 2010a. Marine Knowledge 2020: Marine Data and Observation for Smart and Sustainable Growth. Brussels, 8.9.2010, COM(2010) 461 final.
- EC (European Commission). 2010b. Commission Decision of 1 September 2010 on Criteria and Methodological Standards on Good Environmental Status of Marine Waters (2010/477/EU). *Official Journal of the European Union*, L232: 14–24.
- EC (European Commission). 2012. Blue Growth: Opportunities for Marine And Maritime Sustainable Growth. Brussels, 13.9.2012, COM(2012) 494 final.
- EC (European Commission). 2013a. Proposal for a Directive of the European Parliament and of the Council Establishing a Framework for Maritime Spatial Planning and Integrated Coastal Management. Brussels, 12.3.2013, COM(2013) 133 final.
- EC (European Commission). 2013b. Regulation (EU) No. 1380/2013 of the European Parliament and of the Council of 11 December 2013 on the Common Fisheries Policy, Amending Council Regulations (EC) No. 1954/2003 and (EC) No. 1224/2009 and Repealing Council Regulations (EC) No. 2371/2002 and (EC) No. 639/2004 and Council Decision 2004/585/EC. *Official Journal of the European Union*, L354: 22–61.
- EC (European Commission). 2014. Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions—Innovation in the Blue Economy: Realising the Potential of Our Seas and Oceans for Jobs and Growth. Brussels, 14 March 2014, COM(2014) 254 final/2.
- Edgar, G. J., and Samson, C. R. 2004. Catastrophic decline in mollusc diversity in eastern Tasmania and its concurrence with shellfish fisheries. *Conservation Biology*, 18: 1579–1588.
- EEC (European Economic Community). 1992. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Official Journal of the European Communities*, L206: 7–50.
- Eero, M. 2012. Reconstructing the dynamics of sprat (*Sprattus sprattus*) in the Baltic Sea in the 20th century. *ICES Journal of Marine Science*, 69: 1010–1018.
- Eero, M., Köster, F. W., Plikshs, M., and Thurov, F. 2007. Eastern Baltic cod (*Gadus morhua callarias*) stock dynamics: Extending the analytical assessment back to the mid-1940s. *ICES Journal of Marine Science*, 64: 1257–1271.
- Eero, M., MacKenzie, B. R., Köster, F. W., and Gislason, H. 2011. Multi-decadal responses of a cod (*Gadus morhua*) population to

- human-induced trophic changes, fishing, and climate. *Ecological Applications*, 21: 214–226.
- Ellis, J. R., Dulvy, N. K., Jennings, S., Parker-Humphreys, M., and Rogers, S. I. 2005. Assessing the status of demersal elasmobranchs in UK waters: A review. *Journal of the Marine Biological Association of the United Kingdom*, 85: 1025–1047.
- Engelhard, G. H., and Heino, M. 2004. Maturity changes in Norwegian spring-spawning herring (*Clupea harengus*): Compensatory or evolutionary responses? *Marine Ecology Progress Series*, 272: 245–256.
- Engelhard, G. H., Lynam, C. P., García-Carreras, B., Dolder, P. J., and Mackinson, S. 2015. Effort reduction and the large fish indicator: Spatial trends reveal positive impacts of recent European fleet reduction schemes. *Environmental Conservation*, 42: 227–236.
- Engelhard, G. H., Peck, M. A., Rindorf, A., Smout, S., van Deurs, M., Raab, K., Andersen, K. H., et al. 2014b. Forage fish, their fisheries, and their predators: Who drives whom? *ICES Journal of Marine Science*, 71: 90–104.
- Engelhard, G. H., Pinnegar, J. K., Kell, L. T., and Rijnsdorp, A. D. 2011. Nine decades of North Sea sole and plaice distribution. *ICES Journal of Marine Science*, 68: 1090–1104.
- Engelhard, G. H., Righton, D. A., and Pinnegar, J. K. 2014a. Climate change and fishing: A century of shifting distribution in North Sea cod. *Global Change Biology*, 20: 2473–2483.
- Essington, T. E., Beaudreau, A. H., and Wiedenmann, J. 2006. Fishing through marine food webs. *Proceedings of the National Academy of Sciences of the United States of America*, 103: 3171–3175.
- European Marine Board. 2013. Navigating the Future IV. Position Paper 20 of the European Marine Board, Ostend, Belgium. ISBN: 9789082093100.
- FAO (Food and Agriculture Organization). 2014. *The State of World Fisheries and Aquaculture: Opportunities and Challenges*. Food and Agriculture Organisation, Rome. 223 pp. ISBN 978-92-5-108276-8.
- Fazey, I., Bunse, L., Msika, J., Pinke, M., Preedy, K., Evely, A. C., Lambert, E., et al. 2014. Evaluating knowledge exchange in interdisciplinary and multi-stakeholder research. *Global Environmental Change*, 25: 204–220.
- Fazey, I., Evely, A. C., Reed, M. S., Stringer, L. C., Kruijssen, J., White, I. C. L., Newsham, A., et al. 2012. Knowledge exchange: A review and research agenda for environmental management. *Environmental Conservation*, 40: 19–36.
- Ferretti, F., Myers, R. A., Serena, F., and Lotze, H. K. 2008. Loss of large predatory sharks from the Mediterranean Sea. *Conservation Biology*, 22: 952–964.
- Ferretti, F., Osio, G. C., Jenkins, C. J., Rosenberg, A. A., and Lotze, H. K. 2013. Long-term change in a meso-predator community in response to prolonged and heterogeneous human impact. *Scientific Reports*, 3: 1057.
- Fisheries and Oceans Canada. 2002. *Canada's Ocean Strategy: Our Oceans, Our Future. Policy and Operational Framework for Integrated Management of Estuarine, Coastal and Marine Environments in Canada*. Fisheries and Oceans Canada, Ottawa, Ontario. 42 pp.
- Fletcher, W. J., Chesson, J., Fisher, M., Sainsbury, K. J., Hundloe, T., Smith, A. D. M., and Whitworth, B. 2002. *National ESD Reporting Framework for Australian Fisheries: The "How To" Guide for Wild Capture Fisheries*. FRDC Project 2000/145, Canberra, Australia. 120 pp.
- Foley, J. A. 2011. Can we feed the world and sustain the planet? *Scientific American*, 305: 60–65.
- Fortibuoni, T., Aldighieri, F., Giovanardi, O., Pranovi, F., and Zucchetto, M. 2015. Climate impact on Italian fisheries (Mediterranean Sea). *Regional Environmental Change*, 15: 931–937.
- Fortibuoni, T., Libralato, S., Raicevich, S., Giovanardi, O., and Solidoro, C. 2010. Coding early naturalists' accounts into long-term fish community changes in the Adriatic Sea (1800–2000). *PLoS ONE*, 5: e15502.
- Frank, K. T., Petrie, B., Choi, J. S., and Leggett, W. C. 2005. Trophic cascades in a formerly cod-dominated ecosystem. *Science*, 308: 1621–1623.
- Frank, K. T., Petrie, B., Fisher, J. A. D., and Leggett, W. C. 2011. Transient dynamics of an altered large marine ecosystem. *Nature*, 477: 86–89.
- Fromentin, J.-M., and Powers, J. E. 2005. Atlantic bluefin tuna: Population dynamics, ecology, fisheries and management. *Fish and Fisheries*, 6: 281–306.
- Gilis, C. 1962. I. De Belgische haringcampagne 1961–62 in de sector Sandtietje. II. De Belgische haringvisserij in de sector Sandtietje van 1930 tot 1960. *Studies van het Zeewetenschappelijk Instituut Oostende = Etudes de l'Institut d'Etudes Maritimes d'Ostende, 1962. Zeewetenschappelijk Instituut = Institut d'Etudes Maritimes: Ostend, Belgium*. 56 pp.
- Gilson, G. 1931. *Recherches sur la biologie du hareng "guai" et sur la pêche entre Ostende et le Cap Gris-Nez en 1930-1931*. Annales de l'Institut d'Etudes Maritimes d'Ostende = Annalen van het Zeewetenschappelijk Instituut, 1. Zeewetenschappelijk Instituut = Institut d'Etudes Maritimes: Oostende, Belgium. 71 pp.
- Grabowski, J. N., Brumbaugh, R. D., Conrad, R. F., Keeler, A. G., Opaluch, J. J., Peterson, C. H., Piehler, M. F., et al. 2012. Economic valuation of ecosystem services provided by oyster reefs. *BioScience*, 62: 900–909.
- Greenstreet, S. P. R., Rogers, S. I., Rice, J. C., Piet, G. J., Guirey, E. J., Fraser, H. M., and Fryer, R. J. 2010. Development of the EcoQO for the North Sea fish community. *ICES Journal of Marine Science*, 68: 1–11.
- Griffiths, M. H. 2000. Long-term trends in catch and effort of commercial linefish off South Africa's Cape Province: Snapshots of the 20th century. *South African Journal of Marine Science*, 22: 81–110.
- Harma, C., Brophy, D., Minto, C., and Clarke, M. 2012. The rise and fall of autumn-spawning herring (*Clupea harengus* L.) in the Celtic Sea between 1959 and 2009: Temporal trends in spawning component diversity. *Fisheries Research*, 121–122: 31–42.
- Harnik, P. G., Lotze, H. K., Anderson, S. C., Finkel, Z. V., Finnegan, S., Lindberg, D. R., Liow, L. H., et al. 2012. Extinctions in ancient and modern seas. *Trends in Ecology and Evolution*, 27: 608–617.
- HELCOM. 2007. *Baltic Sea Action Plan adopted on 15 November 2007 in Krakow, Poland*. <http://helcom.fi/baltic-sea-action-plan>. Helsinki, Finland: Helcom (Helsinki Commission for the Protection of the Baltic Marine Environment). 1 p.
- Hobday, A. J. 2011. Sliding baselines and shuffling species: Implications of climate change for marine conservation. *Marine Ecology*, 32: 392–403.
- Holm, P., Marboe, A. H., Poulsen, B., and MacKenzie, B. R. 2010. Marine animal populations: A new look back in time. *In Life in the World's Oceans: Diversity, Distribution, and Abundance*, pp. 1–24. Ed. by A. D. McIntyre. Wiley-Blackwell.
- Horwood, J., O'Brien, C., and Darby, C. 2006. North Sea cod recovery? *ICES Journal of Marine Science*, 63: 961–968.
- Hutchinson, W. F., van Oosterhout, C., Rogers, S. I., and Carvalho, G. R. 2003. Temporal analysis of archived samples indicates marked genetic changes in declining North Sea cod (*Gadus morhua*). *Proceedings of the Royal Society B*, 270: 2125–2132.
- ICES. 2011. *Report of the Study Group on the History of Fish and Fisheries (WGHIST)*, 24–27 October 2011, Cefas, Lowestoft, UK. ICES Document CM 2011/SSGSUE: 11. 40 pp.
- ICES. 2013. *Interim Report of the Working Group on the History of Fish and Fisheries (WGHIST)*, 7–11 October 2013, Panicle, Italy. ICES Document CM 2013/SSGSUE: 08. 48 pp.
- ICES. 2014a. *ICES Strategic Plan 2014–2018*. 21 pp.
- ICES. 2014b. *ICES Advice 2014. Book 6. Section 6.3.3. ECOREGION: North Sea. STOCK: Cod in Subarea IV (North Sea) and Division VIIId (Eastern Channel) and IIIa West (Skagerrak)*. 19 pp.
- Interagency Ocean Policy Task Force. 2010. *Final Recommendations of the Interagency Ocean Policy Task Force July 19, 2010*. The White House Council on Environmental Quality, Executive Office of the

- President of the United States. [www.whitehouse.gov/files/documents/OPTF\\_FinalRecs.pdf](http://www.whitehouse.gov/files/documents/OPTF_FinalRecs.pdf).
- IPCC (Intergovernmental Panel on Climate Change). 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R. K. Pachauri and L. A. Meyer (Eds)]. IPCC, Geneva, Switzerland. 151 pp.
- IUCN Standards and Petitions Subcommittee. 2014. Guidelines for Using the IUCN Red List Categories and Criteria. Version 11. Prepared by the Standards and Petitions Subcommittee. [www.iucnredlist.org/documents/RedListGuidelines.pdf](http://www.iucnredlist.org/documents/RedListGuidelines.pdf).
- Jackson, J. B. C., Kirby, M. X., Berger, W. H., Bjorndal, K. A., Botsford, L. W., Bourque, B. J., Bradbury, R. H., *et al.* 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293: 629–638.
- Jackson, J. B. C., Sala, E., and Alexander, K. (Eds) 2011. *Shifting Baselines: The Past and the Future of Ocean Fisheries*. Island Press, Washington, DC.
- Jennings, S., Smith, A. D. M., Fulton, E. A., and Smith, D. C. 2014. The ecosystem approach to fisheries: Management at the dynamic interface between biodiversity conservation and sustainable use. *Annals of the New York Academy of Sciences*, 1322: 48–60.
- Kittinger, J. N., McClenachan, L. E., Gedan, K. B., and Blight, L. K. (Eds) 2015. *Marine Historical Ecology in Conservation: Using the Past to Manage for the Future*. University of California Press, Berkeley.
- Kittinger, J. N., Pandolfi, J. M., Blodgett, J. H., Hunt, T. L., Jiang, H., Maly, K., McClenachan, L. E., *et al.* 2011. Historical reconstruction reveals recovery in Hawaiian coral reefs. *PLoS ONE*, 6: e25460.
- Klein, E. 2013. Change in nonlinear dynamics and spatial structure of coastal socio-ecological systems: Bay of Fundy as case study. PhD thesis, University of New Hampshire, Durham, USA. 265 pp.
- Köster, F. W., Vinther, M., MacKenzie, B. R., Eero, M., and Plikshs, M. 2009. Environmental effects on recruitment and implications for biological reference points of eastern Baltic cod (*Gadus morhua*). *Journal of Northwest Atlantic Fishery Science*, 41: 205–220.
- Kurlansky, M. 1997. *Cod: A Biography of the Fish that Changed the World*. Walker Publishing, New York. 304 pp.
- Laugen, A. T., Engelhard, G. H., Whitlock, R., Arlinghaus, R., Dankel, D. J., Dunlop, E. S., Eikeset, A. M., *et al.* 2014. Evolutionary impact assessment: Accounting for evolutionary consequences of fishing in an ecosystem approach to fisheries management. *Fish and Fisheries*, 15: 65–96.
- Lescrauwaet, A-K. 2013. Belgian fisheries: Ten decades, seven seas, forty species. PhD thesis, University of Ghent, Belgium. 257 pp.
- Lescrauwaet, A-K., Debergh, H., Vincx, M., and Mees, J. 2010. Fishing in the past: Historical data on sea fisheries landings in Belgium. *Marine Policy*, 34: 1279–1289.
- Levin, P. S., Fogarty, M. J., Murawski, S. A., and Fluharty, D. 2009. Integrated ecosystem assessments: Developing the scientific basis for ecosystem-based management of the ocean. *PLoS Biology*, 7: e1000014.
- Lotze, H. K. 2005. Radical changes in the Wadden Sea fauna and flora over the last 2000 years. *Helgoland Marine Research*, 59: 71–83.
- Lotze, H. K. 2010. Historical reconstruction of human-induced changes in US estuaries. *Oceanography and Marine Biology: An Annual Review*, 48: 267–338.
- Lotze, H. K., Coll, M., Magera, M. A., Ward-Paige, C., and Airoldi, L. 2011. Recovery of marine animal populations and ecosystems. *Trends in Ecology and Evolution*, 26: 595–605.
- Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., Kidwell, S. M., *et al.* 2006. Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science*, 312: 1806–1809.
- Lotze, H. K., and McClenachan, L. 2013. Marine historical ecology: Informing the future by learning from the past. *In* *Marine Community Ecology and Conservation*, pp. 165–201. Ed. by
- M. D. Bertness, J. F. Bruno, B. R. Silliman, and J. J. Stachowicz. Sinauer Associates, Inc., Sunderland, Connecticut, USA.
- Lotze, H. K., and Milewski, I. 2002. Two Hundred Years of Ecosystem and Food Web Changes in the Quoddy Region, Outer Bay of Fundy. Conservation Council of New Brunswick, Fredericton, New Brunswick, Canada.
- MacKenzie, B. R., Alheit, J., Conley, D. J., Holm, P., and Kinze, C. C. 2002. Ecological hypotheses for a historical reconstruction of upper trophic level biomass in the Baltic Sea and Skagerrak. *Canadian Journal of Fisheries and Aquatic Sciences*, 59: 173–190.
- MacKenzie, B. R., Eero, M., and Ojaveer, H. 2011. Could seals prevent cod recovery in the Baltic Sea? *PLoS ONE*, 6: e18998-1-9.
- MacKenzie, B. R., Meier, H. E. M., Lindegren, M., Neuenfeldt, S., Eero, M., Blenckner, T., Tomczak, M. T., *et al.* 2012. Impact of climate change on fish population dynamics in the Baltic Sea: A dynamical downscaling investigation. *Ambio*, 41: 626–636.
- Mazzoldi, C., Sambo, A., and Riginella, E. 2014. The Clodia database: A long time series of fishery data from the Adriatic Sea. *Scientific Data*, 1: 140018.
- McCaughey, D. J., Pinsky, M. L., Palumbi, S. R., Estes, J. A., Joyce, F. H., and Warner, R. R. 2015. Marine defaunation: Animal loss in the global ocean. *Science*, 347: 1255641.
- McClenachan, L. 2009a. Historical declines of goliath grouper populations in South Florida, USA. *Endangered Species Research*, 7: 175–181.
- McClenachan, L. 2009b. Documenting loss of large trophy fish from the Florida Keys with historical photographs. *Conservation Biology*, 23: 636–643.
- McClenachan, L., Ferretti, F., and Baum, J. K. 2012. From archives to conservation: Why historical data are needed to set baselines for marine animals and ecosystems. *Conservation Letters*, 5: 349–359.
- Mollet, F. M., Kraak, S. B. M., and Rijnsdorp, A. D. 2007. Fisheries-induced evolutionary changes in maturation reaction norms in North Sea sole *Solea solea*. *Marine Ecology Progress Series*, 351: 189–199.
- NOAA. 2015. National Oceanic and Atmospheric Administration/ National Climatic Data Centre/Arctic Oscillation. <http://www.ncdc.noaa.gov/teleconnections/ao/>.
- O'Brien, C. M., Fox, C. J., Planque, B., and Casey, J. 2000. Climate variability and North Sea cod. *Nature*, 404: 142.
- O'Higgins, T., Cooper, P., Roth, E., Newton, A., Farmer, A., Goulding, I. C., and Tett, P. 2014. Temporal constraints on ecosystem management: Definitions and examples from Europe's regional seas. *Ecology and Society*, 19: 46.
- Olsen, E. M., Heino, M., Lilly, G. R., Morgan, M. J., Brattey, J., Ernande, B., and Dieckmann, U. 2004. Maturation trends indicative of rapid evolution preceded the collapse of northern cod. *Nature*, 428: 932–935.
- Pauly, D. 1995. Anecdotes and the shifting baseline syndrome of fisheries. *Trends in Ecology and Evolution*, 10: 430.
- Payne, M. R. 2010. Mind the gaps: A state-space model for analysing the dynamics of North Sea herring spawning components. *ICES Journal of Marine Science*, 67: 1939–1947.
- Penney, A. J., Buxton, C. D., Garratt, P. A., and Smale, M. J. 1989. The commercial marine linefishery. *In* *Oceans of Life off South Africa*, pp. 214–229. Ed. by A. I. L. Payne, and R. J. M. Crawford. Vlaeberg Publishers, Vlaeberg, South Africa.
- Penney, A. J., Mann-Lang, J. B., van der Elst, R. P., and Wilke, C. G. 1999. Long-term trends in catch and effort in the KwaZulu-Natal near-shore linefisheries. *South African Journal of Marine Science*, 21: 51–76.
- Perry, R. I., and Ommer, R. E. 2003. Scale issues in marine ecosystems and human interactions. *Fisheries Oceanography*, 12: 513–522.
- Piha, H., and Zampoukas, N. 2011. Review of Methodological Standards Related to the Marine Strategy Framework Directive Criteria on Good Environmental Status. JRC Scientific and Technical Reports, Joint Research Centre, Ispra, Italy. <http://mcc.jrc.ec.europa.eu/documents/201409161352.pdf>.

- Pikitch, E. K., Santora, C., Babcock, E. A., Bakun, A., Bonfil, R., Conover, D. O., Dayton, P., et al. 2004. Ecosystem-based fishery management. *Science*, 305: 346–347.
- Pinnegar, J. K., and Engelhard, G. H. 2008. The ‘shifting baseline’ phenomenon: A global perspective. *Reviews in Fish Biology and Fisheries*, 18: 1–16.
- Pinnegar, J. K., Jennings, S., O’Brien, C. M., and Polunin, N. V. C. 2002. Long-term changes in the trophic level of the Celtic Sea fish community and fish market price distribution. *Journal of Applied Ecology*, 39: 377–390.
- Pinsky, M. L., and Palumbi, R. S. 2014. Meta-analysis reveals lower genetic diversity in overfished populations. *Molecular Ecology*, 23: 29–39.
- Pope, J. G., and Macer, C. T. 1996. An evaluation of the stock structure of North Sea cod, haddock and whiting since 1920, together with a consideration of the impacts of fisheries and predation effects on their biomass and recruitment. *ICES Journal of Marine Science*, 53: 1157–1169.
- Raicevich, S., and Fortibuoni, T. 2013. Assessing neoextirpations in the Adriatic Sea: An historical ecology approach. *In* CIESM Workshop Monographs, No. 45. Marine Extinctions—Patterns and Processes, pp. 97–111. Ed. by F. Briand. CIESM Publisher, Monaco.
- Raymond, C. M., Fazey, I., Reed, M. S., Stringer, L. C., Robinson, G. M., and Evely, A. C. 2010. Integrating local and scientific knowledge for environmental management. *Journal of Environmental Management*, 91: 1766–1777.
- Riesen, W., and Reise, K. 1982. Macrobenthos of the subtidal Wadden Sea: Revisited after 55 years. *Helgoländer Meeresuntersuchungen*, 35: 409–423.
- Rijnsdorp, A. D., Peck, M. A., Engelhard, G. H., Möllmann, C., and Pinnegar, J. K. 2010. Resolving climate impacts on fish stocks. *ICES Cooperative Research Report No. 301*. 372 pp.
- Rijnsdorp, A. D., and van Leeuwen, P. I. 1996. Changes in growth of North Sea plaice since 1950 in relation to density, eutrophication, beam-trawl effort, and temperature. *ICES Journal of Marine Science*, 53: 1199–1213.
- Roberts, C. M. 2007. *The Unnatural History of the Sea*. Island Press, Washington, DC. 438 pp.
- Rosenberg, A. A., Bolster, W. J., Alexander, K. E., Leavenworth, W. B., Cooper, A. B., and McKenzie, M. G. 2005. The history of ocean resources: Modeling cod biomass using historical records. *Frontiers in Ecology and the Environment*, 3: 84–90.
- Schwerdtner Máñez, K., Holm, P., Blight, L., Coll, M., MacDiarmid, A., Ojaveer, H., Poulsen, B., et al. 2014. The future of the oceans past: Towards a global marine historical research agenda. *PLoS ONE*, 9: e101466.
- Schwerdtner Máñez, K., and Poulsen, B. (Eds). 2016. *Perspectives on Oceans Past—A Handbook on Marine Environmental History*. Springer Publishers, The Netherlands.
- Science Europe. 2013. *Humanities in the Societal Challenges. 12 Compelling Cases for Policymakers*. Science Europe, Brussels, Belgium. 32 pp. [http://wpp.scienceeurope.org/wp-content/uploads/2014/05/SE\\_broch\\_HUM\\_fin\\_web\\_LR.pdf](http://wpp.scienceeurope.org/wp-content/uploads/2014/05/SE_broch_HUM_fin_web_LR.pdf).
- Thurstan, R. H. 2013. Fishing up the past: What historical records can tell us about marine populations today. *Significance*, 10: 30–33.
- Thurstan, R. H., Campbell, A. B., and Pandolfi, J. M. 2016. Nineteenth century narratives reveal historic catch rates for Australian snapper (*Pagrus auratus*). *Fish and Fisheries*, doi:10.1111/faf.12103.
- Thurstan, R. H., Hawkins, J. P., Raby, L., and Roberts, C. M. 2013. Oyster (*Ostrea edulis*) extirpation and ecosystem transformation in the Firth of Forth, Scotland. *Journal of Nature Conservation*, 21: 253–261.
- Toresen, R., and Jakobsson, J. 2002. Exploitation and management of Norwegian spring-spawning herring in the 20th century. *ICES Marine Science Symposia*, 215: 558–571.
- Toresen, R., and Østvedt, O. J. 2000. Variation in abundance of Norwegian spring spawning herring (*Clupea harengus*, Clupeidae) throughout the 20th century and the influence of climatic fluctuations. *Fish and Fisheries*, 1: 231–256.
- Trove, National Library of Australia. 1882, 26 August. “The Far Far North”. *The Sydney Mail and New South Wales Advertiser*. p. 340. <http://www.nla.gov.au/nla.news-article161925043> (viewed 15 October 2015).
- TWSC (Trilateral Wadden Sea Cooperation). 2010. *Trilateral Wadden Sea Cooperation Revised Governance Arrangements*. Adopted on the signing of the 2010 Joint Declaration on the Protection of the Wadden Sea, 17 March 2010. [www.waddensea-secretariat.org](http://www.waddensea-secretariat.org).
- UK Hydrographic Office. Admiralty Charts, Scotland East Coast, Firth of Forth: St Abb’s Head to Edinburgh. Surveyed by Lieutenant F. W. L. Thomas, 1852–1856. [www.ukho.gov.uk](http://www.ukho.gov.uk).
- US Department of Commerce. 2007. *Magnuson-Stevens Fishery Conservation and Management Act (Public Law 94-265)*. As amended by the Magnuson-Stevens Fishery Conservation and Management Reauthorization Act (Public Law 109-479).
- Van der Elst, R. P. 1989. Marine recreational angling in South Africa. *In* *Oceans of Life off Southern Africa*, pp. 164–176. Ed. by A. I. L. Payne, and R. J. M. Crawford. Vlaeberg Publishers, Vlaeberg, South Africa.
- Ward-Paige, C. A., Mora, C., Lotze, H. K., Pattengill-Semmens, C., McClenachan, L., Arias-Castro, E., and Myers, R. A. 2010. Large-scale absence of sharks on reefs in the Greater-Caribbean. *PLoS ONE*, 5: e11968.
- WGGES (Working Group on Good Environmental Status). 2011. *Common Understanding of (Initial) Assessment, Determination of Good Environmental Status (GES) and Establishment of Environmental Targets (Art. 8, 9 & 10 MSFD)*. Version 6–22 November 2011. [www.havochvatten.se/download/18.64f5b3211343cffddb2800018956/1348912772472/common-understanding-final.pdf](http://www.havochvatten.se/download/18.64f5b3211343cffddb2800018956/1348912772472/common-understanding-final.pdf).
- Winker, H., Kerwath, S. E., and Attwood, C. G. 2012. Report on stock assessments of important South African linefish resources. Report of the Linefish Scientific Working Group, LSWG No. 3, 2012. Department of Agriculture, Forestry and Fisheries, Cape Town, South Africa. p. 65.
- Wolff, W. J. 2000. The south-eastern North Sea: Losses of vertebrate fauna during the past 2000 years. *Biological Conservation*, 95: 209–217.
- Worm, B., Barbier, E. B., Beaumont, N., Duffy, J. E., Folke, C., Halpern, B. S., Jackson, J. B. C., et al. 2006. Impacts of biodiversity loss on ocean ecosystem services. *Science*, 314: 787–790.
- Zu Ermgassen, P. S. E., Spalding, M. D., Blake, B., Coen, L. D., Dumbauld, B., Geiger, S., Grabowski, J. H., et al. 2012. Historical ecology with real numbers: Past and present extent and biomass of an imperiled estuarine habitat. *Proceedings of the Royal Society B*, 279: 3393–3400.

Handling editor: Emory Anderson