

tion points out a seasonal variation of male penis length which is not visible when animals are not narcotized. This seasonal variation in penis length leads to a seasonal variation of RPSI ($RPSI = [\text{mean female penis length}/\text{mean male penis length}]^3 \times 100$) what has for consequence to obtain no good correlation between RPS indices of the three species *Nucella lapillus*, *Ocenebra erinacea* and *Hinia reticulata*. Nevertheless VDSI is not different with the two methods and we obtain a good correlation between these indices in the three species. *Nucella lapillus* is the most sensible with TBT while *Hinia reticulata* is the less one. Two sterilized females of the mud-snail *Hinia reticulata* have been discovered but it seems this sterilization is not due to the overgrowth of vas deferens tissues in the vaginal opening. A more complete research will be necessary to observe intermediary stages between stage 4+ (see Stroben *et al.*, 1992) and sterilized females in this species. An abnormality in *Nucella lapillus* in Brest region (West Brittany) has conducted us to discard some animals from the calculation of the indices : aphyallic males and females in polluted areas. This last point was first discovered by Gibbs (1993) in England and a research on this phenomenon is in preparation.

REFERENCES

- GIBBS, P.E., 1993. *J. Mar. Biol. Ass. UK*, 73 : 667-678.
 STROBEN, E., J. OEHLMANN & P. FIORONI, 1992. *Mar. Biol.*, 113 : 625-636.

An experimental analysis of resource allocation in relation to the radiation of life histories within the *Littorina saxatilis* group

R.N. Hughes

School of Biological Sciences, University of Wales, Bangor, Gwynedd, LL57 2UW, Wales, UK

Responses in flesh growth, shell growth and reproductive output to limited food ration were measured in *Littorina arcana*, *L. neglecta* and in *L. saxatilis*. Flesh growth in fertilized *L. saxatilis* was significantly less than in unfertilized snails taken from the same population sample, indicating a trade off in resource allocation to somatic and reproductive functions. Decreasing food ration significantly depressed flesh growth in all species, but had no significant effect on reproductive output. Priority in resource allocation therefore was given to reproduction. The degree of this priority was similar among species, since their responses to food ration were not significantly different. The principle of allocation therefore appears to have been irrelevant in the radiation of life histories in the *L. saxatilis* group. *

Aminotransferases in *Littorina* : is there a functional story ?

S.L. Hull, P.J. Mill and J. Grahame

Department of Pure and Applied Biology, The University of Leeds, Leeds LS2 9JT, UK

The Aat-1 locus of the enzyme aspartate aminotransferase (Aat, E.C. 2.6.1.1) appears to be under some form of differential selection in both *Littorina saxatilis* (Johannesson & Johannesson 1990) and *L. arcana* (Grahame, Mill, Double & Hull 1992). In both species, Aat-1 120 predominates in high shore populations, whereas the Aat-1 100 allele occurs in high frequencies in low shore populations. The aim of this study was to determine if there was any physiological basis for the reported pattern of allozyme variation.