

Community patterns in sandy beaches of Chile: richness, composition, distribution and abundance of species

Patrones comunitarios en playas arenosas de Chile: riqueza, composición, distribución y abundancia de especies

ALEJANDRO BRAZEIRO

Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Alameda 340, Casilla 114-d, Santiago, Chile
E-mail: eabrazai@genes.bio.puc.cl

ABSTRACT

The search for patterns, i.e., statistical arrangement in the data, is a key step to understand the way in which natural systems change in space and time. In order to detect community patterns I analyzed in this study some community traits of forty-five local assemblages of sandy beach invertebrates of Chile. There were identified three main community patterns. (1) Species richness tended to decrease with beach slope (morphodynamic state index), fitting a power model. The variability non-explained by beach slope (about 50%) was independent of grain size and latitude, but a little fraction (9%) was explained by sampling effort. Beach length seemed also to be related with this unexplained variability, but in a non-linear way. (2) The species composition of the local assemblages tended to be a non-random nested subgroup of the species composition of the richer assemblages, i.e., the species composition fitted a nested pattern. (3) The species abundance and distribution were positively correlated, which indicate that wide distributed species tended also to be locally abundant. This is the first time that the "nested species composition pattern" (Nº 2) and the "interspecific abundance-distribution pattern" (Nº 3) are documented for sandy beach fauna. These three patterns were statistically evident at both regional ($\sim 2 \times 10^2$ km) and geographical ($\sim 2 \times 10^3$ km) scales, suggesting that they are scale-independent along the evaluated range. Much work is still needed to explain how sandy beach communities are structured, in the present study only the first step has been taken, i.e., the detection of patterns, leaving open the question about the underline processes and mechanisms.

Key words: abundance-distribution relationship, Chile, morphodynamic, nestedness, pattern, sandy beaches, spatial scale, species composition, species richness.

RESUMEN

La búsqueda de patrones, i.e., arreglos estadísticos en los datos, constituye un paso clave en el proceso de entendimiento de la variabilidad espacial y temporal de los sistemas naturales. Con el objeto de identificar patrones comunitarios, en este trabajo se analizaron algunos atributos comunitarios de cuarenta y cinco ensambles locales de invertebrados de playas de arena de Chile. Se identificaron tres principales patrones comunitarios. (1) La riqueza de especies tendió a decrecer con la pendiente de la playa (índice del estado morfodinámico), ajustando un modelo potencial. La variabilidad no explicada por la pendiente de la playa (alrededor de 50%) fue independiente del tamaño de grano y latitud, pero una pequeña fracción (9%) fue explicada por el esfuerzo de muestreo. La longitud de la playa también pareció estar asociada con esta variación no explicada, aunque en una forma no lineal. (2) La composición de especies de los ensambles locales tendió a ser un subgrupo anidado, no aleatorio, de la composición de especies de los ensambles más diversos, i.e., la composición de especies siguió un patrón de anidamiento. (3) La distribución y abundancia de las especies se correlacionaron positivamente, lo que indicó que las especies ampliamente distribuidas tendieron además a ser las más abundantes localmente. Esta es la primera vez que se documenta en playas de arena la ocurrencia de "anidamiento en la composición de especies" (patrón Nº 2) y de "relación interespecífica positiva entre distribución y abundancia" (patrón Nº 3). Estos tres patrones fueron estadísticamente evidentes, tanto a escala regional ($\sim 2 \times 10^2$ km) como geográfica ($\sim 2 \times 10^3$ km), lo que sugiere que son independientes de la escala en el rango evaluado. Aún hace falta mucho trabajo para explicar como se estructuran las comunidades de playas de arena, en el presente trabajo solo se ha dado el primer paso, i.e., la detección de patrones, permaneciendo abierta la interrogante respecto a los procesos y mecanismos subyacentes.

Palabras clave: relación distribución-abundancia, Chile, morfodinámica, anidamiento, patrón, playas de arena, escala espacial, composición de especies, riqueza de especies.

(Received March 13, 1998; accepted December 21, 1998; managed by Carlos Moreno)

INTRODUCTION

Ecology deals with the variability of natural systems. The standard scientific approach to explain this variability involve the following steps: (1) identify patterns of variation, (2) infer possible processes and mechanisms capable of generating the patterns detected, (3) propose explicative hypothesis, and finally (4) evaluate the hypothesis. Thus, the search for patterns, i.e., statistical arrangement in the data, is a key step to understand the way in which natural systems change in space and time. Furthermore, as was claimed by Shrader-Frechette & McCoy (1993), in the absence of strict empirical laws the foundations of much of population and community ecology will have to be laid in broad statistical generalizations.

The spatial variability of species richness (e.g. Hutchinson 1959, Ricklefs & Schluter 1993) and composition (e.g. Patterson & Atmar, 1986), and the interspecific variability of abundance (e.g. Preston 1948, 1962) and distribution (e.g. Brown 1984, Hanski 1982), are among the principal topics of community ecology. In this paper, these community attributes were investigated in assemblages of intertidal invertebrates of sandy beaches of Chile, in order to detect patterns of variation. The focus was placed on four main topics: (1) the relationship between local species richness and physical factors, (2) the degree of nestedness in the local species composition, (3) the interspecific relationship between abundance and distribution, and (4) the effect of spatial scale on the detection of community patterns.

The theme of species richness of sandy beaches has been treated in several studies carried out around the world (e.g. Defeo et al. 1992, Borzone et al. 1996, McLachlan et al. 1996), and also in Chile (e.g. Jaramillo 1978, Jaramillo & Gonzalez 1991, Jaramillo & McLachlan 1993). In these studies, local species richness has been related to physical factors such as sediment type (e.g. McLachlan et al. 1981, Defeo et al. 1992), intertidal slope (e.g. McLachlan et al. 1981, Defeo et al. 1992) and wave exposure (Eleftheriou & Nicholson 1975). Nowa-

days it is known that beach slope, grain size and exposure are correlated, and they contribute to determine the morphodynamic state of the beach, which explains great part of the local variability of species richness (e.g. McLachlan 1990, McLachlan et al. 1993). However, a considerable amount of variation remains without explanation, justifying therefore the search for new patterns of species richness in sandy beach assemblages. The first aim of this paper was to determine how local species richness is related with beach slope (morphodynamic state index), and then to evaluate how the variation non explained by the slope is affected by others factors, such as sampling effort, latitude, grain size, and beach length.

Nestedness take place when the composition of the poorer assemblages tend to be a non-random nested subgroup of the composition of the richer ones (Patterson & Atmar 1986). Nestedness implies that the distribution of species among localities is predictable, which has important implications from the ecological and management point of view. Although the problem of nestedness has been recently pointed out (eighties), patterns of nestedness have been broadly documented in the literature (e.g. Patterson & Atmar 1986, Wright & Reeves 1992, Cook 1995). In sandy beaches assemblages the local variability of species composition has been poorly studied, and it is to date unknown if nestedness occur in such habitats. The second objective of this study was to assess the degree of nestedness in the local species composition of sandy beach assemblages of intertidal invertebrates.

One of the more widely supported generalizations in ecology is that the local abundance and distribution of species tend to be positively related (e.g. Brown 1984, Gaston 1994, Gaston et al. 1997). This pattern has been documented in a variety of taxa, habitats, geographic regions, and spatial scales (for recent revision see Brown 1984, Gaston 1994, 1996). However, the relationship between the abundance and distribution of species has never been studied in sandy beach macrofauna. Thus, the third aim was to evaluate how the distribution and abun-

dance of sandy beach species of invertebrates are related.

The description of patterns is the description of variability, and its quantification depends on the scale of observation (Levin 1992). Consequently, the concepts of pattern and scale are strongly connected, in the sense that different patterns (aspects of the system) could emerge depending on the scale of observation (e.g. Wiens 1989). Thus, the fourth objective was to analyze how the spatial scale employed affect the detection of community patterns. I examined two spatial scale, named geographical ($\sim 2 \times 10^3$ km) and regional ($\sim 2 \times 10^2$ km).

In summery, the main objective of the present study was to detect community patterns in sandy beach assemblages, in order to encourage future investigations focused on particular problems. To this end, I analyzed published and unpublished data that include biotic and abiotic information from forty-five sandy beaches of Chile.

METHODS

Database

The database includes biotic and abiotic information from forty-five exposed sandy beaches of Chile (between 20° and 40° S). The information was obtained from published data of thirty three beaches, and from surveys carried out on twelve exposed sandy beaches of central Chile (see Appendix 1). Twelve sandy beaches of central Chile (32°61'-34°40' S) were sampled during low tides of April - May of 1997, following a systematic approach. For the propose of this study, a sandy beach locality is defined as a sandy coastal belt of 100 m in length, where the along-beach variability of physical characteristics is assumed to be relatively low. Along ten transects (10 m apart), five sampling units (SU) per transect were obtained with a core of 0.02 m² and 30 cm in depth. The first SU was collected on the drift line (1-3 m above), the second and third were obtained between the drift and effluent lines, and the fourth and fifth were taken under the effluent line. SU's were sieved through 1-mm meshes, and the

retained organisms were stored in 10% formalin.

The published information included in the database was also obtained from snapshot surveys, carried out in summer - early autumn, during low tides. In all studies, biological information was obtained taking sampling cores along several transects, and sieving sand through 1-mm meshes. Surface of sampling cores varied between 0.03 and 0.1 m², number of transects between 1 and 6, and total sampled area between 0.9 and 4.8 m². Retained organisms were stored in 5-10% formalin until sorting in laboratory.

The database contains information classified by beach of the following variables: (1) species richness, (2) species composition, (3) species density, (4) latitude, (5) beach slope (measured following Emery 1961), (6) mean grain size, (7) beach length and (8) sampling effort (total area sampled). The whole database, that includes forty-five localities distributed from Iquique (20° S) to Valdivia (40° S), was used in the geographical scale analysis ($\sim 2 \times 10^3$ km). The reduced database, that includes the twelve localities sampled in central Chile (32°61'-34°40' S), was used in the regional scale analysis ($\sim 2 \times 10^2$ km).

Statistical analyses

Evaluating sampling effort adequacy:

The number of species that could be found in a given area depends on the sampling effort (e.g. Sanders 1969). Thus, before any analysis with species richness (ideally during pilot sampling), it is important to evaluate if the performed sampling effort (sampled area in benthic communities) was adequate to estimate the number of species occurring in a given area.

To assess the sampling effort adequacy, species-area curves were made for each of the twelve localities surveyed in central Chile. In contrast with the approach followed by Jaramillo *et al.* (1995), the curves were generated by cumulating species by transects, and not by sampling stations. I preferred this alternative procedure because sandy beach species show evident zonation

Patterns of species richness:

At regional and geographical scales the species richness of sandy intertidal assemblages decreased significantly with intertidal slope, following in both cases a similar power model (Fig. 2). It is important to point out that beach slope is an index of beach morphodynamic state. Beaches with flat slopes tend to be in the dissipative extremity of the morphodynamic gradient, while steepest beaches tend to be in the reflective margin (Wright & Short 1983). Therefore, the "richness-slope pattern" detected in this study imply a "richness-morphodynamic pattern". The scale independence of this relationship was also suggested by others regional (e.g. Jaramillo & Gonzalez 1991, Defeo et al. 1992, Jaramillo 1994, Borzone et al. 1996) and geographical studies (McLachlan et al. 1981, McLachlan et al. 1996). Thus, it is categorically demonstrated that the richness-slope relationship is the most conspicuous and relevant community pattern in sandy beach habitats.

Although the "richness-slope pattern" has been well documented (see also McLachlan et al. 1993, Jaramillo & McLachlan 1993, Jaramillo et al. 1993), the processes and mechanisms responsible of its generation remain unknown, but some hypotheses have been proposed. McLachlan et al. (1993) suggested that an increasing number of species is excluded from the beach as swash climate (see McArdle & McLachlan 1991, 1992) become harsher as beach slope become steeper. It was also suggested that, because of its type of coastal circulation (i.e., circulation cells), primary productivity is higher in flat beaches (i.e., dissipatives), allowing therefore that many species were capable to sustain viable populations in this type of beach (McLachlan 1990). These hypotheses have not been properly tested yet.

Although beach slope explained a considerable fraction of the variance of species richness at geographical scale (52% see Fig. 2), a significant proportion remains without explanation. This remained variability was analyzed in relation to sam-




Fig. 2: Relationship between species richness and beach slope at both geographic and regional scales. The fitted models are shown in the upper right hand of the graph.

Relación entre la riqueza de especies y la pendiente de la playa a escala geográfica y regional. Los modelos ajustados se muestran en la parte superior derecha del gráfico.

tion of species richness non-explained by beach slope was also independent of beach length and grain size.

The absence of correlation between beach length and species richness residuals was an unexpected result. It is widely known that islands support fewer species than continental areas of similar surface (MacArthur & Wilson 1967). Likewise, several studies on landscape ecology have found that patches located in continuous large areas tend to sustain more species than patches of comparable area located in small fragments of habitat (e.g. Thomas et al. 1992). The "rescue effect" (i.e., immigration reduce extinction rate, Brown & Kodric-Brown 1981) has been evoked as the responsible process. Thus, it had been suspected before the analysis that local assemblages immersed in longer beaches could maintain higher diversity, but simple linear regression analysis did not show any significant trend ($r=0.1$, $P=0.54$) (Fig. 3a). But when the relation was separately analyzed along the range 0.2-2 km of length, species richness residuals tended to increase with beach length ($r=0.39$, $P=0.058$) (Fig. 3b). This suggests that species richness and beach length could be related, but not in an obvious way. This relationship seemed to be non-linear and scale dependent, but is still needed much work to determine how beach length affect species richness.

Another unexpected result was the fate that sandy beach macrofauna did not present the general trend of diversity increase towards low latitudes (e.g. Brown & Gibson 1983, Rosenzweig 1992, 1995). This result agree with observations carried out by McLachlan et al. (1996), who stated that latitude is a negligible factor in determining local species richness in sandy beaches of Australia. Sandy beach macrofauna seems to be another exception of the general latitudinal pattern of species richness (e.g. Santelices 1980).

Nested species composition

The local species composition in insular systems and fragmented habitats (e.g. lakes, patch forest, mountain tips) tend in many cases to be nested, i.e., the species

assemblages of poorer localities tend to be a nested, non-random subgroup of the species assemblages of the richer ones (Patterson & Atmar 1986). Examples of nested species composition have been broadly reported in the recent ecological literature (e.g. Soulé et al. 1992, Patterson 1990, Wrigth & Reeves 1992, Doak & Mills 1994), but there are not published antecedents of this pattern in sandy beach fauna.

The results obtained in the present study demonstrated that sandy beach macrofauna is not an exception. At both geographical and regional scale, the distribution of species among localities tended to be nested. The temperatures of the geographical and regional matrices were 10.97° and 11.09° respectively, and its probabilities, calculated by 500 montecarlo-randomization, were 4.4×10^{-36} and 2.7×10^{-6} respectively (Fig. 4). These results constitute the first evidence of nested species composition in sandy beach macrofauna.

Three major hypotheses capable to explain the pattern of nestedness have been proposed. (1) The local extinction probabilities of the species of the regional pool are nested (e.g. Patterson & Atmar 1986, Patterson 1990, Worthen & Rohde 1996). (2) The species probabilities of local colonization are nested (e.g. Patterson & Atmar 1986, Worthen & Rohde 1996). (3) The environmental tolerances of species are nested (e.g. Patterson & Brown 1991). Given the importance that beach morphodynamic have to species richness (see Fig. 2), the interspecific variability in the tolerance to the environmental conditions generated by different morphodynamic states, could play an important role in the generation of the pattern of nestedness. Therefore, I suggest that the distribution of the sandy beach macrofauna of Chile follow a nested pattern because the species capability to live throughout the morphodynamic gradient is nested.

Interspecific abundance and range of distribution

The range of distribution (i.e., number of localities occupied) and the average lineal

density (regarding only the occupied localities) of the sandy beach macrofauna of Chile were directly correlated, both at regional ($r=0.63$, $P=0.038$) and geographical ($r=0.76$, $P<0.01$) scales (Fig. 5). In the regional-scale analysis all species followed the general trend, but in the geographical analysis, two outliers species were observed, the amphipod *Tylos spinulosus* and the polychaete *Nephtys lamelibranchiata*. When extensive study areas were analyzed, like in the geographical-scale analysis, it is probable that the species pool include species whose geographical ranges overlap only a little part of the study area. In such cases, the distribution would be underestimated, but not the abundance. Thus, abundant but poor distributed species, that generate noise in the relationship, could appear simple by methodological fails.

But in general, the species with the widest range of distribution tended also to be the most abundant locally, regardless the spatial scale of analysis (Fig. 5). This pattern is known as the "interspecific abundance-range relationship" (Gaston et al. 1997), and has been widely documented in the ecological literature (e.g. Hanski 1982, Brown 1984, Gaston & Lawton 1990, Warren & Gaston 1997), becoming one of the most general pattern in ecology (Gaston et al. 1997). The interspecific relationship between abundance and distribution has never been investigated before in sandy beaches, indeed, this the first time the "interspecific abundance-range relationship" is documented in this system.

Following the nomenclature introduced by Hanski (1983), the species of wide distribution and high abundance are named "core species", whereas the species of restricted distribution and low abundance are known as "satellite species". In both regional and geographical-scale analyses there were detected the same four "core species", being them *Emerita analoga*, *Orchestoidea tuberculata*, *Excirrolana hirsuticauda* and *E. braziliensis* (Fig. 5). This indicates that the consistence between the regional and geographical pattern was not only in the general trend (i.e. positive relationship between abundance and distri-

bution), but also in the identity of the common and rare species.

Many hypotheses have been proposed to explain the "interspecific abundance-range relationship". These involve different mechanisms, such as sampling artifact, geographical distribution, niche breadth, resource availability, metapopulation dynamics, density dependent habitat selection, and vital rates (see Gaston 1994 and Gaston et al. 1997 for revisions). Although the most plausible mechanisms have already been identified, none of the hypotheses have irrefutable evidences (e.g. Gaston




Fig. 5: Interspecific relation between lineal density and range of distribution in sandy beach macrofauna of Chile, at both geographical (upper) and regional (below) scales. The species indicated by an asterisk (1: *Tylos spinulosus* and 2: *Nephtys lamelibranchiata*) were considered outliers and were not included in the lineal regression analysis at geographical scale. The linear models fitted are shown in the upper right hand of the graphs. Ehir: *Excirrolana hirsuticauda*, Ebr: *E. braziliensis*, Otu: *Orchestoidea tuberculata* and Ean: *Emerita analoga*.

Relación interespecífica entre la densidad lineal y el rango de distribución en la macrofauna de playas de arena de Chile, a escala geográfica (arriba) y regional (abajo). Las especies indicadas con un asterisco (1: *Tylos spinulosus* y 2: *Nephtys lamelibranchiata*) fueron consideradas "outliers", y no fueron incluidas en el análisis de regresión lineal a escala geográfica. Los modelos lineales ajustados se muestran en la parte superior derecha de los gráficos.

- GASTON KJ & JH LAWTON (1990) Effects of scale and habitat on the relationship between regional distribution and local abundance. *Oikos* 58: 329-335.
- GASTON KJ, TM BLACKBURN & H LAWTON (1997) Interspecific abundance-range size relationship: an appraisal of mechanism. *Journal of Ecology* 66:579-601.
- HANSKI I (1982) Dynamics of regional distribution: the core and satellite species hypothesis. *Oikos* 38: 210-221.
- HUTCHINSON GE (1959) Homage to Santa Rosalia, or why are there so many kinds of animals? *The American Naturalist* 93:145-159.
- JARAMILLO E (1978) Zonación y estructura de la comunidad macrofaunística en playas de arena del Sur de Chile (Mehuín, Valdivia). *Studies of Neotropical Fauna and Environment* 13:71-92.
- JARAMILLO E (1987) Sandy beach macroinfauna from the Chilean coast: zonation patterns and zoogeography. *Vie Milieu* 37:165-174.
- JARAMILLO E (1994) Patterns of species richness in sandy beaches of south America. *South African Journal of Zoology* 29:227-234.
- JARAMILLO E & M GONZALEZ (1991) Community structure and zonation of the macroinfauna along a dissipative-reflective range of beach category in south Chile. *Studies of Neotropical Fauna and Environment* 26 193-212.
- JARAMILLO E & A McLACHLAN (1993) Community and population response of the macroinfauna to physical factors over a range of exposed sandy beaches in south-central Chile. *Estuarine Coastal and Shelf Science* 37:615-624.
- JARAMILLO E, A McLACHLAN & P COETZEE (1993) Intertidal zonation patterns of macroinfauna over a range of exposed sandy beaches in south-central Chile. *Marine Ecology Progress Series* 101:105-118.
- JARAMILLO E, A McLACHLAN & J DUGAN (1995) Total sample area and estimate of species richness in exposed sandy beaches. *Marine Ecology Progress Series* 101:105-108.
- LEVIN SA (1992) The problem of pattern and scale in ecology. *Ecology* 73:1943-1967.
- MACARTHUR RH & EO WILSON (1967) *Island biogeography*. Princeton University Press, Princeton. 203 pp.
- McARDLE SB & A McLACHLAN (1991) Dynamics of the swash zone and effluent line on sandy beaches. *Marine Ecology Progress Series* 76:91-99.
- McARDLE SB & A McLACHLAN (1992) Sand beach ecology: swash features relevant to the macrofauna. *Journal of Coastal Research* 8:398-407.
- McLACHLAN A (1990) Dissipative beaches and macrofauna communities on exposed intertidal sands. *Journal of Coastal Research* 6:57-71.
- McLACHLAN A, T WOOLDRIDGE & AH DYE (1981) The ecology of two sandy beaches in southern Africa. *South African Journal of Zoology* 16:219-231.
- McLACHLAN A, E JARAMILLO, TE DONN & F WESSELS (1993) Sandy beach macrofauna communities and their control by the physical environment: a geographical comparison. *Journal of Coastal Research* 15:27-38.
- McLACHLAN A & E JARAMILLO (1995) Zonation on sandy beaches. *Oceanography and Marine Biology Annual Review* 33:305-335.
- McLACHLAN A, A RUYCK & N HACKING (1996) Community structure on sandy beaches: patterns of richness and zonation in relation to tide range and latitude. *Revista Chilena de Historia Natural* 69:451-467.
- PATTERSON BD (1990) On the temporal development of nested subset patterns of species composition. *Oikos* 59:330-342.
- PATTERSON BD & W ATMAR (1986) Nested subset and the structure of insular mammalian fauna and archipelagos. *Biological Journal of Linnean Society* 28:65-82.
- PATTERSON BD & JH BROWN (1991) Regionally nested patterns of species composition in granivorous rodent assemblages. *Journal of Biogeography* 18:395-402.
- PRESTON FW (1948) The commonness, and rarity, of species. *Ecology* 29:254-283.
- PRESTON FW (1962) The canonical distribution of commonness and rarity: part I. *Ecology* 43:185-215.
- ROSENZWEIG ML (1992) Species diversity gradients: we know more or less than we thought. *Journal of Mammalogy* 73:715-730.
- ROSENZWEIG ML (1995) *Species diversity in space and time*. Cambridge University Press, Cambridge. 436pp.
- RICKLEFS RE & D SCHLUTER (1993) Species diversity: regional and historical influences. In: Ricklefs RE & D Schluter (eds) *Species diversity in ecological communities. Historical and geographical perspectives*: 350-364. The University of Chicago Press, Chicago.
- SANCHEZ M, JC CASTILLA & O MENA (1982) Variaciones verano - invierno de la macrofauna de arena en playa Morrillos (Norte Chico, Chile). *Studies on Neotropical Fauna and Environment* 17:31-49.
- SANDERS HL (1969) Benthic marine diversity and the stability-time hypothesis. *Brookhaven Symposium of Biology* 22:71-81.
- SANTELICES B (1980) Phytogeographic characterization of the temperate coast of Pacific South America. *Phycologia* 19:1-12.
- SHORT AD (1996) The role of wave height, period, slope, tide range and embaymentisation in beach classifications: a review. *Revista Chilena de Historia Natural* 69:598-604.
- SHORT AD & LD WRIGTH (1983) Physical variability of sandy beaches. In: McLachlan A & T Erasmus (eds) *Sandy beaches as ecosystems*: 133-144. The Hague, The Netherlands, Junk.
- SOULE ME, AC ALBERTS & DT BOLGER (1992) The effects of habitat fragmentation on chaparral plants and vertebrates. *Oikos* 63:39-47.
- SRADER-FRECHETTE KS & ED McCOY (1933) *Methods in Ecology: Strategies for Conservation*. Cambridge University Press, Cambridge.
- THOMAS CD, JA THOMAS & MS WARREN (1992) Distribution of occupied and vacant butterfly habitats in fragmented habitats. *Oecologia* 92: 563-567.
- WARREN PH & JK GASTON (1997) Interspecific abundance occupancy relationships: a test of mechanisms using microcosms. *Journal of Animal Ecology* 66:730-742.
- WIENS JA (1989) Spatial scaling in ecology. *Functional Ecology* 3:385-397.
- WORTHEN B & K ROHDE (1996) Nested subset analyses of colonization-dominated communities: metazoan ectoparasites of marine fishes. *Oikos* 75:471-478.
- WRIGTH DH & JH REEVES (1992) On the meaning and measurement of nestedness of species assemblages. *Oecologia* 92:416-428.

APPENDIX 2

The geographical pool of sandy beach species of Chile

El ensamble geográfico de especies de playas arenosas de Chile

Code	Species	Code	Species
1	<i>Ampelisca</i> sp (amphipod)	15	<i>Chaetilia paucidens</i> (isopod)
2	<i>Bathyporeiapus magallanicus</i> (amphipod)	16	<i>Excirolana braziliensis</i> (isopod)
3	<i>Hurape</i> sp (amphipod)	17	<i>Excirolana hirsuticuada</i> (isopod)
4	<i>Orchestoidea tuberculata</i> (amphipod)	18	<i>Excirolana monodi</i> (isopod)
5	<i>Phoxocephalopsis mehuinensis</i> (amphipod)	19	<i>Macrochiridotea aff lilianae</i> (isopod)
6	<i>Tryphosella schelenbergi</i> (amphipod)	20	<i>Macrochiridotea mehuinensis</i> (isopod)
7	<i>Tylos spinulosus</i> (amphipod)	21	<i>Macrochiridotea setifer</i> (isopod)
8	<i>Emerita analoga</i> (decapod)	22	<i>Euzonus heterocirrus</i> (polychaete)
9	<i>Lepidopa chilensis</i> (decapod)	23	<i>Hemipodus</i> sp (polychaete)
10	<i>Mesodesma donacium</i> (bivalve)	24	<i>Nephtys impressa</i> (polychaete)
11	<i>Bellia picta</i> (decapod)	25	<i>Nephtys monilibranchiata</i> (polychaete)
12	<i>Phalerisidia maculata</i> (insect)	26	<i>Polydora</i> sp (polychaete)
13	<i>Candancia</i> sp (copepod)	27	<i>Scoletepis chilensis</i> (polychaete)
14	<i>Ocypode gaudichaudii</i> (decapod)		