



Pergamon

Marine Pollution Bulletin, Vol. 36, No. 3, pp. 241–247, 1998
 Crown Copyright © 1998 Published by Elsevier Science Ltd
 All rights reserved. Printed in Great Britain
 0025-326X/98 \$19.00+0.00

PII: 0025-326X(97)00015-0

Organotin Compounds in Liver Tissue of Harbour Porpoises (*Phocoena phocoena*) and Grey Seals (*Halichoerus grypus*) from the Coastal Waters of England and Wales

R. J. LAW*†, S. J. BLAKE*, B. R. JONES* and E. ROGAN†

*Centre for Environment, Fisheries and Aquaculture Science, CEFAS Burnham Laboratory, Remembrance Avenue, Burnham on Crouch, CM0 8HA, UK

†Department of Zoology and Animal Ecology, University of Cork, Lee Maltings, Cork, Ireland

Common (harbour) porpoises (*Phocoena phocoena*) are small cetaceans which are widely distributed throughout the temperate and sub-Arctic waters in the Northern Hemisphere. They are essentially coastal animals, generally bottom-feeding on small schooling fish such as herring or anchovy, although they are also sighted offshore in deep-water areas (Martin, 1990; Northridge *et al.*, 1995). They are under threat in a number of areas around the North Atlantic as a result of the scale of incidental capture in fishing gear (Clausen and Andersen, 1988; Hammond *et al.*, 1995; International Whaling Commission, 1996; NMFS, 1992; Read and Gaskin, 1990; Trippel *et al.*, 1996). Grey seal (*Halichoerus grypus*) populations occur in the Baltic Sea and the NE and NW Atlantic Ocean, with the greater part of the NE Atlantic population being found around the British Isles (Bonner, 1989). Grey seals feed on a wide variety of fish, with some invertebrates including cephalopods, sandeels (*Ammodytidae*) and gadoid fish being of prime importance around the UK (Hammond and Prime, 1990; Hammond *et al.*, 1994a; Hammond *et al.*, 1994b; Prime and Hammond, 1990). The major part of the NE Atlantic population is found around Scotland and NE England (Bonner, 1989), and is currently increasing in numbers (Hiby *et al.*, 1996).

Tributyltin (TBT) compounds have been used extensively as the active component of antifouling paints for ships and marine structures since the 1960s. Their use on small craft <25 m in length has been restricted in many countries since the late 1980s as TBT caused shell thickening and spat failure in oysters, and imposex in gastropods (Alzieu, 1991; Bryan *et al.*, 1987; Waite *et al.*, 1991). At that time the open sea was not considered to be at risk, and the use of TBT has

continued on large vessels. However, a recent study in the southern North Sea has demonstrated the presence of imposex in whelks (*Buccinum undatum*) as a result of TBT contamination from shipping (Ten Hallers-Tjabbes *et al.*, 1994), causing its local extinction within the Dutch Wadden Sea (Cadée *et al.*, 1995). The imposition of wider controls on the use of TBT-based antifouling on shipping is currently under discussion within the International Maritime Organisation. Following release to the environment, TBT degrades by dealkylation to form dibutyl- and monobutyltin (DBT and MBT). These compounds are also used industrially as stabilizers for chlorinated polymers, and as catalysts for silicones and polyurethane foams (Evans and Karpel, 1985). Triphenyl tin compounds (as the acetate and hydroxide) are used in Great Britain in the control of potato blight, with 52 t of fentin being applied in 1994 (Garthwaite *et al.*, 1995).

Many studies of organotin contamination have been conducted, but to date few have included determinations in the tissues of marine mammals. As top predators, cetaceans and pinnipeds accumulate high concentrations of lipophilic and persistent organic compounds through their diet, due to a low metabolic capacity for degradation both in the mammals themselves and components of their food-chain (Suzuki *et al.*, 1992; Tanabe and Tatsukawa, 1992). TBT is bioaccumulative, with an octanol–water partition coefficient in the range 10^3 – 10^4 (Laughlin and Lindén, 1987) and the ability of cetaceans to metabolise TBT is thought to be low as a result of the limited activity of some cytochrome P450 enzymes within these animals (Kannan *et al.*, 1996; Tanabe *et al.*, 1988). This suggests that butyltin compounds can accumulate in marine mammals, and this has been demonstrated in bottlenose dolphins from the Italian coast (Kannan *et al.*, 1996) and the USA (Kannan *et al.*, 1997a), porpoises from the Baltic Sea (Kannan and Falandysz, 1997), and seals, Steller sea lions, and a range of cetacean species from Japan and the north Pacific Ocean (Iwata *et al.*, 1994; Iwata *et al.*, 1995; Kim *et al.*, 1996). No comparable information has yet been presented for the United Kingdom. In this study, concentrations of MBT, DBT and TBT have been determined in the livers of porpoises and grey seals stranded or bycaught around England and Wales, in order to establish the current levels of contamination by these compounds, and to assess the need for further work.

Samples of liver tissue were taken from 29 porpoises and five seals bycaught or stranded around England and Wales from 1992 to 1996 according to previously established protocols (Kuiken and Baker, 1993; Law, 1994), with only those carcasses considered freshly dead or slightly decomposed being sampled. The tissues were stored in glass at -20°C prior to analysis. Previous studies have shown no measurable breakdown of TBT to have occurred after 6 months storage at this temperature. After thawing and homogenization, the

†Corresponding author.

samples were analysed for butyltin compounds using a method established for use in a long-term monitoring programme operated by the Burnham on Crouch laboratory and validated during that study (Waldock *et al.*, 1989; Waldock and Waite, 1994). Briefly, sub-samples were extracted by shaking with 0.1% sodium hydroxide and methanol after the addition of a known quantity of an internal (surrogate) standard (tripropyltin chloride) used for quantification. The organotin compounds were then back-extracted into *n*-hexane, and converted to their respective hydrides with sodium borohydride. They were analysed using capillary gas chromatography with flame-photometric detection at 610 nm. The recovery of TBT relative to the internal standard was 99.6% ($n = 13$, s.d. = 5.6%) and the limit of detection was around $10 \mu\text{g kg}^{-1}$ wet weight. The performance of the method was evaluated by routine analysis of an in-house quality control sample (a wet oyster tissue derived from our own studies) within each sample batch. Over the last 2 years this has been analysed 90 times for DBT (mean value $34 \pm 13 \mu\text{g kg}^{-1}$) and TBT (mean value $226 \pm 110 \mu\text{g kg}^{-1}$); MBT is not routinely determined. In a recent European certification exercise for a reference mussel sample (BCR477) our data were: MBT, 2.03 mg kg^{-1} ; DBT, 1.35 mg kg^{-1} ; TBT, 2.16 mg kg^{-1} ; compared with the assigned values of 1.96, 1.58 and 2.14 mg kg^{-1} , respectively. Figure 1 shows the distribution of sampling sites around England and Wales, and gas chromatograms of a standard solution and sample extract are given as Fig. 2.

The results of analyses are given in Table 1, and other data reported in the literature are listed in Table 2. All data are given as micrograms per kilogram on a wet weight basis. Triphenyltin is also detected using our methodology (although with less sensitivity than for butyltins), but none of these samples contained detectable concentrations (limit of detection about $100 \mu\text{g kg}^{-1}$ wet weight).

Although the use of TBT-based antifouling paints on deep-sea vessels results in some direct inputs offshore, the major inputs (including those from shipping) occur in coastal areas. Industrial inputs of MBT and DBT also occur to coastal areas, and concentrations of these compounds will be supplemented by breakdown of TBT after its release. From the data on TBT concentrations in blubber (Iwata *et al.*, 1994, 1995; Table 2), it is clear that organotin residues can be found at low levels in the blubber of dolphins, porpoises, and a beaked whale from offshore locations in the Pacific Ocean, the Bay of Bengal and the Japan Sea, although they were not detectable in a minke whale from the Antarctic Ocean. In the Southern Hemisphere, minke whales feed mainly upon euphausiids in temperate and polar waters (Martin, 1990), and are apparently not exposed to significant concentrations of organotin compounds in their diet. The highest concentrations of organotins were, however, found in finless porpoises inhabiting the coastal waters and rivers of Japan, believed to feed mainly upon sandlance (*Ammodytes*) at the seabed (Martin, 1990). Comparable concentrations have also been found in the blubber of bottlenose




Fig. 1 Map of England and Wales, showing the stranding locations of porpoises (circles) and grey seals (triangles).

dolphins (*Tursiops truncatus*) from the coasts of Italy and the USA (Kannan *et al.*, 1996, 1997a; Table 2), which feed on a wide variety of fish, crustaceans and squid (Evans, 1987).

Organotin compounds have been detected in all the tissues and organs of the finless porpoise (Iwata *et al.*, 1995), with the highest concentrations occurring in the liver. Muscle contained the major burden of TBT (71%), whilst MBT and DBT were retained predominantly in the blubber and liver (41% and 50%, respectively; Table 2 in Iwata *et al.*, 1995). In overall terms, muscle, blubber, and liver retained almost 90% of the total organotin body burden. In Steller sea lions, 40% of the total organotin burden was in the liver, but 26% accumulated in hair, implying that these animals (and presumably also seals) may eliminate a considerable proportion of ingested organotins through the shedding of fur (Kim *et al.*, 1996). Organotin compounds were also found in the brain of finless porpoises (Iwata *et al.*, 1995) and bottlenose dolphins (Kannan *et al.*, 1997a), indicating that these compounds can pass through the blood-brain barrier, possibly by binding to sulphhydryl

residues as for methylmercury (Kannan and Falandysz, 1997).

Concentrations of total butyltins (Σ BTs) in porpoise liver from the present study (Table 1) were within the range 22–640 $\mu\text{g kg}^{-1}$ wet weight, lower than those reported in finless porpoises from Japan (1120–10200 $\mu\text{g kg}^{-1}$ wet weight; Table 2) and bottlenose dolphins from the Adriatic Sea (1200 and 2200 $\mu\text{g kg}^{-1}$ wet weight) or the USA (110–11340 $\mu\text{g kg}^{-1}$ wet weight). On average, only about 20% of the total butyltin in these individuals was present as TBT, with DBT the major component. The livers from two neonatal harbour porpoises from the Baltic Sea yielded concentrations at the lower end of the range of the samples analysed here, at 18 and 27 $\mu\text{g kg}^{-1}$ wet weight (Kannan and Falandysz, 1997), and three pygmy sperm whales from the USA exhibited similar Σ BT levels to those in UK porpoises (350–410 $\mu\text{g kg}^{-1}$ wet weight). Concentrations of total butyltins in the liver tissues of grey seals were lower than those seen in the porpoises (Table 1; ND to 22 $\mu\text{g kg}^{-1}$ wet weight), similar to that reported for a Larga seal from Hokkaido, Japan




Fig. 2 Gas chromatograms (flame-photometric detection) of (a) a standard solution of organotin compounds, and (b) the extract of liver from a harbour porpoise (SW1992/202; Table 1). The capillary column used was a 25 m \times 0.32 mm ID fused silica column coated with 5% phenylmethylsilicone; the column temperature was raised after injection from 40 to 225°C at 15°C min⁻¹. MBT, monobutyltin; DBT, dibutyltin; TPT, tripropyltin; TBT, tributyltin; TPhT, triphenyltin; all as the hydrides.

(4.4 $\mu\text{g kg}^{-1}$) and lower than concentrations in Steller sea lions from the same area (170–460 $\mu\text{g kg}^{-1}$). This suggests that the seals have either a lower intake of organotins from their diet or a higher catabolic capacity for these compounds than sea lions or small cetaceans.

The data reported here demonstrate that low-level organotin contamination in marine mammal predators of upper trophic levels extends to the United Kingdom in addition to those areas previously reported, and tend to suggest that these compounds may be ubiquitous in coastal areas frequented by shipping. However, the significance of these results is harder to assess. The presence of organotins at relatively high concentrations in exposed animals suggests a limited catabolic capacity for such compounds in both the mammals themselves and their prey, although biomagnification seems to be relatively modest (1.8 times in the finless porpoise, Iwata *et al.*, 1995; 0.2 to 7.5-fold in the Ganges river dolphin *Platanista gangetica*, Kannan *et al.*, 1997b; 1.0 to 6.8-fold in the bottlenose dolphin, Kannan *et al.*, 1997a). Preferential enrichment of organotins in the

liver may be associated with the presence of metal-binding proteins such as glutathione, and could indicate the operation of a detoxification mechanism (Kannan *et al.*, 1996). The occurrence of DBT as the major component in liver, whereas TBT predominates in blubber, also indicates that some active metabolism does occur in all the marine mammal species studied to date.

The reported effects of butyltins in mammals include immunosuppression (Vos *et al.*, 1984; Snoeij *et al.*, 1989), and it has been suggested that a number of mass mortalities in marine mammal populations in European and North American waters in recent years resulting from morbillivirus infections may have been rendered more severe in animals whose immune function had been impaired by environmental pollutants (Dietz *et al.*, 1989; Heide-Jørgensen *et al.*, 1992; Ross *et al.*, 1995). Interest in this area has mainly centred on organochlorine contaminants (including PCBs, dieldrin and DDT and its metabolites) which are accumulated to high concentrations in the blubber of many marine mammals (Hall *et al.*, 1992), but high

TABLE 1
Concentrations of organotin compounds in livers of porpoises and grey seals ($\mu\text{g kg}^{-1}$ wet weight).

| Ref. no. | Location | Date found | Sex | Age | Length (cm) | TBT | DBT | MBT | Σ BTs |
|-------------------------|---|-------------------|-----|-----|-------------|-----|-----|------|--------------|
| <i>Harbour porpoise</i> | | | | | | | | | |
| SW1992/202 | Sunderland, Tyne & Wear | 13/10/92 | F | ND | 133 | 180 | 350 | 110 | 640 |
| SW1993/12 | Cambois, Northumberland | 27/1/93 | M | ND | 144 | 39 | 150 | 63 | 252 |
| SW1994/32 | Westward Ho!, Devon | 4/3/94 | M | 2 | 127 | 21 | 99 | <10 | 120 |
| SW1994/45A | Rossall Point, Lancashire | 2/4/94 | M | 8 | 151 | 9 | 25 | 19 | 53 |
| SW1994/53 | Morfa Dyffryn, Gwynedd | 10/4/94 | F | 3 | 167 | 19 | 46 | 9 | 74 |
| SW1994/68 | 6 miles off Walton pier, Essex | 29/4/94 | F | ND | 112 | 53 | 150 | 50 | 253 |
| SW1994/80 | Port Talbot, Glamorgan | 6/5/94 | M | 4 | 122 | 23 | 82 | 16 | 121 |
| SW1994/99 | Tywyn, Gwynedd | 2/6/94 | M | 1 | 120 | <10 | 14 | 8 | 22 |
| SW1994/108 | Weston-Super-Mare, Avon | 25/6/94 | M | 0 | 75 | <10 | 13 | 100 | 113 |
| SW1994/148 | Sandsend, Whitby, N. Yorkshire | 1/9/94 | M | ND | 130 | 40 | 160 | 94 | 294 |
| SW1994/153 | Ynyslas, Dyfed | 7/9/94 | M | ND | 153 | 21 | 97 | 26 | 144 |
| SW1994/171 | Martello Bay, Clacton, Essex | 22/11/94 | M | ND | 146 | 57 | 210 | <10 | 267 |
| SW1994/175 | Marazion, Cornwall | 4/12/94 | F | ND | NR | 8 | 71 | 49 | 128 |
| SW1994/185 | Bosham Harbour, W. Sussex | 30/12/94 | F | 5 | 145 | 36 | 460 | 130 | 626 |
| SW1995/52 | Seaton sluice, Teeside | 20/4/95 | M | 3 | NR | 41 | 140 | <9 | 181 |
| SW1995/55 | Seaton Sands, Teeside | 1/5/95 | M | 8 | 153 | 47 | 190 | 14 | 251 |
| SW1995/76 | Newport, Norfolk | 1995 ¹ | M | 0 | NR | 47 | 58 | <7 | 105 |
| SW1995/78 | Filey Brigg, N. Yorkshire | 7/7/95 | M | ND | NR | 120 | 440 | 18 | 578 |
| SW1995/85 | Gorleston, Norfolk | 5/7/95 | F | 0 | 86 | 19 | 71 | 7 | 97 |
| SW1995/86 | Trearddur Bay, Anglesey, Gwynedd | 7/7/95 | F | 5 | 172 | 46 | 210 | 9 | 265 |
| SW1995/94 | St. Mary's Island, Whitley Bay, Tyne & Wear | 31/7/95 | F | ND | 112 | 33 | 79 | 8 | 120 |
| SW1995/102 | Aberystwyth, Dyfed | 1/10/95 | F | ND | 167 | 14 | 150 | 16 | 180 |
| SW1995/120A | Snettisham, Norfolk | 5/10/95 | F | ND | 107 | 16 | 120 | <10 | 136 |
| SW1995/126 | Withernsea, E. Yorkshire | 13/10/95 | F | 0 | 95 | 64 | 150 | <390 | 214 |
| SW1996/2 | Robin Hood's Bay, N. Yorkshire | 3/1/96 | M | 5 | 140 | 41 | 21 | <7 | 62 |
| SW1996/27(1) | Praa Sands, Cornwall | 22/1/96 | M | ND | 153 | 39 | 240 | 12 | 291 |
| SW1996/29 | Skegness, Lincolnshire | 25/1/96 | M | >1 | 126 | 42 | 91 | <6 | 133 |
| SW1996/30 | Blyth, Northumberland | 26/1/96 | M | 2 | 117 | 32 | 65 | <6 | 97 |
| SW1996/37 | Kessingland, Suffolk | 7/2/96 | M | ND | 137 | 76 | 300 | 15 | 391 |
| <i>Grey seal</i> | | | | | | | | | |
| SS1993/243 | Colwyn Bay, Clwyd | 25/12/93 | F | ND | 176 | 6 | 5 | <10 | 11 |
| SS1994/63 | Poppitt Sands, Cardigan | 24/5/94 | F | ND | 141 | <4 | <5 | <3 | nd |
| SS1994/277 | Tywyn, Gwynedd | 7/10/94 | F | ND | 179 | <4 | 6 | 14 | 20 |
| SS1995/80 | Broad Haven, Pembrokeshire | 13/7/95 | M | ND | 224 | <4 | <4 | 11 | 11 |
| SS1995/143 | River Thames | 11/9/95 | M | ND | 226 | <4 | 11 | 11 | 22 |

¹Precise date not known.

TBT, tributyltin; DBT, dibutyltin; MBT, monobutyltin; Σ BTs, sum of MBT, DBT and TBT.
nd, not detected; ND, not determined; NR, not reported.

concentrations of organotins could have an additive effect (Kannan *et al.*, 1997a). TBT (though not DBT or MBT) also acts as an endocrine disrupter in gastropods, causing an increase in testosterone levels which leads to the development of imposex (Bettin *et al.*, 1996; Spooner *et al.*, 1991), and could interact with the endocrine system in marine mammals. Whilst the levels of organotins in porpoises and seals stranded around England and Wales are lower than some of those reported for small cetaceans from other areas, further study is needed of the possible toxic effects of these compounds and the risk that their accumulation poses

to these animals. Data are also needed for other coastal mammals in the United Kingdom, such as the common seal (*Phoca vitulina*) and the bottlenose dolphin.

Paul Jepson, Dr John Baker, Dr Thijs Kuiken and Vic Simpson are acknowledged for conducting post-mortem examinations and collecting the tissue samples used in this study. The tissue analyses were funded by the Ministry of Agriculture, Fisheries and Food as part of its programme of marine environmental research. The pathological investigations were funded by the Department of Environment, Transport and the Regions as part of their commitment to the ASCOBANS agreement. We also wish to acknowledge the staff of the Natural History Museum, Stella Turk (Cornish Wildlife Trust) and Rod Penrose (Marine Environmental Monitoring) for the

TABLE 2

Concentrations of organotin compounds in blubber and liver tissues of marine mammals reported in the literature ($\mu\text{g kg}^{-1}$ wet weight). Key as for Table 1. Where a range of years is indicated, the date of sampling of individual animals was not given in the original paper.

| Species | Location | Year | Sex | Tissue | TBT | DBT | MBT | ΣBTs | Ref. ¹ |
|-----------------------------|----------------------------------|---------|----------------|---------|---------|----------|----------|--------------------|-------------------|
| Dall's porpoise | NE North Pacific | 1987 | M | Blubber | 4.7 | <1 | <14 | 4.7 | 1 |
| Dall's porpoise | NW North Pacific | 1987 | M | Blubber | 6.2 | <2 | <20 | 6.2 | 1 |
| Common dolphin | NW North Pacific | 1987 | M | Blubber | 15 | 4.3 | <20 | 19 | 1 |
| Atlantic spotted dolphin | Southeastern USA | 1991-94 | M | Blubber | 80 | 12 | 110 | 200 | 2 |
| Atlantic spotted dolphin | Southeastern USA | 1991-94 | M | Liver | 76 | 380 | 170 | 630 | 2 |
| Finless porpoise | Pacific coast of Chiba, Japan | 1981 | M | Blubber | 36 | <5 | 91 | 130 | 1 |
| Finless porpoise | Pacific coast of Chiba, Japan | 1981 | M | Liver | 200 | 790 | 130 | 1120 | 3 |
| Finless porpoise | Seto-inland Sea, Japan | 1985 | M | Blubber | 460 | 74 | 210 | 740 | 1 |
| Finless porpoise | Seto-inland Sea, Japan | 1985 | M | Liver | 1100 | 6100 | 3000 | 10200 | 3 |
| Finless porpoise | Western coast of Nagasaki, Japan | 1992 | M | Blubber | 31 | <5 | 64 | 95 | 1 |
| Finless porpoise | South China Sea | 1990 | M | Blubber | 6.7 | <1 | <14 | 6.7 | 1 |
| Finless porpoise | Ise Bay, Japan | 1994 | F | Blubber | 120 | 20 | 73 | 213 | 3 |
| Finless porpoise | Ise Bay, Japan | 1994 | F | Liver | 810 | 1800 | 680 | 3290 | 3 |
| Harbour porpoise | Baltic Sea, coast of Poland | 1991 | F | Liver | NR | NR | NR | 18 ² | 4 |
| Harbour porpoise | Baltic Sea, coast of Poland | 1991 | F | Liver | NR | NR | NR | 27 ² | 4 |
| Bottlenose dolphin | NW Adriatic Sea, coast of Italy | 1992 | M | Blubber | 41 | 16 | 55 | 110 | 5 |
| Bottlenose dolphin | NW Adriatic Sea, coast of Italy | 1992 | M | Blubber | 14 | 9 | 25 | 48 | 5 |
| Bottlenose dolphin | NW Adriatic Sea, coast of Italy | 1992 | M | Blubber | 210 | 42 | 65 | 320 | 5 |
| Bottlenose dolphin | NW Adriatic Sea, coast of Italy | 1992 | M | Liver | 250 | 800 | 150 | 1200 | 5 |
| Bottlenose dolphin | NW Adriatic Sea, coast of Italy | 1992 | M | Liver | 400 | 1600 | 200 | 2200 | 5 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | M | Blubber | 240 | 310 | 78 | 630 | 2 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | M ³ | Liver | 5.8-770 | 450-8300 | 120-2260 | 570-11340 | 2 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | F ³ | Liver | 26-170 | 290-1570 | 100-760 | 420-2500 | 2 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | M ⁴ | Liver | 22 | 230 | 68 | 320 | 2 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | F ⁴ | Liver | 94-130 | 67-720 | 310-360 | 1120-1160 | 2 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | M ⁵ | Liver | 40-50 | 54-370 | 44-290 | 140-710 | 2 |
| Bottlenose dolphin | Southeastern USA | 1989-94 | F ⁵ | Liver | 10-110 | 70-880 | 32-350 | 110-1260 | 2 |
| Killer whale | Pacific coast of Taiji, Japan | 1986 | M | Blubber | 12 | 3.4 | <20 | 15 | 1 |
| Spinner dolphin | Bay of Bengal | 1990 | M | Blubber | 2.0 | <1 | <14 | 2.0 | 1 |
| Minke whale | Antarctic Ocean | 1985 | M | Blubber | <1 | <1 | <10 | nd | 1 |
| Ginkgo-toothed beaked whale | Japan Sea | 1993 | M | Blubber | 33 | 10 | <20 | 43 | 1 |
| Pygmy sperm whale | Southeastern USA | 1991-94 | M | Liver | 5-12 | 240-290 | 86-160 | 350-410 | 2 |
| Largha seal | Western coast of Hokkaido, Japan | 1992 | M | Blubber | 4.4 | <1 | <14 | 4.4 | 1 |
| Steller sea lion | Coast of Hokkaido, Japan | 1994 | M | Blubber | 5.6 | 5.2 | 7.1 | 18 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1994 | M | Liver | 20 | 89 | 63 | 170 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1995 | M | Liver | 16 | 130 | 100 | 250 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1995 | F | Liver | 22 | 120 | 52 | 190 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1995 | F | Liver | 19 | 98 | 67 | 180 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1995 | F | Liver | 26 | 110 | 100 | 240 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1995 | F | Liver | 16 | 51 | 110 | 180 | 6 |
| Steller sea lion | Coast of Hokkaido, Japan | 1995 | F | Liver | 85 | 370 | <6.6 | 460 | 6 |

¹1. Iwata *et al.* (1994); 2. Kannan *et al.* (1997a); 3. Iwata *et al.* (1995); 4. Kannan and Falandysz (1997); 5. Kannan *et al.* (1996); 6. Kim *et al.* (1996).

²Neonates.

³Adults.

⁴Juveniles.

⁵Calves.

Dall's porpoise, *Phocoenoides dalli*; Spinner dolphin, *Stenella longirostris*; Common dolphin, *Delphinus delphis*; Minke whale, *Balaenoptera acutorostrata*; Atlantic spotted dolphin, *Stenella frontalis*; Ginkgo-toothed beaked whale, *Mesoplodon ginkgodens*; Finless porpoise, *Neophocaena phocaenoides*; Pygmy sperm whale, *Kogia breviceps*; Harbour porpoise, *Phocoena phocaena*; Largha seal, *Phoca largha*; Bottlenose dolphin, *Tursiops truncatus*; Steller sea lion, *Eumetopias jubatus*; Killer whale, *Orcinus orca*.

reporting and collection of seal and cetacean carcasses used in this study. The authors would also like to thank Jackie Reed (CEFAS Burnham Laboratory) for the preparation of Fig. 1.

- Alzieu, C. Environmental problems caused by TBT in France: assessment, regulation, prospects. *Marine Environmental Research*, 1991, **32**, 7–17.
- Bettin, C., Oehlmann, J. and Stroben, E. TBT-induced imposex in marine neogastropods is mediated by an increasing androgen level. *Helgoländer Wissenschaftliche Meeresuntersuchungen*, 1996, **50**, 299–317.
- Bonner, W. N. (1989) *The Natural History of Seals*. Christopher Helm Ltd., Bromley, UK.
- Bryan, G. W., Gibbs, P. E., Burt, G. R. and Hummerstone, L. G. The effects of tributyltin (TBT) accumulation on adult dogwhelks, *Nucella lapillus*: long-term field and laboratory experiments. *Journal of the Marine Biological Association of the United Kingdom*, 1987, **67**, 525–544.
- Cadee, G. C., Boon, J. P., Fischer, C. V., Mensink, B. P. and Ten Hallers-Tjabbes, C. C. Why the whelk (*Buccinum undatum*) has become extinct in the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 1995, **34**, 337–339.
- Clausen, B. and Andersen, S. Evaluation of bycatch and health status of the harbour porpoise (*Phocoena phocoena*) in Danish waters. *Danish Review of Game Biology*, 1988, **13**, 1–20.
- Dietz, R., Heide-Jørgensen, M.-P. and Härkönen, T. Mass deaths of harbor seals (*Phoca vitulina*) in Europe. *Ambio*, 1989, **18**, 258–264.
- Evans, C. J. and Karpel, S. Organotin compounds in modern technology. *Journal of Organometallic Chemistry Library*, 1985, **16**, 178–217.
- Evans, P. G. H. (1987) *The Natural History of Whales and Dolphins*. Christopher Helm Ltd., Bromley, UK.
- Garthwaite, D. G., Thomas, M. R. and Hart, M. (1995) *Pesticide Usage Survey Report 127: Arable Farm Crops in Great Britain 1994*. Ministry of Agriculture, Fisheries and Food, London.
- Hall, A. J., Law, R. J., Wells, D. E., Harwood, J., Ross, H. M., Kennedy, S., Allchin, C. R., Campbell, L. A. and Pomeroy, P. P. Organochlorine levels in common seals (*Phoca vitulina*) which were victims and survivors of the 1988 phocine distemper epizootic. *Science of the Total Environment*, 1992, **115**, 145–162.
- Hammond, P. S. and Prime, J. H. (1990) The diet of British grey seals, *Halichoerus grypus*. In *Population Biology of Sealworm Pseudoterranova decipiens in Relation to its Intermediate and Seal Hosts*, ed. W. D. Bowen. Department of Fisheries and Oceans, Ottawa, Ontario, Canada. *Canadian Bulletin of Fisheries and Aquatic Sciences* **222**, 243–254.
- Hammond, P. S., Hall, A. J. and Prime, J. H. The diet of grey seals around Orkney and other island and mainland sites in north-eastern Scotland. *Journal of Applied Ecology*, 1994a, **31**, 340–350.
- Hammond, P. S., Hall, A. J. and Prime, J. H. The diet of grey seals in the Inner and Outer Hebrides. *Journal of Applied Ecology*, 1994b, **31**, 737–746.
- Hammond, P. S., Benke, H., Berggren, P., Borchers, D. L., Buckland, S. T., Collet, A., Heide-Jørgensen, M.-P., Heimlich-Boran, S., Hiby, A. R., Leopold, M. F. and Øien, N. (1995) Distribution and abundance of the harbour porpoise and other small cetaceans in the North Sea and adjacent waters. *Final report. LIFE 92-2/UK/027*. Sea Mammal Research Unit, St. Andrew's, UK.
- Heide-Jørgensen, M.-P., Härkönen, T., Dietz, R. and Thompson, P. M. Retrospective of the 1988 European seal epizootic. *Diseases of Aquatic Organisms*, 1992, **13**, 37–62.
- Hiby, L., Duck, C., Thompson, D., Hall, A. and Harwood, J. (1996) Seal stocks in Great Britain. *Natural Environment Research Council News* January 1996, pp. 20–22.
- Iwata, H., Tanabe, S., Miyazaki, N. and Tatsukawa, R. Detection of butyltin compound residues in the blubber of marine mammals. *Marine Pollution Bulletin*, 1994, **28**, 607–612.
- Iwata, H., Tanabe, S., Mizuno, T. and Tatsukawa, R. High accumulation of toxic butyltins in marine mammals from Japanese coastal waters. *Environmental Science and Technology*, 1995, **29**, 2959–2962.
- International Whaling Commission (1996) Report of the Scientific Committee. 15. Small Cetaceans. *Report of the International Whaling Commission* **46**, 84–92.
- Kannan, K., Corsolini, S., Focardi, S., Tanabe, S. and Tatsukawa, R. Accumulation pattern of butyltin compounds in dolphin, tuna, and shark collected from Italian coastal waters. *Archives of Environmental Contamination and Toxicology*, 1996, **31**, 19–23.
- Kannan, K. and Falandysz, J. Butyltin residues in sediment, fish, fish-eating birds, harbour porpoise and human tissues from the Polish coast of the Baltic Sea. *Marine Pollution Bulletin*, 1997, **34**, 203–207.
- Kannan, K., Senthilkumar, K., Loganathan, B. G., Takahashi, S., Odell, D. K. and Tanabe, S. Elevated accumulation of tributyltin and its breakdown products in bottlenose dolphins (*Tursiops truncatus*) found stranded along the U.S. Atlantic and Gulf coasts. *Environmental Science and Technology*, 1997a, **31**, 296–301.
- Kannan, K., Senthilkumar, K. and Sinha, R. K. Sources and accumulation of butyltin compounds in Ganges river dolphin *Platanista gangetica*. *Applied Organometallic Chemistry*, 1997b, **11**, 223–230.
- Kim, G. B., Lee, J. S., Tanabe, S., Iwata, H., Tatsukawa, R. and Shimazaki, K. Specific accumulation and distribution of butyltin compounds in various organs and tissues of the Steller sea lion (*Eumetopias jubatus*): comparison with organochlorine accumulation pattern. *Marine Pollution Bulletin*, 1996, **32**, 558–563.
- Kuiken, T. and Baker, J. R. (1993) *Guidelines for the Postmortem and Tissue Sampling of Cetaceans*. Zoological Society of London, London.
- Laughlin, R. B. Jr. and Lindén, O. Tributyltin — contemporary environmental issues. *Ambio*, 1987, **16**, 252–256.
- Law, R. J. (1994) (Compiler). *Collaborative UK Marine Mammal Project: Summary of Data Produced 1988–1992*. Fisheries Research Technical Report, Ministry of Agriculture, Fisheries and Food, Directorate of Fisheries Research, Lowestoft (97).
- Martin, A. R. (1990) *Whales and Dolphins*. Salamander Books Ltd., London.
- NMFS (1992) Harbour porpoise in eastern North America: status and research needs. National Marine Fisheries Service, *Northeast Fisheries Science Center Reference Document* 92-06. NOAA/NMFS/NEFSC Woods Hole, MA 02543.
- Northridge, S. P., Tasker, M. L., Webb, A. and Williams, J. M. Distribution and relative abundance of harbour porpoises (*Phocoena phocoena* L.), white-beaked dolphins (*Lagenorhynchus albirostris* Gray), and minke whales (*Balaenoptera acutorostrata* Lacepède) around the British Isles. *ICES Journal of Marine Science*, 1995, **52**, 55–66.
- Prime, J. H. and Hammond, P. S. The diet of grey seals from the south-western North Sea assessed from analyses of hard parts found in faeces. *Journal of Applied Ecology*, 1990, **27**, 435–447.
- Read, A. J. and Gaskin, D. E. Changes in growth and reproduction of harbour porpoises, *Phocoena phocoena*, from the Bay of Fundy. *Canadian Journal of Fisheries and Aquatic Sciences*, 1990, **47**, 2158–2163.
- Ross, P. S., de Swart, R. L., Reijnders, P. J. H., Loveren, H. V., Vos, J. G. and Osterhaus, A. D. M. E. Contaminant-related suppression of delayed-type hypersensitivity and antibody responses in harbour seals fed herring from the Baltic Sea. *Environmental Health Perspectives*, 1995, **103**, 162–167.
- Snøeij, N. J., Penninks, A. H. and Seinen, W. Thymus atrophy and immunosuppression induced by organotin compounds. *Archives of Toxicology*, 1989, **13**, 171–174.
- Spooner, N., Gibbs, P. E., Bryan, G. W. and Goad, L. J. The effect of tributyltin upon steroid titres in the female dogwhelk, *Nucella lapillus*, and the development of imposex. *Marine Environmental Research*, 1991, **32**, 37–49.
- Suzuki, T., Matsuda, R. and Saito, Y. Molecular species of tri-n-butyltin compounds in marine products. *Journal of Agricultural and Food Chemistry*, 1992, **40**, 1437–1443.
- Tanabe, S. and Tatsukawa, R. (1992) Chemical modernization and vulnerability of cetaceans: increasing toxic threat of organochlorine contaminants. In *Persistent Pollutants in Marine Ecosystems*, eds C. H. Walker and D. R. Livingstone. Pergamon Press, Oxford, pp. 161–177.
- Tanabe, S., Watanabe, S., Kan, H. and Tatsukawa, R. Capacity and mode of PCB metabolism in small cetaceans. *Marine Mammal Science*, 1988, **4**, 103–124.
- Ten Hallers-Tjabbes, C. C., Kemp, J. F. and Boon, J. P. Imposex in whelks (*Buccinum undatum*) from the open North Sea: relation to shipping traffic intensities. *Marine Pollution Bulletin*, 1994, **28**, 311–313.
- Trippel, E. A., Wang, J. Y., Strong, M. B., Carter, L. S. and Conway, J. D. Incidental mortality of harbour porpoise (*Phocoena*

- phocoena*) by the gill-net fishery in the lower Bay of Fundy. *Canadian Journal of Fisheries and Aquatic Sciences*, 1996, **53**, 1294–1300.
- Vos, J. G., De Klerk, A., Krajnc, E. I., Kruizinga, W., Van Ommen, B. and Rozing, J. Toxicity of bis(tri-*n*-butyltin)oxide in the rat. II. Suppression of thymus-dependent immune responses and of parameters of non-specific resistance after short-term exposure. *Toxicology and Applied Pharmacology*, 1984, **75**, 387–408.
- Waite, M. E., Waldock, M. J., Thain, J. E., Smith, D. J. and Milton, S. M. Reductions in TBT concentrations in UK estuaries following legislation in 1986 and 1987. *Marine Environmental Research*, 1991, **32**, 89–111.
- Waldock, M. J. and Waite, M. E. The performance of an analytical method for determination of TBT during a six year monitoring programme. *Journal of Organometallic Chemistry*, 1994, **8**, 649–658.
- Waldock, M. J., Waite, M. E., Miller, D., Smith, D. J. and Law, R. J. (1989) The determination of total tin and organotin compounds in environmental samples. Aquatic Environment Protection: Analytical Methods, Ministry of Agriculture, Fisheries and Food, Directorate of Fisheries Research, Lowestoft (4).
-