



Royal Netherlands Institute for Sea Research

This is a postprint of:

Martínez Martínez, J., Boere, A., Gilg, I., Lent, J.W.M. van, Witte, H., Bleijswijk, J.D.L. van & Brussaard, C.P.D., (2015) New lipid envelope-containing dsDNA virus isolates infecting *Micromonas pusilla* reveal a separate phylogenetic group *Aquatic Microbial Ecology*, 74, 17-28

Published version: dx.doi.org/10.3354/ame01723

Link NIOZ Repository: www.vliz.be/nl/imis?module=ref&refid=245848

[Article begins on next page]

The NIOZ Repository gives free access to the digital collection of the work of the Royal Netherlands Institute for Sea Research. This archive is managed according to the principles of the [Open Access Movement](#), and the [Open Archive Initiative](#). Each publication should be cited to its original source - please use the reference as presented.

When using parts of, or whole publications in your own work, permission from the author(s) or copyright holder(s) is always needed.

New lipid envelop-containing dsDNA virus isolates infecting *Micromonas pusilla* reveal a separate phylogenetic group.

5 Joaquín Martínez Martínez ^{1,4*}, Arjan Boere¹, Ilana Gilg⁴, Jan WM van Lent², Harry J
Witte¹, Judith DL van Bleijswijk¹, Corina PD Brussaard^{1,3}

¹Department of Biological Oceanography, Royal Netherlands Institute for Sea
Research, PO Box 59, NL-1790 AB Den Burg, The Netherlands

10 ²Laboratory of Virology, Department of Plant Sciences, Wageningen University,
Droevendaalsesteeg 1, 6708 PB, Wageningen, the Netherlands.

³Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics,
University of Amsterdam, Amsterdam, The Netherlands

⁴Present address: Bigelow Laboratory for Ocean Sciences, 60 Bigelow Drive, East
15 Boothbay, 04544 ME, USA.

*Corresponding author: jmartinez@bigelow.org

20 **Abstract**

Viral infection of phytoplankton has major implications for biochemical and energy cycles, community dynamics, and microbial evolution in the marine environment. The non-bloom forming picoplankter *Micromonas pusilla*, a significant component of the plankton community worldwide, is known to be susceptible to
25 infection by both dsDNA and dsRNA viruses. Logically, comprehensive knowledge of the ecology of *M. pusilla* requires a better understanding of the diversity and infection mechanisms of their viruses. Here, we investigated 19 new *M. pusilla*-specific viruses (MpVs) isolated from different locations and years. We performed partial characterization of those MpVs including structural characteristics, genome
30 size, phylogenetic analysis based on partial DNA polymerase gene sequences, host range, and stability at different temperatures and upon exposure to chloroform. Combined, these characteristics allowed classification of the MpVs into two groups. Exposure to chloroform led to loss of infectivity by all MpVs in one group, which suggests the presence of an outer lipid envelop. In addition, all except one of the
35 members in that group formed a monophylogenetic clade that was distinct from all other MpV isolates. The distinctive characteristics of the two MpV groups suggest different infection strategies, which may have important implications for the ecology of both host and virus populations in the environment. Knowledge gained from our study adds value to the MpV isolates as a scientific resource as it will aid in
40 developing and testing in the laboratory new hypotheses about the ecological and biogeochemical implications of *M. pusilla* viral infection in the environment.

Keywords

Micromonas pusilla; Virus diversity; Phycodnaviridae; NCLDV; Characterization;
Lipids

45 Introduction

Viruses are the most abundant biological entities in marine environments (Bergh et al. 1989, Danovaro et al. 2001) and are major players in the microbial food web. Viruses facilitate the movement of nutrients from organisms to pools of dissolved and nonliving particulate organic matter, a process termed the viral shunt
50 (Wilhelm & Suttle 1999). The viral shunt affects microbial turnover rates and hence energy and material fluxes. Furthermore, viruses have a profound effect on microbial population dynamics and in shaping microbial evolution mainly through DNA or RNA transduction (Thingstad 2000, Brüssow et al. 2004, Martínez Martínez et al. 2006, Rohwer & Vega-Thurber 2009).

55 Viral infection constitutes a significant source of phytoplankton mortality in environmental aquatic communities and can even be responsible for the demise of large phytoplankton populations within a time scale of days (Bratbak et al. 1993, Brussaard et al. 1996, Evans et al. 2003, Baudoux et al. 2006, Martínez Martínez et al. 2012). Viruses have been isolated that infect many taxa of marine eukaryotic
60 phytoplankton including *Micromonas pusilla* (Mayer & Taylor 1979, Cottrell & Suttle 1991, Brussaard et al. 2004) a member of the class Prasinophyceae, which is considered to be the most primitive in the green lineage from which all other green algae and land plant classes have risen (Sym & Pienaar 1993). *M. pusilla* is a purely planktonic, naked, highly motile, non-blooming unicellular picoflagellate (1–3 µm in
65 diameter) and is ubiquitous in coastal and oceanic marine waters throughout

temperate and cold oceanic regions where it can occur as a prominent constituent of the picoplankton community (Manton & Parke 1960, Zingone et al. 1999, Not et al. 2004, Šlapeta et al. 2006). Predicted climate change outcomes such as increased water column stratification and reduced nutrient concentrations in ocean surface waters can favor picoplankton growth (Schaum et al. 2013), and thus *M. pusilla* dynamics may be useful indicators of ecosystem change.

The *M. pusilla* virus originally isolated from seawater samples collected in the Strait of Georgia, British Columbia (Mayer & Taylor 1979) was the first algal virus ever isolated. Many other dsDNA *M. pusilla* viruses have subsequently been isolated from numerous distant geographic locations (Waters & Chan 1982, Cottrell & Suttle 1991, Sahlsten 1998, Zingone et al. 2006, Bellec et al. 2009). Additionally, a dsRNA MpV *M. pusilla* virus (Mimovirus MpRV) member of the family Reoviridae was isolated from a Norwegian coastal seawater sample using *M. pusilla* strain LAC38 (Brussaard et al. 2004). All *M. pusilla* viruses isolated so far are icosahedral and large in particle size (90-95 nm, MpRV; 100–135 nm, dsDNA *M. pusilla* viruses). The dsDNA *M. pusilla* viruses belong to the algal virus family Phycodnaviridae, which is part of a group of viruses known as nucleocytoplasmic large dsDNA viruses (NCLDVs) that replicate in the cytoplasm and in some cases partly in the nucleus of their eukaryotic host cells (Iyer et al. 2001). Research stemming from independent studies of *M. pusilla*-specific viruses has revealed significant genetic diversity within dsDNA *M. pusilla* viruses (Cottrell & Suttle 1991, 1995, Bellec et al. 2014), some knowledge of their infection strategy and effect on host physiology (Mayer & Taylor 1979, Waters & Chan 1982), variation in strain specificity (Sahlsten 1998, Brussaard et al. 2004), and evidence for significant cospeciation between these viruses and their hosts in addition to host switches (Bellec et al. 2014). Furthermore, investigations of

their dynamics in the environment have shown seasonal and spatial variation in abundance (Cottrell & Suttle 1991, Sahlsten 1998, Zingone et al. 1999) and have indicated the importance of viruses as mortality agents of *M. pusilla* (Evans et al. 2003). Better understanding of the ecology of *M. pusilla* under current and future
95 oceanic conditions is directly linked to better insight into the diversity and infection mechanisms of specific viruses as drivers of the dynamics and evolution of *M. pusilla*. In the era of “cultivation-independent omic” methods, having well characterized model virus-host systems in culture is still essential to investigate the environmental parameters that affect viral replication success and specificity as well as to aid in
100 understanding the mechanisms of virus-host interactions and their ecological and biogeochemical consequences through hypothesis driven experimental manipulation.

In this study, we report the partial characterization including host range, genome sizing, phylogenetic reconstruction based on the DNA polymerase B gene, heat-inactivation experiments, and effect of exposure to chloroform for nineteen new
105 dsDNA *M. pusilla* virus isolates from several locations throughout the North Sea and Dutch coastal waters, the Mediterranean Sea and the English Channel. Our findings add to the accumulating scientific knowledge on this group of interesting viruses and reveal new aspects of *M. pusilla* virus diversity with potentially significant implications for the ecology of the *M. pusilla*-virus system.

110 **Materials and methods**

Algal cultures

Single strain cultures of the *Prasinophyceae M. pusilla* (Butcher) Manton and Parke, were obtained from the Roscoff Culture Collection (RCC strains – clades specified in Table 1), the National Center for Marine Algae and Microbiota at

115 Bigelow Laboratory for Ocean Sciences (strain CCMP 1545 – clade C) and the
Culture Collection at the Marine Research Center of Göteborg University (strain LAC
38 – clade A) (Table 1). The algal cultures were maintained at 15 °C and kept at mid-
exponential growth phase (approx. $1-2 \times 10^6$ cells ml⁻¹) by periodically transferring
5–10% (v/v) culture in a fresh 1:1 mixture of f/2-Si medium (Guillard 1975) and
120 enriched artificial seawater (ESAW) (Cottrell & Suttle 1991). Light (50-100 μ mol
photons m⁻² s⁻¹) was supplied by fluorescence tubes under a light-dark cycle of 16:8
hours. Cell abundances were calculated by flow cytometry (FCM) on fresh samples as
described by Marie et al. (Marie et al. 1999) using a FACScalibur flow cytometer
(Becton Dickinson, Franklin Lakes, NJ), equipped with an air-cooled laser providing
125 15 mW at 488 nm and with standard filter set-up. Trigger was set on red chlorophyll
autofluorescence.

Virus isolation and propagation

Lytic viruses infectious to *M. pusilla* were isolated during spring and summer
of different years from seawater samples originating from several geographical
130 locations in the North Sea, the Dutch coastal waters, the English Channel and the
Mediterranean Sea (Table 2). Isolation was conducted by adding filtered (GF/F filters,
Whatman, GE Healthcare Europe GbmH, Germany) seawater to exponentially
growing *M. pusilla* strains CCMP1545 and LAC38 cultures (10% v/v) in clear
borosilicate tubes. The cultures were incubated for up to 14 days under the light and
135 temperature conditions described above, and were inspected daily for lysis. Lysis was
determined by colour comparison to non-inoculated control cultures. Lysed cultures
were filtered through 0.2 μ m pore size polyestersulfone (PES) filters (Minisart®
High-flow, Sartorius, Rotterdam, the Netherlands) and used to reinfect fresh
exponentially growing algal cultures. *M. pusilla* virus isolates were made clonal

140 (henceforth referred to as MpVs) by end-point dilution as described by Brussaard et al. (2004). Virus strain MpVSP1 was provided by Dr. C. Suttle, UBC, Canada, and originated from waters off Southern California (Cottrell & Suttle 1991).

Virus production was quantified using flow cytometry. For FCM analysis, samples were fixed with glutaraldehyde (0.5% final concentration, EM grade, Sigma-
145 Aldrich) during 30 min at 4 °C followed by freezing in liquid nitrogen and storage at -80 °C until analysis. Thawed samples were diluted in TE 10:1 buffer (10 mM Tris-base, 1 mM EDTA, pH 8.0) and stained with the nucleic acid-specific dye SYBR Green I (Life Technologies, Grand Island, NY) as described by Brussaard (2004). The identity of the MpV population discriminated by FCM after infection and
150 subsequent lysis was verified by comparison to MpV-free cultures at mid-exponential and stationary growth phase, which showed no virus-like particles in the “MpV-region” (Supplementary Figure S1).

Transmission electron microscopy (TEM)

Aliquots of virus lysates (20 ml, filtered through 0.2 µm-pore size filters) were
155 concentrated by ultracentrifugation (142,000 ×g for 2 h at 8°C, in a Centrikon T-1080 Ultracentrifuge, with a TFF55.38 rotor, Kontron Instruments). The viral pellets were resuspended in 100 µl of TE 10:1 buffer. The concentrated suspensions were fixed with formaldehyde and centrifuged onto electron microscope nickel grids with carbon-coated formvar film in a Beckmann airfuge for 20 min at 30 psi. The grids
160 were negatively stained with 2% uranyl acetate and viewed in a Philips CM12 transmission electron microscope.

Additionally, we used TEM to screen thin sections from *M. pusilla* cells infected (4–12 h after infection) with two randomly selected isolates for each host to confirm that the virus particles observed by FCM and TEM in lysate samples were

165 indeed infectious to *M. pusilla*. Infected algal cells were fixed by addition of
glutaraldehyde to a final concentration of 0.1% and after 30 min, cells were pelleted
by centrifugation at 3,200 ×g for 5 min. Cells were resuspended in 2% gelatin,
pelleted again and the pellet was cut in small cubes of approximately 0.5 mm. These
samples were further fixed for 2 h in 2% glutaraldehyde/3% paraformaldehyde in
170 phosphate/citrate buffer (0.1M Na₂HPO₄·2H₂O and 9.7 mM citric acid ,pH 7.2)
containing 2.5 mM CaCl₂, washed 6x10 min in phosphate/citrate buffer and fixed with
1% osmium tetroxide in phosphate/citrate buffer for 1h. Samples were washed,
dehydrated in a graded series of ethanol (50 to 100%) and infiltrated with LR White
(London Resin company) Samples were transferred to gelatin capsules filled with
175 resin and polymerized for 48 h at 55°C. Sections were cut using a Leica Ultracut S
ultramicrotome and stained with 2% uranyl acetate and lead citrate (Reynolds) before
examination in a Philips CM12 transmission electron microscope. Average capsid
sizes were determined by measuring 10 virus particles for each isolate.

Genome size

180 Freshly produced MpV lysates were partially purified by removing cell debris
and bacteria by centrifugation (7,500 × g for 30 min at 4°C, using a fixed-angle rotor
F-34-6-38 in a 5810R centrifuge, Eppendorf). The genome size of the individual MpV
isolates was determined by Pulse Field Gel Electrophoresis (PFGE) as described in
Baudoux and Brussaard (2005). Briefly, the clarified supernatant was decanted and
185 viral particles were pelleted by ultracentrifugation (142,000 ×g for 2 h at 8°C, in a
Centrikon T-1080 Ultracentrifuge, with a TFF55.38 rotor, Kontron Instruments) and
resuspended in 150 µl SM buffer (0.1 M NaCl, 8 mM MgSO₄·7 H₂O, 50 mM Tris-
HCl, 0.0005% (w/v) glycerine). Plugs (at least 3 per virus isolate) were prepared by
mixing equal volumes of virus concentrate and molten 1.5% (w/v) InCert agarose

190 (Cambrex Bioscience, Rockland, ME, USA) in plastic molds. Once hardened, the
plugs were incubated overnight at 30°C in 800 µl of lysis buffer (250mM EDTA, 1%
SDS (v/v), 1 mg ml⁻¹ proteinase K, Sigma-Aldrich, Zwijndrecht, The Netherlands).
After decanting, the digestion buffer the plugs were washed in TE 10:1 buffer (10
mM Tris–Base, 1 mM EDTA, pH 8.0). The plugs were stored at 4°C in TE 20:50 (20
195 mM Tris, 50 mM EDTA, pH 8.0) until analysis. Plugs were loaded alongside DNA
Lambda ladder plugs (Bio-Rad, Richmond, CA) onto 1% SeaKem GTG agarose gels
(Cambrex Bioscience, Rockland, ME USA) prepared in 1 × TBE gel buffer (90 mM
Tris–Borate and 1 mM EDTA, pH 8.0). Samples were electrophoresed in a Bio-Rad
CHEF DR-II Cell unit filled with 0.5x TBE buffer (45 mM Tris–Borate and 0.5 mM
200 EDTA, pH 8.0), at 6 V/cm with pulse ramps of 20 to 45 s at 14°C for 22 h. Gels were
subsequently stained for at least 1 h with SYBR Green I (1 × 10⁴ of commercial
solution, Molecular Probes, Eugene, OR) and visualized in a FluorS imager (Bio-Rad
Instrument). Genome sizes were estimated based on the migration of each individual
genomic DNA band compared to a molecular size marker (note that this method
205 offers only approximate size values and complete genome sequencing is required for
accurate genome sizing).

Viral DNA purification

Fresh MpV lysates (25 ml) were centrifuged at 10,000 × g for 30 minutes at
4°C to remove most bacteria and cellular debris. The clarified lysate was incubated at
210 4°C overnight with NaCl (2 M final concentration) and PEG6000 (final concentration
10 wt%) (CalBiochem, San Diego, CA, USA). The viruses were subsequently
pelleted from the supernatant by centrifugation at 9,000 × g for 25 minutes at 4°C, the
supernatant was decanted and the pellet was air-dried for ~10 minutes. DNA was
extracted by adding 500 µl of prewarmed lysis buffer (0.5% SDS, 20 µg/ml

215 proteinase K) to the pellet, and incubating at 55°C for 30 minutes. Then 80 µl of 5M
NaCl and 100 µl CTAB-buffer (10% hexadecyltrimethyl ammonium bromide
[CTAB] in 0.7 M NaCl) were added and the mixture was incubated again at 65°C for
15m, after which 500 µl of Chloroform : Isoamylalcohol (24:1) was added and mixed
well. The samples were then centrifuged at 20,000 × g for 5 minutes and aqueous
220 phase containing the DNA was transferred to a new tube. Finally, DNA was
precipitated using a standard isopropanol precipitation method and resuspended in 50 µl
of TRIS (50mM, pH8). DNA yield and quality were checked using a NanoDrop®
ND-1000 (Thermo Scientific, Wilmington, DE, USA).

PCR amplification and sequencing of viral DNA polymerase B gene fragments

225 Virus DNA templates were diluted 5,000-fold in molecular-grade water and
DNA polymerase fragments were amplified from each template using previously
described PCR and thermocycling conditions with AVS1-AVS2 primers (Chen &
Suttle 1995a, b) modified by adding a M13 tag (5'-CAC GAC GTT GTA AAA CGA
C [primer]-3') to both primers. Two separate PCRs were run, one with the AVS1-
230 M13 and AVS2, and one with the AVS1 and AVS2-M13 primers. Finally, the PCR
amplicons were cleaned up using Genscript PCR Clean-up kit (Genscript Inc.,
Piscataway Township, NJ, USA) according to the manufacturer's recommendations,
prior to BigDye cycle sequencing on an ABI prism 310 Genetic analyser at the Royal
Netherlands Institute for Sea Research. *M. pusilla* virus DNA polymerase sequences
235 generated in this study were deposited into GenBank (accession numbers in Table 2).

DNA sequences from our MpV isolates were aligned with an additional 35 *M.*
pusilla virus DNA polymerase B partial sequences available through GenBank and
previously published (Chen and Suttle (1996) Bellec et al. (2014)). Five *Ostreococcus*
virus and two *Bathycoccus* virus DNA polymerase B partial sequences (Bellec et al.

240 2014) were included as an outgroup. The sequences were initially aligned using the
ClustalX algorithm (Larkin et al., 2007) within MEGA version 6 (Tamura et al.,
2013). The alignment was manually refined and a sequence mask was applied to
retain regions of unambiguous alignment; only those positions were included in
subsequent phylogenetic analyses. The alignment was subjected to a Bayesian
245 Inference (BI, generally more suitable for phylogenetic reconstruction of highly
similar sequences) using MrBayes, version 3.2.1 (Ronquist et al. 2012). We used the
GTR model of substitution (Lanave et al. 1984; Rodriguez et al. 1990, Tavaré 1986),
considering invariants and a gamma-shaped distribution of the rates of substitution
among sites. We used 4 chains of 1,000,000 generations and trees were sampled every
250 100 generations using a Markoff Chain Monte Carlo (MCMC) analysis. Chain
parameters appeared to be stationary after one hundred thousand sampled trees; the
first 200,000 trees (20%) were discarded as burn-in for the tree topology and posterior
probability. The consensus tree was viewed and edited in MEGA version 6 (Tamura
et al., 2013).

255 **Host range**

Susceptibility of 17 *M. pusilla* strains, originated from different geographical
locations (Table 1), to the 20 MpV isolates was determined using fresh cultures and
virus lysates. Briefly, each MpV isolate was added to exponentially growing cultures
of each *M. pusilla* strain (10% v/v, in triplicate). The inoculated cultures were gently
260 mixed daily to encourage virus adsorption and prevent host cells sedimentation. Host
growth for each *M. pusilla*-MpV combination was monitored daily over a 14 day
period. Lysis was determined by colour comparison to non-inoculated control cultures
and virus production was verified by FCM. FCM analysis of non-infected controls
proved the absence of a MpV-like group in these cultures. Cultures that had not lysed

265 14 days after the addition of an MpV inoculum were considered resistant to that virus strain.

In addition, a strain of the prasinophyte *Ostreococcus tauri* RCC475 (obtained from the Roscoff Culture Collection) was included in the study to test if any of the MpV isolates was able to infect a different, but closely related, phytoplankton species.

Thermal stability and sensitivity to chloroform

To determine the ability of the viral isolates to remain infectious after exposure to freezing temperatures, duplicate 0.5 ml 0.2 µm-filtered aliquots of each virus lysate were stored (without addition of any cryoprotectant) at -196°C, -80°C, and 275 -20°C for 24 h. The aliquots were thawed at room temperature immediately prior to their addition to 5 ml of exponentially growing host cultures (either *M. pusilla* strain CCMP 1545 or LAC38 used for original isolation) in borosilicate tubes. To determine viral stability at higher temperatures (4°C, 15-40°C in 5°C increments, 50°C and 60°C), duplicate 0.5 ml aliquots of each virus lysate were incubated at the specified 280 temperatures for 10 min in a waterbath, followed by 5 min cooling on ice prior to addition to 5 ml of the appropriate host culture. Duplicate virus-free cultures of both host strains served as negative controls. Inoculated and virus-free cultures were incubated under the normal culturing conditions described above and monitored daily for lysis over a 14-day period.

285 MpV lysate infectivity was also tested following 10 min incubation with 10% and 50% (v/v) chloroform. The chloroform was separated from solution by centrifugation at 4,000 × g for 5 min. The aqueous phases containing the viruses were transferred to clean tubes. These tubes were left with lids open at 4°C overnight to allow for evaporation of any remaining chloroform. Exponentially growing cultures

290 (specific *M. pusilla* strain used for isolation) were inoculated with the treated viruses
(10% v/v). The degree of infectivity was determined by end-point dilution with three
replicates and twelve 10-fold dilution levels for each virus strain. Host samples that
received the addition of 10% v/v of fresh culture medium or untreated MpVs served
as controls. The experiment was performed using 96-well plates (approximately 200
295 μ l volume per well) and borosilicate tubes (in 5 ml volume assays) to rule out possible
container or volume effects. Viral production in the last dilution that caused lysis was
confirmed by FCM.

Results

Ultrastructure analysis

300 TEM of thin sections of *M. pusilla* cells (strains CCMP1545 and LAC38, used
for original MpV isolations (Table 2)) from cultures inoculated with viruses (4-12
hours post-infection) revealed degradation of nuclear material and the accumulation
of virus particles in the cytoplasm, which was not present in non-inoculated cultures
(Figure 1A-D). TEM analysis of lysed cultures revealed free virus particles uniform in
305 shape, size, and staining intensity that were comparable to those in the thin section
photographs. Viruses isolated using either *M. pusilla* strain (MpVs) were tailless, with
icosahedral symmetry as suggested by their hexagonal outline, and approximately
125 \pm 6 nm in diameter. The virus particles showed a thick outer layer surrounding an
electron-dense inner core (Figure 1E).

310 Genome size and type

PFGE analysis revealed MpV genomes around 200 Kb (Table 2; Figure S2).
The MpVs isolated using *M. pusilla* strain CCMP1545 (CCMP1545-MpV isolates
hereafter) had smaller genomes, on average 191 \pm 4 Kb, compared to isolates

obtained using *M. pusilla* strain LAC38 (LAC38-MpV isolates hereafter), which
315 were, on average, 206 ± 6 Kb. Since our cultures were not axenic, bands smaller than
48.5 Kb seen in some of the samples likely correspond to phages with small capsid
and genome sizes present in the lysate.

Nucleic acid extraction and direct PCR amplification of the DNA polymerase
B gene (considered a good phylogenetic marker for most members of the NCLDV
320 group (Chen & Suttle 1996, Yutin et al. 2009) confirmed that all virus isolates in this
study contained dsDNA genomes (see below).

Phylogeny

Partial sequences of the DNA polymerase B gene amplified using previously
designed AVS1-AVS2 primers (Chen & Suttle 1995a, b) are highly conserved at the
325 nucleotide level among all our MpV isolates. Indeed, several of the newly isolated
MpVs have nearly identical partial DNA polymerase B gene sequences. Nonetheless,
our Bayesian inference showed strong support (BI posterior probability = 1) for two
major clades. One, which contained all the new LAC38-MpV isolates except
MpV38T and appeared to be monophyletic, and a second clade that contained all
330 remaining MpV, OtV, and OIV sequences from this and other studies. However, the
phylogenetic relationships in the second clade were poorly resolved (BI posterior
probability = 0.58). Based on the partial DNA polymerase B gene sequence, MpV38T
appears phylogenetically closer to a number of MpVs from the English Channel
(MicAV27, MicAV28, MicAV31, MicAV32, MicAV34) (Bellec et al. 2014) that
335 were isolated using a *M. pusilla* host that belonged to genetic clade A, like host strain
LAC38 used for MpV38T isolation. Two of our CCMP1545-MpVs (MpV41T and
MpV05T) were monophyletic to eight *M. pusilla* viruses isolated by Cottrell and Suttle
from a variety of geographical locations (1991, 1995) (grey labels, Figure 2) and to a

subset of MpVs isolated in 2009 (Bellec et al. 2014) from English Channel waters,
340 using clade C *M. pusilla* hosts, (MicCV21, MicCV22, MicCV23, MicCV28,
MicCV36). Another smaller clade revealed that two other 1545-MpVs (MpV40T and
MpV02T) were highly similar to MicC497V2. The remainder of our 1545-MpV
isolates showed no clear affiliation with any previously-isolated virus. MpV40T and
MpV41T were isolated from a single seawater sample collected in 2006 from the
345 exact same location from where the MicC-viruses originated.

Host range

The *M. pusilla* strains included in this study showed variable susceptibility to
infection, but none was resistant to all MpV isolates (Table 1). In contrast, the closely
related prasinophyte *Ostreococcus tauri* strain RCC475 was resistant to infection by
350 all of the 20 MpVs tested (data not shown). Multiple MpV isolates were capable of
infecting the same host strains, yet their specificity was highly variable and ranged
from MpVs capable of infecting as many as 16 of the 17 *M. pusilla* strains tested (e.g.
MpV40T or MpV12T) to as few as 4 *M. pusilla* strains (Mediterranean isolates,
MpV38T and MpV39T). Other than a reduced host range for the Mediterranean
355 isolates, no clear pattern in host susceptibility or virus infectivity was observed based
on the geographical origin and time of isolation of either the host strains or the
viruses. However, it is noteworthy that none of the CCMP1545-MpV isolates was
able to infect *M. pusilla* strain LAC38 and vice versa (Table 1).

Thermal stability and sensitivity to chloroform

360 All of our MpV isolates remained infectious after freezing at -20°C, -80°C and
-196°C for 24 hours, and slow thawing at room temperature. However, LAC38-MpV
isolates were more sensitive to high temperatures and lost infectivity following

exposure to temperatures above 35°C, while heat inactivation of CCMP1545-MpVs only occurred above 40°C (Table 3).

365 Treatment of the MpV isolates with either 10% or 50% chloroform resulted in complete loss of infectivity of all LAC38-MpV isolates, while none of the CCMP1545-MpV isolates lost infectivity (Table 3).

Discussion

All MpV isolates described here appeared to be assembled in the cytoplasm of the host and showed similar hexagonal morphology and particle diameter (125±6 nm). These characteristics are comparable to those of earlier published *M. pusilla* viruses (Mayer & Taylor 1979, Cottrell & Suttle 1991, Zingone et al. 2006). However, we were able to separate the new MpVs into two distinct groups based on patterns in: average genome size (larger genome size for LAC38-MpVs isolates); susceptibility to viral infection of *M. pusilla* strains CCMP1545 and LAC38 (viruses that lysed one of this host strains could not lyse the other host strain and vice versa); sensitivity to different temperatures treatments (LAC38-MpV isolates more sensitive to high temperature exposure); and loss of infectivity following exposure to chloroform only by LAC38-MpV isolates. The same two virus groups could be resolved based on the phylogeny of their partial DNA polymerase B gene sequences, with the exception of MpV38T that was phylogenetically apart from the rest of LAC38-MpVs isolates. Overall, the phylogenetic grouping depending on the host strain and clade is consistent with that reported by Bellec et al. (2014). Some of our MpV isolates for *M. pusilla* strain CCMP1545 (clade C) are phylogenetically close to six dsDNA *M. pusilla* viruses isolated from the English Channel (Bellec et al. 2014) and eight dsDNA *M. pusilla* viruses isolated from other distant geographical

locations (Cottrell & Suttle 1991, Chen & Suttle 1996), one of which, MpVSP1, was included for comparison in our characterization study. The six English Channel *M. pusilla* viruses were isolated using *M. pusilla* strain RCC834 and the other eight *M. pusilla* viruses were isolated and propagated using *M. pusilla* strain Plymouth27 (obtained from the algae culture collection at the University of Texas at Austin). Interestingly, *M. pusilla* strain Plymouth27 was the initial isolate from which strains RCC834 (Roscoff Culture Collection, Station Biologique Roscoff) and CCMP1545 (National Center for Marine Algae and Microbiota, Bigelow Laboratory) originated (information obtained from culture collection sites). Furthermore, *M. pusilla* strain RCC834 was also included in our host range analysis but it showed different susceptibility to our MpV strains than CCMP1545. Indeed, RCC834 was susceptible to infection by eight of the twelve LAC38-MpVs. Changes in susceptibility to viral infection by two distinct host cultures originating from a common isolate have also been reported for the phytoplankton species *Emiliana huxleyi* (Allen et al. 2007). It is likely that susceptibility differences observed for the two *M. pusilla* strains with a common origin, as for *E. huxleyi*, are due to changes in the host, adaptive or not, induced by slight changes in culture conditions under which the algae is maintained for long periods of time (Lakeman et al. 2009). The idea of changes in the host that can affect host range is also supported by the results from a recent study of genomic data that suggests slower evolutionary divergence of prasinoviruses than that of their hosts (Moreau et al. 2010).

MpVs belonging to the two groups described here coexist in the environment as several strains were isolated from the same water sample (e.g. MpV02T and -14T; MpV40T, -41T, -42T and -43T). However, we do not have data about the relative abundance and dynamics of members of both MpV groups that co-occur in the

environment. Previous studies have investigated seasonal abundance and dynamics of *M. pusilla* and their viruses in geographically distant locations throughout coastal waters of the Pacific and Atlantic Oceans (Cottrell & Suttle 1991), Scandinavian waters (Sahlsten 1998) and the Mediterranean Sea (Zingone et al. 1999). All three studies reported the presence of dsDNA *M. pusilla* viruses at all times of the year, yet they varied among locations and temporally at the same location. In particular, Sahlsten (1998) investigated the abundance of *M. pusilla* viruses infectious to different *M. pusilla* host strains including strains LAC38, CCMP1545, and CCMP491 (a reclone of CCMP1545), and found that viruses infectious to *M. pusilla* strain LAC38 were numerically dominant at all time points. Host range analysis in that study is in accordance with our findings since MpV isolates infectious to *M. pusilla* strain LAC38 appear innocuous to strains CCMP1545 and CCMP491, and vice versa, under the specific culture and environmental conditions set during these studies. What determines such specificity remains unanswered, especially given the fact that isolates obtained with *M. pusilla* strain LAC38 (clade A) can infect other clade C hosts and CCMP1545-MpVs can lyse clade A hosts other than LAC38. Further genomic and phenotypic characterization of both *M. pusilla* strains might aid in figuring out this conundrum. Unfortunately, no other parameters in Sahlsten's paper are comparable to those in our study to allow us to place their *M. pusilla* virus isolates into either group as defined here.

Combined, the distinct effect of chloroform exposure and maintenance of infectivity at high temperature exposures differed between the MpV groups in this study, suggesting different infection efficiencies under various environmental conditions and even possibly different propagation strategies. To our knowledge, this is the first study that reports the distinction between dsDNA *M. pusilla* viruses either

susceptible or resistant to chloroform exposure. Maintenance of infectivity after exposure to chloroform by all of the members in the CCMP1545-MpVs group suggests that they lack a lipid membrane surrounding their capsids, i.e. they are non-
440 enveloped viruses. The opposite argument is the case for LAC38-MpVs. To date, our understanding of the mechanisms of infection in prasinoviruses (dsDNA *M. pusilla* viruses) is limited. In the initial report of *M. pusilla* viruses isolation, Mayer and Taylor (1979) showed an infection mechanism similar to what is known for non-enveloped NCLDV members of the genera Chlorovirus and Phaeovirus (van Etten et
445 al. 2002, Wilson & Allen 2009). Briefly, virus adhesion to the host cell surface is followed by fusion of a viral inner-membrane to the host membrane, allowing entry of the virus genome into the host, and leaving an empty extracellular capsid. Progeny particles form in the cytoplasm of the host and are released through localized rupture of the host's cell membrane. This infection mechanism is consistent with more recent
450 observations on two *M. pusilla* virus isolates from Mediterranean waters (Zingone et al. 2006). Our TEM analysis did not provide information regarding the entry mechanisms of MpVs in this study, but shows MpV accumulation in the cytoplasm and rupture of the host's cell membrane. Yet, variations of infection strategies exist among members of the NCLDV group in particular regarding viral entry to and exit
455 from the host cell (Iyer et al. 2001, Law et al. 2006, Mackinder et al. 2009, Mutsafi et al. 2013). Such differences occur even within NCLDV families. For example the double membrane enveloped vaccinia viruses (family Poxviridae, animal viruses) can occur in three infectious forms with different structures, abundance and roles (Smith et al. 2002). The vaccinia viruses display a non-fusogenic outer membrane dissolution
460 allowing the inner envelope to fuse with the host's plasma membrane and releasing an intact virion core into the host's cytoplasm (Law et al. 2006, Doceul et al. 2010). *E.*

huxleyi-virus EhV-86, another member of the NCLDV group, has a different infection mechanism than that described for other algal viruses (Chloroviruses, Phaeoviruses and Prasinoviruses). EhV-86's entry and exit strategies are more similar to those of
465 animal NCLDVs. EhV-86 has a lipid membrane-enveloped capsid and although it is still not clear whether it enters its host via an endocytotic or an envelope fusion mechanism, an intact viral nucleoprotein core with its capsid reaches the host's cytoplasm. EhV-86 progeny is released via a budding mechanism during which the virions become again enveloped with host's plasma membrane (Mackinder et al.
470 2009). Based on our results, it is tempting to hypothesize that MpVs infectious to *M. pusilla* strain CCMP1545, which retained infectivity after treatment with chloroform, might be non-enveloped and follow an infection mechanism as previously described for other dsDNA *M. pusilla* virus isolates. On the contrary, the chloroform-sensitive viruses that infect *M. pusilla* strain LAC38 might carry lipid membranes around their
475 capsids and have entry and exit strategies similar to the ones described for EhV-86.

Furthermore, the loss of infectivity at temperatures above 35°C by viruses that infect *M. pusilla* strain LAC38, compared to infectivity maintenance up to 40°C by MpVs specific to *M. pusilla* strain CCMP1545, is in agreement with literature since the 1950s that indicates that non-enveloped viruses are usually more heat-resistant
480 than viruses with a lipid bilayer envelope (Nims & Plavsic 2013). The different sensitivity of MpVs with and without lipid envelopes to heat suggests variable ecological niche adaptation between both groups. Although it is unlikely that viruses get exposed to such high temperatures in the environment, the result may indicate differential ability among both MpV groups to handle longer term exposure to high
485 temperatures, hence allowing niche differentiation and variable propagation success

throughout changing environmental conditions, e.g. one MpV group could be more successful during warmer seasons and the other in colder seasons.

Finally, it is worth noting that the genomes of all the MpVs infectious to *M. pusilla* CCMP1545 are less than 200 Kb in size (average 191 ± 4 Kb), while the
490 MpVs that can infect *M. pusilla* strain LAC38 carry genomes of 206 ± 6 Kb on average, as estimated by PFGE. Full genome sequencing of several representatives from each group would conclusively determine whether the observed differences in genome sizes correspond to a deletion or insertion of one or several genes of ecological significance. It could also shed light on alternative propagation
495 mechanisms as suggested by the differential characteristics we observed among the two MpV groups in this study.

Combined, the results in our study add to the known complexity of the *M. pusilla*-virus system (dsDNA and dsRNA *M. pusilla* viruses) by revealing a greater than previously thought diversity of dsDNA *M. pusilla* viruses, based overall on
500 phylogenetic distinction, average genome size, and the indications of the existence of enveloped and non-enveloped dsDNA *M. pusilla* viruses. In addition, the new chloroform-sensitive MpV isolates infectious to *M. pusilla* strain LAC38 suggest that different infection mechanisms may exist among dsDNA *M. pusilla* viruses. Our study also highlights the importance of basic characterization of cultured viruses to
505 reveal traits and key features that might not have been revealed by genome sequencing of a few isolates. Isolation and maintenance of host-virus systems in the laboratory provides extremely valuable resources for the investigation of major plankton-virus ecology questions, biogeochemical consequences of viral infection, microbial evolution and unveiling their potential for translational and applied science.
510 Often, thorough testing of hypotheses cannot be achieved by cultivation-independent

approaches, in particular in the case of viruses, due to the lack of universal phylogenetic markers and the limitations in appropriate methods for in situ field measurements. Full benefit from these model systems can only be achieved through comprehensive knowledge of the virus characteristics.

515 **Acknowledgements**

We thank Anne-Claire Baudoux for her help with virus isolation, Anita Stoker, Bart Kay, Pim Steinberg, and Marco Flohil for their help in the laboratory and processing samples. TEM work was done at the Wageningen EM Center (WEMC), Wageningen University, The Netherlands. This work was part of the MICROVIR
520 project, which was supported by the division for Earth and Life Sciences Foundation (ALW), with financial aid from the Netherlands Organization for Scientific Research (NWO).

525

530

References

- 535 Allen MJ, Martínez Martínez J, Schroeder DC, Somerfield PJ, Wilson WH (2007)
Use of microarrays to assess viral diversity: from genotype to phenotype.
Environ Microbiol 9:971-982
- Baudoux A-C, Brussaard CPD (2005) Characterisation of different viruses infecting
the marine harmful algal bloom species *Phaeocystis globosa*. *Virology*
341:80-90
- 540 Baudoux A-C, Noordeloos AAM, Veldhuis MJW, Brussaard CPD (2006) Virally
induced mortality of *Phaeocystis globosa* during two spring blooms in
temperate coastal waters. *Aquat Microb Ecol* 44:207-217
- Bellec L, Clerissi C, Edern R, Foulon E, Simon N, Grimsley N, Desdevises Y (2014)
Cophylogenetic interactions between marine viruses and eukaryotic
545 picophytoplankton. *BMC Evol Biol* 14:59
- Bellec L, Grimsley N, Moreau H, Desdevises Y (2009) Phylogenetic analysis of new
Prasinoviruses (Phycodnaviridae) that infect the green unicellular algae
Ostreococcus, *Bathycoccus* and *Micromonas*. *Environ Microbiol Rep* 1:114-
123
- 550 Bergh O, Borsheim KY, Bratbak G, Heldal M (1989) High abundance of viruses
found in aquatic environments. *Nature* 340:467-468
- Bratbak G, Egge JK, Heldal M (1993) Viral mortality of the marine alga *Emiliana*
huxleyi (Haptophyceae) and termination of algal blooms. *Mar Ecol Prog Ser*
93:39-48
- 555 Brüssow H, Canchaya C, Hardt W-D (2004) Phages and the evolution of bacterial
pathogens: from genomic rearrangements to lysogenic conversion. *Microbiol*
Mol Biol Rev 68:560-602
- Brussaard CPD (2004) Optimization of procedures for counting viruses by flow
cytometry. *Appl Environ Microbiol* 70:1506-1513
- 560 Brussaard CPD, Kempers RS, Kop AJ, Riegman R, Heldal M (1996) Virus-like
particles in a summer bloom of *Emiliana huxleyi* in the North Sea. *Aquat*
Microb Ecol 10:105-113
- Brussaard CPD, Noordeloos AAM, Sandaa RA, Heldal M, Bratbak G (2004)
Discovery of a dsRNA virus infecting the marine photosynthetic protist
565 *Micromonas pusilla*. *Virology* 319:280-291
- Chen F, Suttle CA (1995a) Amplification of DNA-polymerase gene fragments from
viruses infecting microalgae. *Appl Environ Microbiol* 61:1274-1278
- Chen F, Suttle CA (1995b) Nested Polymerase chain reaction with three highly
degenerate primers for amplification and identification of DNA from related
570 organisms. *Biotechniques* 18:609 et seq.
- Chen F, Suttle CA (1996) Evolutionary relationships among large double-stranded
DNA viruses that infect microalgae and other organisms as inferred from
DNA polymerase genes. *Virology* 219:170-178
- 575 Cottrell MT, Suttle CA (1991) Wide-spread occurrence and clonal variation in viruses
which cause lysis of a cosmopolitan, eukaryotic marine phytoplankter,
Micromonas pusilla. *Mar Ecol Prog Ser* 78:1-9
- Cottrell MT, Suttle CA (1995) Genetic diversity of algal viruses which lyse the
photosynthetic picoflagellate *Micromonas pusilla* (Prasinophyceae). *Appl*
Environ Microbiol 61:3088-3091
- 580 Danovaro R, Dell'anno A, Trucco A, Serresi M, Vanucci S (2001) Determination of
virus abundance in marine sediments. *Appl Environ Microbiol* 67:1384-1387

- Doceul V, Hollinshead M, van der Linden L, Smith GL (2010) Repulsion of superinfecting virions: a mechanism for rapid virus spread. *Science* 327:873-876
- 585 Evans C, Archer SD, Jacquet S, Wilson WH (2003) Direct estimates of the contribution of viral lysis and microzooplankton grazing to the decline of a *Micromonas* spp. population. *Aquat Microb Ecol* 30:207-219
- Guillard RRL (1975) Culture of phytoplankton for feeding marine invertebrates. In: Smith WL, Chanley MH (eds) *Culture of marine invertebrate animals*. Plenum Press, New York, p 29-60
- 590 Iyer L, Aravind L, Koonin E (2001) Common origin of four diverse families of large eukaryotic DNA viruses. *J Virol* 75:11720 - 11734
- Lakeman MB, von Dassow P, Cattolico RA (2009) The strain concept in phytoplankton ecology. *Harmful Algae* 8:746-758
- 595 Law M, Carter GC, Roberts KL, Hollinshead M, Smith GL (2006) Ligand-induced and nonfusogenic dissolution of a viral membrane. *Proc Natl Acad Sci U S A* 103:5989-5994
- Mackinder LCM, Worthy CA, Biggi G, Hall M, Ryan KP, Varsani A, Harper G, Wilson WH, Brownlee C, Schroeder DC (2009) A unicellular algal virus, *Emiliana huxleyi* virus 86, exploits an animal-like infection strategy. *J Gen Virol* 90:2306-2316
- 600 Manton I, Parke M (1960) Further observations on small green flagellates with special reference to possible relatives of *Chromulina pusilla* Butcher. *J Mar Biol Assoc U.K.* 39:275-298
- 605 Marie D, Brussaard CPD, Partensky F, Vaulot D (1999) Enumeration of phytoplankton, bacteria and viruses in marine samples. . In: Robinson J.P. ZD, P.N. Dean, A. Orfao and 4 others (ed) *Current protocols in cytometry*. John Wiley & Sons, Chichester, p 11.11.11-11.11.15
- Martínez Martínez J, Norland S, Thingstad F, Schroeder DC, Bratbak G, Wilson WH, 610 Larsen A (2006) Variability in microbial population dynamics between similarly perturbed mesocosms. *J Plankton Res* 28:783-791
- Martínez Martínez J, Schroeder DC, Wilson WH (2012) Dynamics and genotypic composition of *Emiliana huxleyi* and their co-occurring viruses during a coccolithophore bloom in the North Sea. *FEMS Microbiol Ecol* 81:315-323
- 615 Mayer JA, Taylor FJR (1979) A virus which lyses the marine nanoflagellate *Micromonas pusilla*. *Nature* 281
- Moreau H, Piganeau G, Desdevises Y, Cooke R, Derelle E, Grimsley N (2010) Marine prasinovirus genomes show low evolutionary divergence and acquisition of protein metabolism genes by horizontal gene transfer. *J Virol* 620 84:12555-12563
- Mutsafi Y, Shimoni E, Shimon A, Minsky A (2013) Membrane assembly during the infection cycle of the giant Mimivirus. *PLoS Pathog* 9:e1003367
- Nims RW, Plavsic M (2013) Intra-family and inter-family comparisons for viral susceptibility to heat inactivation. *J Microb Biochem Tec* 5:136-141
- 625 Not F, Latasa M, Marie D, Cariou T, Vaulot D, Simon N (2004) A single species, *Micromonas pusilla* (Prasinophyceae), dominates the eukaryotic picoplankton in the Western English Channel. *Appl Environ Microbiol* 70:4064-4072
- Rohwer F, Vega-Thurber R (2009) Viruses manipulate the marine environment. *Nature* 459:207-212

- 630 Sahlsten E (1998) Seasonal abundance in Skagerrak-Kattegat coastal waters and host specificity of viruses infecting the marine photosynthetic flagellate *Micromonas pusilla*. *Aquat Microb Ecol* 16:103-108
- Schaum E, Rost B, Millar AJ, Collins S (2013) Variation in plastic responses of a globally distributed picoplankton species to ocean acidification. *Nature Clim Change* 3:298-302
- 635 Šlapeta J, López-García P, Moreira D (2006) Global dispersal and ancient cryptic species in the smallest marine eukaryotes. *Mol Biol Evol* 23:23-29
- Smith GL, Vanderplasschen A, Law M (2002) The formation and function of extracellular enveloped vaccinia virus. *J Gen Virol* 83:2915-2931
- 640 Sym SD, Pienaar RN (1993) Further observations on *Trichocystis*, a subgenus of *Pyramimonas* (Prasinophyceae, Chlorophyta). *Phycologia* 32:338-350
- Thingstad TF (2000) Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnol Oceanogr* 45:1320-1328
- 645 van Etten JL, Graves MV, Muller DG, Boland W, Delaroque N (2002) Phycodnaviridae - large DNA algal viruses. *Arch Virol* 147:1479-1516
- Waters RE, Chan AT (1982) *Micromonas pusilla* virus: the virus growth cycle and associated physiological events within the host cells; host range mutation. *J Gen Virol* 63:199-206
- 650 Wilhelm SW, Suttle CA (1999) Viruses and nutrient cycles in the sea. Viruses play critical roles in the structure and function of aquatic food webs. *Bioscience* 49:781-788
- Wilson WH, Allen MJ (2009) Giant viruses and their genomes. In: Feng Z, Long M (eds) *Viral genomes: Diversity, properties and parameters*. Nova Science Publishers, Inc
- 655 Yutin N, Wolf Y, Raoult D, Koonin E (2009) Eukaryotic large nucleo-cytoplasmic DNA viruses: Clusters of orthologous genes and reconstruction of viral genome evolution. *Virol J* 6:223
- Zingone A, Natale F, Biffali E, Borra M, Forlani G, Sarno D (2006) Diversity in morphology, infectivity, molecular characteristics and induced host resistance between two viruses infecting *Micromonas pusilla*. *Aquat Microb Ecol* 45:1-14
- 660 Zingone A, Sarno D, Forlani G (1999) Seasonal dynamics in the abundance of *Micromonas pusilla* (Prasinophyceae) and its viruses in the Gulf of Naples (Mediterranean Sea). *J Plankton Res* 21:2143-2159
- 665

670 **Figures**

Figure 1. Representative transmission electron micrographs of thin sections of healthy and infected *M. pusilla* cells and free viruses.

(A) *M. pusilla* strain CCMP1545 uninfected; (B) *M. pusilla* strain CCMP1545 infected with MpV02T; (C) *M. pusilla* strain LAC38 uninfected; (D) *M. pusilla* strain LAC38 infected with MpV03T; (E) free MpV06T particles from a *M. pusilla* strain LAC38 culture lysate. Arrows point at some MpVs within infected cells.

Figure 2. Phylogenetic tree based on DNA polymerase family B gene fragments.

Phylogenetic reconstruction was based on nucleotide sequences using Bayesian Inference. Numbers indicate branch support as posterior probabilities. In **bold** are novel MpV isolates from this study, isolated with *M. pusilla* strain CCMP1545 (▲) or with *M. pusilla* strain LAC38 (●). In **grey** are MpV isolates from Cottrell and Suttle (1991), which were isolated with *M. pusilla* Plymouth 27 (from which CCMP1545 originated). Not highlighted are MpVs and other picoeukaryote viruses (OIV and OtV, *Ostreococcus* spp.; BpV and BatV, *Bathycoccus* spp.) from Bellec et al. 2014. Letters A, B, and C within the names of these isolates indicate *M. pusilla* host clade. GenBank accession number for DNA polymerase sequences previously reported can be found within the respective references.

Tables

690 **Table 1. Host strain specificity of 19 new MpV isolates from this study and MpVSP1 (Cottrell and Suttle 1991).**

Clear boxes indicate no lysis; grey boxes indicate lysis; • denotes MpV from this host strain. Geographical origins: Pacif = Pacific Ocean; Med = Mediterranean Sea; Sarg =

Sargasso Sea; Sout = Southern Ocean; Nort = North Sea; Engl = English Channel;

695 Balt = Baltic Sea; Duct = Dutch Coastal waters.

Table 2. *M. pusilla*-virus isolates data including genome size as estimated in this study and GenBank accession number for their partial DNA polymerase B sequences.

Table 3. Sensitivity of MpV isolates to exposure to a wide range of temperature and to chloroform.

700

Grey boxes indicate lysis by the MpV isolate after the specified treatment. Clear boxes indicate loss of viral infectivity.

Supplementary Information

Supplementary Figure S1. Representative flow cytometry plot showing virus and bacteria populations.

705

(A) *M. pusilla* strain CCMP1545 infected with MpV40T; (B) *M. pusilla* strain CCMP1545 uninfected. Discrimination of groups was based on green fluorescence and side scatter signatures.






Supplementary Figure S2. PFGE photographs.

710


The MpV isolate loaded into each well is indicated at the top of the lane. λ symbols indicate wells loaded with DNA Lambda ladder plugs. Variable amounts of each MpV isolate were randomly loaded onto different gels to increase confidence when estimating genome size across samples that were electrophoresed separately. The arrows mark the ~ 200 Kb bands identified as MpV genomes. Bands < 48.5 Kb are

715

likely from phages present in some of the cultures. **Non-MpV** denotes wells that were loaded with samples other than MpV isolates and were not relevant for this study.



BpV178
BatV3



0.1

Table 1. Host strain specificity of 19 new MpV isolates from this study and MpVSP1 (Cottrell and Suttle 1991).

			<i>Micromonas pusilla</i> -viruses (MpVs)																			
Host Clade	Origin	Strain	Nor	Nor	Nor	Nor	Nor	Pac	Eng	Eng	Nor	Nor	Nor	Dut	Dut	Dut	Dut	Nor	Med	Med	Eng	Eng
			02T	04T	05T	07T	13T	SP1	40T	41T	03T	06T	08T	09T	10T	11T	12T	14T	38T	39T	42T	43T
<i>Micromonas pusilla</i> strains	A	Pac	RCC299																			
	A	Med	RCC449																			
	A	Sar	RCC450																			
	A	Sou	RCC658																			
	A	Med	RCC372																			
	A	Nor	RCC692																			
	?	Sou	RCC803																			
	B	Med	RCC434																			
	B	Eng	RCC418																			
	B	Eng	RCC461																			
	C	Med	RCC498																			
	C	Eng	RCC465																			
	C	Eng	RCC834																			
	C	Nor	RCC629																			
	C	Bal	RCC373																			
	C	Eng	CCMP1545	•	•	•	•	•	•	•												
	A	Bal	LAC38											•	•	•	•	•	•	•	•	•

Clear boxes indicate no lysis; grey boxes indicate lysis; • denotes MpV from this host strain. Geographical origins: Pac = Pacific Ocean; Med = Mediterranean Sea; Sar = Sargasso Sea; Sou = Southern Ocean; Nor = North Sea; Eng = English Channel; Bal = Baltic Sea; Dut = Dutch Coastal waters.

Table 2. *M. pusilla*-virus isolates' data including genome size as estimated in this study and GenBank accession number for their partial DNA polymerase B sequences.


MpV strain	Geographical origin	Isolation date	Isolation host	Genome size (Kbp)	GenBank Accession #
02T	North Sea 54 29'N; 04 38'E	July 2003		195	KJ955609
04T	North Sea 59 34'N; 00 30'E	July 2003		194	KJ955611
05T	North Sea 57 83'N; 07 42'E	July 2003		189	KJ955612
07T	North Sea 56 43'N; 06 00'E	July 2003	<i>M. pusilla</i> CCMP1545	186	KJ955614
13T	North Sea 59 00'N; 00 00'E	July 2003		187	KJ955620
40T	English Channel 48 45'N; 03 57'W	April 2006		194	KJ955624
41T	English Channel 48 45'N; 03 57'W	April 2006		194	KJ955625
SP1	Pacific Ocean 27 50'N; 97 04'W	October 1990		186	U32975*
03T	North Sea 55 46'N; 02 56'E	July 2003		212	KJ955610
06T	North Sea 57 83'N; 07 42'E	July 2003		200	KJ955613
08T	North Sea 56 43'N; 06 00'E	July 2003		208	KJ955615
09T	Dutch Coast 52 10'N; 04 16'E	April 2003		204	KJ955616
10T	Dutch Coast 52 10'N; 04 16'E	April 2003		210	KJ955617
11T	Dutch Coast 52 10'N; 04 16'E	April 2003	<i>M. pusilla</i> LAC38	203	KJ955618
12T	Dutch Coast 52 10'N; 04 16'E	May 2003		200	KJ955619
14T	North Sea 54 29'N; 04 38'E	July 2003		209	KJ955621
38T	Mediterranean Sea 41 40'N; 02 48'W	March 2006		205	KJ955622
39T	Mediterranean Sea 41 40'N; 02 48'W	March 2006		207	KJ955623
42T	English Channel 48 45'N; 03 57'W	April 2006		204	KJ955626
43T	English Channel 48 45'N; 03 57'W	April 2006		205	KJ955627

* Chen and Suttle 1996

Table 3. Sensitivity of MpV isolates to exposure to a wide range of temperatures and to chloroform.


MpVs	Host <i>M. pusilla</i> CCMP1545								Host <i>M. pusilla</i> LAC38													
	02T	04T	05T	07T	13T	40T	41T	SP1	03T	06T	08T	09T	10T	11T	12T	14T	38T	39T	42T	43T		
Treatment	Temperature	-196 °C	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey						
		-80 °C	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey						
		-20 °C	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey						
		4-35 °C	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey						
		40 °C	Grey	White																		
		>40 °C	White	White	White	White	White	White	White	White	White	White	White	White	White	White						
Chloroform	10%	Grey	Grey	Grey	Grey	Grey	Grey	Grey	White	White	White	White	White	White	White	White	White	White	White	White	White	
	50%	Grey	Grey	Grey	Grey	Grey	Grey	Grey	White	White	White	White	White	White	White	White	White	White	White	White	White	

Clear boxes indicate loss of infectivity. Grey boxes indicate that the MpV isolate caused lysis of the host after the specified treatment.




Supplementary Figure S1. Representative flow cytometry plots showing virus and bacteria populations.


(A) *M. pusilla* strain CCMP1545 infected with MpV40T; (B) *M. pusilla* strain CCMP1545 uninfected. Plots from uninfected cultures in late stationary phase (example given) and in exponential phase look the same. Discrimination of groups was based on green fluorescence and side scatter signatures.




Filename: PFGE 2h 1min d 10-3-06




Filename: PFGE 1.5h 1min 30-3-06



Filename: PFGE A 30min 15sec 6-4-06



Filename: PFGE 2h 1min 28-4-06



Filename: PFGE 35min 1.30min 19-5-06

Supplementary Figure S2. PFGE photographs.

The MpV isolate loaded into each well is indicated at the top of the lane. λ symbols indicate wells loaded with DNA Lambda ladder plugs. Variable amounts of each MpV isolate were randomly loaded onto different gels to increase confidence when estimating genome size across samples that were electrophoresed separately. The arrows mark the ~ 200 Kb bands identified as MpV genomes. Bands < 48.5 Kb are likely from phages present in some of the cultures. **Non-MpV** denotes wells that were loaded with samples other than MpV isolates and were not relevant for this study.