



Royal Netherlands Institute for Sea Research

This is a postprint of:

Maat, D.S., van Bleijswijk, J.D.L., Witte, H.J. & Brussaard, C.P.D. (2016). Virus production in phosphorus-limited *Micromonas pusilla* stimulated by a supply of naturally low concentrations of different phosphorus sources, far into the lytic cycle. *FEMS Microbiology Ecology*, 92, fiw136

Published version: [dx.doi.org/10.1093/femsec/fiw136](https://doi.org/10.1093/femsec/fiw136)

Link NIOZ Repository: www.vliz.be/nl/imis?module=ref&refid=281168

[Article begins on next page]

The NIOZ Repository gives free access to the digital collection of the work of the Royal Netherlands Institute for Sea Research. This archive is managed according to the principles of the [Open Access Movement](#), and the [Open Archive Initiative](#). Each publication should be cited to its original source - please use the reference as presented.

When using parts of, or whole publications in your own work, permission from the author(s) or copyright holder(s) is always needed.

1 **Manuscript for FEMS microbiology ecology**

2

3 **Virus production in phosphorus limited *Micromonas pusilla* stimulated by a supply of**
4 **naturally low concentrations of different phosphorus sources, far into the lytic cycle**

5

6 Douwe S Maat, Judith DL van Bleijswijk, Harry J Witte and Corina PD Brussaard

7

8 NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and
9 Biogeochemistry, and Utrecht University, P.O. Box 59, 1790 AB Den Burg, Texel, The
10 Netherlands.

11

12 Correspondence: Douwe Maat, Landsdiep 4, 1797 SZ, 't Horntje. Tel: +31 222 369482; Fax:
13 +31 319674. e-mail: douwe.maat@nioz.nl

14

15 Running title: Algal viruses stimulated by P-uptake during infection

16

17 Keywords: phytoplankton virus, phosphorus limitation, latent period, burst size, organic
18 phosphorus, remineralization

19

20 **Subject Category:** Microbe-microbe and microbe-host interactions

21

22 **Abstract**

23 Earlier studies show that the proliferation of phytoplankton viruses can be inhibited by
24 depletion of soluble reactive phosphorus (SRP; orthophosphate). In natural marine waters,
25 phytoplankton P-availability is, however, largely determined by the supply rate of SRP (e.g.
26 through remineralization) and potentially by the source of P as well (i.e. the utilization of
27 soluble non-reactive P; SNP). Here we show how a steady low supply of P (mimicking

28 natural P-recycling) to virally infected P-limited *Micromonas pusilla* stimulates virus
29 proliferation. Independent of the degree of P-limitation prior to infection (0.32 and 0.97 μ_{\max}
30 chemostat cultures), SRP-supply resulted in 2-fold higher viral burst sizes (viruses lysed host
31 cell⁻¹) as compared to no addition (P-starvation). Delaying these spikes during the infection
32 cycle showed that the added SRP was utilized for extra *M. pusilla* virus (MpV) production far
33 into the lytic cycle (18h post infection). Moreover, P-limited *M. pusilla* utilized several SNP-
34 compounds with high efficiency and with the same extent of burst size stimulation as for SRP.
35 Finally, addition of virus-free MpV lysate (representing a complex SNP-mixture) to newly
36 infected cells enhanced MpV production, implicating host associated alkaline phosphatase
37 activity, and highlighting its important role in oligotrophic environments.

38

39 **Introduction**

40 Phosphorus (P) is an important macronutrient for all organisms. As a component of
41 lipids, sugars and nucleic acids it is involved in the structuring, metabolism and reproduction
42 of cells. In many marine systems, relative shortage of this element is responsible for the
43 limitation of phytoplankton productivity and biomass (Ruttenberg 2003; Dyhrman *et al.*
44 2007). The ecological importance of P-limitation of phytoplankton is expected to increase as a
45 consequence of enhanced vertical stratification of many oceanic regions due to global climate
46 change (Karl *et al.* 1997; Sarmiento *et al.* 2004; Behrenfeld *et al.* 2006).

47 Reduced availability of soluble reactive phosphorus (SRP; orthophosphate) has been
48 shown to lower the impact of viral infection on phytoplankton mortality (Bratbak *et al.* 1993;
49 Wilson *et al.* 1996; Bratbak *et al.* 1998; Clasen and Elser 2007; Maat *et al.* 2014). Thus far
50 viral infection studies which tested the effects of nutrient availability only focused on the
51 effect of P-depletion or P-starvation. Although these conditions may represent the end of
52 phytoplankton spring blooms (Ly *et al.* 2014), they do not simulate the natural P-limiting

53 growth conditions that can be found in oligotrophic waters. Phytoplankton P-limitation is not
54 solely the result of low P-concentration, but is also defined by the supply rate of low
55 concentrations of P (Harris 1986). In P-limited open ocean regions, phytoplankton growth
56 completely depends on P-recycling (Benitez-Nelson 2000; Benitez-Nelson and Buesseler
57 1999; Karl and Bjorkman 2002; Dyrhman *et al.* 2007). We hypothesize that a low supply of
58 SRP, simulating P-turnover as by remineralization, stimulates virus production under
59 oligotrophic conditions. Under such conditions virally induced phytoplankton mortality rates
60 in the natural environment might be less affected by P-limitation than thus far conveyed
61 (Bratbak *et al.* 1993; Wilson *et al.* 1996; Bratbak *et al.* 1998; Clasen and Elser 2007; Maat *et*
62 *al.* 2014).

63 The extent of P-limitation depends on the capability of phytoplankton to utilize
64 alternative P-sources, such as short chain polyphosphates (SCPs), phosphomonoesters
65 (PMEs) and nucleotides (Cembella *et al.* 1984b; Casey *et al.* 2009), collectively called soluble
66 non-reactive phosphorus (SNP; Benitez-Nelson 2000). The utilization of SNP requires special
67 enzymes, such as extracellular alkaline phosphatases (APs; Cembella *et al.* 1984b, Dyrhman
68 & Palenik 2003). These enzymes hydrolyze organic P-compounds and release the
69 orthophosphates which can then be assimilated. In this way phytoplankton are able to utilize
70 all sorts of molecules, ranging from nucleotides and phosphosugars to large compounds such
71 as phospholipids and phytin (Cembella *et al.* 1984a). The concentration of SNP in the open
72 ocean can be up to five times higher than the SRP-fraction (Karl 2002) and thus a significant
73 fraction of primary production is thought to be sustained by SNP (Benitez-Nelson and
74 Buesseler 1999; Lomas *et al.* 2010). To date, there is no published data available on the
75 possible utilization of SNP-compounds by virally infected cells and the effects on virus
76 proliferation. We hypothesize that SNP-compounds can be utilized by the host cell during the
77 viral infection cycle, thereby also enhancing viral production.

78 Here we examined three research questions: i) does a steady supply of low
79 concentrations of SRP to virally infected P-limited phytoplankton stimulate virus proliferation
80 as compared to no addition, and to what extent is this dependent on the degree of P-limitation,
81 ii) how far into the infection cycle can SRP-supply be utilized for virus production, and iii) is
82 virus production positively affected by a low, but continuous supply of SNP?

83

84 **Materials and Methods**

85

86 *Culturing*

87 Experiments were performed with the picoeukaryotic photoautroph *Micromonas*
88 *pusilla*, pre-cultured axenically in P-limited chemostats at two different growth rates,
89 representing different levels of P-limitation. This ubiquitous phytoplankton species is found in
90 coastal and oceanic regions worldwide (Not *et al.* 2004; Slapeta *et al.* 2006), copes well with
91 low P-availability (Maat *et al.* 2014), and is readily infected by lytic viruses (Mayer 1977;
92 Cottrell and Suttle 1991; Baudoux and Brussaard 2008). Prior to culturing, all glassware and
93 tubing was rinsed with 0.1 M HCl and ultrapure water before autoclaving. Axenic cultures of
94 *M. pusilla* (Mp-Lac38; culture collection Marine Research Center, Goteborg University) were
95 grown in 5 l P-limited chemostats at 15°C (using a Lauda Ecoline StarEdition RE104 water
96 bath, and pumping water through water jackets of the borosilicate vessels). The cultures were
97 gently stirred by a glass clapper above a magnetic stirrer, moving at 15 RPM. Irradiance at
98 100 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ was supplied by 18W/965 OSRAM daylight spectrum fluorescent
99 tubes (QSL-2100; Biospherical Instruments Inc., San Diego, USA) in a light dark cycle of
100 16:8 h. Phosphate was the limiting nutrient (Na_2HPO_4 , 0.25 μM final concentration) in filter-
101 sterilized (0.1 μm Sartopore Midicap filter, Sartorius A.G. Germany) f/2 medium (Guillard
102 and Ryther 1962) which was based on aged, nutrient-poor ocean water and modified to

103 contain 40 μM NaNO_3 and 0.01 μM Na_2SeO_3 (Cottrell and Suttle 1991). The concentration of
104 SRP in the aged sea water was below the detection limit of 0.02 μM . Final SRP-
105 concentrations of the medium were always verified using a colorimetric assay (see below).
106 The chemostat dilution rate was adjusted to obtain two different growth rates, representing
107 different degrees of P-limitation. To allow comparison to P-replete growth conditions
108 (exponential growth, $\mu_{\text{max}} = 0.72 \text{ d}^{-1}$), separate dilution rates were set to the highest possible
109 while avoiding washout (i.e. near-maximum growth rate of $0.97\mu_{\text{max}}$; MacIntyre & Cullen
110 2005) and to $0.32\mu_{\text{max}} \text{ d}^{-1}$ to allow testing of a stronger P-limitation. After at least 5 volume
111 changes, steady state conditions (constant algal abundance) were maintained for another 3
112 months (demonstrating long term consistency). Algal alkaline phosphatase activity (APA) and
113 photosynthetic capacity (Fv/Fm) were stable over this period and SRP-concentrations in the
114 culture vessels were always below the detection limit ($< 0.02 \mu\text{M}$).

115 Lysate of the lytic dsDNA virus MpV-08T (virus culture collection, Royal
116 Netherlands Institute for Sea Research) was depleted in SRP by 3 recurring virus infection
117 cycles of axenic algal host, cultured under P-starved culture conditions (batch cultures derived
118 from the $0.97\mu_{\text{max}}$ P-limited chemostats, deprived of SRP and thus P-starved) prior to the
119 experiments. Relatively high volumes of lysate of 10-15% v/v were used because the MpV
120 stock under P-starved conditions (quantified by flow cytometry, see below) was relatively low
121 and methods to concentrate the viruses (i.e., ultracentrifugation and tangential flow filtration)
122 before addition led to decreased MpV infectivity (in a variable manner). The SRP-
123 concentration in the lysates used for the experiments was below the limit of detection (0.02
124 μM) and considered zero. Subsamples of glutaraldehyde-fixed (0.25% final concentration;
125 Sigma-Aldrich, St. Louis, USA) algal cultures and viral lysates were regularly checked for
126 axenity (containing no contaminating organisms) by epifluorescence microscopy (Axioplan,
127 Oberkochen, Germany) using the DNA-stain DAPI (4',6-diamidino-2-phenylindole,

128 dihydrochloride; Life Technologies Ltd. Paisley, UK) in combination with 0.2 μm pore-size
129 black polycarbonate filters (Whatman, Maidstone, UK) according to Porter & Feig
130 (1980). The cultures and lysates were shown to be axenic at all times.

131

132 *Experimental set-up*

133 The viral infection experiments were performed with cultures that were collected from
134 the P-limited steady-state chemostats. However, dilution of the cultures strongly complicates
135 the study of one-step virus growth cycle as the virus-host contact rate will immediately
136 change and non-growing infected host cells will be washed out. To allow for optimal infection
137 (i.e. one-step infection allowing for one-step virus growth cycle), the dilution of the cultures
138 was stopped at the moment of infection and P was thereafter (during the infection cycle)
139 supplied (spiked) by hand at fixed intervals and with similar overall rates as under chemostat
140 culturing. This way we were able to keep the cells in a constant state of P-limitation and thus
141 maintain P-limited growth conditions for the non-infected cultures. Besides this P-spiked
142 treatment, control cultures were taken from the same chemostats that (i) were deprived of SRP
143 upon viral infection (P-starved) and (ii) received directly upon infection 4 μM SRP
144 (Na_2HPO_4) final concentration in order to overcome P-limitation (P-enriched). Additionally,
145 P-replete control cultures (f/2 with $\text{Na}_2\text{HPO}_4 = 36 \mu\text{M}$, $\text{NaNO}_3 = 882 \mu\text{M}$) were grown at 1.0
146 μ_{max} (0.72 d^{-1}) for 4 weeks in a semi-continuous fashion according to the turbidostat principle
147 (regulated abundance of cells by daily dilution), facilitating non-limited cell physiology and
148 constant exponential growth (MacIntyre & Cullen 2005). Under these optimal growth
149 conditions, we expected the cultures to produce the maximum viral burst size (Maat *et al.*
150 2014).

151 For the viral infection experiments, duplicate subcultures were incubated in Erlenmeyer flasks
152 under light and temperature conditions identical to the chemostats and inoculated with 0.2 μm

153 filtered (polyethersulfone membrane filtration, Sartopore Midicap, Sartorius A.G. Goettingen,
154 Germany) axenic MpV-08T lysate at a virus : host cell ratio of 10. Virus infectivity was close
155 to 100%, as determined by comparing the most probable number of infective MpV (MPN
156 endpoint dilution; Suttle 1993) with flow cytometry MpV total counts (Fig. S1). The non-
157 infected control cultures received an equal volume of 0.1 μm filtered (polyethersulfone
158 membrane filtration, Sartopore Midicap, Sartorius A.G. Goettingen, Germany) seawater with
159 an SRP-concentration below the detection limit. Algal and viral abundance samples were
160 generally taken every 6 hours or after longer time intervals later in the infection cycle. Algal
161 samples were analyzed fresh and virus samples (1 ml) were fixed with glutaraldehyde-fixed
162 (EM-grade, 0.5% final concentration; Sigma-Aldrich, St. Louis, USA) for 15-30 min, flash
163 frozen in liquid nitrogen and stored at -80°C . Monitoring of the non-infected controls
164 continued until the infected cultures were almost completely lysed. When lysis seemed to be
165 delayed, sampling of the non-infected controls was also extended with at least another time
166 point.

167

168 *Addition of SRP to infected, P-limited M. pusilla*

169 The cultures (derived from the respective chemostats) were spiked with SRP at concentrations
170 that allowed similar growth as under 0.97 and $0.32\mu_{\text{max}}$ chemostat culturing. The first 12 h
171 post infection (p.i.) spiking took place every hour (10.5 and $2.6 \text{ amol P cell}^{-1} \text{ hour}^{-1}$, for the
172 0.97 and $0.32\mu_{\text{max}}$ cultures, respectively), but thereafter (until full lysis of culture) the
173 frequency was reduced to every 6 h as the viral latent periods (the time until first release of
174 progeny viruses) were largely covered. The volume of added SRP-stock ($[\text{Na}_2\text{HPO}_4]=104.2$
175 μM) was adapted to the time period in between spiking, and at all times corrected for the
176 change in cell abundance due to growth (non-infected controls) or cell lysis (infected
177 cultures). This correction for cell abundance ensured that the non-infected controls and the

178 infected cells obtained the same amount of SRP per algal cell. For example, the total
179 concentration of spiked P to the $0.97\mu_{\max}$ roughly doubled per day for the non-infected
180 controls, while it reduced for the lysing virally infected cultures according to the number of
181 cells remaining. The replicate of the $0.32\mu_{\max}$ P-enriched treatment failed for technical
182 reasons.

183 Finally, to study the possible effects of P-limitation on viral genome production as a
184 causal factor for potentially delayed and decreased viral production, the intracellular
185 production of viral genomes in the $0.97\mu_{\max}$ cultures was monitored by quantifying MpV
186 DNA polymerase gene copies (*DNApol*). The total MpV *DNApol* abundance per lysed host
187 cell was calculated by dividing the increase in concentration of MpV *DNApol* copy number
188 by the maximum decline in host cells over the entire one-step infection cycle (lysed host
189 cells).

190

191 *Delayed addition of SRP to infected, P-limited M. pusilla*

192 To test how far into the virus growth cycle the supplied SRP can still be utilized by the host
193 for virus production, the $0.97\mu_{\max}$ cultures received SRP-additions at similar concentrations as
194 for Experiment 1, but the administration was delayed for 1, 6, 12, 18, 24, 36 or 48 h p.i. To
195 ensure that the potential effects on virus growth characteristics originated from the timing of
196 spiking rather than the amount of P, the total amount of SRP at the first spiking event was
197 increased with the period of delay and corrected for the remaining volume and *M. pusilla*
198 abundance (thus equaling cellular P-quota). Neither the infected nor the non-infected controls
199 were negatively affected in their growth rate or Fv/Fm by this relatively large input of SRP
200 (Fig. S2).

201

202 *Addition of SNP to infected, P-limited M. pusilla*

203 To determine whether SNP-availability affects viral proliferation in infected *M. pusilla* in a
204 similar manner as SRP, the 0.97 μ_{\max} cultures were spiked with different SNP-compounds at
205 the same frequency and final P-concentration as for Experiment 1. The compounds used were
206 disodium-glycerolphosphate hydrate (GP; Sigma-Aldrich, G6501), disodium adenosine 5'-
207 monophosphate (AMP; Sigma-Aldrich, 01930) and pentasodium tripolyphosphate
208 hexahydrate (PP; Sigma-Aldrich, T5633).

209 As viral lysate is an important source of SNP (Gobler *et al.* 1997; Haaber and
210 Middelboe 2009), we tested whether the addition of the P-starved lysate that we added to the
211 viral infection experiments was responsible for part of the virus production in our
212 experiments. For this, 15% v/v virus-free MpV lysate (0.02 μm pore-size Whatman Anodisc
213 25, UK) was added to P-limited 0.97 μ_{\max} cultures at T0. Note that all treatments received the
214 same amount of lysate at the start of the experiment, which enables comparison across the
215 different treatments.

216

217 *Enumeration of algae, viruses and viral genomes*

218 Algal abundances were analyzed on fresh samples by flow cytometry (Marie *et al.* 1999)
219 using a BD Accuritm C6 cytometer (BD Biosciences, San Jose, Ca, USA), triggered on
220 chlorophyll *a* red autofluorescence. Viral abundance samples were samples were thawed,
221 diluted with TE buffer and stained with SYBRGreen I (final concentration of 0.5 x 10⁻⁴ of the
222 commercial stock; Life Technologies Ltd., Paisley, UK) according to Brussaard (2004).
223 Following a 10 minute 80°C heat incubation, the samples were analyzed using a benchtop BD
224 FacsCalibur (BD Biosciences, San Jose, USA) with the trigger set on green fluorescence. All
225 flow cytometry data were analyzed using CYTOWIN 4.31 (Vaulot 1989). Viral burst size was
226 determined by dividing the number of newly produced viruses (released by the host cells) by
227 the maximum decline in host cells, i.e. number of lysed host cells.

228 Quantitative PCR was based on Brown *et al.* (2007), but included a heat treatment according
229 to Short and Short (2008), new primers specific for strain MpV-08T, and the use of a
230 calibration curve to allow absolute quantification of the number of *DNApol* copies μl^{-1} . Total
231 MpV genome abundance, i.e. the sum of intra- and extracellular viral genomes in the sample,
232 was determined on 1 ml samples that were stored at -20°C in 2 ml cryovials (Greiner bio-one
233 GmbH, Frickenhausen, Germany). One day before analysis the samples were thawed, diluted
234 1:5 in freshly prepared deionized water (18.2 M Ω), sonicated (MSE Soniprep 150, UK) for 3
235 $\times 10$ s at amplitude 8, followed by a 5 minute heat treatment at 80°C and storage at -80°C .
236 This treatment was sufficient to break down all cells and viruses, as ascertained by flow
237 cytometry (no cells or viruses could be detected anymore by the protocols described above).
238 The new forward (MpV08T_qF1: 5'-ATGGAAATATCGAAGGTATTA-3') and reverse
239 (MpV08T_qR1: 5'-ACCATATATCGAGTTCATTG-3') primers targeted the viral DNA
240 polymerase gene and produce a product of 220 bp. PCR reactions of 20 μl , contained 1 μl
241 template, 1u Picomax polymerase, 1x Picomax buffer (including 1.5 mM MgCl_2), 200 μM
242 dNTP's, 0.2 μM of each primer, 1 mg/ml BSA and 0.2 \times SYBRgreen. After each run, a
243 melting curve was constructed from a 5 s scan during stepwise increments of temperature
244 from 72 to 95°C . All experimental samples were run along a calibration curve (constructed in
245 duplicates from 7 samples increasing from 10 copies to 10 million copies of the target) made
246 from purified MpV product of newly designed strain-specific forward (MpV08T_Fa; 5'-
247 AAGGGIGCITATTACACACC-3') and reverse (MpV08T_Rt: 5'-
248 GGCTTITTGAAIAGTGCACT-3') primers, amplifying a fragment from 5 bases downstream
249 of AVS1 to 46 bases upstream of AVS2 (AVS primers by Chen and Suttle 1995) producing a
250 589 bp product that includes the target of the qPCR primers. The calibration curve was spiked
251 with 10 million copies non target SPUD-A DNA (Nolan *et al.* 2006; 5'-
252 AACTTGGCTTTAATGGACCTCCAATTTTGAGTGTGCACAAGCTATGGAA

253 CACCACGTAAGACATAAAAACGGCCACATATGGTGCCATGTAAGGATGAATGT-3').

254 The efficiency of the qPCR reactions of experimental and calibration samples with primer
255 pair MpV08T_qF1 -_qR1 was 96.1% ($R^2 = 0.994$).

256

257 *Inorganic P-concentrations, algal physiology and statistics*

258 Concentrations of SRP were determined colorimetrically as described by Hansen and
259 Koroleff (1999). Samples were filtered (0.2 μm , FP 30/0.2 CA-S Whatman, Dasser,
260 Germany) into clean screw cap vials and stored at -20°C until analysis. The lower limit of
261 detection was 0.02 μM . All chemostat cultures in steady-state gave undetectable
262 concentrations, demonstrating that all added P (in the medium) was directly taken up by the
263 cells (this is standard practice for axenic phytoplankton cultures; e.g. Veldhuis & Admiraal
264 1987; Yao *et al.* 2011). The SRP-concentrations were measured at the beginning and end of
265 all infection experiments and were all shown to be below the limit of detection, with of course
266 the exception for the P-replete and P-enriched treatments.

267 As an indicator of P-limitation within the steady state algal cultures (Healey and
268 Hendzel 1979; Beardall *et al.* 2001), alkaline phosphatase activity (APA) was determined
269 fluorometrically according to Perry (1972). To a glass cuvette containing 2 ml of culture, 250
270 μl of 3-O-methylfluorescein phosphate (MFP; Sigma-Aldrich, M2629) was added to a final
271 concentration of 595 μM . Emission at 510 nm was measured on a Hitachi F2500
272 Fluorescence spectrophotometer (Hitachi Instruments, San Jose, CA, USA) for 60 s with an
273 excitation wavelength of 430 nm. The rate of MFP conversion was determined from a
274 standard curve of 3-O-methylfluorescein (Sigma-Aldrich, M7004). As expected, APA
275 increased with the extent of P-limitation, showing zero for the P-replete semi-continuous
276 cultures and 9.5 ± 0.9 and 31 ± 0.1 $\text{amol P cell}^{-1} \text{ s}^{-1}$ for the 0.97 and $0.32 \mu_{\text{max}}$ P-limited cultures,
277 respectively.

278 Photosynthetic capacity F_v/F_m was determined by PAM fluorometry (Water-PAM,
279 Walz, Germany). Samples (2 ml) were dark-adapted at culture temperature (15°C) for 15
280 minutes before analysis. All measurements were carried out with the same Water-PAM
281 settings, whereby 0.2 μm filtered sea water was used as a blank. After determining the
282 minimal (F_0) and maximal fluorescence (F_m), the variable fluorescence (F_v) was calculated
283 as $F_m - F_0$ (see Maxwell & Johnson 2000).

284 Statistics were carried out using SigmaplotTM 12.0 (Systat software Inc, Chicago, IL,
285 USA). One-way ANOVAs were used for testing the differences between the treatments, but
286 when the assumptions were not met non-parametric (Kruskall-Wallis or Mann-Whitney) tests
287 were used. Values in tables and figures are average \pm standard deviation (s.d.).

288

289 **Results**

290

291 *Addition of SRP to infected, P-limited M. pusilla*

292 While growth in the P-starved cultures ceased, SRP-addition allowed continued
293 growth of the non-infected *M. pusilla* cultures (Fig. 1). The extent of P-limitation determined
294 the growth rates, i.e., the 0.32 and 0.97 μ_{max} P-spiked cultures maintained growth at 0.30 \pm 0.0
295 and 0.68 \pm 0.0 d⁻¹, while this was 0.76 \pm 0.0 and 0.53 \pm 0.1 d⁻¹ after receiving a surplus of 4 μM
296 (SRP-enriched). The level of limitation (0.97 and 0.32 μ_{max} ; starved, spiked or enriched) did
297 not seem to affect the lysis dynamics of the infected cultures (Fig. 1b). All P-controlled
298 cultures showed slightly slower lysis at the end of the first day post infection (18-24h)
299 compared to the P-replete treatments (Fig. 1b).

300 The period to the release of the first progeny viruses (i.e. latent period of MpV) in the
301 0.97 μ_{max} P-spiked algal host cultures was of similar length as in the P-enriched and P-replete
302 cultures, i.e. 6-12 h (Fig. 1c). The MpV latent periods in all 0.32 μ_{max} cultures were prolonged

303 to 12-18 h. Independent of the effect on the latent period, the rates of increase in extracellular
304 progeny viruses varied for the different treatments, i.e. rates reduced with increasing P-stress
305 (from 1.7 to 0.8-1.1×10⁷ MpV ml⁻¹ h⁻¹ under replete and enriched conditions, and from 0.4-
306 0.8 to 0.2 MpV ml⁻¹ h⁻¹ for the spiked compared to the P-starved treatment) MpV burst sizes
307 of the P-starved 0.32μ_{max} cultures were not statistically different from the 0.97μ_{max} treatment
308 (average of 75±1; Kruskal Wallis ANOVA on ranks, p=0.667, n=2; Table 1). Independent
309 (pilot) experiments with the same set-up and virus-host model systems (0.97 and 0.32μ_{max}
310 SRP-starved cultures, n=15 in total) gave similar results, i.e. average 74 ±3 MpV lysed host
311 cell⁻¹ (data not shown). Likewise, the semi-continuously grown P-replete cultures showed
312 good replication with an average MpV burst size of 310 ±26 when including the results of
313 other independent experiments (n=12 over a period of 2 years). Furthermore, the 0.97μ_{max} P-
314 starved treatment in all three experiments in the current study did not show significant
315 differences in MpV latent period or burst size (average of 75±2; Kruskal Wallis ANOVA on
316 ranks, p=0.333, total n=6; Table 1), illustrating that reproduction for our experiments was
317 good. Independent of the degree of P-limitation (0.97 vs 0.32μ_{max}), spiking of the infected
318 cultures with SRP led to significantly increased MpV burst sizes compared to the P-starved
319 cultures (on average a doubling; ANOVA, p=0.003; Table 1). However, the burst sizes of the
320 SRP-spiked cultures were still lower than for the P-enriched and P-replete treatments
321 ANOVA, p=0.015; Table 1). The enriched cultures still showed lower burst sizes than the
322 replete treatment (i.e., 44 and 22% lower for the 0.32 and 0.97μ_{max} culture, respectively;
323 Table 1).

324 Similarly to the MpV burst sizes in the 0.97 μ_{max} spiked and starved treatments, the
325 MpV *DNApol* copy number was also strongly affected by the strength of P-limitation (Fig. 2),
326 i.e., respectively 7- and 11-fold lower than P-replete. Although the temporal dynamics of
327 MpV *DNApol* copy number and viruses over the infection cycle were the same, the total

328 number of viral *DNApol* cell⁻¹ was 3, 2 and 7-fold higher than the viral burst sizes of the 0.97
329 μ_{\max} P-starved, P-spiked and P-replete treatment, respectively (Table 1). Hence, under P-
330 replete conditions the ratio of viral genomes to viruses was higher than under P-spiked and P-
331 starved conditions.

332

333 *Delayed addition of SRP to infected, P-limited M. pusilla*

334 Delaying the addition (spiking) of SRP led to a lag in growth of the non-infected control
335 cultures, (i.e., 0.22 ± 0.0 d⁻¹ until being spiked, compared to 0.61 ± 0.1 d⁻¹ for the continuously
336 spiked culture), but growth quickly recovered upon spiking. The temporal dynamics of the
337 infected cultures were not affected in comparison to the regularly spiked cultures (Fig. 3).
338 Delayed SRP-addition up to 18 h p.i. still gave comparable MpV production and burst sizes
339 compared to continuous spiking from 0 h p.i. (Fig. 1 and 3, Table 1). Starting the additions of
340 SRP after 24 h p.i., gave similar results as for the P-starved cultures (Mann-Whitney Rank
341 sum test, $p < 0.001$, $n \geq 8$; Figure 3c, Table 1). The 16:8h light-dark synchronized growth of *M.*
342 *pusilla* may theoretically have influenced the results of the delayed spiking experiment.
343 However, at the moment of the 18h p.i. SRP-addition the cultures were already in the dark for
344 5 h while still showing similar virus growth kinetics as the 0-12h p.i. SRP-supply treatments.
345 Furthermore, the 24 h p.i. treatment started in the light and still showed lower MpV yield.
346 Light-dark cycle did thus not affect the outcome.

347

348 *Addition of SNP to infected, P-limited M. pusilla*

349 Supplying the $0.97\mu_{\max}$ P-limited non-infected control cultures with the SNP-compounds GP,
350 AMP and PP resulted in maximum growth rates, similar as with SRP (0.72 ± 0.1 d⁻¹; average
351 over 48h experiment). Independent of SNP-source, all infected algal cultures showed
352 comparable lysis dynamics and MpV latent periods similar to SRP-spiked treatment (Fig. 4b

353 and c; Table 1). All SNP-compounds stimulated MpV production (as compared to SRP-
354 starved), PP even to the same degree as found for the SRP-spiked treatment (Fig. 4c).
355 Although it took longer (about 24 h), the maximal yield of MpV under GP and AMP spiking
356 was not significantly different from the PP and SRP-treatments (Fig. 4c). Consequently, MpV
357 burst sizes for the PP, GP, and AMP spiked treatments were not different from SRP-spiked
358 (Table 1).

359 The addition of virus-free lysate, to test whether lysate itself was a significant source
360 of rapidly bioavailable SRP and subsequently MpV production in all infected cultures, led to a
361 burst size slightly higher than the P-starved cultures (Fig. 4d; Table 1). This result was tested
362 once more in a separate experiment (n=2) with similar results. Overall, average burst size
363 increase was 16 ± 5 MpV host cell⁻¹ (t-test, n=4, p=0.003). This increase was clearly lower than
364 the increase in MpV burst size obtained for the other SRP-treatments, as the added total
365 amount of P in the lysate was also lower than that for the other treatments.

366

367 **Discussion**

368

369 In this study we tested the effects of low supply rates of SRP (simulating natural
370 remineralization conditions) and SNP (simulating e.g. release of cell content by viral lysis) on
371 the virus growth cycle of P-limited *M. pusilla*. Spiking with SRP during the infection cycle
372 distinctly increased the viral burst sizes (no effect on latent period) around 2-fold as compared
373 to the P-starved treatment. Since the algal host cells were still P-limited, the burst sizes were
374 not as high as those under P-replete or P-enrichment conditions (this study; Maat *et al.* 2014).
375 Additionally, the strength of P-limitation, obtained by adapting the algal host in P-limited
376 chemostats to different growth rates prior to infection (i.e. 0.97 and low 0.32 μ_{\max}), affected
377 the burst size upon SRP-spiking, with the strongest increase for the 0.97 μ_{\max} cultures.

378 Viruses are strictly dependent on their host for the substrate, enzymes and energy that
379 are involved in virus replication, and hence the physiological state of the host can be expected
380 to drive the outcome of infection. This was best illustrated by P-enrichment (addition of 4 μM
381 SRP to P-limited cultures), whereby SRP was no longer limiting *per se* and the difference in
382 burst size between the 0.97 and 0.32 μ_{max} pre-cultured *M. pusilla* was solely due to a constraint
383 on the host's physiology prior to infection. Potential damage of photosynthetic machinery
384 and/or additional components of the energy metabolism under P-limitation could be such
385 constraint (Theodorou *et al.* 1991; Graziano *et al.* 1996). Previous studies have shown the
386 dependence of viral replication on photophosphorylation for several phytoplankton species
387 (Padan *et al.* 1970; Waters and Chan 1982; Vanetten *et al.* 1983; Juneau *et al.* 2003; Baudoux
388 and Brussaard 2008). The strength of P-limitation at steady state (prior to MpV infection)
389 affected the Fv/Fm of *M. pusilla* negatively, i.e. 0.49 for 0.32 μ_{max} cultures as compared to
390 0.61 for 0.97 μ_{max} and 0.66 for P-replete cultures (Maat *et al.* 2014). Upon infection with
391 MpV, SRP-spiking, at least partly, counteracted the strong decline in Fv/Fm observed under
392 P-starved conditions (Supplement Fig. S3a). Spiking with P could thus stimulate host energy
393 metabolism and hence viral proliferation. These data furthermore affirm the interplay of viral
394 infection and environmental factors such as light and nutrient limitation on the photosynthetic
395 capacity of phytoplankton (Fig S3b; Kimmance *et al.* 2014).

396 However, the supplied P could also be (partly) allocated to the actual production of the
397 viruses inside the host cell, e.g. synthesized into nucleic acids. Viruses have been found to
398 recycle host nucleotides and it has been argued that the relative size of a phytoplankton
399 genome is a good predictor of the viral burst size (Paul *et al.*, 2002; Brown *et al.*, 2006;
400 Brown and Bidle, 2014). Assuming non-limiting growth conditions of the host, a MpV
401 genome of 208 kb (Martínez Martínez *et al.* 2015) and *M. pusilla* host genome of 22 Mbp
402 (Worden *et al.* 2009), the predicted maximum viral burst size for our virus-host system would

403 then be 106 MpV lysed host cell⁻¹, which is around 3-fold lower than experimentally recorded
404 under P-spiked, enriched and replete conditions (this study; 141 – 330 MpV lysed host cell⁻¹).
405 Thus at least part of the viral nucleotides has to have been the product of *de novo* synthesis.

406 Furthermore, there was a strong, 10-fold, overproduction of MpV *DNApol* cell⁻¹ (as
407 compared to MpV burst size) under P-replete conditions. Such overproduction of MpV
408 genomes in comparison to released MpV progeny has been suggested to be a way to increase
409 the speed of viral genome packaging in the dense host cytoplasm (Brown *et al.* 2007;
410 Weidmann *et al.* 2011). Alternatively, Nissimov *et al.* (2015) argued that overproduction of
411 genome copy numbers for *Emiliana huxleyi* viruses was due to a limitation in the availability
412 of components of the protein capsid or lipid membrane. As the capsid of MpV-08T is also
413 surrounded by a lipid membrane (Maat *et al.* 2016), both capsid proteins or lipids may
414 potentially have been limiting MpV-08T production. Under P-limiting and P-starved
415 conditions, both the viral burst sizes and the genome copy number per cell were lower than
416 under P-replete conditions. It is likely that the availability of P affects the transcription of
417 viral genomes directly, either because of its function in compounds that supply energy to the
418 necessary enzymatic activity (Theodorou *et al.* 1991) or as element in viral nucleotides
419 (besides potential recycling from the host; Wikner *et al.* 1993, Brown & Bidle 2014).
420 Remarkably, the overproduction of MpV *DNApol* cell⁻¹ compared to MpV burst size was only
421 2- and 3-fold under P-starved and P-limiting conditions, respectively. The packaging
422 efficiencies (defined as the number of viruses divided by the number of virus genome copies)
423 under P-starved and P-limiting conditions were thus higher than under P-replete conditions
424 (47, 35 and 15%, respectively).

425 Delayed SRP-spiking of the 0.97 μ_{\max} cultures up to 18 h p.i. gave similar viral burst
426 sizes as spiking from the start of infection. *M. pusilla* was thus able to utilize the supplied
427 SRP for additional MpV progeny far into the lytic cycle. Zheng and Chisholm (2012) showed

428 that P-addition to P-depleted and infected *Prochlorococcus* sp. led to a decrease in the
429 transcription of a virally encoded gene that is likely involved in the uptake of P. Effects on the
430 viral burst size were however not investigated in their study. Phytoplankton P-uptake during
431 the virus infection cycle has been the subject of discussion in several theoretical studies, as it
432 strongly affects the modelling results of the impact of viruses in natural nutrient limited
433 systems (Rhodes and Martin 2010; Jover et al. 2014). Overall, our results demonstrate that it
434 is key for future research to consider SRP-supply rate, i.e. P-turnover rate under natural
435 conditions, as a significant driver of virus proliferation and thus to our understanding to what
436 extent viruses structure host population dynamics.

437 Viral lysis rates of different phytoplankton species (including *M. pusilla*) under natural
438 oligotrophic conditions are found to be at least comparable to grazing rates (Cottrell and
439 Suttle 1995; Baudoux et al. 2007; Mojica et al. 2014). Besides, Lønborg *et al.* (2013) showed
440 that a high percentage of the dissolved cellular material released upon viral cell lysis of axenic
441 *M. pusilla* is readily bioavailable. Such a steady supply of labile organic matter stimulates
442 bacterial production and through remineralization SRP is constantly brought back into the
443 system (Gobler *et al.* 1997; Brussaard *et al.* 2005; Haaber and Middelboe 2009). Our data
444 illustrate that under P-limitation a low supply of SRP during the infection cycle positively
445 affects the production of MpV. Therefore the impact of viruses on phytoplankton mortality in
446 oligotrophic ecosystems may be higher than previously anticipated (Wilson *et al.* 1996;
447 Bratbak *et al.* 1998; Clasen and Elser 2007; Maat *et al.* 2014).

448 Not only SRP, but also SNP-compounds were found to be efficiently utilized by
449 axenic *M. pusilla*. Growth and viral production dynamics of the non-infected and infected
450 SNP-spiked cultures were similar as under SRP-supply. The ecological potential is substantial
451 since SNP in the oligotrophic open ocean can be up to 5 times higher than SRP (Karl 2002).
452 The more readily available SNP-compounds such as PP and SCPs may accordingly not only

453 be important stimulants of phytoplankton physiology and growth (Cembella 1984a), but also
454 of virus proliferation within infected algal host. The conversion of SNPs to SRP is likely to be
455 catalyzed by APs and other enzymes (Chróst & Siuda 2002, Dyhrman & Palenik 2003). We
456 show that *M. pusilla* APA in P-limited chemostats increases with decreasing growth rate. This
457 increase continues when the chemostat pumps are stopped (P-deprivation leading to
458 starvation), also in virally infected cultures (Maat et al. 2016; Fig. S4). Additionally, by
459 testing the effect of lysate as source of SRP, we found that even relatively small amounts of
460 lysate (of more complex composition) can directly stimulate growth of axenic non-infected *M.*
461 *pusilla* as well as viral production in infected cultures. Literature suggests that around 38-49%
462 of total P in the viral lysates is transferred back into the dissolved phase, whereby enzymatic
463 activity from heterotrophic bacteria is suggested to be responsible for the conversion of total P
464 to SRP (Gobler *et al.* 1997; Haaber and Middelboe 2009; Lønborg and Álvarez-Salgado
465 2012). We speculate that algal APs in our axenic algal cultures, potentially in combination
466 with activity of other enzymes (e.g. nucleases), associated to lysed *M. pusilla* cells could have
467 performed the same process (Chróst & Siuda 2002, Dyhrman & Palenik 2003). We were able
468 to still detect APA in a week old (axenic) P-starved cultures of *M. pusilla* (data not shown).
469 This was probably due to APs bound to *M. pusilla* cell debris (lysed cells). Although some
470 phytoplankton species excrete free APs to the environment (Chróst & Siuda 2002), we did not
471 find APA in 0.2 μm filtered (lysed and non-lysed) *M. pusilla* cultures. Viral lysis-induced
472 release of host cellular content seems an ecologically interesting source of bioavailable SNP
473 that can be directly utilized and as such has the potential to stimulate MpV production of
474 newly infected (but not yet lysed) neighboring cells. The reduction of P-limitation induced by
475 viral lysis and the subsequent stimulation of viral production implies a positive feedback
476 process that maintains regeneration in P-limited oligotrophic ecosystems. We do however,
477 realize that we used axenic phytoplankton cultures in our study, while in the natural

478 environment heterotrophic bacteria may compete for the same sources of SNP (and even SRP
479 that is made available by phytoplankton extracellular APA). Løvdal and coworkers (2008)
480 showed that biomass-specific uptake of SRP and SNP was similar for phytoplankton and
481 heterotrophic bacteria during an *Emiliana huxleyi* bloom in the coastal North Sea, which
482 suggests that products of viral lysis are indeed available for phytoplankton and subsequently
483 algal virus proliferation. In contrast, Hartmann *et al.* (2011) showed lower biomass specific
484 uptake for small protists than for bacteria in the North Atlantic subtropical gyre, suggesting
485 that the outcome of competition may depend on the (trophic) state of the marine system.
486 However, the experimental incubations in their study were carried out in the dark, which
487 could also have led to an underestimation of phytoplankton P-uptake (Nalewajko & Lee
488 1983). We recommend forthcoming studies to test the influence of potentially competing
489 heterotrophic bacteria. Alternatively to uptake of dissolved P, photosynthetic protists can
490 display considerable bacterivory in the oligotrophic Atlantic Ocean (Hartmann *et al.* 2011,
491 2012). Also *M. pusilla* strains have been shown to be bacterivores, particularly under low P-
492 conditions (Gonzalez *et al.* 1993; McKie-Krisberg & Sanders 2014) and as such potentially
493 providing an additional source of P (next to SRP and SNP) that may be utilized to promote
494 MpV production upon infection. Studies describing the effects of P-availability on
495 phytoplankton proliferation thus far, merely focused on the effects of SRP-depletion (low
496 ambient SRP-concentrations at the moment of infection). We show here that besides low and
497 relatively constant provision of P, also the source of P appears of ecological relevance for
498 algal virus ecology. It will be interesting to test different algal species (including larger-sized
499 and theoretically more vulnerable to nutrient limitation) for their potential to utilize P (SRP-
500 and SNP-sources) during viral infection.

501

502 **Funding**

503 This work was supported by the Royal Netherlands Institute for Sea Research (NIOZ), an
504 institute of the Netherlands Organisation for Scientific Research (NWO) through a grant to
505 C.B. Furthermore, the authors acknowledge additional support of the European Commission
506 7th Framework Programme project MaCuMBA (Marine Microorganisms: Cultivation
507 Methods for Improving their Biotechnological Applications) contract no. 311957, and the
508 Division for Earth and Life Sciences (ALW) as part of the NWO project CHARLET under
509 contract no. 839.10.513.

510

511 **Acknowledgements**

512 We thank Reinhoud de Blok for technical assistance and Jennifer E. Welsh for assisting with
513 proofreading.

514

515 **Conflict of Interest**

516 The authors declare no conflict of interest

517

518 **References**

519

520 Baudoux A-C, Veldhuis MJ, Witte HJ, Brussaard CPD. Viruses as mortality agents of
521 picophytoplankton in the deep chlorophyll maximum layer during IRONAGES III.

522 *Limnol Oceanogr* 2007;**52**:2519-2529.

523 Baudoux A-C, Brussaard CPD. Influence of irradiance on virus-algal host interactions. *J*

524 *Phycol* 2008;**44**:902-908.

525 Beardall J, Young E, Roberts S. Approaches for determining phytoplankton nutrient
526 limitation. *Aquatic Sciences* 2001;**63**:44-69.

527 Behrenfeld MJ, O'Malley RT, Siegel DA, *et al.* Climate-driven trends in contemporary ocean
528 productivity. *Nature* 2006;**444**:752-755.

529 Benitez-Nelson CR, Buesseler KO. Variability of inorganic and organic phosphorus turnover
530 rates in the coastal ocean. *Nature* 1999;**398**:502-505.

531 Benitez-Nelson CR. The biogeochemical cycling of phosphorus in marine systems. *Earth Sci*
532 *Rev* 2000;**51**:109-135.

533 Bratbak G, Egge JK, Heldal M. Viral mortality of the marine alga *Emiliania huxleyi*
534 (Haptophyceae) and termination of algal blooms. *Mar Ecol Prog Ser* 1993;**93**:39-48.

535 Bratbak G, Jacobsen A, Heldal M, *et al.* Virus production in *Phaeocystis pouchetii* and its
536 relation to host cell growth and nutrition. *Aquat Microb Ecol* 1998;**16**:1-9.

537 Brown CM, Bidle KD. Attenuation of virus production at high multiplicities of infection in
538 *Aureococcus anophagefferens*. *Virology* 2014;**466**:71-81.

539 Brown CM, Lawrence JE, Campbell DA. Are phytoplankton population density maxima
540 predictable through analysis of host and viral genomic DNA content? *J Mar Biol*
541 *Assoc UK* 2006;**86**:491-498.

542 Brown CM, Campbell DA, Lawrence JE. Resource dynamics during infection of *Micromonas*
543 *pusilla* by virus MpV-Sp1. *Environ Microbiol* 2007;**9**:2720-2727.

544 Brussaard C, Mari X, Bleijswijk JV, *et al.* A mesocosm study of *Phaeocystis globosa*
545 (Prymnesiophyceae) population dynamics: II. Significance for the microbial
546 community. *Harmful algae* 2005;**4**:875-893.

547 Brussaard CPD. Optimization of procedures for counting viruses by flow cytometry. *Appl*
548 *Environ Microb* 2004;**70**:1506-1513.

549 Brussaard CPD, Wilhelm SW, Thingstad F, *et al.* Global-scale processes with a nanoscale
550 drive: the role of marine viruses. *ISME J* 2008;**2**:575-578.

551 Casey JR, Lomas MW, Michelou VK, *et al.* Phytoplankton taxon-specific orthophosphate (Pi)
552 and ATP utilization in the western subtropical North Atlantic. *Aquat Microb Ecol*
553 2009;**58**:31-44.

554 Cembella AD, Antia NJ, Harrison PJ. The utilization of inorganic and organic phosphorus
555 compounds as nutrients by eukaryotic microalgae - a multidisciplinary perspective:
556 part 1. *CRC Cr Rev Microbiol* 1984a;**10**:317-391.

557 Cembella AD, Antia NJ, Harrison PJ. The utilization of inorganic and organic phosphorus
558 compounds as nutrients by eukaryotic microalgae - a multidisciplinary perspective:
559 part 2. *CRC Cr Rev Microbiol* 1984b;**11**:13-81.

560 Chen F, Suttle CA. Amplification of DNA polymerase gene fragments from viruses infecting
561 microalgae. *Appl Environ Microbiol* 1995;**61**:1274-1278.

562 Chróst RJ, Siuda W. Ecology of microbial enzymes in lake ecosystems. In: Burns RG, Dick
563 RP (eds.) *Enzymes in the environment: activity, ecology, and applications*. CRC Press,
564 New York; 2002, p. 39-84.

565 Clasen JL, Elser JJ. The effect of host *Chlorella* NC64A carbon : phosphorus ratio on the
566 production of *Paramecium bursaria Chlorella* Virus. *Freshw Biol* 2007;**52**:112-122.

567 Cottrell MT, Suttle CA. Wide-spread occurrence and clonal variation in viruses which cause
568 lysis of a cosmopolitan, eukaryotic marine phytoplankter, *Micromonas pusilla*. *Mar*
569 *Ecol-Prog Ser* 1991;**78**:1-9.

570 Cottrell MT, Suttle CA. Dynamics of a lytic virus infecting the photosynthetic marine
571 picoflagellate *Micromonas pusilla*. *Limnol Oceanogr* 1995;**40**:730-739.

572 Dyrhrman ST, Ammerman JW, Van Mooy BAS. Microbes and the Marine Phosphorus Cycle.
573 *Oceanography* 2007;**20**:110-116.

574 Dyhrman ST, Palenik B. Characterization of ectoenzyme activity and phosphate-regulated
575 proteins in the coccolithophorid *Emiliania huxleyi*. *J Plankton Res* 2003; **25**:1215-
576 1225.

577 Gobler CJ, Hutchins DA, Fisher NS, *et al.* Release and bioavailability of C, N, P, Se, and Fe
578 following viral lysis of a marine chrysophyte. *Limnol Oceanogr* 1997;**42**:1492-1504.

579 Gonzalez JM, Sherr BF, Sherr E. Digestive enzyme activity as a quantitative measure of
580 protistan grazing: the acid lysozyme assay for bacterivory. *Mar Ecol-Prog Ser*
581 1993;**100**:197-206.

582 Graziano LM, Roche J, Geider RJ. Physiological responses to phosphorus limitation in batch
583 and steady-state cultures of *Dunaliella tertiolecta* (chlorophyta): a unique stress
584 protein as an indicator of phosphate deficiency. *J Phycol* 1996;**32**:825-838.

585 Guillard RR, Ryther JH. Studies of marine planktonic diatoms. 1. *Cylotella nana hustedt*, and
586 *Detonula convervacea* (cleve) gran. *Can J Microbiol* 1962;**8**:229.

587 Guo PX, Peterson C, Anderson D. Prohead and DNA-GP3_dependent ATPase Activity of the
588 DNA Packaging protein GP16 of Bacteriophage-PHI-29. *J Mol Biol* 1987;**197**:229-
589 236.

590 Haaber J, Middelboe M. Viral lysis of *Phaeocystis pouchetii*: implications for algal
591 population dynamics and heterotrophic C, N and P cycling. *ISME J* 2009;**3**:430-441.

592 Hansen HP, Koroleff F. Determination of nutrients. In: Grasshoff K, Kremling, K., Erhardt,
593 M. (ed). *Methods of Seawater Analysis 3.ed.* Wiley: Weinheim, Germany; 1999, p.
594 159-228.

595 Harris G. The concept of limiting nutrients. In: Harris G (ed) *Phytoplankton ecology-*
596 *structure, function and fluctuation.* Chapman and Hall, London; 1986, p. 137-164.

597 Hartmann M, Grob C, Scanlan DJ, *et al.* Comparison of phosphate uptake rates by the
598 smallest plastidic and aplastidic protists in the North Atlantic subtropical gyre. *FEMS*
599 *Microbiol Ecol* 2011;**78**:327-335.

600 Hartmann M, Grobb C, Tarran GA, *et al.* Mixotrophic basis of Atlantic oligotrophic
601 ecosystems. *PNAS* 2012;**109**:5756-5760.

602 Healey F, Hendzel L. Indicators of phosphorus and nitrogen deficiency in five algae in
603 culture. *Journal of the Fisheries Board of Canada* 1979;**36**:1364-1369.

604 Jover LF, Effler TC, Buchan A, *et al.* The elemental composition of virus particles:
605 implications for marine biogeochemical cycles. *Nat Rev Microbiol* 2014;**12**:519-528.

606 Juneau P, Lawrence JE, Suttle CA, *et al.* Effects of viral infection on photosynthetic processes
607 in the bloom-forming alga *Heterosigma akashiwo*. *Aquat Microb Ecol* 2003;**31**:9-17.

608 Karl D, Letelier R, Tupas L, *et al.* The role of nitrogen fixation in biogeochemical cycling in
609 the subtropical North Pacific Ocean. *Nature* 1997;**388**:533-538.

610 Karl D, Bjorkman, KM. Dynamics of DOP. In: Hansell D, Carlson, CA (eds.)
611 *Biogeochemistry of Marine Dissolved Matter*. Academic: London, 2002, 249-366.

612 Kimmance SA, Allen MJ, Pagarete A, Martinez Martinez J, Wilson WH: Reduction in
613 photosystem II efficiency during a virus-controlled *Emiliana huxleyi* bloom. *Mar*
614 *Ecol Prog Ser* 2014;**495**:65-76.

615 Lomas MW, Burke AL, Lomas DA, *et al.* Sargasso Sea phosphorus biogeochemistry: an
616 important role for dissolved organic phosphorus (DOP). *Biogeosciences* 2010;**7**:695-
617 710.

618 Lønborg C, Álvarez-Salgado XA. Recycling versus export of bioavailable dissolved organic
619 matter in the coastal ocean and efficiency of the continental shelf pump. *Global*
620 *Biogeochem Cycles* 2012;**26**.

621 Lønborg C, Middelboe M, Brussaard CP. Viral lysis of *Micromonas pusilla*: impacts on
622 dissolved organic matter production and composition. *Biogeochemistry* 2013;**116**:231-
623 240.

624 Løvndal T, Eichner C, Grossart H-P, *et al.* Competition for inorganic and organic forms of
625 nitrogen and phosphorous between phytoplankton and bacteria during an *Emiliania*
626 *huxleyi* spring bloom. *Biogeosciences* 2008;**5**:371-383.

627 Ly J, Philippart CJM, Kromkamp JC. Phosphorus limitation during a phytoplankton spring
628 bloom in the western Dutch Wadden Sea. *Journal of Sea Research* 2014;**88**:109-120.

629 Maat DS, Crawford KJ, Timmermans KR, *et al.* Elevated CO₂ and phosphate limitation favor
630 *Micromonas pusilla* through stimulated growth and reduced viral impact. *Appl*
631 *Environ Microb* 2014;**80**:3119-3127.

632 Maat DS, Bale NJ, Hopmans EC, Damsté JSS, Schouten S, Brussaard CPD. Increasing P-
633 limitation and viral infection impact lipid remodeling of the picophytoplankter
634 *Micromonas pusilla*. *Biogeosciences* 2016;**13**:1667-1676

635 MacIntyre HL, Cullen JJ. Using cultures to investigate the physiological ecology of
636 microalgae. In: Anderson RA (ed) *Algal culturing techniques*, Elsevier Academic
637 press, Amsterdam; 2005, p. 287-327.

638 Marie D, Brussaard CPD, Thyrhaug R, *et al.* Enumeration of marine viruses in culture and
639 natural samples by flow cytometry. *Appl Environ Microb* 1999;**65**:45-52.

640 Martínez Martínez JM, Boere A, Gilg I, *et al.* New lipid envelope-containing dsDNA virus
641 isolates infecting *Micromonas pusilla* reveal a separate phylogenetic group. *Aquat*
642 *Microb Ecol* 2015;**74**:17-28.

643 Mayer JA. Viral infection in marine Prasinophycean alga, *Micromonas pusilla*. *J Phycol*
644 1977;**13**:44-44.

645 McKie-Krisberg ZM, Sanders RW. Phagotrophy by the picoeukaryotic green alga
646 *Micromonas*: implications for Arctic Oceans. *The ISME journal*. 2014;**8**:1953-1961

647 Murray AG, Jackson GA. Viral Dynamics - A model of the effects of size, shape, motion and
648 abundance of single-celled planktonic organisms and other particles. *Mar Ecol-Prog*
649 *Ser* 1992;**89**:103-116.

650 Nalewajko C, Lee K. Light stimulation of phosphate uptake in marine phytoplankton. *Mar*
651 *Biol* 1983;**74**:9-15.

652 Nissimov JI, Napier JA, Allen MJ, Kimmance SA. Intragenus competition between
653 coccolithoviruses: an insight on how a select few can come to dominate many.
654 *Environ. Microbiol* 2016;**18**:133-145.

655 Nolan T, Hands RE, Ogunkolade W, Bustin SA. SPUD : A quantitative PCR assay for the
656 detection of inhibitors in nucleic acid preparations. *Anal Biochem* 2006;**351**:308-310

657 Not F, Latasa M, Marie D, *et al.* A single species, *Micromonas pusilla* (Prasinophyceae),
658 dominates the eukaryotic picoplankton in the western English channel. *Appl Environ*
659 *Microbiol* 2004;**70**:4064-4072.

660 Padan E, Ginzburg D, Shilo M. Reproductive cycle of cyanophage LPP1-G in *Plectonema*
661 *boryanum* and its dependence on photosynthetic and respiratory systems. *Virology*
662 1970;**40**:514.

663 Paul JH, Sullivan MB, Segall AM, *et al.* Marine phage genomics. *Comp Biochem Physiol*,
664 *Part B: Biochem Mol Biol* 2002;**133**:463-476.

665 Porter KG, Feig YS. The use of DAPI for identifying and counting aquatic microflora. *Limnol*
666 *Oceanogr* 1980;**25**:943-948.

667 Rhodes CJ, Martin AP. The influence of viral infection on a plankton ecosystem undergoing
668 nutrient enrichment. *J Theor Biol* 2010;**265**:225-237.

669 Ruttenberg K. The global phosphorus cycle. *Treatise on geochemistry* 2003;**8**:585-643

670 Sarmiento JL, Slater R, Barber R, *et al.* Response of ocean ecosystems to climate warming.
671 *Glob Biogeochem Cycles* 2004;**18**:1-23.

672 Short SM, Short CM. Diversity of algal viruses in various North American freshwater
673 environments. *Aquat Microb Ecol* 2008;**51**:13-21.

674 Slapeta J, Lopez-Garcia P, Moreira D. Global dispersal and ancient cryptic species in the
675 smallest marine eukaryotes. *Mol Biol Evol* 2006;**23**:23-29.

676 Suttle CA. Enumeration and isolation of viruses In: In Kemp PF, Sherr, B. F., Sherr, E. F. *et*
677 *al.* (eds.) *Current Methods in Aquatic Microbial Ecology*. Lewis Publishers Boca
678 Raton, Florida; 1993, p. 121-134.

679 Theodorou ME, Elrifi IR, Turpin DH, *et al.* Effects of phosphorus limitation on respiratory
680 metabolism in the green alga *Selenastrum minutum*. *Plant physiol* 1991;**95**:1089-1095

681 Vanetten JL, Burbank DE, Xia Y, *et al.* Growth cycle of a virus, PBCV-1, that infects
682 *Chlorella*-like algae. *Virology* 1983;**126**:117-125.

683 Vaultot D (1989). CYTOPC: Processing software for flow cytometric data. Signal and Noise
684 2:8. [http://www.sb-roscoff.fr/Phyto/index.php?option=com_content&task=view&id=7](http://www.sb-roscoff.fr/Phyto/index.php?option=com_content&task=view&id=72&Itemid=123)
685 [2&Itemid=123](http://www.sb-roscoff.fr/Phyto/index.php?option=com_content&task=view&id=72&Itemid=123).

686 Waters RE, Chan AT. *Micromonas pusilla* virus - the virus growth-cycle and associated
687 physiological events within the host-cells - host range mutation. *J Gen Virol*
688 1982;**63**:199-206.

689 Weidmann M, Sall AA, Manuguerra J-C, *et al.* Quantitative analysis of particles, genomes
690 and infectious particles in supernatants of haemorrhagic fever virus cell cultures. *Virol*
691 *J* 2011;**8**:81.

692 Wilson WH, Carr NG, Mann NH. The effect of phosphate status on the kinetics of
693 cyanophage infection in the oceanic cyanobacterium *Synechococcus sp.* WH7803. *J*
694 *Phycol* 1996;**32**:506-516.

695 Zheng Q, Chisholm SW. Marine viruses exploit their host's two-component regulatory system
696 in response to resource limitation. *Current Biology* 2012;**22**:124-128.

697

698

1 **Titles and legends to figures**

2

3 Figure 1: Temporal dynamics of *Micromonas pusilla* and viruses upon SRP addition to
4 infected cells. Abundances of non-infected (a) and virally infected (b) *Micromonas pusilla*
5 cultures, as well as the virus MpV (c) over time under different levels of P-limitation. The
6 viral infection experiments were carried out in Erlenmeyer flasks with cultures derived from
7 P-limited chemostats at 0.97 and 0.32 μ_{\max} . Spiked cultures received SRP at an hourly rate
8 until 12h post infection to maintain chemostat P-supply (thereafter at a 6h rate until all cells
9 lysed). P-starved control cultures did no longer receive P-supply. Enriched cultures were
10 spiked with 4 μM SRP to overcome the P-limitation. P-replete cultures were taken along as
11 positive control (non-limited cell physiology). Sampling times were 0, 3, 6, 12, 18, 24, 32, 36,
12 48 and 72 h post infection). All treatments, except the 0.32 μ_{\max} enriched are averages of
13 duplicate cultures (\pm standard deviation).

14

15 Figure 2: Viral genome production represented by total MpV *DNApol* abundance over time
16 (SRP-addition). Spiked cultures received SRP at a 1-6 hourly rate to maintain chemostat P-
17 supply. Grey bars represent dark (night) period. Sampling times were 0, 6, 12, 18, 24, 32, 36,
18 48 and 72 h post infection). All treatments are averages of duplicate cultures (\pm standard
19 deviation).

20

21 Figure 3: Temporal dynamics of *Micromonas pusilla* and viruses in Experiment 2.
22 Abundances of non-infected (a) and virally infected (b) *Micromonas pusilla* cultures, as well
23 as the virus MpV (c) under delayed SRP-addition. The viral infection experiments were
24 carried out in Erlenmeyer flasks with cultures derived from P-limited chemostats (0.97 μ_{\max}).
25 SRP was supplied similar to Experiment 1 but with a 6, 12, 18, 24, 30, 36 and 48h delay,

26 respectively (for proper comparison the concentration SRP added was corrected for the
27 delay). As controls P-limited cultures received either SRP-addition alike Experiment 1
28 (starting 1h p.i.) or not receiving SRP-addition (P-starved). Sampling times were on 0, 6, 12,
29 18, 24, 32, 36, 48, 54 and 72 h post infection). All treatments are averages of duplicate
30 cultures (\pm standard deviation).

31

32 Figure 4: Temporal dynamics of *Micromonas pusilla* and viruses in Experiment 3.

33 Abundances of non-infected (a) and virally infected (b) *Micromonas pusilla* cultures, as well
34 as the virus MpV (c) under SNP-supply. The viral infection experiments were carried out in
35 Erlenmeyer flasks with cultures derived from P-limited chemostats ($0.97\mu_{\max}$). The used SNP-
36 compounds glycerophosphate (GP; closed triangle), adenosine monophosphate (AMP; open
37 circle) and polyphosphate (PP; open triangle) were compared with an SRP-spiked and starved
38 control (closed circle and square, respectively) whereby the supply rates were similar as in
39 Experiment 1. One-time supply (at T0) of 15% v/v virus-free and P-starved MpV lysate (d)
40 was added as additional control treatment to test the potential stimulating effect on MpV
41 production by SNP-compounds in the lysate (note here the different scale of the Y-axis).
42 Sampling times were 0, 6, 12, 18, 24, 32, 36, 48, 60 and 72 h post infection). All treatments
43 are averages of duplicate cultures (\pm standard deviation).

44




45

Table 1: Viral burst size, i.e. the number of newly produced *Micromonas pusilla* viruses (MpV) or viral genomes (*DNApol*) per lysed host cell under the different P-treatments. Experiment 1 demonstrates the effect of spiking the infected cultures with soluble reactive phosphorus (SRP) at a rate similar to their chemostat growth rate (0.97 and $0.32\mu_{\max}$) or a surplus of P (enrichment). Experiment 2 demonstrates the influence of delayed SRP-spiking (from 1 to 48 h post infection) of the $0.97\mu_{\max}$ cultures. Experiment 3 demonstrates the effect of spiking with soluble non-phosphorus (SNP) compounds: Adenosine monophosphate (AMP), Glycerophosphate (GP) and Polyphosphate (PP) and filtered (virus-free) lysate. Average ($n=2$) \pm standard deviations (s.d.).

Treatment	MpV (host cell ⁻¹)	<i>MpV-DNApol</i> (host cell ⁻¹)
<i>Experiment 1 (SRP-spiked)</i>		
$1.0 \mu_{\max}$ P-replete	330 ± 16	2221 ± 174
$0.97\mu_{\max}$ P-enriched	256 ± 4	
$0.32\mu_{\max}$ P-enriched	186^{\dagger}	
$0.97\mu_{\max}$ P-spiked	160 ± 8	341 ± 26
$0.32\mu_{\max}$ P-spiked	141 ± 4	
$0.97\mu_{\max}$ P-starved	74 ± 1	210 ± 34
$0.32\mu_{\max}$ P-starved	76 ± 11	
<i>Experiment 2 (delayed SRP-spike)</i>		
$0.97\mu_{\max} + 1\text{h}$	154 ± 17	
$0.97\mu_{\max} + 6\text{h}$	150 ± 18	
$0.97\mu_{\max} + 12\text{h}$	156 ± 8	
$0.97\mu_{\max} + 18\text{h}$	153 ± 6	
$0.97\mu_{\max} + 24\text{h}$	110 ± 32	
$0.97\mu_{\max} + 30\text{h}$	90 ± 6	
$0.97\mu_{\max} + 36\text{h}$	86 ± 1	
$0.97\mu_{\max} + 48\text{h}$	78 ± 2	
$0.97\mu_{\max}$ P-starved	77 ± 1	
<i>Experiment 3 (SNP-spiked)</i>		
$0.97\mu_{\max} + \text{SRP}$	171 ± 3	
$0.97\mu_{\max} + \text{AMP}$	165 ± 20	
$0.97\mu_{\max} + \text{GP}$	162 ± 2	
$0.97\mu_{\max} + \text{PP}$	177 ± 1	
$0.97\mu_{\max} + \text{lysate}$	94 ± 3	
$0.97\mu_{\max}$ P-starved	75 ± 2	

[†] Replicate failed for technical reasons

Figure 1



- $0.32\mu_{\max}$ starved
- $0.97\mu_{\max}$ starved
- ▼— $0.32\mu_{\max}$ spiked
- △— $0.97\mu_{\max}$ spiked
- $0.32\mu_{\max}$ enriched
- $0.97\mu_{\max}$ enriched
- ◆— P-replete

Figure 2







Figure 3



- 1 h
- 6 h
- ▼ 12 h
- △ 18 h
- 24 h
- 30 h
- ◆ 36 h
- ◇ 48 h
- ▲ P-starved

Figure 4








Figure S1



Virus abundances determined by flow cytometry (FCM) and most probable number (MPN) dilution assay to determine the percentage of infective viruses (MPN/ FCM). Virus infectivity in the lysates was considered 'close to 100%' (error bars show 95% confidence intervals).

Figure S2




Figure S2: The effects of a single SRP-addition (to $0.97\mu_{\max}$ culture) of $300 \mu\text{mol cell}$ at T24 versus frequent SRP-additions of $38 \mu\text{mol cell h}$ every 6h on (A) the growth of cells and (B) the Fv/Fm over 48 h. Note that the total P additions over 48 h are less than what the $0.97\mu_{\max}$ culture would receive to sustain growth at chemostat growth rate (which would be $504 \mu\text{mol cell}$).

Figure S3





Figure S3a: Photosynthetic efficiency (Fv/Fm) of the SRP-spiked (triangles) and SRP-starved (circles) 0.97 μ_{\max} *M. pusilla* cultures during the infection experiment for the infected (closed symbols) and non-infected (open symbols) cultures. r.u. stands for relative units.

Figure S3b: Fv/Fm relative to the non-infected 0.97 μ_{\max} SRP spiked culture to distinguish between viral infection and P-starvation on host Fv/Fm. Hence the other treatments show the effects on Fv/Fm of viral lysis (0.97 μ_{\max} spiked, infected), P-starvation (0.97 μ_{\max} P-starved) and infection + P-starvation (0.97 μ_{\max} starved infected). At T24 the reduction in Fv/Fm by the respective treatments were 11, 18 and 33%. r.u. stands for relative units.

Figure S4




Figure S4: Alkaline phosphatase activity (APA) during an infection experiment of the 0.32 (circles) and $0.97\mu_{\text{max}}$ *M. pusilla* cultures. APA in the non-infected (open symbols) as well as in the infected (closed symbols) start increasing when the dilution of the cultures is stopped (T0).