

## PHOTOPROTECTION AND XANTHOPHYLL-CYCLE ACTIVITY IN THREE MARINE DIATOMS<sup>1</sup>

Céline Dimier, Federico Corato, Ferdinando Tramontano, and Christophe Brunet<sup>2</sup>

Stazione Zoologica “Anton Dohrn,” Villa Comunale, 80121 Naples, Italy

Light is one of the most important factors affecting marine phytoplankton growth, and its variability in time and space strongly influences algal performance and success. The hypothesis tested in this work is that the activity of the xanthophyll cycle and the development of nonphotochemical quenching could be considered a functional trait of algal diversity. If this hypothesis is true, a relationship must exist between fast-activated pigment variations linked to photoprotective behavior and the ecology of the species. This assumption was tested on three diatoms: *Skeletonema marinoi* Sarno et Zingone, *Thalassiosira rotula* Meunier, and *Chaetoceros socialis* Lauder. These three diatoms occupy different ecological niches. Strains of these diatoms were subjected to five changes in irradiance. Xanthophyll-cycle activity, quantum yield of fluorescence, and electron transport rate were the main parameters determined. There were marked interspecific differences in xanthophyll-cycle activity, and these differences were dependent on the light history of the cells. *Chaetoceros socialis* responded efficiently to changing irradiance, which might relate to its dominance during the spring bloom in some coastal areas. In contrast, *T. rotula* responded with a slower photoprotection activation, which seems to reflect its more offshore ecological distribution. The photo-response of *S. marinoi* (a late-winter coastal species blooming in the Adriatic Sea) was light-history dependent, becoming photoinhibited under high light when acclimated to low light, but capable of reaching a high photoprotection level when acclimated to moderate light. Our hypothesis on the photoprotection capacity as a functional trait in microalgae seems to be validated given the results of this study.

**Key index words:** diatoms; NPQ; photoacclimation; quantum yield of fluorescence; xanthophyll cycle

**Abbreviations:** Ax, antheraxanthin;  $\beta$ -car,  $\beta$ -carotene; chl *a*, chlorophyll *a*; chl *c*, chlorophyll *c*; Dd, diadinoxanthin; DES, de-epoxidation state (= Dt/[Dt + Dd] ratio); Dt, diatoxanthin; *F*, instantaneous fluorescence value; *F*<sub>m</sub>, maximal fluorescence in dark-adapted state; *F*'<sub>m</sub>, maximal fluorescence in light-adapted state; *F*<sub>o</sub>, minimal

fluorescence in dark-adapted cells; *F*'<sub>q</sub>/*F*'<sub>m</sub>, effective quantum yield of fluorescence (light-adapted cells); *F*<sub>s</sub>, fluorescence intensity at steady state at given actinic level (RLC measurement); Fuco, fucoxanthin; *F*<sub>v</sub>/*F*<sub>m</sub>, quantum yield of fluorescence (dark-adapted cells); HL, high light; LHC, light-harvesting complex; LL, low light; ML, moderate light; NPQ, nonphotochemical quenching; PAM, pulse amplitude modulated; PFD, photon flux density; RLC, rapid light curve; SV<sub>m</sub>, Stern–Volmer expression for NPQ calculation; Vx, violaxanthin; Zx, zeaxanthin

The photoacclimation process in marine phytoplankton covers a large range of temporal scales and consists of many biological modifications, from molecular to morphological (Falkowski and La Roche 1991, Prézelin et al. 1991), which drive variations in the ecophysiological behavior of a species. At oversaturating irradiances, excess photons absorbed by the light-harvesting complexes (LHCs) can be dissipated (e.g., through thermal dissipation occurring as nonphotochemical quenching [NPQ]). This phenomenon involves xanthophyll-cycle activity (Ruban et al. 2004), corresponding to the reversible de-epoxidation of violaxanthin (Vx) into zeaxanthin (Zx) through antheraxanthin (Ax) for green algae and higher plants (Demmig et al. 1988). In chl *c*-containing algae, the photoprotective cycle involves the conversion of the monoepoxide diadinoxanthin (Dd) into the de-epoxidized diatoxanthin (Dt; Arsalane et al. 1994, Olaizola et al. 1994), even though the Vx, Ax, and Zx pigments are present in low quantity in this taxon (Lohr and Wilhelm 1999, 2001). Recent studies on diatoms indicate that Dt is responsible for NPQ (Jakob et al. 2001, Lavaud et al. 2002a,b,c, 2003), which can reach very high values in *Phaeodactylum tricornutum* Bohlin, much higher than the values reported for terrestrial plants (Lavaud et al. 2002b, 2004).

Comparative studies showed that the maximal value of nonphotochemical quenching (NPQ) is taxon dependent (Casper-Lindley and Björkman 1998) and can differ markedly even among different species of, for example, diatoms (Lavaud et al. 2004). However, relationships between photoprotective response diversity and algal ecological properties

<sup>1</sup>Received 31 March 2006. Accepted 23 March 2007.

<sup>2</sup>Author for correspondence: e-mail brunet@szo.it.

remained hypothetical. Recently, some studies revealed an adaptation of the photosynthetic properties of algae to the ecological niche in which they grow. Strzepek and Harrison (2004) showed that acclimation capacities in two diatoms depended on biochemical characteristics of their photosynthetic apparatus, which in turn are related to the ecosystem properties. In our study, we wanted to test the assumption that the capacity of short-term pigment variations, activated as a photoprotective process, depends on the ecological characteristics of the species (i.e., on the ecological niche properties). For instance, the short-term photoresponse is probably more often activated in highly mixed water columns (e.g., coastal) than in more hydrodynamically stable (e.g., oceanic) ecosystems, and this characteristic could therefore be one factor determining the competitive performance and the success of some algae. In relation to this hypothesis, the short-term photoresponse would act as a functional component of phytoplankton diversity.

Diatoms were chosen as models since they generally dominate algal communities in coastal ecosystems, accounting for ~40% of the marine primary production (Armbrust et al. 2004), and they present a high level of morphological, ecological, and molecular diversity (Medlin and Kaczmarska 2004). The short-term photoprotective capacities, mainly controlled by xanthophyll-cycle activation and development of NPQ as well as variations of fluorescence parameters, were followed during five different light-shift experiments. A comparative study of the dynamics of  $V_x$ ,  $A_x$ , and  $Z_x$  among the three diatoms was also undertaken to test their potential role in Dd-Dt xanthophyll-cycling intensity, since these pigments could be intermediary pigments in the photoprotective pathway activation (Lohr and Wilhelm 2001).

Previous photophysiological studies were generally carried out using very strong photon flux densities (PFDs; e.g.,  $>1000 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; Casper-Lindley and Björkman 1998, Lavaud et al. 2002b, 2003) to dissect the main traits of the photoresponse. In this study, to test the “ecological-physiological relationship” assumption, we used three diatoms representative of different ecological niches, applying PFDs ranging from 40 to  $400 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ , which correspond to values often experienced by cells during their growth in coastal ecosystems. The three species used as models are *Skeletonema marinoi*, *Chaetoceros socialis*, and *Thalassiosira rotula*. The first species blooms in the coastal area of the Adriatic Sea in the late winter (Totti and Artegiani 2001, Sarno et al. 2005), while the second co-occurs with other species during the spring bloom in some coastal areas (Marshall and Ranasinghe 1989, Totti and Artegiani 2001), and the third grows in more offshore ecosystems (Syvertsen 1977). Different photoresponses were determined, depending mainly on the species but also on the

growth PFD acclimation of the cells. According to the light condition and the species, the xanthophyll cycle was quickly activated (e.g., exponential synthesis of Dt), followed a slow and linear activation, or was almost inactivated when photoinhibition occurred.

## MATERIALS AND METHODS

**Algal cultures and experimental design.** Nonaxenic strains of *S. marinoi*, *T. rotula*, and *C. socialis*, isolated from the Adriatic Sea (*S. marinoi*) or from the Tyrrhenian Sea (*T. rotula* and *C. socialis*), were provided by the Laboratory of Ecophysiology (Stazione Zoologica “A. Dohrn,” Naples, Italy). Cells of *S. marinoi* are about 7–8  $\mu\text{m}$  diameter, forming long chains. *Thalassiosira rotula* is a chain-forming species with cell diameter of 25  $\mu\text{m}$ , while *C. socialis* forms colonies with cells of about 8–9  $\mu\text{m}$  diameter.

The three species were cultured semicontinuously at 20°C in 3 L Pyrex glass bottles (Microglass, Naples, Italy) containing natural sterile seawater amended with f/2 nutrients (Guillard and Rytner 1962), with a 12:12 light:dark (L:D) photoperiod at low light (LL,  $40 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) or moderate light (ML,  $100 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ). Light was provided with white fluorescent tubes (58W/840; Phillips, Paris, France), and irradiance was measured with a laboratory PAR sensor (4  $\Pi$ , QSL-2100; Biospherical Instruments Inc., San Diego, CA, USA). Before the beginning of the experiments, cultures were kept in exponential growth phase for at least 10 generations by repeatedly diluting them with fresh medium. Experiments were performed twice independently and consisted of the following light changes: the LL-cultured cells were shifted to 200 or 400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  and the ML-cultured cells were shifted to 40, 200, or 400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ . During each experiment, one flask was kept at the same irradiance as growth conditions. Experiments lasted 3 or 4 h and samples for pigment, variable fluorescence, and flow cytometry measurements were taken at high time frequency.

**Pigment content.** Ten milliliters of culture was filtered through 25 mm GF/F filters (Whatman, Maidstone, UK) and immediately stored in liquid nitrogen for later pigment analysis (Casotti et al. 2005). Briefly, pigment filters were extracted in 3 mL 100% methanol, and 500  $\mu\text{L}$  of  $1 \text{ mol} \cdot \text{L}^{-1}$  ammonium acetate was added to the 1 mL pigment extract for 5 min before the analysis by a Hewlett Packard series 1100 HPLC (Hewlett-Packard, Wilmington, NC, USA). A 3  $\mu\text{m}$  C<sub>8</sub> BDS column (ThermoHypersil, Runcorn, UK) was used, and the mobile phase was composed of two solvents: (A) methanol and aqueous ammonium acetate (70:30), and (B) methanol. Pigments were also measured at 440 nm using a photodiode array detector (model DAD series 1100; Hewlett Packard), which gave the 400 to 700 nm spectrum for each detected pigment. Pigments were detected by fluorometry (series 1100 fluorometer; Hewlett Packard), using a 410 nm excitation wavelength and a 665 nm emission wavelength, and quantified using standards from the VKI (Water Quality Institute, Horsholm, Denmark).

**Fluorescence measurements.** Variable fluorescence was measured using a Phyto-PAM (pulse-amplitude-modulated) fluorometer (Heinz Walz GmbH, Effeltrich, Germany). Quantum yield of fluorescence for the light-adapted ( $F'_q/F'_m$ , with  $F'_q = F'_m - F$ ) and for the 15 min dark-adapted samples ( $F_v/F_m$ , with  $F_v = F_m - F_o$ ) was determined through the measurements of the minimum fluorescence level,  $F$  (or  $F_o$ ), and the maximum fluorescence level,  $F'_m$  (or  $F_m$ ), respectively. The latter corresponds to the maximum fluorescence measured after a saturation pulse of bright red light (655 nm,  $2400 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) applied for 450 ms. The pulse

was saturating since the increase of its duration did not increase the fluorescence yield in any of the three species studied.

The NPQ was estimated during the rapid light curve (RLC) measurement performed three or four times per experiment, as follows: After the 15 min dark adaptation,  $F_v/F_m$  was estimated, and cells were exposed to 10 levels of increasing actinic light (from 8 to 1500  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ , 2 min at each PFD level). At the end of each illuminated step, the quantum yield of fluorescence was measured ( $F'_q/F'_m = [F'_m - F'_s]/F'_m$ ), and the relative electron transport rate (rETR) was estimated as follows:

$$\text{rETR} = (F'_q/F'_m) \cdot \text{PFD} \cdot 0.5 \quad (1)$$

where PFD was the incident irradiance ( $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ), and 0.5 corresponded to the assumption that half of the absorbed light is distributed to PSII. The RLCs were fitted with the equation of Eilers and Peeters (1988) to retrieve the photosynthetic parameters  $\text{rel}\alpha_{\text{etr}}$ ,  $E_k$ , and  $\text{rETR}_{\text{max}}$ . The NPQ was calculated at each light level by the Stern–Volmer expression (Krause and Weis 1991):

$$\text{NPQ} = \text{SV}_m = (F'_q/F'_m) - 1 \quad (2)$$

where  $\text{SV}_m$  is the Stern–Volmer quenching coefficient of the maximum fluorescence yield, and  $F'_m$  and  $F_m$  are the maximum fluorescence of light-adapted (after each light level) and dark-adapted samples (measured at the beginning of the RLC), respectively (Villareal 2004).

## RESULTS

### *Acclimation of Chaetoceros socialis to changing PFD.*

The ML-growing cells of *C. socialis* ( $100 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) had a lower  $F_v/F_m$  and photosynthetic parameter values than the LL-growing cells (Table 1) and presented a greater capacity for energy dissipation as revealed by the higher  $\text{NPQ}_{\text{max}}$  (5.60 vs. 1.12; Fig. 1, a and b). This finding was in relation to the 4-fold greater xanthophyll pool (Dd + Dt) in the ML-acclimated cells (Fig. 2, a and b). The light-limitation experienced by cells growing under  $40 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ , as revealed by the  $E_k$  value (Table 1), might be a determinant for the increase of the chl *a*  $\cdot \text{cell}^{-1}$  content with respect to the ML condition (Table 2). The accessory pigments versus chl *a* ratio did not change between the two PFD-growing conditions, except that  $\beta\text{-car}/\text{chl } a$  doubled in the ML-growing cells (Table 2).

Photoresponses of ML-growing cells to the high PFD condition ( $400 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) were

characterized by an increase in the  $\text{rETR}_{\text{max}}$  and  $E_k$  (by 40%) and a decrease in  $\text{NPQ}_{\text{max}}$  (–44%; Fig. 1a). It is noteworthy that a quenching of  $F_m$  occurred (first point of the evolution of maximum fluorescence parameter from RLC; Fig. 1c) between the start ( $T_o$ ) and the end of the experiment (240 min), which could indicate dark-NPQ. A possible cause could be chlororespiration (Jakob et al. 2001) generating a transthylakoid pH gradient, allowing the maintenance of some xanthophylls in the de-epoxidized state (Villareal 2004). Indeed, the diatoxanthin pool was high since it followed a 3-fold increase mainly in the first minutes after the shift from ML to HL (Fig. 2a). This Dt increase, mainly due to de novo synthesis would be responsible for the strong quenching of  $F$  and  $F'_m$  ( $P < 0.01$ ; Fig. 2c).

Photoresponse of LL-growing cells to high PFD was characterized by a strong decrease in PSII operating efficiency ( $F'_q/F'_m$ ; Fig. 2d), which could be related to xanthophyll-cycle activity (Fig. 2b). Diatoxanthin synthesis was slower than for the ML-growing cells shifted to HL ( $k = 0.37$  and  $1.61 \text{ min}^{-1}$ , respectively) and was primarily due to the conversion of preexisting Dd, followed by a de novo synthesis of Dd and Dt. A development of dark-NPQ was also revealed by the quenching of  $F_m$  between  $T_o$  and the end of experiment (180 min) from RLC measurement (Fig. 1d).

The photobiological properties of ML-growing cells shifted to low PFD remained rather similar, except for the 50% reduction in diatoxanthin and the 30% increase in chl *a*  $\cdot \text{cell}^{-1}$  (data not shown).

*Acclimation of Skeletonema marinoi to changing PFD.* The maximum PSII photochemical efficiencies, the relative photosynthetic efficiencies, and the  $\text{NPQ}_{\text{max}}$  were almost similar in ML- and LL-growing cells ( $\approx 0.65$ ,  $0.266 \mu\text{mol e}^- \cdot \mu\text{mol photons}^{-1}$ , and 0.8 to 1.2, respectively; Table 1 and Fig. 3, a and b). The light-saturation of cells at  $100 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  with respect to their light limitation at  $40 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  (see  $E_k$  values, Table 1) could be a determinant for the decrease in cellular chl *a* quota and fucoxanthin (Fuco; Table 2). The latter probably induced a lowering of cell absorption capacity, which might account for the reduction of  $\text{rETR}_{\text{max}}$  (Table 2), together with the 2- to 3-fold greater pool of Dd and Dt in

TABLE 1. Fluorescence and photosynthetic parameters of the three diatoms grown under low ( $40 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) and moderate ( $100 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) photon flux density (PFD) conditions.

	$F_v/F_m$	$F'_q/F'_m$	$E_k$ ( $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ )	$\text{rETR}_{\text{max}}$ ( $\mu\text{mol e}^- \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ )	$\text{rel}\alpha$ ( $\mu\text{mol e}^- \cdot \mu\text{mol photons}^{-1}$ )
<i>Chaetoceros socialis</i> PFD: 40	0.70	0.72	98.9	28.9	0.292
<i>C. socialis</i> PFD: 100	0.56	0.57	77	18.7	0.243
<i>Skeletonema marinoi</i> PFD: 40	0.65	0.63	160.5	42.8	0.266
<i>S. marinoi</i> PFD: 100	0.64	0.66	118.4	31.5	0.266
<i>Thalassiosira rotula</i> PFD: 40	0.66	0.67	93.7	25.9	0.277
<i>T. rotula</i> PFD: 100	0.50	0.36	67.7	14.4	0.213




FIG. 1. Evolution of NPQ (a, b) and  $F_s$  and  $F_m'$  (c, d) from an upward shift in irradiance, from rapid light curve measurements performed on *Chaetoceros socialis*. Cells were shifted to 400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  from growth in medium light (100  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; a, c) and low light (40  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; b, d). All the data are means of two values. Numbers on the graph represent the time (min) after the light shift.

ML-growing cells compared to LL-growing cells (Fig. 4, a and b).

The shift in ML-growing cells to the high PFD condition (400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) induced a strong accumulation of Dt (Fig. 4a) correlated to a reduction of PSII operating efficiency ( $P < 0.05$ ; Fig. 4c). The lowering of  $\text{NPQ}_{\text{max}}$  with time (Fig. 3a) might be related to the strong decrease of  $F_m$  (first point of the RLC, Fig. 3c). This absence of  $F_m$  recovery could indicate the development of dark-NPQ. Photoinhibition could be another hypothesis, but the minimal fluorescence  $F$  did not increase (Fig. 4c), as generally observed in such cases (Müller et al. 2001). On the contrary, the LL-growing cells shifted to the high PFD condition showed a low Dt synthesis while Dd accumulated (Fig. 4b). This pigment response might account for the small decrease in the PSII operating efficiency (Fig. 4d). The increase in the minimal level of fluorescence ( $F_0$  [+14%], data not shown) might indicate the closure

of the reaction centers related to photoinhibition (Krause et al. 1990). The RLC measurement revealed an increase in  $\text{NPQ}_{\text{max}}$  (Fig. 3c) and the absence of dark-NPQ (Fig. 3d).

The photobiological properties of ML-growing cells shifted to low PFD remained rather similar, except for a 42% increase of chl  $a \cdot \text{cell}^{-1}$  (data not shown).

#### Acclimation of *Thalassiosira rotula* to changing PFD.

The ML-growing cells of *T. rotula* (100  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) were light-saturated ( $E_k = 67.7 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ) in contrast to cells growing at low PFD ( $E_k = 93.7 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ). This feature was responsible for the greater xanthophyll pigment pool (Fig. 6, a and b), as well as the low PSII quantum yield of fluorescence (Table 1) and the higher  $\text{NPQ}_{\text{max}}$  (Fig. 5, a and b) compared to the low PFD condition. The ML-growing cells appeared to be already photoinhibited. Therefore, when shifted to high PFD (400  $\mu\text{mol}$




FIG. 2. Evolution of xanthophyll ratios (a, b) and  $F$ ,  $F_m$ , and  $F_q/F_m'$  (c, d) from an upward shift in irradiance on *Chaetoceros socialis*. Cells were shifted to 400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  from growth in medium light (100  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; a, c) and low light (40  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; b, d). Pigment ratios in  $\text{mol} \cdot 10 \text{ mol chl } a^{-1}$ . All the data are means of two values.

TABLE 2. Mean ( $n = 6$ ) and standard deviation (in brackets) of chl  $a \cdot \text{cell}^{-1}$  ( $\text{mol chl } a \cdot \text{cell}^{-1}$ ) and pigment ratios ( $\text{mol pigment} \cdot 10 \text{ mol chl } a^{-1}$ ) for the three species after 10 generations of acclimation at 40 (LL) or 100 (ML)  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ .

	Chl $a \cdot \text{cell}^{-1}$	$\beta\text{-car}/\text{chl } a$	$(Vx + Ax + Zx)/\text{chl } a$	Fucox/chl $a$
<i>Chaetoceros socialis</i> PFD: 40	$8.86 \times 10^{-16}$ ( $0.9 \times 10^{-16}$ )	0.356 (0.025)	0.173 (0.095)	5.33 (0.48)
<i>C. socialis</i> PFD: 100	$7.26 \times 10^{-16}$ ( $0.8 \times 10^{-16}$ )	0.602 (0.080)	0.183 (0.179)	5.94 (0.75)
<i>Skeletonema marinoi</i> PFD: 40	$6.49 \times 10^{-16}$ ( $0.7 \times 10^{-16}$ )	0.477 (0.053)	0.338 (0.061)	6.78 (0.648)
<i>S. marinoi</i> PFD: 100	$4.23 \times 10^{-16}$ ( $0.7 \times 10^{-16}$ )	0.637 (0.244)	0.708 (0.206)	8.84 (3.77)
<i>Thalassiosira rotula</i> PFD: 40	$3.65 \times 10^{-14}$ ( $1.8 \times 10^{-14}$ )	0.227 (0.027)	0.156 (0.016)	4.18 (0.45)
<i>T. rotula</i> PFD: 100	$4.04 \times 10^{-14}$ ( $0.9 \times 10^{-14}$ )	0.253 (0.017)	0.135 (0.043)	4.27 (0.21)

$\beta\text{-car}$ ,  $\beta$ -carotene; Vx, violaxanthin; Zx, zeaxanthin; Ax, antheraxanthin; Fucox, fucoxanthin; LL, low light; ML, moderate light.

photons  $\cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ), they did not accumulate a large amount of Dt (Fig. 6a), which in turn implied a slight decrease of  $F_q/F_m'$  (Fig. 6c). The  $\text{NPQ}_{\text{max}}$  was strongly reduced with time (Fig. 5a), which might be related to development of dark-NPQ as suggested by the strong decrease of  $F_m$  (first point of the RLC, Fig. 5c). On the contrary, the LL-growing cells shifted to the high PFD condition slowly synthesized Dt (Fig. 6b), which was not correlated to the rapid decrease of  $F_m'$  ( $P > 0.05$ ; Fig. 6d). The low  $\text{NPQ}_{\text{max}}$  was reduced with time (Fig. 5b) due to dark-NPQ and/or photoinhibition

(see Fig. 5d), the latter being revealed by the increase of  $F_s$  along the RLC (+56%).

The ML-growing cells shifted to low PFD revealed a quick relaxation of PSII. The quenching of  $F_m$  and the dark-NPQ disappeared after 7 min, while Dt/chl  $a$  was reduced by 50% (data not shown).

#### DISCUSSION

Our results highlight a strong variability of photo-responses developed by these three centric diatoms. The high photophysiological diversity among the




FIG. 3. Evolution of NPQ (a, b) and  $F_s$  and  $F_m'$  (c, d) from an upward shift in irradiance, from rapid light curve measurements performed on *Skeletonema marinoi*. Cells were shifted to 400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  from growth in medium light (100  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; a, c) and low light (40  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; b, d). All the data are means of two values. Numbers on the graph represent the time (min) after the light shift.

three species does not match their phylogenetic relationships (Medlin and Kaczmarek 2004), since *S. marinoi* and *T. rotula* are closely related, whereas *C. socialis* is more distant. The ecological properties of the species could represent one of the factors responsible for photophysiological differences. Responses varied from a quick and efficient photoacclimation to photoinhibition, and the physiological processes activated by cells concern xanthophyll cycling and NPQ as well as dark-NPQ related to the formation of a transthylakoidal  $\Delta\text{pH}$  in the dark, for instance by chlororespiration (Nixon 2000, Villareal 2004). Intraspecific photoresponse variability mainly depends on the light history of the species (i.e., on the growth PFD condition) as already observed for other species (Casper-Lindley and Björkman 1998). The higher content of diatoxanthin in ML-growing cells strongly increases the further development of NPQ as observed in *C. socialis* and *T. rotula*, in which the  $\text{NPQ}_{\text{max}}$  increased 5 and

10 times, respectively, compared to the LL-growing cells.

*Photoprotective behavior as a functional trait in diatoms.* In *T. rotula*, the low capacity for short-term photoprotective acclimation prohibits this species from coping with rapidly increasing irradiance when grown under low PFD. Its very low capacity of NPQ ( $\text{NPQ}_{\text{max}} = 0.17$ ) probably determines photoinhibition of cells under high PFD, as revealed by the increase in  $F_o$  (+154%), reflecting the closure of reaction centers (Müller et al. 2001, Voronova et al. 2002). This species when acclimated under low- or moderate-PFD conditions exhibits the slowest Dt synthesis under high PFD, which might reflect a slow formation of a transthylakoidal  $\Delta\text{pH}$ . A low cytochrome *b<sub>6</sub>f* content could be hypothesized to be responsible for this feature, as found in another oceanic *Thalassiosira* species (*T. oceanica* Hasle, Strzepek and Harrison 2004). The presence of NPQ in ML-growing cells of *T. rotula* and the stability of




FIG. 4. Evolution of xanthophyll ratios (a, b) and  $F$ ,  $F_m'$ , and  $F_q/F_m'$  (c, d) from an upward shift in irradiance on *Skeletoneema marinoi*. Cells were shifted to  $400 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  from growth in medium light ( $100 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; a, c) and low light ( $40 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; b, d). Pigment ratios in  $\text{mol} \cdot 10 \text{ mol chl } a^{-1}$ . All the data are means of two values.

Dt/chl *a* during transfer to  $400 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  suggest that these processes act on a longer-term scale, as already seen in studies carried out on different *Thalassiosira* species (Meyer et al. 2000, Lavaud et al. 2004, Strzepek and Harrison 2004). *Thalassiosira rotula* is also peculiar in that it was the only one of the three species studied that does not appear to have a significant change in the most common long-term acclimation processes, such a decrease in chl *a* content or an increase in the accessory pigments linked to photoprotection (Dd,  $\beta$ -car, Vx, Ax, or Zx; see below). This species appears, therefore, to be able to cope with "limited" increasing light by activating slow physiological responses. This physiological property fits well with the low physical motions present in the offshore, seasonally stratified ecosystems. The very fast recovery to the low-light state of PSII is in agreement with this hypothesis since these very large cells may also sink to escape from the very high-light level of the surface in such stable water columns.

There is marked interspecific diversity in the capacity of NPQ development, as already shown for species belonging to different groups (Casper-

Lindley and Björkman 1998) and among diatoms (Lavaud et al. 2004). The highest value reported for *C. socialis* ( $\text{NPQ}_{\text{max}} = 5.6$ ) is in the range of the highest values reported by other authors on diatoms and other microalgal groups (Casper-Lindley and Björkman 1998, Lavaud et al. 2004, Ruban et al. 2004). The quick activation of the xanthophyll cycle and the strong accumulation of Dt in cells exposed to high PFD are responsible for this high NPQ and probably contribute to the dark-NPQ measured in this species. *Chaetoceros socialis* had the greatest physiological plasticity of the three species studied, being able to efficiently acclimate to high PFD when grown under either low or moderate PFD. We interpret this plasticity as a functional trait that contributes to its ecological success in the turbulent environment where it grows. This species dominates the algal community during the spring bloom in the northern Adriatic Sea (Totti and Artegiani 2001) and in coastal waters of the northwest Atlantic (Marshall and Ranasinghe 1989). *Chaetoceros* as a genus might be seen as having high photophysiological plasticity since other species have the same acclimation behavior with respect to light variations




FIG. 5. Evolution of NPQ (a, b) and  $F_s$  and  $F'_m$  (c, d) from an upward shift in irradiance, from rapid light curve measurements performed on *Thalassiosira rotula*. Cells were shifted to 400  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  from growth in moderate light (100  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; a, c) and low light (40  $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; b, d). All the data are means of two values. Numbers on the graph represent the time (min) after the light shift.

(*C. muelleri* Lemmerm., Olaizola and Yamamoto 1994; *C. gracilis* Pant., Kashino and Kudoh 2003).

In contrast, *S. marinoi* does not show high capacity for energy dissipation, which is also strongly dependent on the light history of the cells. The LL-growing cells are not capable of accumulating high amounts of Dt when transferred under high light, probably leading to photoinhibition. In the LL-growing cells, the lack or the delay of Dt synthesis could be due to the low content of Dd ( $\sim 0.065 \text{ mol Dd} \cdot \text{mol chl } a^{-1}$ ) most probably strongly attached to FCPs and therefore not really convertible. A low thylakoid lumen pH due to the delay of photoprotection might inactivate the de-epoxidase (Olaizola et al. 1994) because of conformational change of the enzyme (Emanuelsson et al. 2003). When the new synthesis of Dd occurs (i.e., after 30 min), efficient photoprotection is thus prevented. On the other hand, the ML-growing cells activate a strong synthesis of Dt, reaching  $0.80 \text{ mol} \cdot \text{mol chl } a^{-1}$  under high PFD. This

response is relatively slow due to the delay of about 30 min before the Dd de novo synthesis (Fig. 4a), which is needed when this pigment reaches the minimal value of  $0.065 \text{ mol} \cdot \text{mol chl } a^{-1}$ . This value, identical for both LL- and ML-growing cells of *S. marinoi*, could correspond to a nonconvertible Dd pool. A lag of 30 min would be the time needed to activate the photoprotective biosynthetic pathway, leading to the synthesis of new Dd molecules. The high Dt content could be responsible for the development of dark-NPQ and the nonreversibility of  $F'_m$  in the dark (see Fig. 3c). This dark-NPQ is a bias for the interpretation of NPQ measurement based on  $F_m$ , the maximal fluorescence of dark-adapted cells. The maintenance of a  $\Delta\text{pH}$  in the dark could be an advantage under fluctuating irradiance, allowing the rapid development of energy dissipation upon exposure to oversaturating PFD (Lavaud et al. 2002c). Over long timescales, in ML-growing cells, the strong increase in Dd and Dt is accompanied by an






FIG. 6. Evolution of xanthophyll ratios (a, b) and  $F$ ,  $F_m'$ , and  $F_q/F_m'$  (c, d) from an upward shift in irradiance on *Thalassiosira rotula*. Cells were shifted to  $400 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  from growth in medium light ( $100 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; a, c) and low light ( $40 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; b, d). Pigment ratios in  $\text{mol} \cdot 10 \text{ mol chl } a^{-1}$ . All the data are means of two values.

important reduction in chl *a* content, indicating a probable decrease in the number of thylakoids. A reduction in the antenna size is not proposed since  $\alpha_{\text{rel}}$  is the same between LL- and ML-growing cells. These changes in pigment content act as a relevant acclimation process, optimizing the maximum PSII photochemical efficiency (see parameters in Table 1). The physiological plasticity of *S. marinoi* is “intermediate” between *T. rotula* and *C. socialis*. This feature is in agreement with the ecological characteristics of this species growing in the coastal area of the Adriatic Sea (Totti and Artegiani 2001). It dominates the late winter algal bloom, characterized by very low-light conditions, and can be present—without dominating—in the springtime community, when light increases.

**Pool of accessory xanthophyll pigments.** Low variations in the concentration of Vx, Ax, and Zx at our experimental timescale (<4 h) support findings by Lohr and Wilhelm (1999) that these pigments have no direct function in energy dissipation. At long-term scales, they might have a structural role, for instance, by decreasing the thylakoid membrane fluidity (Havaux and Gruszecki 1993). The higher




FIG. 7. Relationship between the  $(Vx + Ax + Zx + \beta\text{-car})/Dd$  ratio measured for LL- and ML-acclimated cells for the three species (*Chaetoceros socialis*, *Skeletonema marinoi*, and *Thalassiosira rotula*). Data are means of two values. The regression was significant ( $P < 0.05$ ;  $r^2 = 0.98$ ;  $y = 0.72x + 0.56$ ). Ax, antheraxanthin;  $\beta\text{-car}$ ,  $\beta$ -carotene; Vx, violaxanthin; Zx, zeaxanthin.

content of these pigments (per mol chl *a*), together with  $\beta$ -car, Dd, and Dt, in the ML-growing cells with respect to LL-growing cells corresponds to their role as intermediaries in the photoprotective biosynthetic pathway (Lohr and Wilhelm 1999, 2001). The significant relationship between the  $(Vx + Ax + Zx + \beta\text{-car})/\text{Dd}$  ratio estimated in the ML- and LL-growing cells (Fig. 7) reveals that the quantity of precursor pigments is linked to the quantity of Dd, and vice versa. The slope of the linear regression (Fig. 7) being  $<1$  indicates that when this ratio increases, the differences between the LL- and ML-growing cells tend to be smaller, perhaps indicating a saturation of the loci for these pigments in the PSII. The low capacity for changing the xanthophyll content inside the PSII between the two PFD conditions could be responsible for the slow photo-response capacity exhibited by the algae, for instance *T. rotula*. It appears that the faster the kinetics of the xanthophyll cycle, the lower the  $(Vx + Ax + Zx + \beta\text{-car})/\text{Dd}$  ratio (reaching values  $<1$ , Fig. 7), indicating a trend toward net Dd accumulation. This could be a key factor in quickly activating the conversion from Dd to Dt, which is one of the main short-term processes of photoprotection of PSII.

The authors thank the Laboratory of Ecophysiology of the SZN and F. Esposito for sharing the three strains of diatoms, as well as R. Casotti for her flow cytometry experience. Wiebe Kooistra is kindly acknowledged for the critical reading of the manuscript. The two referees and L. Franklin are acknowledged for all their comments and suggestions to improve this manuscript. C. D. was supported by a PhD grant from SZN. This paper has been produced within the framework of the Marine Biodiversity and Ecosystem Functioning (MARBEF) Network of Excellence, which is funded in the Community's Sixth Framework Programme (contract no. GOCE-CT-2003-505446). This publication is contribution number MPS-07027 of MARBEF.

- Armbrust, E. V., Berges, J. A., Bowler, C., Green, B. R., Martinez, D., Putnam, N. H., Zhou, S., et al. 2004. The genome of the diatom *Thalassiosira pseudonana*: ecology, evolution and metabolism. *Science* 306:79–86.
- Arsalane, W., Rousseau, B. & Duval, J. C. 1994. Influence of the pool size of the xanthophyll cycle on the effects of light stress in a diatom: competition between photoprotection and photoinhibition. *Photochem. Photobiol.* 60:237–43.
- Casotti, R., Mazza, S., Brunet, C., Vantrepotte, V., Ianora, A. & Miralto, A. 2005. Growth inhibition and toxicity of the diatom aldehyde 2-trans-4-trans decadienal on *Thalassiosira weissflogii* (Bacillariophyceae). *J. Phycol.* 41:7–20.
- Casper-Lindley, C. & Björkman, O. 1998. Fluorescence quenching in four unicellular algae with different light-harvesting and xanthophyll-cycle pigments. *Photosynth. Res.* 56:277–89.
- Demmig, B., Winter, K., Kruger, A. & Czygan, F. C. 1988. Zeaxanthin and the heat dissipation of excess light energy in *Nerium oleander* exposed to a combination of high light and water stress. *Plant Physiol.* 87:17–24.
- Eilers, P. H. C. & Peeters, J. C. H. 1988. A model for the relationship between light intensity and the rate of photosynthesis in phytoplankton. *Ecol. Model.* 42:199–215.
- Emanuelsson, A., Eskling, M. & Akerlund, H.-E. 2003. Chemical and mutational modification of histidines in violaxanthin de-epoxidase from *Spinacia oleracea*. *Physiol. Plant.* 119:97–104.
- Falkowski, P. G. & La Roche, J. 1991. Molecular biology in studies of ocean processes. *Int. Rev. Cytol.* 128:261–303.
- Guillard, R. R. R. & Ryther, J. H. 1962. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacea* (Cleve) Gran. *Can. J. Microbiol.* 8:229–38.
- Havaux, M. & Gruszecki, W. I. 1993. Heat- and light-induced chlorophyll *a* fluorescence changes in potato leaves containing high or low levels of the carotenoid zeaxanthin: indications of a regulatory effect of zeaxanthin on membrane fluidity. *Photochem. Photobiol.* 58:607–14.
- Jakob, T., Goss, R. & Wilhelm, C. 2001. Unusual pH-dependence of diadinoxanthin de-epoxidase activation causes chlororespiratory induced accumulation of diatoxanthin in the diatom *Phaeodactylum tricornutum*. *J. Plant Physiol.* 158:383–90.
- Kashino, Y. & Kudoh, S. 2003. Concerted response of xanthophyll-cycle pigments in a marine diatom, *Chaetoceros gracilis*, to shifts in light condition. *Phycol. Res.* 51:168–72.
- Krause, G. H., Somersalo, S., Zumbush, E., Weyers, B. & Laasch, H. 1990. On the mechanism of photoinhibition in chloroplasts. Relationships between changes in fluorescence and activity of photosystem II. *J. Plant Physiol.* 136:472–9.
- Krause, G. H. & Weis, E. 1991. Chlorophyll fluorescence and photosynthesis: the basics. *Annu. Rev. Plant Physiol.* 42:313–49.
- Lavaud, J., Rousseau, B. & Etienne, A. L. 2002a. In diatoms, a thylakoid proton gradient alone is not sufficient to induce a non-photochemical fluorescence quenching. *FEBS Lett.* 253:163–6.
- Lavaud, J., Rousseau, B. & Etienne, A. L. 2003. Enrichment of the light-harvesting complex in diadinoxanthin and implications for the non-photochemical quenching in diatoms. *Biochemistry* 42:5802–8.
- Lavaud, J., Rousseau, B. & Etienne, A. L. 2004. General features of photoprotection by energy dissipation in planktonic diatoms (Bacillariophyceae). *J. Phycol.* 40:130–7.
- Lavaud, J., Rousseau, B., van Gorkom, H. & Etienne, A. L. 2002b. Influence of the diadinoxanthin pool size on photoprotection in the marine planktonic diatom *Phaeodactylum tricornutum*. *Plant Physiol.* 129:1398–406.
- Lavaud, J., van Gorkom, H. J. & Etienne, A. L. 2002c. Photosystem II electron transfer cycle and chlororespiration in planktonic diatoms. *Photosynth. Res.* 74:51–9.
- Lohr, M. & Wilhelm, C. 1999. Algae displaying the diadinoxanthin cycle also possess the violaxanthin cycle. *Proc. Natl. Acad. Sci. U. S. A.* 96:8784–9.
- Lohr, M. & Wilhelm, C. 2001. Xanthophyll synthesis in diatoms: quantification of putative intermediates and comparison of pigment conversion kinetics with rate constants derived from a model. *Planta* 212:382–91.
- Marshall, H. G. & Ranasinghe, J. A. 1989. Phytoplankton distribution along the eastern coast of the USA. VII. Mean cell concentration and standing crop. *Cont. Shelf Res.* 9:153–64.
- Medlin, L. K. & Kaczmarek, I. 2004. Evolution of the diatoms: V. Morphological and cytological support for the major clades and a taxonomic revision. *Phycologia* 43:245–70.
- Meyer, A., Tackx, M. & Daro, N. 2000. Xanthophyll cycling in *Phaeocystis globosa* and *Thalassiosira* sp.: a possible mechanism for species succession. *J. Sea Res.* 43:373–84.
- Müller, P., Li, X.-P. & Niyogi, K. K. 2001. Non-photochemical quenching. A response to excess light energy. *Plant Physiol.* 125:1558–66.
- Nixon, P. J. 2000. Chlororespiration. *Phil. Trans. R. Soc. Lond. B Biol. Sci.* 355:1541–7.
- Olaizola, M., La Roche, J., Kolber, Z. & Falkowski, P. G. 1994. Non-photochemical fluorescence quenching and the diadinoxanthin cycle in a marine diatom. *Photosynth. Res.* 41:357–70.
- Olaizola, M. & Yamamoto, H. Y. 1994. Short-term response of the diadinoxanthin cycle and fluorescence yield to high irradiance in *Chaetoceros muelleri* (Bacillariophyceae). *J. Phycol.* 30:606–12.
- Prézelin, B. B., Tilzer, M. M., Schofield, O. & Haese, C. 1991. The control of the production process of phytoplankton by the

- physical structure of the aquatic environment with special reference to its optical properties. *Aquat. Sci.* 53:136–86.
- Ruban, A. V., Lavaud, J., Rousseau, B., Guglielmi, G., Horton, P. & Etienne, A. L. 2004. The super-excess energy dissipation in diatom algae: comparative analysis with higher plants. *Photosynth. Res.* 82:165–75.
- Sarno, D., Kooistra, W. H. C. F., Medlin, L. K., Percopo, I. & Zingone, A. 2005. Diversity in the genus *Skeletonema* (Bacillariophyceae). II. An assessment of the taxonomy of *S. costatum*-like species, with the description of four new species. *J. Phycol.* 41:151–76.
- Strzepek, R. F. & Harrison, P. J. 2004. Photosynthetic architecture differs in coastal and oceanic diatoms. *Nature* 431:689–92.
- Syvertsen, E. 1977. *Thalassiosira rotula* and *T. gravida*: ecology and morphology. *Nova Hedwigia Beih.* 54:99–112.
- Totti, C. & Artegiani, A. 2001. Phytoplankton time series in the northern Adriatic Sea: the Senigallia transect (1988–1994). *Arch. Oceanogr. Limnol.* 22:107–12.
- Villareal, T. A. 2004. Single-cell amplitude modulation fluorescence measurements of the giant diatom *Ethmodiscus* (Bacillariophyceae). *J. Phycol.* 40:1052–61.
- Voronova, E. N., Volkova, E. V., Kazimirko, Y. V., Chivkunova, O. B., Merzlyak, M. N., Pogosyan, S. I. & Rubin, A. B. 2002. Response of the photosynthetic apparatus of the diatom *Thalassiosira weissflogii* to high irradiance light. *Russ. J. Plant Physiol.* 49:311–9.