



## Phylogenetics and biogeography of the Balkan ‘sand gobies’ (Teleostei: Gobiidae): vulnerable species in need of taxonomic revision

MAARTEN P. M. VANHOVE<sup>1\*</sup>, ALCIBIADES N. ECONOMOU<sup>2</sup>, STAMATIS ZOGARIS<sup>2</sup>,  
MAARTEN H. D. LARMUSEAU<sup>1,3,4</sup>, SOFIA GIAKOUMI<sup>2</sup>, ELENI KALOGIANNI<sup>2</sup>,  
FILIP A. M. VOLCKAERT<sup>1</sup> and TINE HUYSE<sup>1</sup>

<sup>1</sup>Laboratory of Animal Diversity and Systematics, Department of Biology, Katholieke Universiteit Leuven, Ch. Deberiotstraat 32, B-3000 Leuven, Belgium

<sup>2</sup>Institute of Inland Waters, Hellenic Centre for Marine Research, PO Box 712, 190 13, Anavyssos, Greece

<sup>3</sup>Laboratory of Forensic Genetics and Molecular Archaeology, UZ Leuven, Kapucijnenvoer 33, B-3000 Leuven, Belgium

<sup>4</sup>Department of Human Genetics, Campus Gasthuisberg, Katholieke Universiteit Leuven, Herestraat 49, B-3000 Leuven, Belgium

Received 30 June 2011; revised 27 July 2011; accepted for publication 27 July 2011

Within the Atlantic–Mediterranean region, the ‘sand gobies’ are abundant and widespread, and play an important role in marine, brackish, and freshwater ecosystems. They include the smallest European freshwater fish, *Economidichthys trichonis*, which is threatened by habitat loss and pollution, as are several other sand gobies. Key to good conservation management is an accurate account of the number of evolutionary significant units. Nevertheless, many taxonomic and evolutionary questions remain unresolved within the clade, and molecular studies are lacking, especially in the Balkans. Using partial 12S and 16S mitochondrial ribosomal DNA sequences of 96 specimens of at least eight nominal species (both freshwater and marine populations), we assess species relationships and compare molecular and morphological data. The results obtained do not support the monophyly of *Economidichthys*, suggesting the perianal organ to be a shared adaptation to hole-brooding rather than a synapomorphy, and urge for a taxonomic revision of *Knipowitschia*. The recently described *Knipowitschia montenegrina* seems to belong to a separate South-East Adriatic lineage. *Knipowitschia milleri*, an alleged endemic of the Acheron River, and *Knipowitschia* cf. *panizzae*, are shown to be very closely related to other western Greek *Knipowitschia* populations, and appear conspecific. A distinct Macedonian–Thessalian lineage is formed by *Knipowitschia thessala*, whereas *Knipowitschia caucasica* appears as an eastern lineage, with populations in Thrace and the Aegean. The present study combines the phylogeny of a goby radiation with insights on the historical biogeography of the eastern Mediterranean, and identifies evolutionary units meriting conservation attention. © 2011 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2012, 105, 73–91.

**ADDITIONAL KEYWORDS:** *Economidichthys* – ichthyology – *Knipowitschia* – Mediterranean – *Pomatoschistus*.

### INTRODUCTION

The Balkans are considered as Europe’s prime centre of endemism and a hotspot of biodiversity (Kryštufek

& Reed, 2004). Their glacial refugium role renders the Balkans, as well as other regions (e.g. the Iberian and Italian peninsulae), particularly interesting both in a biogeographical and a conservation framework, leading to a highly diverse fauna (Durand, Persat & Bouvet, 1999a). Nevertheless, their biogeographical history and the processes leading to this high

\*Corresponding author.  
E-mail: maarten.vanhove@bio.kuleuven.be

diversity are poorly investigated or understood (Kryštufek & Reed, 2004). An important mechanism underlying species richness, and hence a useful target in evolutionary research and biogeography, is adaptive radiation, as exemplified by phylogenetic work on fishes such as the Neotropical monophyletic goby tribe Gobiosomatini (seven-spined gobies) (Rüber, Van Tassell & Zardoya, 2003; Taylor & Hellberg, 2005; Larmuseau *et al.*, 2011). The goby family (Teleostei, Perciformes, Gobiidae) represents one of the most (if not the most) speciose fish families worldwide, with about 1950 described species (Nelson, 2006). Gobies have adapted to a wide range of habitats, including marine, brackish, and freshwater, and show an impressive range of mating behaviours (Miller, 1986; Kottelat & Freyhof, 2007). The eastern Mediterranean, in particular the Balkans and the Black and Caspian basins, hosts the highest diversity of gobies in Europe, including many endemic species (Kottelat & Freyhof, 2007; Kovačić & Patzner, 2011), dominated in freshwater by the so-called 'sand gobies'. The Balkan 'sand gobies' present a useful framework for studying patterns of diversity and evolution because of: (1) their species richness and degree of endemism; (2) the importance of species radiations in evolutionary research; (3) the understudied biogeography of the eastern Mediterranean in general (Zogaris, Economou & Dimopoulos, 2009); and (4) the usefulness of freshwater fishes as biogeographical tools (Hugueny, 1989; Heino, 2001). Additional intrinsic features (e.g. perianal organ; conservation importance) render them scientifically interesting. Surprisingly, considering their prevalence, abundance and potential with respect to biogeographical studies in the Balkans, molecular studies on these species are still rare.

The monophyletic sand goby clade (Huysse, Van Houdt & Volckaert, 2004), morphologically characterized by features of the head canals and vertebrae, includes the genera *Pomatoschistus* Gill, 1863, *Knipowitschia* Iljin, 1927, *Economidichthys* Bianco, Bullock, Miller & Roubal, 1987, and *Gobiusculus* Duncker, 1928. *Orsinigobius* Gandolfi, Marconato & Torricelli, 1986, and possibly *Hyrceanogobius* Iljin, 1928, are synonymous with *Knipowitschia* (Miller, 2004b). These small benthic species occupy parts of the eastern Atlantic Ocean, the Mediterranean Sea, and the Balkan and Ponto-Caspian continental regions (Miller, 2004c). The diversity of sand gobies is underestimated, as illustrated by numerous recent descriptions (Kovačić, 2005; Kovačić & Šanda, 2007; Miller & Šanda, 2008; Miller, 2009) and new records (Van Neer *et al.*, 1999; Stefanni, 2000; Turan *et al.*, 2005; Ozen, Irmak & Bilecenoglu, 2007; Mejri *et al.*, 2009). Several sand gobies deserve conservation attention as a result of anthropogenic threats, such as


wetland degradation, water extraction, land reclamation, structural changes of water bodies, introduction of predators, and pollution (Economidis, 1995; Maitland & Crivelli, 1996; Kottelat & Freyhof, 2007; Özcan, 2009). Furthermore, because endangered species are important for environmental assessment and for prioritization in conservation management (Smith & Darwall, 2006), the freshwater sand gobies, with their wide habitat range and high level of endemism, might be considered as useful indicator species for monitoring the conservation of Mediterranean inland aquatic ecosystems. Scientific attention could perhaps render them attractive enough to be regarded as flagship species for the Mediterranean ichthyofauna and ambassadors for wetlands in the region.

Biodiversity surveys and conservation measures are seriously hampered by ambiguities in species delineations and assignments (Bickford *et al.*, 2006; Economou *et al.*, 2007). Based on morphology and morphometrics, many controversies exist about sand goby taxonomy, such as the systematic validity of taxa diagnosed by characters that may exhibit intraspecific phenotypic plasticity (e.g. reduction in head canals and squamation; Kovačić, 2005). The status of nominal genera currently recognized is likely to change (Kottelat & Freyhof, 2007). Molecular studies could help to clarify the relationships and evolutionary history of sand gobies (Penzo *et al.*, 1998; Huysse *et al.*, 2004; Miller & Šanda, 2008). Phylogenetic analyses allow us to pinpoint some issues of taxonomic interest, and to identify evolutionary significant units (ESUs) for conservation. Although the definition of this concept, which should represent the minimal unit to address conservation matters, is somewhat problematic (Vogler & DeSalle, 1994; Crandall *et al.*, 2000; Fraser & Bernatchez, 2001), ESUs coupled with historical biogeography have already been defined in Mediterranean sand gobies (Stefanni & Thorley, 2003).

## MATERIAL AND METHODS

### SPECIMEN COLLECTION

We sampled Economidis' goby *Economidichthys pygmaeus* (Holly, 1929), Trichonis goby *Economidichthys trichonis* Economidis & Miller, 1990, Caucasian dwarf goby *Knipowitschia caucasica* (Kawrajsky in Berg 1916), Miller's dwarf goby *Knipowitschia milleri* (Ahnelt & Bianco, 1990), Thessaly goby *Knipowitschia thessala* (Vinciguerra, 1921), Adriatic dwarf goby *Knipowitschia pannizae* (Verga, 1841) (*sensu* the Evinos population identified by Ahnelt & Bianco, (1990); hereafter referred to as *Knipowitschia* cf. *panizae*), Montenegro dwarf goby *Knipowitschia*



**Figure 1.** Map showing sand goby sampling localities, except for the location on the River Morača. More details on sampling sites and species are provided in Table 1.

*montenegrina* Kovačić and Šanda, 2007 and marbled goby *Pomatoschistus marmoratus* (Risso, 1810). Valid taxon names (according to currently accepted descriptions and identifications), status, sampling locations, habitats, number of individuals sequenced, and

GenBank accession numbers are shown in Figure 1 and Table 1. Taxon and author names are used *sensu* Eschmeyer (2010). Gobies were collected using hand and seine nets and backpack electrofishing in a range of freshwater and brackish habitats. Specimens were

**Table 1.** Populations included in the phylogenetic analyses, including their species ascription, collection location, hydrographic drainage basin, habitat description, IUCN Red List category (Smith & Darwall, 2006), and number of sequenced specimens

| Species  | Site number | Location                         | Drainage basin | Habitat description                     | Number of sequences | GenBank accession number                        |
|--|-------------|----------------------------------|----------------|---|---------------------|---|
| <i>Economidichthys pygmaeus</i>                                    | 1           | Kryoneri Springs                 | Evinos         | Freshwater karstic springs with reedbed | 3                   | JN388369-72                                     |
| (freshwater)   | 2           | Kalamas Delta                    | Kalamas        | Deltaic freshwater channel              | 2                   | JN388373-5                                      |
| Least concern  | 3           | Lake Trichonis                   | Acheloo        | Large freshwater lake-shore             | 4                   | JN388376-7                                      |
|  | 4           | Skala Springs                    | Louros         | Spring-fed freshwater channel           | 2                   | JN388378-80                                     |
|  | 5           | Agios Georgios Springs           | Arachthos      | Spring-fed freshwater stream            | 2                   | JN388381-2                                      |
|  | 6           | Chiliadou Springs                | Mornos         | Spring-fed freshwater stream            | 4                   | JN388383-4                                      |
|  | 7           | Astakos Springs                  | Astakos        | Spring-fed brackish stream              | 2                   | JN388385-6                                      |
|  | 8           | Vlychos Springs, Vonitsa Lagoon  | Vlychos        | Spring-fed brackish channel             | 2                   | JN388387-8                                      |
|  | 9           | Aitoliko Stream                  | Acheloo        | Spring-fed freshwater creek             | 8                   | GQ398131-2                                      |
| <i>Economidichthys trichonis</i><br>(freshwater) Endangered        | 3           | Lake Trichonis, site Myrtias     | Acheloo        | Large deep freshwater lake -shore       | 4                   | Larmuseau<br><i>et al.</i> (2010)<br>JN388364-8 |
| <i>Knipowitschia caucasica</i><br>(euryhaline)                     | 10          | Lake Vidra (Karystos Wetland)    | Rigia          | Semi-natural freshwater lake            | 2                   | JN388321-2                                      |
|  | 11          | Rigia Stream                     | Rigia          | Brackish river-mouth                    | 2                   | JN388321-2                                      |
| <i>Knipowitschia</i> sp.<br>(brackish water)                       | 12          | Kalogria Laspoloutra             | Prokopos       | Artificial pool beside lagoon           | 2                   | JN388306-7                                      |
|  | 13          | Kalogria Prokopos Lagoon         | Prokopos       | Brackish marsh                          | 2                   | JN388308-9                                      |
| <i>Knipowitschia</i> cf. <i>panizzae</i><br>(freshwater, brackish) | 14          | Evinos Lagoon (Klisova)          | Evinos         | Brackish lagoon                         | 2                   | JN388310-2                                      |
| <i>Knipowitschia</i> sp.<br>(brackish water)                       | 15          | Aitoliko Lagoon                  | Acheloo        | Brackish lagoon                         | 2                   | JN388313-5                                      |
|  | 16          | Acheloo Estuary                  | Acheloo        | Brackish lagoon                         | 1                   | JN388316,<br>JN399059                           |
| <i>Knipowitschia caucasica</i><br>(euryhaline)                     | 17          | Evos River, site Feres           | Evos           | Freshwater canalized river              | 2                   | JN388323-4                                      |
|  | 18          | Strymon River, Strymoniko bridge | Strymon        | Riparian riverine marshes               | 2                   | JN388319-20                                     |

|    |   |   |                   |  |    |   |
|----|---|---|-------------------|--|----|---|
| 19 | <i>Knipowitschia milleri</i><br>(freshwater to slightly<br>brackish) Critically<br>endangered | Vosvozis Stream near Lake<br>Mitrkou                | Filiouris         | Lowland freshwater river                         | 2  | JN388325-7  |
| 20 | <i>Knipowitschia thessala</i><br>(freshwater) Endangered                                      | Acheron Delta                                       | Acheron           | Brackish channel                                 | 12 | GQ398133-7<br>Larmuseau<br><i>et al.</i> (2010)<br>JN388303-5<br>JN388317-8 |
| 21 |   | Enipeas tributary                                   | Pinios            | Small lowland river                              | 1  | JN388317-8  |
| 22 |   | Pinios River, main course, near<br>Farkadona bridge | Pinios            | Large lowland river                              | 4  | JN388317-8  |
| 32 | <i>Knipowitschia montenegrina</i>   | Morača River, site Ponari<br>(Montenegro)           | Morača            | Deltaic channel, near estuary                    | 3  | JN388301-2  |
| 23 | <i>Pomatoschistus marmoratus</i>  | Livadaki  | Karystos<br>Bay   | Marine beach                                     | 2  | JN388331-3  |
| 24 | (marine and brackish water)   | Chiliadou estuary                                   | Mornos            | Marine beach                                     | 2  | JN388357-8  |
| 25 |   | Kryoneri estuary                                    | Evinos            | Marine beach                                     | 2  | JN388353-6  |
| 26 |   | Evinos estuary                                      | Evinos            | River-mouth                                      | 2  | JN388359-60   |
| 27 |   | Drepano beach                                       | Kalamas           | Marine beach                                     | 2  | JN388361-3  |
| 28 |   | Acheloos estuary                                    | Acheloos          | Marine beach, outflow of<br>brackish lagoon      | 8  | JN388342-52   |
| 29 |   | Plakias Bay   | Southern<br>Crete | Marine sandy beach                               | 2  | JN388334-7  |
| 30 |   | Frangokastello cove                                 | Southern<br>Crete | Marine sandy beach                               | 2  | JN388338-41   |
| 31 |   | Giorgiopoulis stream-mouth                          | Kourna            | Brackish spring-fed stream;<br>near stream-mouth | 2  | JN388328-30   |

Site numbers correspond to the map (Fig. 1), excluding number 32, which is the only site outside Greece. For details on the hydrographic basins of most sites, see Economou *et al.* (2007). Note that *K. caucasica* is not mentioned in Smith & Darwall (2006) and that *K. montenegrina* was only described subsequently; hence, no conservation status is mentioned.

preserved in 99% ethanol until further processing. Voucher specimens are part of the reference collection of the Hellenic Centre for Marine Research (Institute of Inland Waters, Anavyssos, Greece) and are available from the authors on request.

#### EXTRACTION AND AMPLIFICATION OF MITOCHONDRIAL 12S AND 16S rDNA

DNA was extracted using the NucleoSpin Tissue Kit (Macherey-Nagel) in accordance with the manufacturer's instructions. Polymerase chain reaction (PCR) was performed using a GeneAmp PCR system 2700 thermocycler (Applied Biosystems). The reaction volume of 25 µL contained 1 × PCR buffer (Eurogentec), 200 µM of dNTPs (Amersham Pharmacia Biotech), 2 mM of MgCl<sub>2</sub> (50 mM) (Eurogentec), 0.5 U of *Taq*-polymerase (Eurogentec), 0.8 µM of each primer (Eurogentec), 1 µL of template DNA, topped up with milli-Q water (Millipore). Primers used were 16SH 5'-CCGGTCTGAACTCAGATCACGT-3', 16SL 5'-CGCCTGTTTATCAAAAACAT-3' (Palumbi *et al.*, 1991), 12SH 5'-TGACTGCAGAGGGTGACGGGGCG GTGTGT-3' and 12SL 5'-AAAAAGCTTCAAACCTGG GATTAGATACCCCACTAT-3' (Kocher *et al.*, 1989).

After an initial denaturation of 3 min at 95 °C, samples were subject to 35 cycles of 30 s at 95 °C, 30 s at 52 °C (16S) or 64 °C (12S) and 50 s at 72 °C. After a final elongation of 7 min at 72 °C, samples were cooled down to 4 °C. PCR products were purified with NucleoFast (Macherey-Nagel) in accordance with the manufacturer's instructions. They were sequenced using the same primers as above, with the Big Dye Terminator 3.1 kit (Applied Biosystems), applying a 1/8 dilution of the Big Dye Terminator sequencing protocol. Finally the products were run on an ABI PRISM 3130 Avant Genetic Analyser automated sequencer (Applied Biosystems).

#### SEQUENCE ALIGNMENT AND PHYLOGENETIC ANALYSIS

Sequences available from other sand goby species were retrieved from GenBank (Table 2). *Padogobius bonelli* (Bonaparte, 1846) (synonymous with *Padogobius martensii*) and *Padogobius nigricans* (Canestrini, 1867) were selected as outgroup taxa. As has been extensively shown, these species belong to the same clade as, for example, *Gobius* Linnaeus, 1758 and *Millerigobius* Bath, 1973 and are firmly distinct from the sand gobies (Penzo *et al.*, 1998; Huyse *et al.*, 2004; Giovannotti

**Table 2.** GenBank sequences of sand gobies with habitat description and sampling location

| Species  | Location                           | GenBank accession number | Reference                        |
|--|------------------------------------|--------------------------|----------------------------------|
| <i>Pomatoschistus quagga</i> (Heckel, 1837) (marine)               | Northern Adriatic                  | AF067264 and -77         | Penzo <i>et al.</i> (1998)       |
| <i>Knipowitschia punctatissima</i> (freshwater)                    | Veneto Streams basin (Italy)       | AF067260 and -73         |                                  |
| <i>Pomatoschistus minutus</i> (Pallas, 1770) (brackish, marine)    | Venice Lagoon (Italy)              | AJ616816 and -33         | Huyse <i>et al.</i> (2004)       |
| <i>Pomatoschistus microps</i> (Krøyer, 1838) (brackish, marine)    | Ostend (North Sea, Belgium)        | AJ616813 and -30         |                                  |
| <i>Pomatoschistus marmoratus</i> (brackish, marine)                | Ostend (North Sea, Belgium)        | AJ616811 and -28         |                                  |
| <i>Pomatoschistus canestrinii</i> (freshwater, brackish)           | Venice Lagoon (Italy)              | AJ616809 and -26         |                                  |
| <i>Pomatoschistus lozanoi</i> (de Buen, 1923) (brackish, marine)   | Dalmatia (Croatia)                 | AJ616818 and -35         |                                  |
| <i>Pomatoschistus norvegicus</i> (Collett, 1902) (marine)          | North Sea (Belgium)                | AJ616815 and -32         |                                  |
| <i>Pomatoschistus pictus</i> (Malm, 1865) (marine)                 | Bergen (Norway)                    | AJ616814 and -31         |                                  |
| <i>Knipowitschia panizzae</i> (freshwater, brackish)               | Bergen (Norway)                    | AJ616817 and -34         |                                  |
| <i>Gobiusculus flavescens</i> (Fabricius, 1779) (brackish, marine) | Po Delta and Venice Lagoon (Italy) | AJ616806 and -22-4       |                                  |
| <i>Padogobius bonelli</i> (freshwater)                             | Trondheim (Norway)                 | AJ616836 and -19         |                                  |
| <i>Padogobius nigricans</i> (freshwater)                           | Sentino River (Italy)              | EF218637 and -52         | Giovannotti <i>et al.</i> (2007) |
|  | Tevere River (Italy)               | AF067257 and -70         | Penzo <i>et al.</i> (1998)       |

*et al.*, 2007; Vanhove *et al.*, 2011). Following the incongruence-length difference test (Farris *et al.*, 1995) carried out for the same regions by Huyse *et al.* (2004), 12S and 16S rDNA fragments were concatenated into a single dataset. Sequences were aligned by MUSCLE, version 3.8 (Edgar, 2004) using default distance measures and sequence weighting schemes. The resulting alignment was trimmed with trimAl, version 1.2b (Capella-Gutiérrez, Silla-Martínez & Gabaldón, 2009), making use of an automated trimming algorithm optimized for subsequent maximum likelihood (ML) phylogenetic tree reconstruction. jModelTest, version 0.1.1 (Guindon & Gascuel, 2003; Felsenstein, 2005; Posada, 2008) was used to estimate the optimal model of molecular evolution, using a ML optimized starting tree. The TPM2uf+I+Γ model was selected based on the corrected Akaike information criterion (AICc) (Hurvich & Tsai, 1989). For subsequent implementation in phylogenetic software, we chose the GTR+I+Γ model (gamma shape parameter: 0.48; proportion of invariable sites: 0.58) (Tavaré, 1986; Rodríguez *et al.*, 1990), representing the commonly available one with the highest AICc score. A model-averaged phylogeny demonstrated that hardly any nodes displayed phylogenetic uncertainty because of model selection. Hence, we feel confident that the slight overparametrization will not influence our analyses. Under this model, a ML search was carried out in PhyML, version 3.0 (Guindon & Gascuel, 2003), assessing nodal support through 1000 bootstrap samples using the nearest-neighbour interchange branch swapping algorithm. The model was also used in Bayesian analysis, implemented in MrBayes, version 3 (Huelsenbeck & Ronquist, 2001; Ronquist & Huelsenbeck, 2003). Posterior probabilities were calculated over  $4 \times 10^6$  generations. At that point, stationarity of the Markov chain was reached, as indicated by a standard deviation of split frequencies of 0.005, by the absence of a trend in the log probabilities plotted over generations, and by a potential scale reduction factor (Gelman & Rubin, 1992) converging to 1. The Markov chain was sampled every other 100 generations. We discarded 25% of the samples as 'burn-in'. For calculation of ML genetic distances according to the optimized model, and for phylogeny reconstruction using the maximum parsimony (MP) method, PAUP\* 4.01b (Swofford, 2001) was used, with the PaupUp interface (Calendini & Martin, 2005). A heuristic search was performed using 1000 replicates of nearest-neighbour interchange branch swapping; gaps were treated as a fifth character.


Because the monophyly of *Economidichthys* is under discussion, we performed constraint analyses to assess whether enforcing monophyly on this genus significantly changed the likelihood of the tree. We used the Shimodaira–Hasegawa (Shimodaira & Hasegawa,

1999) and Kishino–Hasegawa topological tests (Kishino & Hasegawa, 1989) (one-sided, in view of the inclusion of the ML tree; Goldman, Anderson & Rodrigo, 2000; Schmidt, 2009), implemented in PAUP\*.

To assess the phylogenetic content of the dataset, a likelihood-mapping analysis (Strimmer & von Haeseler, 1997) was carried out in TREE-PUZZLE, version 5.2 (Schmidt *et al.*, 2002). Using the same software, a likelihood-ratio-test (LRT) was performed to test the molecular clock hypothesis, and a clock-constrained tree was constructed by quartet puzzling, with the parameters resulting from jModelTest. Based on mutation rates of mitochondrial rDNA of  $0.5\text{--}0.56\% \times 10^{-6}$  years (sicydiine gobies; Keith *et al.*, 2011),  $0.6\% \times 10^{-6}$  years (*Mugilogobius* Smitt, 1900 and *Pandaka* Herre, 1927 gobies; Mukai, Suzuki & Nishida, 2004),  $0.7\% \times 10^{-6}$  years (the cyprinid *Telestes souffia* (Risso, 1827); Costedoat & Gilles, 2009),  $1\% \times 10^{-6}$  years (Centropominae; Tringali *et al.*, 1999),  $1.11\% \times 10^{-6}$  years (*Syngnathus* Linnaeus, 1758 pipefish; Wilson, 2006),  $1.4\% \times 10^{-6}$  years ('sand gobies'; Huyse *et al.*, 2004) and  $1.3\text{--}1.6\% \times 10^{-6}$  years (Blenniidae; Almada *et al.*, 2009), we used constant mutation rates of 0.5% and  $1.5\% \times 10^{-6}$  years to draw a calibrated tree. 12S and 16S are relatively conservative markers compared to several other mitochondrial (mt)DNA genes (Wan *et al.*, 2004) and poikilotherm animals often display relatively slow evolutionary rates (Johns & Avise, 1998). However, because we are dealing (over a smaller phylogenetic range) with small-bodied taxa with a short generation time, relatively higher mutation rate estimations are more realistic (Martin & Palumbi, 1993; Gillooly *et al.*, 2005; Wilson, 2006). Hence, we did not take unconventionally low values (such as  $0.23\% \times 10^{-6}$  years for gymnotiforms and mormyriiforms; Alves-Gomes, 1999) into account. To check whether there was any noise left in the dataset as a result of unstable characters, we partitioned the data in ten mutation rate categories using TIGER, version 1.0 (Cummins & McInerney, 2011) and removed the fastest evolving category with the StripBin script. Likelihood-mapping was subsequently performed on the resulting alignment. Alignment files were converted using ALTER, version 1.2 (Glez-Peña *et al.*, 2010) and trees were drawn with FigTree, version 1.3 (<http://tree.bio.ed.ac.uk/software/figtree>) and MEGA, version 5 (Tamura *et al.*, 2011).

## RESULTS

Ninety-six specimens were sequenced, belonging to at least eight nominal species: *K. montenegrina*, *K. cf. panizzae*, *K. caucasica*, *K. milleri*, *K. thessala*, *E. pygmaeus*, *E. trichonis*, and *P. marmoratus*. The length of



**Figure 2.** Likelihood-mapping analysis of the dataset (815 bp sequences of 12S and 16S rDNA for all sand gobies) demonstrating the strong phylogenetic content.


the amplified products was approximately 444 and 584 bp for 12S and 16S rDNA, respectively. After trimming, a total of 815 bp were retained in the dataset. This data matrix contained 204 variable sites of which 145 were identified as parsimony-informative. Fifty-five unique haplotypes were identified. Conspecific individuals originating from distinct geographical populations often differed in one to four positions (except for rather 'distant' *P. marmoratus* populations from the Adriatic, Euboea, and Crete). The MP tree had a consistency index of 0.56, a retention index of 0.81 and a tree length of 501. Likelihood-mapping showed that 95.1% of quartets were fully ('tree-like') and 2.4% partly ('network-like') resolved, whereas 2.6% remained unresolved (Fig. 2). After removal of the fastest evolving 30 nucleotide positions, this changed to 91.3%, 3.7%, and 5%, respectively. Hence, the resolution of the dataset did not decrease as a result of fast evolving sites. The LRT showed that a clock-like evolution of the sequences should not be rejected at a significance level of 0.05. A clock-constrained quartet puzzling tree with timescales corresponding to two various mutation rates is shown in Figure 3. Using the upper ( $1.5\% \times 10^{-6}$  years) and lower ( $0.5\% \times 10^{-6}$  years) assumed mutation rates, the first divergence event, leading to the major lineages of sand gobies, took place between 2.9 and 9 Mya). A first diversification within the different genera is estimated to date back to 2–6 Mya.

*Knipowitschia* appear paraphyletic in all analyses (Fig. 4). However, except for *Knipowitschia punctatissima* (Canestrini, 1864), which did not cluster with its alleged congeners but rather with *Pomatoschistus canestrinii* (Ninni, 1883), all taxa belonging to the genus are consistently grouped together. This *Knipowitschia* cluster is estimated to have diverged between 0.6 and 2.8 Mya. It is subdivided into distinct lineages with appreciable statistical support (posterior probability of at least 95 under Bayesian


inference, or a bootstrap value of at least 75 in all other analyses): (1) *K. montenegrina* from the Morača River in Montenegro, perhaps being part of a south-eastern Adriatic lineage; (2) *K. panizzae* from the Adriatic basin (Po Delta and Venice Lagoon); (3) an Ionian clade: *Knipowitschia* sp. (Aitoliko Lagoon, Acheloos Lagoon, Prokopos Lagoon at Kalogria), *K. cf. panizzae* (Evinos Lagoon) and *K. milleri* (Acheron Delta); (4) the Macedonian–Thessalian *K. thessala* from the Pinios River; and (5) the Thracian–Aegean *K. caucasica* (Strymon, Evros and Vosvozis Rivers) and *K. caucasica* from the island of Euboea (Rigia Stream and Lake Vidra). Within the Ionian *Knipowitschia* lineage, the Aitoliko, Acheloos, Kalogria, and Evinos (*K. cf. panizzae*) populations never differed by more than 0.6% from one another, and never more than 1.3% from *K. milleri* (see the Supporting information, Table S1). Populations of *K. caucasica* did not differ more than 0.8% gamma-corrected GTR distance from each other. *Knipowitschia montenegrina* was separated by 2.1–2.9% from the other representatives in the main *Knipowitschia* cluster, Adriatic *K. panizzae* by 1.8–2.9%, and *K. thessala* by 1.3–2.8%.

Monophyly of *Economidichthys* is not supported in any of the analyses. Although enforcing monophyly of the genus significantly reduced the length of the MP tree to 487, the Kishino–Hasegawa test proved the constraint tree to be significantly worse at  $P < 0.001$  ( $P < 0.08$  for the Shimodaira–Hasegawa test). The genetic distance between its two species is 4.4–6.0%, which exceeds the distance between the various *Knipowitschia* species (except for *K. punctatissima*). *Economidichthys pygmaeus* sequences are very similar (and, after alignment trimming, mostly identical) throughout the range, never exceeding a 0.3% GTR+I+ $\Gamma$  distance, except for the population from Chiliadou Springs (Mornos Delta, Corinthian Gulf) which differed by 1.5–1.7%. The split between *E. pygmaeus* from the Mornos Basin and its conspecifics is timed at 0.8–2.3 Mya.

Although *Pomatoschistus* is paraphyletic, all *Pomatoschistus* spp. cluster together in all analyses, apart from *Pomatoschistus quagga* and *P. canestrinii*. *Gobiusculus flavescens* is a member of the main *Pomatoschistus* clade (confirming Huyse *et al.*, 2004). Within *P. marmoratus*, three lineages are apparent: (1) an Euboean–Cretan cluster; (2) the representatives from the Venice Lagoon and from Drepano beach, the northernmost Greek location (located near the Greek–Albanian border); and (3) all other *P. marmoratus* samples from mainland western Greece. The most recent common ancestor of eastern Mediterranean *P. marmoratus* dates back 0.7–2.1 Mya. *Pomatoschistus marmoratus* from Venice Lagoon differed by up to 2.4% from its Greek conspecifics, with the minimal distance being 1.3–1.4% (to gobies from



**Figure 3.** Clock-constrained quartet puzzling tree based on 815 bp of 12S and 16S rDNA sequences for the sand gobies under study. Maximum likelihood distances are indicated, and a timescale is provided, calibrated assuming a strict molecular clock of  $1.5\% \times 10^{-6}$  years, respectively. For additional explanation regarding the evolutionary events indicated, see text.



**Figure 4.** Cladogram constructed for the 12S and 16S rDNA sequences (815 bp) of the sand gobies. Statistical support for each node is shown: BI posterior probability/ML bootstrap/puzzle support/MP bootstrap. Clades not recovered in a particular analysis are marked with ‘-’. Branch lengths indicate the expected number of substitutions per site under Bayesian inference.

Drepano). Most Greek *P. marmoratus* sequences were less than 0.5% apart, except for the Euboean (Livadaki, Karystos Bay) and Cretan (Plakias, Frangokastello, Giorgiopoulos) populations, which were more similar to each other than to specimens from mainland western Greece.

## DISCUSSION

The sand gobies are a monophyletic clade within the Gobiidae (Huysse *et al.*, 2004), including many endemics in the eastern Mediterranean basin. Many species are threatened by man and, for *E. pygmaeus* and *K. thessala*, the type localities have been destroyed and their populations lost (Economidis, 1991; Maitland & Crivelli, 1996). Despite the urgent need for their protection, conservation efforts are hindered by a lack of taxonomic knowledge (Economou *et al.*, 2007). Indeed, complexes of so-called cryptic species can be problematic in conservation policy (Wan *et al.*, 2004; Bickford *et al.*, 2006). Larmuseau *et al.* (2008) have shown that molecular work may assist in identifying morphologically cryptic sand gobies. Apart from species identification, molecular phylogeny/phylogeography is necessary to identify conservation units allowing management of the genetic diversity within species (Wan *et al.*, 2004) and to demarcate biogeographical boundaries in aquatic biomes (Riddle *et al.*, 2008; Costedoat & Gilles, 2009), with both representing crucial steps in conservation planning and resource management. We can pinpoint some lineages that merit additional taxonomic and conservational scrutiny because genetic distances can be compared to those between recognized species (for an example in *Rutilus rutilus* (Linnaeus, 1758), see Larmuseau *et al.*, 2009), although this of course depends on the quality of current morphological taxonomy (Johns & Avise, 1998). Based on previous work on gobies using mitochondrial rDNA (Harada *et al.*, 2002; Huysse *et al.*, 2004; Mukai *et al.*, 2004), corrected pairwise distances between closely-related species appear to be 2–4% at least. With this cut-off in mind, we might suggest some sand gobies whose taxonomy merits further scrutiny. Because a thorough revision is hence essential for these taxa, we will focus on lineages rather than species to see how major sand goby lineages fit into the historical biogeography of the Balkans.

Partial 12S and 16S mitochondrial ribosomal DNA sequences were used to clarify the phylogenetic relationships of taxa belonging to *Economidichthys* and *Knipowitschia*. The various analyses show a similar picture. *Economidichthys* spp. appear as a basal, paraphyletic group within the consistently monophyletic sand gobies. Most taxa assigned to *Knipowitschia* and *Pomatoschistus*, respectively, strongly cluster

together. The only exceptions are the well-supported clade formed by *P. canestrinii* and *K. punctatissima*, and *P. quagga* as sister to the remaining representatives of *Knipowitschia*.

## PHYLOGENETIC STATUS OF THE GENUS *ECONOMIDICHTHYS*

*Economidichthys* is endemic to western Greece. The widespread *E. pygmaeus* occurs in rivers, lowland streams, and freshwater bodies along the western Greek mainland, the island of Lefkas, and Lake Trichonis. This lake and the adjacent Lake Lysimachia are the only water bodies hosting *E. trichonis*. This is the smallest European freshwater fish and, in its larval stage, the smallest free-living European vertebrate (Miller, 2004a; Kottelat & Freyhof, 2007). The position of the genus *Economidichthys* based on molecular data was not resolved in the study of Huysse *et al.* (2004), which only included *E. pygmaeus*. The genus appears to be ancestral within the sand gobies, although many basal nodes are unresolved. Gamma-corrected genetic distances situate *E. trichonis* closest to its alleged congener. However, none of the phylogenetic analyses support monophyly of the genus. The two species of the genus differ more from each other than do species in other sand goby genera (a minimum of 4.4% pairwise genetic distance between *E. pygmaeus* and *E. trichonis* versus a maximum of 2.9% between the species in the *Knipowitschia* clade).

The most remarkable autapomorphy of *Economidichthys* is the perianal organ, a highly-derived organ unique among teleosts (Miller, 1990). However, both species exhibit substantial embryological and developmental differences, which might be adaptations to different lifestyles. Embryonic and larval developmental patterns have long been suggested to be evolutionarily conservative (and phylogenetically informative) compared to highly progressive adult patterns (Ahlstrom & Moser, 1976; Moser *et al.*, 1984; Strauss, 1992; Strathmann, 2000). Because of the dissimilar development of both *Economidichthys* species, Daoulas *et al.* (1993) hypothesized the perianal organ to be a shared adaptation (rather than a synapomorphy) to hole-brooding, with an antiseptic role. Although eggs are always more vulnerable to infections when deposited in sheltered locations (Potts, 1984), it is argued that both *Economidichthys* spp. experience this risk to a higher extent than other hole-brooders. They are almost unique among teleosts in using reed cavities as spawning sites, which are poorly aerated holes of organic origin. Many gobies possess accessory testis-associated structures, most often termed seminal vesicles or sperm duct glands, whose species-specific structure has been hypothesized to depend on ecological or physiological

conditions rather than on phylogeny (Fishelson, 1991). These are mostly present in males only (Cole & Hoese, 2001), in contrast to the perianal organ. However, both sexes in *Gobiodon* Bleeker, 1856 and *Paragobiodon* Bleeker, 1873 exhibit seminal vesicles that are suggested to include an antiseptic function. Species in both of these genera brood on corals (Mazzoldi, Petersen & Rasotto, 2005), and hence, similar to *Economidichthys*, on organic substrates.

*Aphia minuta* (Risso, 1810) and *Crystallogobius linearis* (Düben, 1845) are Atlantic–Mediterranean gobies known to be similar to *E. trichonis* with respect to biology and egg morphology (Caputo *et al.*, 2003; La Mesa *et al.*, 2005). Comparing the 12S and 16S rDNA sequences of those species (GenBank accession numbers EF218623, EF218635, EF218638, and EF218650; Giovannotti *et al.*, 2007), however, showed a Kimura-two-parameter (Kimura, 1980) genetic distance of approximately 16% and 13%, respectively, to *E. trichonis*, refuting sister-taxa affinities between those taxa (and showing that developmental characters can be homoplasious as well). Hence, to the best of our current knowledge, there are no known sister taxa to *Economidichthys*. This fact, together with the existence of only two lineages belonging to this genus (and this despite our thorough sampling) suggests that (past) extinction events might have been important in this group.

An exception to the generally very low genetic diversity within *E. pygmaeus* are the specimens caught in the Chiliadou Springs, off the Mornos River, at its easternmost distribution and less than 100 km from the nearest *E. pygmaeus* sampling location. They differ from the other *E. pygmaeus* populations by 1.5–1.7%, which is, by approximately two-fold, the highest pairwise distance found within the Thracian–Aegean *K. caucasica* or within the Ionian *Knipowitschia*. A difference between Chiliadou and western Greek populations was also found in *Valencia letourneuxi* (Sauvage, 1880) (Cyprinodontiformes, Valenciidae) (Vogiatzi *et al.*, 2009). Although this does not suffice to raise this population to the species level (see above), it is remarkable that it also deviates phenotypically from other *E. pygmaeus* [e.g. in (head) size parameters and in a colour pattern with reduced vertical stripes; A. N. Economou, S. Zogaris, pers. observ.] to the extent that it might have erroneously been assigned to *Knipowitschia* several times (Economou *et al.*, 1999). A geological event possibly isolating the Mornos basin from other drainages home to *E. pygmaeus* is the alternation between marine and freshwater (the so-called Corinth Lake) conditions in the Corinthian Gulf during the past million years (Moretti *et al.*, 2003; Zelilidis, 2003; Lykousis *et al.*, 2007; Lykousis, 2009). Approximately 0.9 Mya, the then lacustrine Gulf was flooded by the sea (West-

away, 1996). This corresponds to the estimated timing of the separation of the Mornos population (0.8–2.3 Mya) because the actual mutation rate of the genes is probably towards the higher edge of the range used (see above). In view of the relatively high risk of anthropogenic extinction of often unique local sand goby populations (Miller, 1990), this Mornos population deserves to be treated with extra conservational focus as an ESU because it represents the most phylogenetically distinct *E. pygmaeus* population.

#### NEED FOR TAXONOMIC REVISION OF THE PARAPHYLETIC *KNIPOWITSCHIA*

*Knipowitschia* has representatives in the Caspian, Black, Aegean, Ionian, and Adriatic basins, many of which are local freshwater endemics. Based on 12S and 16S rDNA sequence data, Penzo *et al.* (1998) and Huyse *et al.* (2004) demonstrated the paraphyly of the genus. The additional species sequenced here fall within the *Knipowitschia* clade, so the genus appears valid, with the exception of *K. punctatissima* that clusters with *P. canestrinii* in all analyses. It should be noted that the affiliation of *K. punctatissima* has been under discussion before, and the absence of the longitudinal row and suborbital papillae led to its classification under the newly-described *Orsinigobius*. Subsequently, morphological similarities with *Knipowitschia* led to the synonymization of *Orsinigobius* with *Knipowitschia* (which appears to be unjustified based on the present results), and the absent characters were considered as secondary losses (Economidis & Miller, 1990; Penzo *et al.*, 1998). The genetic divergence among the various *Knipowitschia* species is limited (2.9% at most). Furthermore, the genetic distances within the Ionian populations are almost equal to those between Ionian populations and *K. thessala* or *K. caucasica*. The fact that the level of inter- and intraspecific distances is similar, confusing current species delineation, indicates the existence of a cryptic species complex (Nelson, 1999), and *Knipowitschia* is certainly in need of a revision.

The recently described *Knipowitschia montenegrina* clusters with its Adriatic congener *K. panizzae*. Before being raised to species status, its representatives were assigned to *K. panizzae*, although it is morphologically not its most reminiscent species (Kovačić & Šanda, 2007). In view of the genetic distance of at least 2.1% to its congeners, the morphological species is genetically confirmed. It is, however, difficult to explain how isolation, either through geographical distance or vicariant barriers, could have provoked speciation here (Skoulikidis *et al.*, 2009; Zogaris *et al.*, 2009). Perhaps *K. montenegrina* and *K. panizzae* stem from separate Danubian colonization events. Mostly on the basis of mtDNA markers or allozyme

genetics, other Balkan fish populations were suggested to stem from Pleistocene river capture from the Danube to the southern (Durand *et al.*, 1999b, 2003; Šanda, Lusková & Vukić, 2005) and northern Adriatic (Tsigenopoulos & Berrebi, 2000; Šlechtová *et al.*, 2004; Papa & Schreiber, 2005).

The 0.5–1.3% genetic distance of *K. milleri* to its only other Ionian congeners *Knipowitschia* sp. and *K. cf. panizzae* is low, indicative of intraspecific rather than interspecific variation. Their respective larvae cannot be reliably distinguished using morphology (Miller, Madurell & Economou, 2004). This could suggest that divergence was indeed recent enough for genetic mixing to occur. Bearing in mind that few Ionian *Knipowitschia* populations are adequately studied morphologically, and that phenotypic variation is observed in *Knipowitschia* spp. (Miller, 2004b; Kovačić, 2005), it is uncertain whether the distinctive features of the Acheron *K. milleri* are real or merely represent intraspecific variation. In view of the low genetic distances, all Ionian representatives of *Knipowitschia* appear to be conspecific with *K. milleri*. River confluences as a result of recent marine regression in the Ionian region (Chronis, Piper & Anagnostou, 1991), and a drop of salinity in the coastal zone, provided occasions for sympatry and could explain the homogeneity in Ionian *Knipowitschia* (and *E. pygmaeus*) populations (e.g. for chub; Durand *et al.*, 1999b). Because Kovačić (2008) asserts that *K. panizzae* in the Adriatic cannot be distinguished from *K. caucasica*, a thorough study of *Knipowitschia* in the Adriatic and Ionian regions is necessary to determine whether there are different species present, and whether certain endemics are limited in range and should as such be treated as ESUs. Caution is still required before forging species entities (or in this case, lumping taxa) in the Ionian and South-East Adriatic ecoregions. It would be interesting to include other poorly known *Knipowitschia* spp. from these regions in the analysis, such as *Knipowitschia croatica* Mrakovic, Kerovec, Misetic & Schneider, 1996, *Knipowitschia mrakovici* Miller, 2009, *Knipowitschia radovici* Kovačić, 2005, and *Knipowitschia goerneri* Ahnelt, 1991, an endemic to the island of Corfu.

#### BIOGEOGRAPHICAL AND HISTORICAL IMPLICATIONS FOR THE BALKANS

Five lineages of *Knipowitschia* are recognized in the present study: south-eastern Adriatic, Adriatic, Ionian, Macedonian–Thessalian and Thracian–Aegean. These insights provide support for the historical view that the south-eastern Adriatic, Ionian, Macedonian–Thessalian, and Thracian–Aegean (or northern Aegean) regions have a distinct evolutionary history reflecting ichthyogeographical boundaries

(Bianco, 1986; Bănărescu, 2004; Economou *et al.*, 2007; Zogaris *et al.*, 2009). As such, each region requires separate conservation planning strategies. We reiterate that more work, such as in the Western and Eastern Aegean, may reveal the historical affinities and biogeographies among these and the Thracian ecoregions. *Knipowitschia* spp., being tiny fish, survive in small populations in minute wetlands on several Aegean islands. They may represent one of the few widespread genera, suitable as biogeographical indicators, in the remarkably species-depauperate insular realm of these ecoregions.

Several studies have invoked Lago Mare-phases during the Messinian Salinity Crisis (MSC) to explain the presence of Paratethyan *Knipowitschia* in the Mediterranean basin (Miller, 1990; Huyse *et al.*, 2004). If the genus is this old, a multitude of 'diversification routes' could have led to its current diversity: two Lago Mare phases (Clauzon *et al.*, 2005), independent Paratethyan connections (Sonnenfeld, 1974; Sakiņ & Yaltırak, 2005), and several colonization routes for Danubian freshwater fishes towards the Adriatic or Aegean (Doadrio & Carmona, 2003; Levy, Doadrio & Almada, 2009). However, even the slowest molecular clock dates *Knipowitschia* diversification in this region well after the MSC or Lago Mare phases. We suggest that Pleistocene glaciation cycles drove diversification within *Knipowitschia*, as has been proposed for *Pomatoschistus* (Gysels *et al.*, 2004; Huyse *et al.*, 2004), and allowed the spread and diversification of freshwater fishes in the region (Miller & Šanda, 2008). However, this alone does not explain the range of the various *Knipowitschia* species. Indeed, clustering of *K. thessala* with the Ionian lineage appears to favour Pleistocene river capture events rather than a spread through the Adriatic and Ionian seas (Durand *et al.*, 1999b). Both Thracian–Aegean *K. caucasica* and Macedonian–Thessalian *K. thessala* could stem from originally Paratethyan river captures (Economidis & Bănărescu, 1991). Geographically, *K. thessala* interrupts northern and southern Thracian–Aegean *K. caucasica*. This might result from relatively later Danubian colonization via the Axios River (Durand *et al.*, 1999b). Lastly, concerning the Thracian–Aegean *Knipowitschia* lineage, clustering of the Euboean and Thracian–Aegean specimens could be the result of dispersal through a freshened Aegean Sea approximately 0.2 Mya, which corresponds to the molecular dating. Affinities between Euboea and Thrace have also been found for two cyprinid genera (Zardoya, Economidis & Doadrio, 1999; Tsigenopoulos & Berrebi, 2000).

The presence of both of the *Economidichthys* spp. and an endemic *Knipowitschia* lineage in the Ionian region leads to a higher local biodiversity compared to

the other study areas. This supports the general agreement that Ionian drainages have long been isolated from other European (including Greek) drainages. Together with the glacial refugium role of the western Balkans and with the presence of ancient lakes in the Acheloos drainage (e.g. Lake Trichonis: Economou *et al.*, 1994), this isolation shaped a unique freshwater ichthyofauna with a high degree of endemism (Economou *et al.*, 2007; Skoulikidis *et al.*, 2009) deserving special attention for conservation.

#### COMPARISON WITH MARINE TAXA: *P. MARMORATUS*

The genetic clusters of the marine *P. marmoratus*, namely (1) the islands of Euboea and Crete; (2) Drepano and the Adriatic; and (3) western Greece, confirm the biogeographical boundary between the Ionian and Aegean in the eastern Mediterranean Sea (Spalding *et al.*, 2007). In view of the timing, Pleistocene events must have played a role in structuring the region's marbled goby populations. Separate monophyletic clusters of Aegean and Adriatic marine sand gobies, as a result of glaciation events, were predicted by Stefanni & Thorley (2003) based on *Pomatoschistus minutus* molecular data. The fact that the Drepano specimens (near the Kalamas Basin and the Greek–Albanian border) cluster with the Adriatic *P. marmoratus* could point to a correlation between genetic and geographical distances. The genetic distances found in *P. marmoratus* throughout the sampling area do not exceed those found in freshwater gobies on a much smaller spatial scale (e.g. within *Knipowitschia*), suggesting that dispersal barriers in the marine realm are not as sharp as in freshwater systems.

#### PERSPECTIVES

The results obtained in the present study indicate that the natural range, phylogeny, and phylogeography of Balkan sand gobies, both marine and freshwater, reflect ichthyogeographical boundaries proposed on the basis of fish assemblage composition (Bianco, 1986; Bănărescu, 2004; Economou *et al.*, 2007; Zogaris *et al.*, 2009). Taxonomic uncertainties were highlighted in *Knipowitschia* and *Economidichthys*, and ancient fish dispersal pathways previously mentioned in the literature were recognized in the goby radiation by linking its phylogeny to the region's historical biogeography. On the scale of single or adjacent river drainages, *Knipowitschia* is understudied, necessitating further molecular and morphological work to formally describe species demarcations. With the exception of one taxon, it appears as a monophyletic genus. Its low within-clade genetic variability might represent recent or ongoing speciation

in a cryptic species complex, and its phylogeny reflects the Balkan aquatic ecoregions. Because the geographical resolution of our sampling design was rather high, additional molecular work should focus on other regions, such as the northern Balkans along the Adriatic coastline, and the Ponto-Caspian basin. Indeed, approximately 12 endemic sand gobies remain to be phylogenetically positioned (Eschmeyer, 2010). The retrieval of a basal polytomy in several studies on this system, current higher taxon coverage, the fast divergence in other gobies (Akihito *et al.*, 2000), and the high phylogenetic content of the dataset point to a fast radiation process. In view of classical problems with mitochondrial gene trees (Shaw, 2002; Rubinoff, 2006), additional morphological work is recommended (Avice, 2000), together with nuclear sequences to assess phenomena such as hybridization, and for a more reliable assignment of ESUs *sensu* Moritz (1994).

To increase resolution, parasite molecular data might be useful because parasites are known to sometimes complement (genetic) host data (Nieberding & Olivieri, 2007; Blakeslee, Byers & Lesser, 2008; Barson *et al.*, 2010). Preliminary molecular results on *Gyrodactylus* von Nordmann, 1832 flatworms on various *E. pygmaeus* populations show a higher genetic diversity than their host, pointing to a Pleistocene isolation event in the Kalamas River (the northernmost border of its range) (M. P. M. Vanhove, B. Geens, pers. observ.). As such, this approach looks very promising for generating additional information on the current study system.

#### ACKNOWLEDGEMENTS

The authors dedicate this article to Professor Peter J. Miller, in acknowledgement of his enormous and inspiring contribution to the study of the Greek ichthyofauna. Fieldwork was partly financed with a travel grant to M.P.M.V. from BeNCoRe (Belgian Network for Coastal Research; ENCORA coordination action contract number FP6-2004-Global-3–518120). We are grateful to D. Mrdak and V. Pešić (Department of Biology, University of Montenegro) for help with the collection of *Knipowitschia* (Joint Research and Technology Programme 'Scientific cooperation and technology transfer for the development of a fish-based assessment method of surface waters ecological status'). A. Vidalis (Biodiversity East) produced Figure 1; S. Janssens (Laboratory of Plant Systematics, K.U.Leuven) provided useful phylogenetic suggestions; and B. Geens assisted in collecting the *Gyrodactylus* data from *E. pygmaeus* hosts. We thank L. Vardakas, Y. Chatzinikolaou, V. Tachos, D. Kommatas, Y. Koutsikos, and Y. Kapakos for their participation in the sampling campaigns. The

anonymous reviewers contributing greatly to the improvement of the manuscript are gratefully acknowledged. M.P.M.V and T.H. benefit from a PhD and a postdoctoral fellowship, respectively, of the Research Council – Flanders (FWO-Vlaanderen), and M.H.D.L. benefits from a postdoctoral position of the K.U.Leuven (BOF PDM-Kort).

## REFERENCES

- Ahlstrom EH, Moser HG. 1976.** Eggs and larvae of fishes and their role in systematic investigations and in fisheries. *Revue des Travaux de l'Institut des Pêches Maritimes* **40**: 379–398.
- Ahnelt H, Bianco PG. 1990.** *Orsinigobius milleri* n. sp., a new species of freshwater goby from W-Greece (Pisces: Gobiidae). *Annalen des Naturhistorischen Museums in Wien* **91B**: 1–6.
- Akihito IA, Kobayashi T, Ikeo K, Imanishi T, Ono H, Umehara Y, Hamamatsu CH, Sugiyama K, Ikeda Y, Sakamoto K, Fumihito A, Ohno S, Gojobori T. 2000.** Evolutionary aspects of gobioid fishes based upon a phylogenetic analysis of mitochondrial cytochrome *b* genes. *Gene* **259**: 5–15.
- Almada VC, Robalo JI, Levy A, Freyhof J, Bernardi G, Doadrio I. 2009.** Phylogenetic analysis of peri-Mediterranean blennies of the genus *Salaria*: molecular insights on the colonization of freshwaters. *Molecular Phylogenetics and Evolution* **52**: 424–431.
- Alves-Gomes JA. 1999.** Systematic biology of gymnotiform and mormyrid electric fishes: phylogenetic relationships, molecular clocks and rates of evolution in the mitochondrial rRNA genes. *The Journal of Experimental Biology* **202**: 1167–1183.
- Avise JC. 2000.** *Phylogeography: the history and formation of species*. Cambridge, MA: Harvard University Press.
- Bănărescu PM. 2004.** Distribution pattern of the aquatic fauna of the Balkan Peninsula. In: Griffiths HI, Kryštufek B, Reed JM, eds. *Balkan biodiversity: pattern and process in the European hotspot*. Dordrecht: Kluwer Academic Publishers, 203–217.
- Barson M, Prikrylová I, Vanhove MPM, Huyse T. 2010.** Parasite hybridization in African *Macrogryrodactylus* spp. (Monogenea, Platyhelminthes) signals historical host distribution. *Parasitology* **137**: 1585–1595.
- Bianco PG. 1986.** The zoogeographic units of Italy and the Western Balkans based on cyprinid species ranges (Pisces). *Biologia Gallo-Hellenica* **12**: 291–299.
- Bickford D, Lohman DJ, Sodhi NS, Ng PKL, Meier R, Winker K, Ingram KK, Das I. 2006.** Cryptic species as a window on diversity and conservation. *Trends in Ecology and Evolution* **22**: 148–155.
- Blakeslee AMH, Byers JE, Lesser MP. 2008.** Solving cryptogenic histories using host and parasite molecular genetics: the resolution of *Littorina littorea*'s North American origin. *Molecular Ecology* **17**: 3684–3696.
- Calendini F, Martin JF. 2005.** *PaupUp version 1.0.3.1. A free graphical frontend for PAUP\* Dos software*. Available at: <http://www.agromontpellier.fr/spe/Recherche/JFM/PaupUp/>
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009.** trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972–1973.
- Caputo V, La Mesa M, Candi G, Nisi Cerioni P. 2003.** The reproductive biology of the crystal goby with a comparison to that of the transparent goby. *Journal of Fish Biology* **62**: 375–385.
- Chronis G, Piper DJW, Anagnostou C. 1991.** Late Quaternary evolution of the Gulf of Patras, Greece: tectonism, deltaic sedimentation and sea-level change. *Marine Geology* **97**: 191–209.
- Clauzon G, Suc J-P, Popescu S-M, Marunteanu M, Rubino J-L, Marinescu F, Melinte MC. 2005.** Influence of Mediterranean sea-level changes on the Dacic Basin (Eastern Paratethys) during the late Neogene: the Mediterranean Lago Mare facies deciphered. *Basin Research* **17**: 437–462.
- Cole KS, Hoese DF. 2001.** Gonad morphology, colony demography and evidence for hermaphroditism in *Gobiodon okinawae* (Teleostei, Gobiidae). *Environmental Biology of Fishes* **61**: 161–173.
- Costedoat C, Gilles A. 2009.** Quaternary pattern of freshwater fishes in Europe: comparative phylogeography and conservation perspective. *The Open Conservation Biology Journal* **3**: 36–48.
- Crandall KA, Bininda-Emonds ORP, Mace GM, Wayne RK. 2000.** Considering evolutionary processes in conservation biology. *Trends in Ecology and Evolution* **15**: 290–295.
- Cummins C, McInerney JO. 2011.** A method for inferring the rate of evolution of homologous characters that can potentially improve phylogenetic inference, resolve deep divergence and correct systematic biases. *Systematic Biology*. DOI: 10.1093/sysbio/syr064.
- Daoulas C, Economou AN, Psarras T, Barbieri-Tseliki R. 1993.** Reproductive strategies and early development of three freshwater gobies. *Journal of Fish Biology* **42**: 749–776.
- Doadrio I, Carmona JA. 2003.** Testing freshwater Lago Mare dispersal theory on the phylogeny relationships of Iberian cyprinid genera *Chondrostoma* and *Squalius* (Cypriniformes, Cyprinidae). *Graellsia* **59**: 457–473.
- Durand JD, Persat H, Bouvet Y. 1999a.** Phylogeography and postglacial dispersion of the chub (*Leuciscus cephalus*) in Europe. *Molecular Ecology* **8**: 989–997.
- Durand JD, Templeton AR, Guinand B, Imsiridou A, Bouvet Y. 1999b.** Nested clade and phylogeographic analyses of the chub, *Leuciscus cephalus* (Teleostei, Cyprinidae), in Greece: implications for Balkan Peninsula biogeography. *Molecular Phylogenetics and Evolution* **13**: 566–580.
- Durand J-D, Bianco PG, Laroche J, Gilles A. 2003.** Insight into the origin of endemic Mediterranean ichthyofauna: phylogeography of *Chondrostoma* genus (Teleostei, Cyprinidae). *Journal of Heredity* **94**: 315–328.

- Economidis PS. 1991.** *Check list of freshwater fishes of Greece. Recent status of threats and protection.* Athens: Hellenic Society for the Protection of Nature.
- Economidis PS. 1995.** Endangered freshwater fishes of Greece. *Biological Conservation* **72**: 201–211.
- Economidis PS, Bănărescu PM. 1991.** The distribution and origins of freshwater fishes in the Balkan peninsula, especially in Greece. *Internationale Revue der Gesamten Hydrobiologie* **76**: 257–283.
- Economidis PS, Miller PJ. 1990.** Systematics of freshwater gobies from Greece (Teleostei: Gobiidae). *Journal of Zoology* **221**: 125–170.
- Economou AN, Barbieri R, Daoulas CH, Psarras T, Stoumboudi M, Bertahas I, Giakoumi S, Patsias A. 1999.** *Endangered freshwater fish of western Greece and the Peloponnese – distribution, abundance, threats and measures for protection. Final technical report* [in Greek]. Athens: National Centre for Marine Research.
- Economou AN, Daoulas CH, Psarras TH, Barbieri-Tseliki R. 1994.** Freshwater larval fish from Lake Trichonis (Greece). *Journal of Fish Biology* **45**: 17–35.
- Economou AN, Giakoumi S, Vardakas L, Barbieri R, Stoumboudi M, Zogaris S. 2007.** The freshwater ichthyofauna of Greece – an update based on a hydrographic basin survey. *Mediterranean Marine Science* **8**: 91–166.
- Edgar RC. 2004.** MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**: 1792–1797.
- Eschmeyer WN, ed. 2010.** *Catalog of fishes electronic version.* Available at: <http://research.calacademy.org/ichthyology/catalog/fishcatmain.asp>
- Farris JS, Källersjö M, Kluge AG, Bult C. 1995.** Constructing a significance test for incongruence. *Systematic Biology* **44**: 570–572.
- Felsenstein J. 2005.** *PHYLIP (Phylogeny Inference Package)*, Version 3.6. Distributed by the author. Seattle, WA: Department of Genome Sciences, University of Washington. Available at: <http://evolution.genetics.washington.edu/phylip.html>
- Fishelson L. 1991.** Comparative cytology and morphology of seminal vesicles in male gobiid fishes. *Japanese Journal of Ichthyology* **38**: 17–30.
- Fraser DJ, Bernatchez L. 2001.** Adaptive evolutionary conservation: towards a unified concept for defining conservation units. *Molecular Ecology* **10**: 2741–2752.
- Gelman A, Rubin DB. 1992.** Inference from iterative simulation using multiple sequences. *Statistical Science* **7**: 457–472.
- Gillooly JF, Allen AP, West GB, Brown JH. 2005.** The rate of DNA evolution: effects of body size and temperature on the molecular clock. *Proceedings of the National Academy of Sciences of the United States of America* **102**: 140–145.
- Giovannotti M, Nisi Cerioni P, La Mesa M, Caputo V. 2007.** Molecular phylogeny of the three pedomorphic Mediterranean gobies (Perciformes: Gobiidae). *Journal of Experimental Zoology Part B (Molecular and Developmental Evolution)* **308**: 722–729.
- Glez-Peña D, Gómez-Blanco D, Reboiro-Jato M, Fdez-Riverola F, Posada D. 2010.** ALTER: program-oriented format conversion of DNA and protein alignments. *Nucleic Acids Research* **38** (Suppl. 2): W14–W18.
- Goldman N, Anderson JP, Rodrigo AG. 2000.** Likelihood-based tests of topologies in phylogenetics. *Systematic Biology* **49**: 652–670.
- Guindon S, Gascuel O. 2003.** A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Systematic Biology* **52**: 696–704.
- Gysels ES, Hellemans B, Patarnello T, Volckaert FAM. 2004.** Current and historic gene flow of the sand goby *Pomatoschistus minutus* on the European Continental Shelf and in the Mediterranean Sea. *Biological Journal of the Linnean Society* **83**: 561–576.
- Harada S, Jeon S-R, Kinoshita I, Tanaka M, Nishida M. 2002.** Phylogenetic relationships of four species of floating gobies (*Gymnogobius*) as inferred from partial mitochondrial cytochrome *b* gene sequences. *Ichthyological Research* **49**: 324–332.
- Heino J. 2001.** Regional gradient analysis of freshwater biota: do similar biogeographic patterns exist among multiple taxonomic groups? *Journal of Biogeography* **28**: 69–76.
- Huelsenbeck JP, Ronquist F. 2001.** MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**: 754–755.
- Hugueny B. 1989.** West African rivers as biogeographic islands: species richness of fish communities. *Oecologia* **79**: 236–243.
- Hurvich CM, Tsai C-L. 1989.** Regression and time series model selection in small samples. *Biometrika* **76**: 297–307.
- Huysse T, Van Houdt J, Volckaert FAM. 2004.** Paleoclimatic history and vicariant speciation in the ‘sand goby’ group (Gobiidae, Teleostei). *Molecular Phylogenetics and Evolution* **32**: 324–336.
- Johns GC, Avise JC. 1998.** A comparative summary of genetic distances in the vertebrates from the mitochondrial cytochrome *b* gene. *Molecular Biology and Evolution* **15**: 1481–1490.
- Keith P, Lord C, Lorion J, Watanabe S, Tsukamoto K, Couloux A, Dettai A. 2011.** Phylogeny and biogeography of Sicydiinae (Teleostei: Gobiidae) inferred from mitochondrial and nuclear genes. *Marine Biology* **158**: 311–326.
- Kimura M. 1980.** A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* **16**: 111–120.
- Kishino H, Hasegawa M. 1989.** Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order of the Hominoida. *Journal of Molecular Evolution* **29**: 170–179.
- Kocher TD, Thomas WK, Meyer A, Edwards SV, Paabo S, Villablanca FX, Wilson AC. 1989.** Dynamics of mitochondrial DNA evolution in animals – amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences of the United States of America* **86**: 6196–6200.

- Kottelat M, Freyhof J. 2007.** *Handbook of European freshwater fishes*. Cornol: Kottelat and Berlin: Freyhof.
- Kovačić M. 2005.** A new species of *Knipowitschia* (Gobiidae) from Dalmatia, Croatia. *Cybium* **29**: 275–280.
- Kovačić M. 2008.** The key for identification of Gobiidae (Pisces: Perciformes) in the Adriatic Sea. *Acta Adriatica* **49**: 245–254.
- Kovačić M, Patzner RA. 2011.** North-Eastern Atlantic and Mediterranean gobies. In: Patzner RA, Van Tassell JL, Kovačić M, Kapoor BG, eds. *The biology of gobies*. Enfield: Science Publishers Inc, 177–206.
- Kovačić M, Šanda R. 2007.** A new species of *Knipowitschia* (Perciformes: Gobiidae) from southern Montenegro. *Journal of the National Museum (Prague), Natural History Series* **176**: 81–89.
- Kryštufek B, Reed JM. 2004.** Pattern and process in Balkan biodiversity – an overview. In: Griffiths HI, Kryštufek B, Reed JM, eds. *Balkan biodiversity: pattern and process in the European hotspot*. Dordrecht: Kluwer Academic Publishers, 1–8.
- La Mesa M, Arneri E, Caputo V, Iglesias M. 2005.** The transparent goby, *Aphia minuta*: review of biology and fisheries of a paedomorphic European fish. *Reviews in Fish Biology and Fisheries* **15**: 89–109.
- Larmuseau MHD, Freyhof J, Volckaert FAM, Van Houdt JKJ. 2009.** Matrilinear phylogeography and demographical patterns of *Rutilus rutilus*: implications for taxonomy and conservation. *Journal of Fish Biology* **75**: 332–353.
- Larmuseau MHD, Guelinckx J, Hellemans B, Van Houdt JKJ, Volckaert FAM. 2008.** Fast PCR-RFLP method facilitates identification of *Pomatoschistus* species in the North Atlantic. *Journal of Applied Ichthyology* **24**: 342–344.
- Larmuseau MHD, Huysse T, Vancampenhout K, Van Houdt JKJ, Volckaert FAM. 2010.** High molecular diversity in the rhodopsin gene in closely related goby fishes: a role for visual pigments in adaptive speciation? *Molecular Phylogenetics and Evolution* **55**: 689–698.
- Larmuseau MHD, Vanhove MPM, Huysse T, Volckaert FAM, Decorte R. 2011.** Signature of selection on the rhodopsin gene in the marine radiation of American seven-spined gobies (Gobiidae, Gobiosomatini). *Journal of Evolutionary Biology* **24**: 1618–1625.
- Levy A, Doadrio I, Almada VC. 2009.** Historical biogeography of European leuciscins (Cyprinidae): evaluating the Lago Mare dispersal hypothesis. *Journal of Biogeography* **36**: 55–65.
- Lykousis V. 2009.** Sea-level changes and shelf break prograding sequences during the last 400 ka in the Aegean margins: subsidence rates and palaeogeographic implications. *Continental Shelf Research* **29**: 2037–2044.
- Lykousis V, Sakellariou D, Moretti I, Kaberi H. 2007.** Late Quaternary basin evolution of the Gulf of Corinth: sequence stratigraphy, sedimentation, fault-slip and subsidence rates. *Tectonophysics* **440**: 29–51.
- Maitland PS, Crivelli AJ. 1996.** *Conservation of freshwater fish. Conservation of Mediterranean wetlands* 7. Arles: Tour du Valat.
- Martin AP, Palumbi SR. 1993.** Body size, metabolic rate, generation time, and the molecular clock. *Proceedings of the National Academy of Sciences of the United States of America* **90**: 4087–4091.
- Mazzoldi C, Petersen CW, Rasotto MB. 2005.** The influence of mating system on seminal vesicle variability among gobies (Teleostei, Gobiidae). *Journal of Zoological Systematics and Evolutionary Research* **43**: 307–314.
- Mejri R, Lo Brutto S, Ben Hassine OK, Arculeo M. 2009.** A study on *Pomatoschistus tortonesei* Miller 1968 (Perciformes, Gobiidae) reveals the Siculo–Tunisian Strait (STS) as a breakpoint to gene flow in the Mediterranean basin. *Molecular Phylogenetics and Evolution* **53**: 596–601.
- Miller PJ. 1986.** Gobiidae. In: Whitehead PJP, Bauchot M-L, Hureau J-C, Nielsen J, Tortonese E, eds. *Fishes of the North-Eastern Atlantic and the Mediterranean*, Vol. III. Paris: Unesco, 1019–1085.
- Miller PJ. 1990.** The endurance of endemism – the Mediterranean fresh-water gobies and their prospects for survival. *Journal of Fish Biology* **37**: 145–156.
- Miller PJ. 2004a.** *Economidichthys*. In: Miller PJ, ed. *The freshwater fishes of Europe – Gobiidae 2*. Wiebelsheim: AULA-Verlag GmbH, 439–442.
- Miller PJ. 2004b.** *Knipowitschia*. In: Miller PJ, ed. *The freshwater fishes of Europe – Gobiidae 2*. Wiebelsheim: AULA-Verlag GmbH, 331–337.
- Miller PJ. 2004c.** *Pomatoschistus*. In: Miller PJ, ed. *The freshwater fishes of Europe – Gobiidae 2*. Wiebelsheim: AULA-Verlag GmbH, 279–283.
- Miller PJ. 2009.** A west Balkanian freshwater gobiid fish, *Knipowitschia mrakovcici* sp. nov. (Teleostei: Gobiidae). *Journal of Fish Biology* **74**: 1499–1507.
- Miller PJ, Madurell T, Economou AN. 2004.** *Knipowitschia milleri*. In: Miller PJ, ed. *The freshwater fishes of Europe – Gobiidae 2*. Wiebelsheim: AULA-Verlag GmbH, 399–404.
- Miller PJ, Šanda R. 2008.** A new west Balkanian sand-goby (Teleostei: Gobiidae). *Journal of Fish Biology* **72**: 259–270.
- Moretti I, Sakellariou D, Lykousis V, Micarelli L. 2003.** The Gulf of Corinth: an active half graben? *Journal of Geodynamics* **36**: 323–340.
- Moritz C. 1994.** Defining 'Evolutionarily Significant Units' for conservation. *Trends in Ecology and Evolution* **9**: 373–375.
- Moser HG, Richards WJ, Cohen DM, Fahay MP, Kendall AW, Richardson SL, eds. 1984.** *Ontogeny and systematics of fishes*. La Jolla, California: American Society of Ichthyologists and Herpetologists Special Publication 1.
- Mukai T, Suzuki T, Nishida M. 2004.** Genetic and geographical differentiation of *Pandaka* gobies in Japan. *Ichthyological Research* **51**: 222–227.
- Nelson JS. 1999.** Editorial and introduction: the species concept in fish biology. *Reviews in Fish Biology and Fisheries* **9**: 277–280.
- Nelson JS. 2006.** *Fishes of the world*, 4th edn. New York, NY: John Wiley and Sons.
- Nieberding CM, Olivieri I. 2007.** Parasites: proxies for host genealogy and ecology? *Trends in Ecology and Evolution* **22**: 156–165.

- Özcan G. 2009.** The status of the freshwater gobiid, *Knipowitschia mermere* Ahnelt, 1995: distribution ecology and threats. *Biharean Biologist* **3**: 139–141.
- Ozen O, Irmak E, Bilecenoglu M. 2007.** Occurrence of *Pomatoschistus minutus* (Pallas, 1770) (Pisces: Gobiidae) along the North Aegean coast of Turkey. *Annales, Series Historia Naturalis* **17**: 161–164.
- Palumbi SR, Martin A, Romano S, McMillan WO, Stice L, Grabowski G. 1991.** *The simple fools guide to PCR*. Honolulu: University of Hawaii Press.
- Papa L, Schreiber A. 2005.** Genetic variability in European sculpin, *Cottus gobio* (Pisces: Cottidae): assignment of population samples from the Brenta (Italy) and the Sava (Slovenia) river systems to Central European population groups. *Stuttgarter Beiträge zur Naturkunde Serie A (Biologie)* **678**: 1–15.
- Penzo E, Gandolfi G, Bargelloni L, Colombo L, Patarinello T. 1998.** Messinian salinity crisis and the origin of freshwater lifestyle in Western Mediterranean gobies. *Molecular Biology and Evolution* **15**: 1472–1480.
- Posada D. 2008.** jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* **25**: 1253–1256.
- Potts GW. 1984.** Parental behaviour in temperate marine teleosts with special reference to the development of nest structures. In: Potts GW, Wootton RJ, eds. *Fish reproduction: strategies and tactics*. London: Academic Press, 223–244.
- Riddle BR, Dawson MN, Hadly EA, Hafner DJ, Hickerson MJ, Mantooth SJ, Yoder AD. 2008.** The role of molecular genetics in sculpting the future of integrative biogeography. *Progress in Physical Geography* **32**: 173–202.
- Rodriguez F, Oliver JL, Marin A, Medina JR. 1990.** The general stochastic model of nucleotide substitution. *Journal of Theoretical Biology* **142**: 485–501.
- Ronquist F, Huelsenbeck JP. 2003.** MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Rüber L, Van Tassell JL, Zardoya R. 2003.** Rapid speciation and ecological divergence in the American seven-spined gobies (Gobiidae, Gobiomatini) inferred from a molecular phylogeny. *Evolution* **57**: 1584–1598.
- Rubinoff D. 2006.** Utility of mitochondrial DNA barcodes in species conservation. *Conservation Biology* **20**: 1026–1033.
- Sakinç M, Yaltrak C. 2005.** Messinian crisis: what happened around the northeastern Aegean? *Marine Geology* **221**: 423–436.
- Šanda R, Lusková V, Vukić J. 2005.** Notes on the distribution and taxonomic status of *Gobio gobio* from the Morača River basin (Montenegro). *Folia Zoologica* **54** (Suppl. 1): 73–80.
- Schmidt HA. 2009.** Testing tree topologies. In: Lemey PH, Salemi M, Vandamme A-M, eds. *The phylogenetic handbook: a practical approach to phylogenetic analysis and hypothesis testing*. Cambridge: Cambridge University Press, 381–404.
- Schmidt HA, Strimmer K, Vingron M, von Haeseler A. 2002.** TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. *Bioinformatics* **18**: 502–504.
- Shaw KL. 2002.** Conflict between nuclear and mitochondrial DNA phylogenies of a recent species radiation: what mtDNA reveals and conceals about modes of speciation in Hawaiian crickets. *Proceedings of the National Academy of Sciences of the United States of America* **99**: 16122–16127.
- Shimodaira H, Hasegawa M. 1999.** Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Molecular Biology and Evolution* **16**: 1114–1116.
- Skoulikidis NTH, Economou AN, Gritzalis KC, Zogaris S. 2009.** Rivers of the Balkans. In: Tockner K, Robinson CT, Uehlinger U, eds. *Rivers of Europe*, 1st edn. London: Academic Press, 421–466.
- Šlechtová V, Bohlen J, Freyhof J, Persat H, Delmastro GB. 2004.** The Alps as barrier to dispersal in cold-adapted freshwater fishes? Phylogeographic history and taxonomic status of the bullhead in the Adriatic freshwater drainage. *Molecular Phylogenetics and Evolution* **33**: 225–239.
- Smith KG, Darwall WRT (compilers). 2006.** *The status and distribution of freshwater fish endemic to the Mediterranean Basin*. Gland and Cambridge: IUCN.
- Sonnenfeld P. 1974.** The Upper Miocene evaporite basins in the Mediterranean Region – a study in paleo-oceanography. *International Journal of Earth Sciences* **63**: 1133–1172.
- Spalding MD, Fox HE, Allen GR, Davidson N, Ferdaña ZA, Finlayson M, Halpern BS, Jorge MA, Lombana A, Lourie SA, Martin KD, McManus E, Molnar J, Recchia CA, Robertson J. 2007.** Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *BioScience* **57**: 573–583.
- Stefanni S. 2000.** First record of the Norway goby in the Adriatic Sea. *Journal of Fish Biology* **57**: 828–830.
- Stefanni S, Thorley JL. 2003.** Mitochondrial DNA phylogeography reveals the existence of an Evolutionarily Significant Unit of the sand goby *Pomatoschistus minutus* in the Adriatic (Eastern Mediterranean). *Molecular Phylogenetics and Evolution* **28**: 601–609.
- Strathmann RR. 2000.** Functional design in the evolution of embryos and larvae. *Seminars in Cell & Developmental Biology* **11**: 395–402.
- Strauss RE. 1992.** Developmental variability and heterochronic evolution in poeciliid fishes (Cyprinodontiformes). In: Mayden RL, ed. *Systematics, historical ecology, and North American freshwater*. Stanford, CA: Stanford University Press, 492–514.
- Strimmer K, von Haeseler A. 1997.** Likelihood-mapping: a simple method to visualize phylogenetic content of a sequence alignment. *Proceedings of the National Academy of Sciences of the United States of America* **94**: 6815–6819.
- Swofford DL. 2001.** *PAUP\*: phylogenetic analysis using parsimony (\*and other methods)*, Version 4.0b10. Computer software and manual. Sunderland, MA: Sinauer Associates.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011.** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**: 2731–2739.
- Tavaré S. 1986.** Some probabilistic and statistical problems

- in the analysis of DNA sequences. In: Miura RM, ed. *Some mathematical questions in biology – DNA sequence analysis*. Providence, RI: American Mathematical Society, 57–86.
- Taylor MS, Hellberg ME. 2005.** Marine radiations at small geographic scales: speciation in neotropical reef gobies (*Elacatinus*). *Evolution* **59**: 374–385.
- Tringali MD, Bert TM, Seyoum S, Bermingham E, Bartolacci D. 1999.** Molecular phylogenetics and ecological diversification of the Transisthmian fish genus *Centropomus* (Perciformes: Centropomidae). *Molecular Phylogenetics and Evolution* **13**: 193–207.
- Tsigenopoulos CS, Berrebi P. 2000.** Molecular phylogeny of North Mediterranean freshwater barbs (genus *Barbus*: Cyprinidae) inferred from cytochrome *b* sequences: biogeographic and systematic implications. *Molecular Phylogenetics and Evolution* **14**: 165–179.
- Turan D, Berber S, Topkara ET, Verep B. 2005.** A first record [*Knipowitschia longicaudata* (Kessler, 1877)] for the fish fauna of Lake Manyas. *Turkish Journal of Zoology* **29**: 171–176.
- Van Neer W, Wildekamp RH, Kucuk F, Unlusayin M. 1999.** First inland records of the euryhaline goby *Knipowitschia caucasica* from lakes in Anatolia, Turkey. *Journal of Fish Biology* **54**: 1334–1337.
- Vanhove MPM, Kovačić M, Koutsikos NE, Zogaris S, Vardakas LE, Huyse T, Economou AN. 2011.** First record of a landlocked population of marine *Millerigobius macrocephalus* (Perciformes: Gobiidae): observations from a unique spring-fed karstic lake (Lake Vouliagmeni, Greece) and phylogenetic positioning. *Zoologischer Anzeiger* **250**: 195–204.
- Vogiatzi E, Kalogianni E, Giakoumi S, Magoulas A, Tsigenopoulos C. 2009.** Genetic variation and biogeography of the Corfu toothcarp *Valencia letourneuxi* Sauvage, 1880 (Pisces, Valenciidae) based on microsatellite markers. Poster presentation. Eleventh International Congress on the Zoogeography and Ecology of Greece and Adjacent Regions, Herakleion, Crete, 21–25 September 2009. Book of Abstracts, p. 202.
- Vogler AP, DeSalle R. 1994.** Diagnosing units of conservation management. *Conservation Biology* **8**: 354–363.
- Wan Q-H, Wu H, Fujihara T, Fang S-G. 2004.** Which genetic marker for which conservation genetics issue? *Electrophoresis* **25**: 2165–2176.
- Westaway R. 1996.** Quaternary elevation change of the Gulf of Corinth in Central Greece. *Philosophical Transactions of the Royal Society Series A, Mathematical, Physical and Engineering Sciences* **354**: 1125–1164.
- Wilson AB. 2006.** Interspecies mating in sympatric species of *Syngnathus* pipefish. *Molecular Ecology* **15**: 809–824.
- Zardoya R, Economidis PS, Doadrio I. 1999.** Phylogenetic relationships of Greek Cyprinidae: molecular evidence for at least two origins of the Greek cyprinid fauna. *Molecular Phylogenetics and Evolution* **13**: 122–132.
- Zelilidis A. 2003.** The geometry of fan-deltas and related turbidites in narrow linear basins. *Geological Journal* **38**: 31–46.
- Zogaris S, Economou AN, Dimopoulos P. 2009.** Ecoregions in the Southern Balkans: should their boundaries be revised? *Environmental Management* **43**: 682–697.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Table S1.** GTR+I+ $\Gamma$  pairwise genetic distances (in %) between the haplotypes included in the phylogenetic analysis, for a fragment of 815 bp of 12S and 16S rDNA. The number of times a haplotype was found on a given location is provided in parenthesis. Haplotypes preceded by an asterisk (\*) were retrieved from GenBank. Sequences were available for *Knipowitschia milleri* (GQ398133-7) and *Economidichthys pygmaeus* (GQ398131-2), of which additional specimens were sequenced.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.