

Antarctic macrobenthic communities: A compilation of circumpolar information

Julian Gutt¹, David K. A. Barnes², Susanne J. Lockhart³, Anton van de Putte⁴

1 Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Columbusstr., 27568, Bremerhaven, Germany **2** British Antarctic Survey, High Cross, Madingley Road, CB3 0ET, Cambridge, United Kingdom **3** NOAA Southwest Fisheries Science Center, La Jolla Shore Drive, CA 92037, La Jolla, United States **4** ANTA-BIF, Vautierstraat 29, B-1000 Brussels, Belgium

Corresponding author: *Julian Gutt* (julian.gutt@awi.de)

Academic editor: *L. Penev* | Received 12 December 2012 | Accepted 12 February 2013 | Published 19 February 2013

Citation: Gutt J, Barnes DKA, Lockhart SJ, van de Putte A (2013) Antarctic macrobenthic communities: A compilation of circumpolar information. *Nature Conservation* 4: 1–13. doi: 10.3897/natureconservation.4.4499

Abstract

Comprehensive information on Antarctic macrobenthic community structure has been publicly available since the 1960s. It stems from trawl, dredge, grab, and corer samples as well as from direct and camera observations (Table 1–2). The quality of this information varies considerably; it consists of pure descriptions, figures for presence (absence) and abundance of some key taxa or proxies for such parameters, e.g. sea-floor cover. Some data sets even cover a defined and complete proportion of the macrobenthos with further analyses on diversity and zoogeography. As a consequence the acquisition of data from approximately 90 different campaigns assembled here was not standardised. Nevertheless, it was possible to classify this broad variety of known macrobenthic assemblages to the best of expert knowledge (Gutt 2007; Fig. 1). This overview does not replace statistically sound community and diversity analyses. However, it shows from where which kind of information is available and it acts as an example of the feasibility and power of such data collections. The data set provides unique georeferenced biological basic information for the planning of future coordinated research activities, e.g. under the umbrella of the biology program “Antarctic Thresholds - Ecosystem Resilience and Adaptation” (AnT-ERA) of the Scientific Committee on Antarctic Research (SCAR) and especially for actual conservation issues, e.g. the planning of Marine Protected Areas (MPAs) by the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR).

Keywords

Macrobenthic communities, trawls, dredges, grabs, corers, direct observations (scuba-diving, sea-bed video, sea-bed photography)

Data resources

Data published through GBIF: <http://ipt.biodiversity.aq/resource.do?r=macrobenthos>

Seabed images through Pangaea: <http://www.pangaea.de/> (sample: <http://doi.pangaea.de/10.1594/PANGAEA.702075>)

General description

Additional information: Additional files uploaded: list of references (Table 1–2) and classification of macrobenthic communities (Fig. 1).

Project details

Project title: Antarctic macrobenthic communities: A compilation of circumpolar information.

Personnel: Julian Gutt.

Taxonomic coverage

General taxonomic coverage description (for detailed information see references in Table 1): Macrobenthic communities have been uploaded in the category “vernacularNames”, abbreviations in “taxonRemarks”.

“Sessile suspension feeders and associated fauna” can be dominated by both demosponges, e.g., *Cinachyra* or *Mycale* and hexactinellid (glass) sponges. The most common genus is *Rossella*. The sponges include fast growing genera, such as *Homaxinella* or those that grow slowly, at least during the adult life stage, such as the also common hexactinellid genus *Anoxycalix*. The associated fauna comprises specialised predators, such as nudibranches, asteroids (especially *Acodontaster conspicuus* and *Perknaster fuscus*, which control fast growing *Mycale acerata* populations) and gastropods. Other fauna groups are symbionts, amphipods and other macroorganisms that prefer an epibiotic life-mode (mainly from the echinoderms, such as sedentary holothurians, ophiuroids and crinoids). If space is not monopolised by sponges, then, cnidarians (such as gorgonians, pennatularians, alconarians or hydrocorals), solitary and compound ascidians, and a variety of bryozoans can be most abundant. A recently described population of lithodid crabs is speculated to grow fast due to oceanic warming and was associated with the “mobile deposit feeders, infauna and grazers”. Other mobile epifauna assemblages can be dominated in shallow areas by the asteroid *Acodontaster validus*, by two species of the grazing echinoid *Sterechinus*, a variety of deposit feeding and scavenging ophiuroids and mobile holothurians. The infauna is comparably rare; however, polychaetes and the clams *Yoldia* as well as *Laternula* can reach high densities, especially in shallow muddy sediments. A

general depth gradient exists for biomass and abundances. In addition, very low biomass and abundances are found in shallow habitats that are physically and permanently disturbed by sea-scour, in intermediately deep shelf areas that are scoured by icebergs and in extremely oligotrophic situations under or close to the ice-shelves. Intensively disturbed assemblages can be dominated by very few species, appearing to be almost "monospecific", during recolonisation by pioneers such as the ascidian *Mogula pedunculata*, bryozones like *Cellarinella* and *Cellaria* or the gorgonian *Primnoisis antarctica* or in physically disturbed areas, where only opportunistic mobile species survive. Locally clams of the species *Adamussium colbecki* can live in several layers on top of each other simply due to suitable environmental conditions and low competition. Species can also become very abundant when they are better local competitors for space, such as the demosponge *Cinachyra barbata*, s.l. Recently, fauna-rich vent sites and far poorer seeps have been discovered.

Common names: sessile suspension feeders and associated fauna (SSFA), sessile suspension feeders and associated fauna - predator driven (SSFA-PRED), sessile suspension feeders and associated fauna - dominated by sponges (SSFA-SPO), sessile suspension feeders and associated fauna - dominated by taxa other than sponges (SSFA-OTH), mixed assemblage (MIX), very low biomass or absence of trophic guilds (VLB), "monospecific" (MONO), physically controlled (PHYCO), mobile deposit feeders, infauna and grazers (MOIN), mobile deposit feeders, infauna and grazers - infauna dominated (MOIN-INF), mobile deposit feeders, infauna and grazers - epi-fauna dominated (MOIN-EPI), vent (VENT), and seep (SEEP).

Spatial coverage

General spatial coverage: The study area generally covers almost the entire Southern Ocean, including single ice-shelf covered sites (Fig. 2). The vast majority of information is from shelf areas around the continent at water depth shallower than 800m. Non-ice shelf covered shelf areas can be up to 300km wide or the shelf-edge at 600 to 800m depth can "disappear" beneath the floating ice-shelf. Shallow areas (<50m) are rare because 86% of the coast-line is glaciated or consists of an ice-shelf edge. A non-glaciated coast mainly exists along the Antarctic Peninsula. The coastline is either extremely complex with bays, channels, peninsulas, islands etc. or less structured, especially where it is formed by the ice-shelf. Overdeepened basins (inner-shelf depressions) can reach >1200m water depth. Most islands exist west of the Antarctic Peninsula and along the Scotia Arc linking the Peninsula with the southern tip of South America. The coastal waters are mainly affected by the Antarctic Coastal Current (East Wind Drift), whilst the largest off-shore part of the Southern Ocean is dominated by the Antarctic Circumpolar Current (West Wind Drift) with gyres of different size. Sediments are predominantly poorly sorted but also cobble "fields", bedrock, and pure soft sediments exist. The Antarctic marine ecosystem is shaped by a distinct seasonality of the sea-ice cover affecting a short and intensive primary production in austral summer, by predominantly stable low temperature to which most organisms are thought

to be specifically adapted to, and very little terrestrial run-off. Most of the shelf-inhabiting macrobenthic species are endemic; some taxa reach above-average species richness (Clarke and Johnston 2003). Only few marine habitats are protected, most of which are small. Plans and proposals for large Marine Protected Areas (MPA's), e.g. in East Antarctica, in the Ross and Weddell Seas, exist but require good scientific knowledge and data to be meaningful.

Coordinates: 83°0'0"S and 52°0'0"S Latitude; 180°0'0"W and 180°0'0"E Longitude.

Temporal coverage: March 1, 1956–February 21, 2010.

Methods

Method step description: Attribution of the information from the different sources (for references see Table 1, for hyperlinks see Table 2) to the classified macrobenthic assemblages (Fig. 1) was done to the best of expert knowledge. This was done for the entire data set simultaneously and the results were made publically available for the first time via the database “Antarctic Biodiversity Facility” (ANTABIF). The principal parameter on which these assumptions have been made was biomass or a proxy for biomass such as sea-floor coverage. Some information on benthic functioning is also included directly or indirectly, e.g. predation, competition, succession after ice-berg scouring, epi-biotic life-mode and oligotrophic conditions under ice shelves. The source publications listed (Table 1) comprise descriptions of catches, other observations, and data on fauna and were mainly from historical and modern peer-reviewed articles. Other information sources were sea-bed videos and still images together with associated meta-data (Table 2). All the latter source material has an associated DOI and is available at the database PANGAEA (www.pangaea.de).

Study extent description: Southern Ocean with emphasize on coastal shelf areas and some islands without specific temporal patterns of sampling.

Sampling description: This project aggregates data from various expeditions with a full range of benthic sampling methods, such as grabs, corers, dredges, and trawls as well as non-invasive observations by scuba divers, stationary, towed, or ROV-based still and video-cameras. For detail descriptions see original publications in journals (Table 1) or data repositories (Table 2).

Quality control description: A first version of the classification of the macrobenthic communities had been published in a peer-reviewed journal (Gutt 2007). A modified version had been published in the Antarctic Climate Change and the Environment report (ACCE, Turner et al. 2009). The actual version is depicted in Fig. 1. Data presented here is available at ANTABIF/SCAR-MarBIN and will contribute to the biogeographic atlas project of SCAR and the Census of Marine Life (De Broyer et al. in prep.), <http://atlas.biodiversity.aq/>.

Table 1. References of results and data used for the compilation of information on Antarctic macrobenthic communities presented in this article.

- Arnaud P (1974) Contribution a la bionomie marine benthique des régions antarctiques et subantarctiques. *Téthys* 6: 1–464.
- Azam F, Beers JR, Campbell L, Carlucci AF, Holm-Hansen O, Reis FMH, Karl DM (1979) Occurrence and metabolic activity of organisms under the Ross Ice Shelf, Antarctica, at station J9. *Science* 203: 451–453.
- Bannasch R, Feiler K, Rauschert M (1984) Fortsetzung der biologischen Untersuchungen im Gebiet der sowjetischen Antarktisstation Bellingshausen. *Geodätische und geophysikalische Veröffentlichungen, Reihe 1, Heft 11*: 37–55.
- Barnes DKA (1995a) Sublittoral epifaunal communities at Signy Island, Antarctica. I. The ice-foot zone. *Marine Biology* 121: 555–563.
- Barnes DKA (1995b) Sublittoral epifaunal communities at Signy Island, Antarctica. II. Below the ice-foot zone. *Marine Biology* 121: 565–572.
- Barnes DKA, Clarke A (1995) Epibiotic communities on sublittoral macroinvertebrates at Signy Island, Antarctica. *Journal of the Marine Biological Association of the United Kingdom* 75: 689–703.
- Barry JP, Dayton PK (1988) Current patterns in McMurdo Sound, Antarctica and their relationship to local biotic communities. *Polar Biology* 8: 367–376.
- Barthel D, Gutt J (1992) Sponge associations in the eastern Weddell Sea. *Antarctic Science* 4: 137–150.
- Beaman RJ, O'Brien PE (2009) Collaborative East Antarctic Marine Census (CEAMARC): Post-Survey Report, RSV Aurora Australis Voyage 3, December 2007 - January 2008. *Geoscience Australia Record 2008/05*, 61 pp.
- Bellisio NB, Lopez RB, Tomo AP (1972) Distribucion vertical de la fauna bentonica en tres localidades antarticas: Bahía Esperanza, Isla Petermann y Archipelago Melchior. *Contribucion Instituto Antartico Argentino* 142: 1–89.
- Belyaev GM, Ushakov PV (1957) Certain regularities in the quantitative distribution of the bottom fauna in Antarctic waters. *Doklady Akademii Nauk SSSR* 112: 137–140.
- Bowden DA, Schiaparelli S, Clark MR, Rickard GJ (2011) A lost world? Archaic crinoid-dominated assemblages on an Antarctic seamount. *Deep-Sea Research II* 58: 119–127.
- Bruchhausen PM, Raymond JA, Jacobs SS, DeVries AL, Thorndike EM, DeWitt HH (1979) Fish, crustaceans, and the sea floor under the Ross Ice Shelf. *Science* 203: 449–451.
- Bullivant JS (1967) Ecology of the Ross Sea benthos. *Bulletin of the New Zealand Department of Scientific and Industrial Research* 176: 49–78.
- Castellanos ZJA de (1973) Estratificacion del complejo bentonico de invertebrados en Puerto Paraiso (Antartida). *Contribucion del Instituto Antartico Argentino* 164: 4–23.
- Cattaneo-Vietti R, Chiantore M, Albertelli G (1997) The population structure and ecology of the Antarctic scallop *Adamussium colbecki* (Smith, 1902) at Terra Nova Bay (Ross Sea, Antarctica). *Scientia Marina, Supl.* 2: 15–24.
- Cattaneo-Vietti R, Chiantore M, Gambi MC, Albertelli G, Cormaci M, Di Geronimo I (2000a) Spatial and vertical distribution of benthic littoral communities in Terra Nova Bay. In: Faranda FM, Guglielmo L, Ionora A (Eds) *Ross Sea Ecology*. Springer, Berlin, 503–514.
- Cattaneo-Vietti R, Bavestrello G, Cerrano C, Gaino E, Mazzella L, Pansini M, Sarà M (2000b) The role of sponges in the Terra Nova Bay ecosystem. In: Faranda FM, Guglielmo L, Ionora A (Eds) *Ross Sea Ecology*. Springer, Berlin, 539–549.
- Cerrano C, Bavestrello G, Calcinai B, Cattaneo-Vietti R, Sarà A (2000) Asteroids eating sponges from Tethys Bay, East Antarctica. *Antarctic Science* 12: 425–426.
- Chiantore M, Cattaneo-Vietti R, Berkman PA, Nigro M, Vacchi M, Schiaparelli H, Albertelli G (2001) Antarctic scallop (*Adamussium colbecki*) spatial population variability along the Victoria Land Coast, Antarctica. *Polar Biology* 24: 139–143.
- Cranmer TL, Ruhl HA, Baldwin RJ, Kaufmann RS (2003) Spatial and temporal variation in the abundance, distribution and population structure of epibenthic megafauna in Port Foster, Deception Island. *Deep-Sea Research II* 50: 1821–1842.

- Davis RW, Castellini MA, Horning M, Davis MP (1983) Maintenance of an observation hole through the McMurdo Ice Shelf for winter oceanography. *Antarctic Journal of the United States* 18: 12–14.
- Dayton PK (1979) Observations on growth, dispersal and population dynamics of some sponges in McMurdo Sound, Antarctica. *Colloques internationaux du Centre national de la recherche scientifique* 291: 271–282.
- Dayton PK, Oliver JS (1977) Antarctic soft-bottom benthos in oligotrophic and eutrophic environments. *Science* 197: 55–58.
- Dayton PK, Kooyman GL, Barry JP (1984) Benthic life under thick ice. *Antarctic Journal of the United States* 19: 128.
- Dayton PK, Robillard GA, Paine RT (1970) Benthic faunal zonation as a result of anchor ice at McMurdo Sound Antarctica. In: Holdgate MW (Ed) *Antarctic Ecology*. Academic Press, New York, 244–258.
- Dayton PK, Robillard GA, Paine RT, Dayton LB (1974) Biological accommodation in the benthic community at McMurdo Sound, Antarctica. *Ecological Monographs* 44: 105–128.
- Domack E, Ishman S, Leventer A, Sylva S, Willmott V, Huber B (2005) A chemotrophic ecosystem found beneath Antarctic ice shelf. *EOS* 86 (269): 271–272.
- Drescher HE, Hubold G, Piatkowski U, Plötz J, Voß J (1983) Das biologische Programm der ANTARKTIS-I-Expedition mit FS “Polarstern”. *Reports on Polar Research* 12: 1–34.
- Everitt DA, Poore GCB, Pickard J (1980) Marine benthos from Davis Station, east Antarctica. *Australian Journal of Freshwater Research* 31: 829–836.
- Gallardo VA, Castillo JG, Retamal MA, Yáñez A, Moyano HI, Hermosilla JG (1977) Quantitative studies on the soft-bottom macrobenthic animal communities of shallow Antarctic bays. In: Llano GA (Ed) *Adaptations within Antarctic ecosystems*. Smithsonian Institution, Washington, DC, 361–387.
- Gerdes D, Klages M, Arntz WE, Herman RL, Galéron J, Hain S (1992) Quantitative investigations on macrobenthos communities of the southeastern Weddell Sea shelf based on multibox corer samples. *Polar Biology* 12: 291–301.
- Gruzov EN, Pushkin AF (1970) Bottom communities of the upper sublittoral of Enderby Land and the South Shetlands. In: Holdgate MW (Ed) *Antarctic Ecology*, Vol. 1., Academic Press, London, 235–238.
- Gruzov EN, Propp MV, Pushkin AF (1968) Biological associations of coastal areas of the Davis Sea (based on the observations of divers). *Soviet Antarctic Expedition Information Bulletin* 6(6): 523–533.
- Gutt J, Piepenburg D (1991) Dense aggregations of deep-sea holothurians in the southern Weddell Sea, Antarctica. *Marine Ecology Progress Series* 68: 277–285.
- Gutt J, Starmans A (1998) Structure and biodiversity of megabenthos in the Weddell and Lazarev Seas (Antarctica): ecological role of physical parameters and biological interactions. *Polar Biology* 20: 229–247.
- Gutt J (2007) Antarctic macro-zoobenthic communities: a review and an ecological classification. *Antarctic Science* 19: 165–182.
- Gutt J, Koubbi P, Eléaume M (2007) Mega-epibenthic diversity off Terre Adélie (Antarctica) in relation to disturbance. *Polar Biology* 30: 1323–1329.
- Gutt J, Barratt I, Domack E, Udekem d’Acoz C, Dimmler W, Grémare A, Heilmayer O, Isla E, Janussen D, Jorgensen E, Kock K-H, Lehnert LS, López-González P, Langner S, Linse K, Manjón-Cabeza ME, Meißner M, Montiel A, Raes M, Robert H, Rose A, Sañé-Schepisi E, Saucède T, Scheidat M, Schenke H-W, Seiler J, Smith C (2010) Biodiversity change after climate-induced ice-shelf collapse in the Antarctic. *Deep-Sea Research II* 58: 74–83.
- Hamada E, Numanami H, Naito Y, Taniguchi A (1986) Observation on the marine benthic organisms at Syowa Station in Antarctica using a remotely operated vehicle. *Memoirs of National Institute of Polar Research (Japan)* 40: 289–298.
- Jazdzewski K, Juraz W, Kittel W, Presler P, Sicinski J (1986) Abundance and biomass estimates of the benthic Fauna in Admiralty Bay, King George Island, South Shetland Islands. *Polar Biology* 6: 5–16.
- Jazdzewski K, De Broyer C, Pudlarz M, Zielinski D (2001) Seasonal fluctuations of vagile benthos in the uppermost sublittoral of a maritime Antarctic fjord. *Polar Biology* 24: 910–917.
- Jones CD, Lockhart SJ (2011) Detecting Vulnerable Marine Ecosystems in the Southern Ocean using research trawls and underwater imagery. *Marine Policy* 35(5): 7732–7736.

- Jones DOB (2005) Ecological controls on density, diversity and community structure of polar megabenthos. PhD thesis, Southampton, UK: University of Southampton.
- Kirkwood JM, Burton HR (1988) Macrobenthic species assemblages in Ellis Fjord, Vestfold Hills, Antarctica. *Marine Biology* 97: 445–457.
- Kohnen H (1984) Die Expedition ANTARKTIS-II mit FS 'Polarstern' 1983/84. Reports on Polar Research 19: 1–185.
- Lipps JH, Ronan TE Jr, DeLaca TE (1979) Life below the Ross Ice Shelf, Antarctica. *Science* 203: 447–449.
- Littlepage JL, Pearse JS (1962) Biological and oceanographic observations under an Antarctic ice shelf. *Science* 137: 679–681.
- Lockhart SJ, Jones CD (2008) Biogeographic patterns of benthic invertebrate megafauna on shelf areas within the Southern Ocean Atlantic sector. *CCAMLR Science* 15: 167–192.
- Lovell LL, Trego KD (2003) The epibenthic megafaunal and benthic infaunal invertebrates of Port Foster, Deception Island (South Shetland Islands, Antarctica). *Deep-Sea Research II* 50: 1799–1819.
- Lowry JK (1975) Soft bottom macrobenthic community of Arthur Harbor, Antarctica. *Antarctic Research Series* 23: 1–19.
- Nakajima Y, Watanabe K, Naito Y (1982) Diving observations of the marine benthos at Syowa station, Antarctica. *Memoirs of the National Institute of Polar Research (Japan), Special Issue* 23: 44–54.
- Niemann H, Fischer D, Graffe D, Knittel K, Montiel A, Heilmayer O, Nöthen K, Pape T, Kasten S, Bohrmann G, Boetius A, Gutt J (2009) Biogeochemistry of a low-activity cold seep in the Larsen B area, western Weddell Sea, Antarctica. *Biogeosciences* 6: 2383–2395.
- Oliver JS, Slattery PN (1985) Effects of crustacean predators on species composition and population structure of soft-bodied infauna from McMurdo Sound, Antarctica. *Ophelia* 24: 155–175.
- Oliver JS, Watson DJ, O'Connor EF, Dayton PK (1976) Benthic communities of McMurdo Sound. *Antarctic Journal of the U.S.* 11: 58–549.
- Pabis K, Siciński J, Krymariy M (2011) Distribution patterns in the biomass of macrozoobenthic communities in Admiralty Bay (King George Island, South Shetlands, Antarctic). *Polar Biology* 34: 489–500.
- Piepenburg D, Voß J, Gutt J (1997) Assemblages of sea stars (Echinodermata: Asteroidea) and brittle stars (Echinodermata: Ophiuroidea) in the Weddell Sea (Antarctica) and off Northeast Greenland (Arctic): a comparison of diversity and abundance. *Polar Biology* 17: 305–322.
- Post AL, Hemer MA, O'Brien PE, Roberts D, Craven M (2007) History of benthic colonisation beneath the Amery Ice Shelf, East Antarctica. *Marine Ecology Progress Series* 344: 29–37.
- Post AL, O'Brien PE, Beaman RJ, Riddle MJ, de Santis L (2010a) Physical controls on deep water coral communities on the George V Land slope, East Antarctica. *Antarctic Science* 22: 371–378.
- Post AL, Beaman RJ, O'Brien PE, Eléaume M, Riddle MJ (2011) Community structure and benthic habitats across the George V Shelf, East Antarctica: trends through space and time. *Deep-Sea Research II* 58: 105–118.
- Propp MV (1977) The study of bottom fauna at Haswell Island by scuba diving. In: Holdgate MW (Ed) *Antarctic Ecology*, Vol. 1., Academic Press, London, 239–241.
- Raguá-Gil JM, Gutt J, Clarke A, Arntz WE (2004) Antarctic shallow-water mega-epibenthos: shaped by circumpolar dispersion or local conditions? *Marine Biology* 144: 829–839.
- Rehm P, Hooke RA, Thatje S (2011) Macrofaunal communities on the continental shelf off Victoria Land, Ross Sea, Antarctica. *Antarctic Science* 23: 449–455.
- Riddle MJ, Craven M, Goldsworthy PM, Carsey F (2007) A diverse benthic assemblage 100 km from open water under the Amery Ice Shelf, Antarctica. *Paleoceanography* 22, doi: 10.1029/2006PA001327
- Rogers AD, Tyler PA, Connelly DP, Copley JT, James R, Larter RD, Linse K, Mills RA, Naveira Garabato A, Pancost RD, Pearce DA, Polunin NVC, German CR, Shank T, Boersch-Supan PH, Alker BJ, Aquilina A, Bennett SA, Clarke A, Dinley RJJ, Graham AGC, Green DRH, Hawkes JA, Hepburn L, Hilario A, Huvenne VAI, Marsh L, Ramirez-Llodra E, Reid WDK, Roterman CN, Sweeting CJ, Thatje S, Zwirgmaier K (2012) The discovery of new deep-sea hydrothermal vent communities in the Southern Ocean and implications for biogeography. *PLoS Biology* 10(1): e1001234. doi: 10.1371/journal.pbio.1001234
- Sahade R, Tatián M, Kowalke J, Kühne S, Esnal GB (1998) Benthic faunal associations on soft substrates at Potter Cove, King George Island, Antarctic. *Polar Biology* 19: 85–91.

- Saiz JI, García FJ, Manjón-Cabeza ME, Parapar J, Peña-Cantero A, Saucède T, Troncoso JS, Ramos A (2008) Community structure and spatial distribution of benthic fauna in the Bellingshausen Sea (West Antarctica). *Polar Biology* 31: 735–743 (only stations <1000m considered).
- Sicinski J, Pabis K, Jazdzewski K, Konopacka A, Blazewicz-Paszkwycz M (2011) Macrobenthos of two Antarctic glacial covres: a comparison with non-disturbed bottom areas. *Polar Biology* 35: 355–367.
- Smith CR, Grange LJ, Honig DL, Naudts L, Huber B, Guidi L, Domack E (2012) A large population of king crabs in Palmer Deep on the west Antarctic Peninsula shelf and potential invasive impacts. *Proceedings of the Royal Society B, Biological Sciences* 279: 1017–1026.
- Starmans A, Gutt J, Arntz WE (1999) Mega-epibenthic communities in Arctic and Antarctic shelf areas. *Marine Biology* 135: 269–280.
- Teixidó N, Garrabou J, Arntz WE (2002) Spatial pattern quantification of Antarctic benthic communities using landscape indices. *Marine Ecology Progress Series* 242: 1–14.
- Sumida PYG, Bernardino AF, Stedall VP, Glover AG, Smith CR (2008) Temporal changes in benthic megafaunal abundance and composition across the West Antarctic Peninsula shelf: Results from video surveys. *Deep-Sea Research II* 55: 2465–2477.
- Teixidó N, Garrabou J, Gutt J, Arntz WE (2004) Recovery in Antarctic benthos after iceberg disturbance: trends in benthic composition, abundance, and growth forms. *Marine Ecology Progress Series* 278: 1–16.
- Teixidó N, Garrabou J, Gutt J, Arntz WE (2007) Iceberg disturbance and successional spatial patterns: the case of the shelf Antarctic benthic communities. *Ecosystems* 10: 142–157.
- Voß J (1988) Zoogeographie und Gemeinschaftsanalyse des Makrozoobenthos des Weddellmeeres (Antarktis). *Berichte zur Polarforschung* 45: 1–145.
- Watters G, Bergström B, Gutt J, Petterson J-O, Rosenberg J, Setran A, Valenzuela C (1995) 9. Leg III: Crab and epifaunal surveys of bays, anchorages, and fjords around South Georgia. AMLR 1994/95 Field Season Report, Administrative Report LJ-95–13, National Oceanic and Atmospheric Administration, Southwest Fisheries Science Center, Antarctic Ecosystem Research Group, 113–130.
- Zamorano JH (1983) Zonación y Biomasa de la macrofauna bentónica en Bahía South, Archipiélago de Palmer, Antártica. *Serie científica. Instituto Antártico Chileno* 30: 27–38.

Table 2. Hyperlinks (DataCite DOIs), which provide access to seabed images and metadata from single stations where the images have been taken. The macrobenthos depicted in these images was classified and used for the compilation of information on Antarctic macro-benthic communities presented in this article.

doi: 10.1594/PANGAEA.702075	doi: 10.1594/PANGAEA.770359
doi: 10.1594/PANGAEA.702059	doi: 10.1594/PANGAEA.198690
doi: 10.1594/PANGAEA.702076	doi: 10.1594/PANGAEA.198691
doi: 10.1594/PANGAEA.702077	doi: 10.1594/PANGAEA.198692
doi: 10.1594/PANGAEA.702062	doi: 10.1594/PANGAEA.198693
doi: 10.1594/PANGAEA.702078	doi: 10.1594/PANGAEA.198694
doi: 10.1594/PANGAEA.702064	doi: 10.1594/PANGAEA.198695
doi: 10.1594/PANGAEA.702065	doi: 10.1594/PANGAEA.198696
doi: 10.1594/PANGAEA.702066	doi: 10.1594/PANGAEA.198697
doi: 10.1594/PANGAEA.702067	doi: 10.1594/PANGAEA.198698
doi: 10.1594/PANGAEA.702079	doi: 10.1594/PANGAEA.198699
doi: 10.1594/PANGAEA.702069	doi: 10.1594/PANGAEA.198667
doi: 10.1594/PANGAEA.702070	doi: 10.1594/PANGAEA.198668
doi: 10.1594/PANGAEA.702080	doi: 10.1594/PANGAEA.198669
doi: 10.1594/PANGAEA.702072	doi: 10.1594/PANGAEA.198670
doi: 10.1594/PANGAEA.702073	doi: 10.1594/PANGAEA.198671
doi: 10.1594/PANGAEA.702074	doi: 10.1594/PANGAEA.198672

doi: 10.1594/PANGAEA.198673

doi: 10.1594/PANGAEA.198674

doi: 10.1594/PANGAEA.198675

doi: 10.1594/PANGAEA.198676

doi: 10.1594/PANGAEA.198677

doi: 10.1594/PANGAEA.198678

doi: 10.1594/PANGAEA.198679

doi: 10.1594/PANGAEA.198680

doi: 10.1594/PANGAEA.198681

doi: 10.1594/PANGAEA.198682

doi: 10.1594/PANGAEA.206413

doi: 10.1594/PANGAEA.198683

doi: 10.1594/PANGAEA.198684

doi: 10.1594/PANGAEA.198685

doi: 10.1594/PANGAEA.198686

doi: 10.1594/PANGAEA.198687

doi: 10.1594/PANGAEA.198688

doi: 10.1594/PANGAEA.198689

doi: 10.1594/PANGAEA.250203

doi: 10.1594/PANGAEA.250204

doi: 10.1594/PANGAEA.250205

doi: 10.1594/PANGAEA.250206

doi: 10.1594/PANGAEA.250207

doi: 10.1594/PANGAEA.250208

doi: 10.1594/PANGAEA.250209

doi: 10.1594/PANGAEA.250210

doi: 10.1594/PANGAEA.250211

doi: 10.1594/PANGAEA.250212

doi: 10.1594/PANGAEA.250213

doi: 10.1594/PANGAEA.250214

doi: 10.1594/PANGAEA.250215

doi: 10.1594/PANGAEA.250216

doi: 10.1594/PANGAEA.250217

doi: 10.1594/PANGAEA.250218

doi: 10.1594/PANGAEA.250219

doi: 10.1594/PANGAEA.198643

doi: 10.1594/PANGAEA.198646

doi: 10.1594/PANGAEA.198644

doi: 10.1594/PANGAEA.198645

doi: 10.1594/PANGAEA.198647

doi: 10.1594/PANGAEA.198648

doi: 10.1594/PANGAEA.198649

doi: 10.1594/PANGAEA.198650

doi: 10.1594/PANGAEA.198653

doi: 10.1594/PANGAEA.198654

doi: 10.1594/PANGAEA.198656

doi: 10.1594/PANGAEA.198657

doi: 10.1594/PANGAEA.198655

doi: 10.1594/PANGAEA.198658

doi: 10.1594/PANGAEA.198659

doi: 10.1594/PANGAEA.198660

doi: 10.1594/PANGAEA.198661

doi: 10.1594/PANGAEA.198662

doi: 10.1594/PANGAEA.198663

doi: 10.1594/PANGAEA.198664

doi: 10.1594/PANGAEA.198665

doi: 10.1594/PANGAEA.198666

doi: 10.1594/PANGAEA.757741

doi: 10.1594/PANGAEA.757740

doi: 10.1594/PANGAEA.227308

doi: 10.1594/PANGAEA.227309

doi: 10.1594/PANGAEA.227310

doi: 10.1594/PANGAEA.227311

doi: 10.1594/PANGAEA.666972

doi: 10.1594/PANGAEA.666973

doi: 10.1594/PANGAEA.666974

doi: 10.1594/PANGAEA.666975

doi: 10.1594/PANGAEA.666976

doi: 10.1594/PANGAEA.666977

doi: 10.1594/PANGAEA.666978

doi: 10.1594/PANGAEA.666979

doi: 10.1594/PANGAEA.666980

doi: 10.1594/PANGAEA.666981

doi: 10.1594/PANGAEA.666982

doi: 10.1594/PANGAEA.666983

doi: 10.1594/PANGAEA.666984

doi: 10.1594/PANGAEA.666986

doi: 10.1594/PANGAEA.666991

doi: 10.1594/PANGAEA.666992

doi: 10.1594/PANGAEA.666993

doi: 10.1594/PANGAEA.666994

doi: 10.1594/PANGAEA.666995

doi: 10.1594/PANGAEA.666996

doi: 10.1594/PANGAEA.666997

doi: 10.1594/PANGAEA.227670

doi: 10.1594/PANGAEA.220741

doi: 10.1594/PANGAEA.220742

doi: 10.1594/PANGAEA.220743

doi: 10.1594/PANGAEA.220744

doi: 10.1594/PANGAEA.220745

doi: 10.1594/PANGAEA.220746

doi: 10.1594/PANGAEA.220747

doi: 10.1594/PANGAEA.319912

doi: 10.1594/PANGAEA.319913

doi: 10.1594/PANGAEA.319914	doi: 10.1594/PANGAEA.667026
doi: 10.1594/PANGAEA.319915	doi: 10.1594/PANGAEA.667027
doi: 10.1594/PANGAEA.319917	doi: 10.1594/PANGAEA.667028
doi: 10.1594/PANGAEA.319918	doi: 10.1594/PANGAEA.667029
doi: 10.1594/PANGAEA.319916	doi: 10.1594/PANGAEA.667030
doi: 10.1594/PANGAEA.319919	doi: 10.1594/PANGAEA.667031
doi: 10.1594/PANGAEA.319920	doi: 10.1594/PANGAEA.667032
doi: 10.1594/PANGAEA.319894	doi: 10.1594/PANGAEA.691559
doi: 10.1594/PANGAEA.319895	doi: 10.1594/PANGAEA.691560
doi: 10.1594/PANGAEA.319896	doi: 10.1594/PANGAEA.740036
doi: 10.1594/PANGAEA.319897	doi: 10.1594/PANGAEA.691561
doi: 10.1594/PANGAEA.319898	doi: 10.1594/PANGAEA.691562
doi: 10.1594/PANGAEA.319899	doi: 10.1594/PANGAEA.691563
doi: 10.1594/PANGAEA.319900	doi: 10.1594/PANGAEA.691564
doi: 10.1594/PANGAEA.319901	doi: 10.1594/PANGAEA.691565
doi: 10.1594/PANGAEA.319902	doi: 10.1594/PANGAEA.691566
doi: 10.1594/PANGAEA.319903	doi: 10.1594/PANGAEA.691567
doi: 10.1594/PANGAEA.319904	doi: 10.1594/PANGAEA.691568
doi: 10.1594/PANGAEA.319905	doi: 10.1594/PANGAEA.691569
doi: 10.1594/PANGAEA.319906	doi: 10.1594/PANGAEA.691570
doi: 10.1594/PANGAEA.319907	doi: 10.1594/PANGAEA.691571
doi: 10.1594/PANGAEA.319908	doi: 10.1594/PANGAEA.691572
doi: 10.1594/PANGAEA.319909	doi: 10.1594/PANGAEA.691573
doi: 10.1594/PANGAEA.319910	doi: 10.1594/PANGAEA.691574
doi: 10.1594/PANGAEA.319888	doi: 10.1594/PANGAEA.691575
doi: 10.1594/PANGAEA.319889	doi: 10.1594/PANGAEA.691576
doi: 10.1594/PANGAEA.319890	doi: 10.1594/PANGAEA.691577
doi: 10.1594/PANGAEA.319891	doi: 10.1594/PANGAEA.691578
doi: 10.1594/PANGAEA.319892	doi: 10.1594/PANGAEA.691579
doi: 10.1594/PANGAEA.319893	doi: 10.1594/PANGAEA.691580
doi: 10.1594/PANGAEA.667009	doi: 10.1594/PANGAEA.691581
doi: 10.1594/PANGAEA.667008	doi: 10.1594/PANGAEA.691545
doi: 10.1594/PANGAEA.667010	doi: 10.1594/PANGAEA.691546
doi: 10.1594/PANGAEA.667011	doi: 10.1594/PANGAEA.691547
doi: 10.1594/PANGAEA.667012	doi: 10.1594/PANGAEA.691548
doi: 10.1594/PANGAEA.667013	doi: 10.1594/PANGAEA.691549
doi: 10.1594/PANGAEA.667014	doi: 10.1594/PANGAEA.691550
doi: 10.1594/PANGAEA.667015	doi: 10.1594/PANGAEA.691551
doi: 10.1594/PANGAEA.667016	doi: 10.1594/PANGAEA.691552
doi: 10.1594/PANGAEA.667017	doi: 10.1594/PANGAEA.691553
doi: 10.1594/PANGAEA.667018	doi: 10.1594/PANGAEA.691554
doi: 10.1594/PANGAEA.667019	doi: 10.1594/PANGAEA.691555
doi: 10.1594/PANGAEA.667020	doi: 10.1594/PANGAEA.691556
doi: 10.1594/PANGAEA.667021	doi: 10.1594/PANGAEA.691557
doi: 10.1594/PANGAEA.667022	doi: 10.1594/PANGAEA.691558
doi: 10.1594/PANGAEA.667023	doi: 10.1594/PANGAEA.713323
doi: 10.1594/PANGAEA.667024	doi: 10.1594/PANGAEA.713324
doi: 10.1594/PANGAEA.667025	doi: 10.1594/PANGAEA.713325

doi: 10.1594/PANGAEA.713326

doi: 10.1594/PANGAEA.713327

doi: 10.1594/PANGAEA.713328

doi: 10.1594/PANGAEA.713329

doi: 10.1594/PANGAEA.713330

doi: 10.1594/PANGAEA.713331

doi: 10.1594/PANGAEA.713332

Sessile Ssuspension Feeders with Associated fauna (SSFA)

Mobile deposit feeders, Infauna & Grazers (MOIN)




Figure 1. Classification of Antarctic macro-benthic communities (after Gutt 2007 and Turner et al. 2009).

Datasets

Dataset description

Object name: Darwin Core Archive Antarctic macrobenthic communities: A compilation of circumpolar information

Character encoding: UTF-8

Format name: Darwin Core Archive format

Format version: 1.0

Distribution: <http://ipt.biodiversity.aq/archive.do?r=macrobenthos>




Figure 2. Geographic coverage of the circumpolar distribution of information on Antarctic macrobenthic communities provided by ANTABIF.

Publication date of data: 2012-07-19

Language: English

Licenses of use: This data-set is entitled “Antarctic macrobenthic communities: A compilation of circumpolar information” and has been uploaded to (ANTABIF). The data set has been made available under the Open Data Commons Attribution License: <http://www.opendatacommons.org/licenses/by/1.0/>

Metadata language: English

Date of metadata creation: 2012-07-19

Hierarchy level: Dataset

Acknowledgements

Thanks are due to all who provided data for this compilation.

References

- De Broyer C, Koubbi P, Danis B, David B, Grant S, Griffiths H, Gutt J, Held C, Huettmann F, Post A, Ropert-Coudert Y (in prep.) The CAML / SCAR-MarBIN Biogeography Atlas of the Southern Ocean.
- Clarke A, Johnston NM (2003) Antarctic marine benthic diversity. *Oceanography and Marine Biology: an Annual Review* 41: 47–114.
- Gutt J (2007) Antarctic macro-zoobenthic communities: a review and an ecological classification. *Antarctic Science* 9: 165–182. doi: 10.1017/S0954102007000247
- Turner J, Bindschadler R, Convey P, di Prisco G, Fahrbach E, Gutt J, Hodgson D, Mayewsky P, Summerhayes C (2009) *Antarctic Climate Change and the Environment*. SCAR, Scott Polar Research Institute, Cambridge, 526 pp.