

Diagnoses **Molluscorum novorum**,

AUCTORE H. CROSSE.

1. MUREX LIENARDI.

T. vix aut non rimata, fusiformis, solidula, pallide violaceo-albida; spira elongata, apice obtusulo; sutura late marginata; anfr. 6 sexvaricosi (interdum septemvaricosi), embryonales primi 1 1/2 sublæves, sequentes varicibus rotundatis longitudinaliter muniti, costulis prominulis, plus minusve obsolete squamulatis, varices superantibus transversim cincti, ultimus spira paulo major, varicibus alternatim majoribus et minoribus instructus, varice ultimo, penultimo et antepenultimo et costulis violaceo-fuscis; apertura ovata, parva, intus violacea, in canalem breviusculum, subrecurvum, violaceum desinens; peristoma subcontinuum, violaceum, margine columellari subarcuato, plicis 3 minutis, albicantibus munito, externo intus denticulis 6 armato, ad limbum crenulato, mox extus, occurso varicis ultimi, in alam brevem, pulchre fimbriatam, purpureo-violaceam producto. — Long. 25, diam. maj. 13 1/2 mill. Apertura 7 mill. longa (canali excluso), 4 lata. (Coll. Crosse, Liénard et Rœters van Lennep.)

Habitat ad littus insulæ Mauritiï, parum frequens. (Robillard.)

2. MEROE ROETERSIANA.

T. ovata, transversa, subcompressa, subinæquilateralis (latere antico brevior), antice rotundata, postice obtuse truncato-rotundata, lævigata, pallide roseo-fuscescens, guttulis longitudinalibus, paulo saturatioribus, parum conspicuis suffusa; fossa ligamenti longa, profunda, extus

utrinque striis impressis, subito evanidis circumdata; lunula lanceolata; cardo normalis; pagina interna roseo-albida, in vicinio cicatriculæ posticæ purpureo-violaceo picta; margo externus crassiusculus, tenuiter crenulato-denticulatus. — Diam. antero-post. 52, umbono-margin. 39, crass. 19 1/2 mill. (Coll. Ræters van Lennep.)

Habitat?

3. TEREBRATULA CERNICA.

T. globoso-piriformis, convexa, pellucida, sublævigata, vix concentrice striatula, vitreo-hyalina; valva dorsalis medio convexa, utrinque attenuata, vix obsolete subangulata; foramen ovato-rotundatum, sat magnum; rostrum vix productum, oblique truncatum; valva ventralis valde inflata; margo externus ex utroque latere subflexuosus. — Diam. antero-post. 28, umbono-marg. valvæ dorsalis (e foramine ad marginem) 35, ventralis 33, crass. 20 mill. (Coll. E. Liénard.)

Habitat in mari insulam Mauritiæ alluente. (Teste E. Liénard.)

Obs. Species inter Terebratulam vitream et T. uvam quasi media.

H. C.

Description d'un **Murex** nouveau, provenant de l'île **Maurice**,

PAR E. LIÉNARD.

MUREX CROSSEI.

T. vix obtecte subrimata, breviter pyramidato-fusifor-