

## Air-sea carbon flux at the Belgian Continental Shelf

Pint Steven, Everaert Gert, Gkritzalis Thanos and Vandegehuchte Michiel

Flanders Marine Institute (VLIZ), InnovOcean site, Wandelaarkaai 7, 8400 Oostende, Belgium  
E-mail: [steven.pint@vliz.be](mailto:steven.pint@vliz.be)

Observing the balance of greenhouse gases is an important way to keep track of global change (Steinhoff et al., 2019). One important element in this balance is the atmosphere-water exchange of CO<sub>2</sub> in the ocean. The air-sea CO<sub>2</sub> flux provides insight in how much CO<sub>2</sub> is incorporated in the marine environment (i.e. the sea being a sink for atmospheric CO<sub>2</sub>) or emitted by the marine environment (i.e. the sea being a source). As of 2013, as part of the European research infrastructure “Integrated Carbon Observation System” (ICOS), the Flanders Marine Institute (VLIZ) measures the pCO<sub>2</sub> in the surface layer of the water at the Belgian Continental Shelf. In this study, we used observations of pCO<sub>2</sub> collected at the Thornton buoy; a measuring buoy located at the Thorntonbank, a sandbank approximately 30 km seawards from the coast near Zeebrugge, from February until December 2018. We calculated the air-sea carbon fluxes according to the wind driven turbulence diffusivity model of Nightingale (2000). Our results show a clear seasonality of air-sea carbon flux at the Thornton buoy, with the sea being a carbon sink from February until June switching to a carbon source from July until December. This seasonality is also reported in Gypens et al. (2004 and 2011) and is hypothesized to be driven by temperature, biological processes and the impact of the freshwater plume of the Scheldt river (Gypens et al., 2011). We calculated that the sink was largest in April, while in August, the source was at its maximum. Increasing the amount of pCO<sub>2</sub> observations with the RV Simon Stevin will allow us to further explore the spatial variability of the air-sea carbon flux at the Belgian Continental Shelf.

### References

- Gypens, N., Lancelot, C., & Borges, A. (2004). Carbon dynamics and CO<sub>2</sub> air-sea exchanges in the eutrophied coastal waters of the southern bight of the North Sea: a modelling study. *Biogeosciences Discussions*, 1(1), 561–589. doi: 10.5194/bgd-1-561-2004
- Gypens, N., Lacroix, G., Lancelot, C., & Borges, A. (2011). Seasonal and inter-annual variability of air-sea CO<sub>2</sub> fluxes and seawater carbonate chemistry in the Southern North Sea. *Progress in Oceanography*, 88(1-4), 59–77. doi: 10.1016/j.pocean.2010.11.004
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., ... Upstill-Goddard, R. C. (2000). In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers. *Global Biogeochemical Cycles*, 14(1), 373–387. doi: 10.1029/1999gb900091
- Steinhoff, T., Gkritzalis, T., Lauvset, S. K., Jones, S., Schuster, U., Olsen, A., ... Watson, A. (2019). Constraining the Oceanic Uptake and Fluxes of Greenhouse Gases by Building an Ocean Network of Certified Stations: The Ocean Component of the Integrated Carbon Observation System, ICOS-Oceans. *Frontiers in Marine Science*, 6. doi: 10.3389/fmars.2019.00544

Keywords: Air-sea flux; Carbon dioxide; pCO<sub>2</sub>; North Sea; Belgian Continental Shelf