



Author Correction: Attraction Pheromone of The Benthic Diatom *Seminavis robusta*: Studies on Structure-Activity Relationships

Christine Lembke¹ · Daniel Stettin¹ · Franziska Speck¹ · Nico Ueberschaar¹ · Sam De Decker² · Wim Vyverman² · Georg Pohnert^{1,3}

Published online: 2 March 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Author Correction: Journal of Chemical Ecology (2018) 44:354–363

<https://doi.org/10.1007/s10886-018-0944-2>

The original version of this article unfortunately contained a mistake. The chemical structure of compound **6** in Fig. 1 was incorrect. The tested compound **6** in this study was (3*S*,8*aS*)-3-isobutylhexahydropyrrolo[1,2-*a*]pyrazine-1,4-dione as shown in the corrected version of Fig. 1 here.

The author apologizes for this oversight and for any confusion it may have caused.

The online version of the original article can be found at <https://doi.org/10.1007/s10886-018-0944-2>

✉ Georg Pohnert
Georg.Pohnert@uni-jena.de

¹ Institute for Inorganic and Analytical Chemistry, Bioorganic Analytics, Friedrich-Schiller-Universität Jena, Lessingstrasse 8, D-07743 Jena, Germany

² Laboratory of Protistology and Aquatic Ecology, Department of Biology, University Gent, Krijgslaan 281 S8, 9000 Gent, Belgium

³ Max Planck Institute for Chemical Ecology, Hans-Knöll-Str. 8, D-07745 Jena, Germany

Fig. 1 Compounds tested in this study. The natural pheromone (*S,S*)-1 (gray box) was synthesized stereospecifically, as well as with undefined stereochemical composition (1). Stereoisomers of 1 and 2 were separated by chiral HPLC

