



J. Plankton Res. (2019) 41(4): 375–391. First published online July 15, 2019 doi:10.1093/plankt/fbz026

HORIZONS

Mixotrophic protists and a new paradigm for marine ecology: where does plankton research go now?

KEVIN J FLYNN^{1,*}, ADITEE MITRA¹, KONSTANTINOS ANESTIS², ANNA A ANSCHÜTZ¹, ALBERT CALBET³,
GUILHERME DUARTE FERREIRA³, NATHALIE GYPENS⁴, PER J HANSEN⁵, UWE JOHN^{2,6}, JON LAPEYRA MARTIN⁴,
JOOST S MANSOUR⁷, MAIRA MASELLI⁵, NIKOLA MEDIC⁵, ANDREAS NORLIN¹, FABRICE NOT⁷, PARASKEVI PITTA⁸,
FILOMENA ROMANO⁸, ENRIC SAIZ³, LISA K. SCHNEIDER⁹, WILLEM STOLTE⁹ AND CLAUDIA TRABONI³

¹BIOSCIENCES, SWANSEA UNIVERSITY, SWANSEA SA2 8PP, UK ²ALFRED-WEGENER-INSTITUTE HELMHOLTZ CENTRE FOR POLAR AND MARINE RESEARCH, AM HANDELSHAFEN 12, 27570 BREMERHAVEN, GERMANY ³INSTITUT DE CIÈNCIES DEL MAR (ICM, CSIC), PASSEIG MARÍTIM DE LA BARCELONETA 37-49, 08003 BARCELONA, SPAIN ⁴UNIVERSITÉ LIBRE DE BRUXELLES, ECOLOGIE DES SYSTÈMES AQUATIQUES, CP-221, BOULEVARD DU TRIOMPHE, 1050 BRUSSELS, BELGIUM ⁵MARINE BIOLOGICAL SECTION, UNIVERSITY OF COPENHAGEN, STRANDPROMENADEN 5, DK-3000 HELSINGØR, DENMARK ⁶HELMHOLTZ INSTITUTE FOR FUNCTIONAL MARINE BIODIVERSITY (HIFMB), AMMÄLÄNDER HEERSTR. 231, 23129 OLDENBURG, GERMANY ⁷SORBONNE UNIVERSITY, CNRS, UMR7144, ADAPTATION AND DIVERSITY IN MARINE ENVIRONMENT (AD2M) LABORATORY, ECOLOGY OF MARINE PLANKTON TEAM, STATION BIOLOGIQUE DE ROSCOFF, PLACE GEORGES TEISSIER, 29680, ROSCOFF, FRANCE ⁸INSTITUTE OF OCEANOGRAPHY, HELLENIC CENTRE FOR MARINE RESEARCH, PO BOX 2214, 71003 HERAKLION, GREECE ⁹DELTA RES, BOUSSINESQWEG 1, 2629 HV DELFT, THE NETHERLANDS

*CORRESPONDING AUTHOR: k.j.flynn@swansea.ac.uk

Received March 22, 2019; editorial decision May 24, 2019; accepted May 24, 2019

Corresponding editor: John Dolan

Many protist plankton are mixotrophs, combining phototrophy and phagotrophy. Their role in freshwater and marine ecology has emerged as a major developing feature of plankton research over recent decades. To better aid discussions, we suggest these organisms are termed “mixoplankton”, as “planktonic protist organisms that express, or have potential to express, phototrophy and phagotrophy”. The term “phytoplankton” then describes phototrophic organisms incapable of phagotrophy. “Protozooplankton” describes phagotrophic protists that do not engage in acquired phototrophy. The complexity of the changes to the conceptual base of the plankton trophic web caused by inclusion of mixoplanktonic activities are such that we suggest that the restructured description is termed the “mixoplankton paradigm”. Implications and opportunities for revision of survey and fieldwork, of laboratory experiments and of simulation modelling are considered. The main challenges are not only with taxonomic and functional identifications, and with measuring rates of potentially competing processes within single cells, but with decades of inertia built around the traditional paradigm that assumes a separation of trophic processes between

different organisms. In keeping with the synergistic nature of cooperative photo- and phagotrophy in mixoplankton, a comprehensive multidisciplinary approach will be required to tackle the task ahead.

KEYWORDS: mixotrophy; protist; mixoplankton; phytoplankton; protozooplankton; microbial loop; allometry

INTRODUCTION

Mixotrophy is the combination of autotrophy and heterotrophy in the same organism. The means by which this combination occurs in different organisms varies, but the form that has most fascinated scientists (and captured the public attention; Wyndham, 1951) has been that which combines photo(auto)trophy with carnivory, or as applied to protists, with phago (hetero)trophy. Long considered by most aquatic scientists to be of minor significance, or just a curiosity, we now recognize that protist plankton engaging in photo- and phago-mixotrophy are common and important members of the global plankton community (Leles *et al.*, 2017, 2019; Faure *et al.*, 2019).

Even though photo- and phago-mixotrophy in planktonic protists has been studied by many scientists over many decades in freshwater and marine systems (e.g. Biecheler, 1936; Blackbourn *et al.*, 1973; Bird and Kalf, 1986; Estep *et al.* 1986; Laval-Peuto and Febvre, 1986; Sanders *et al.*, 1990; Sanders, 1991; Jones, 1994; Stoecker *et al.*, 1987; Jeong *et al.*, 2010; Hansen, 2011), appreciation that these organisms play a core role in especially marine ecosystems has been slow to mature. From 2011 to 2013 a series of workshops (funded by the Leverhulme Trust, U.K.) were held in Sweden (Kalmar), UK (Swansea) and the USA (Horn Point), bringing together experts across the field of marine planktonic protists. Outputs from those meetings explored the false dichotomy of the traditional phytoplankton–zooplankton paradigm (Flynn *et al.*, 2013), advantages of acquired phototrophy (Flynn and Hansen, 2013), the role of mixotrophy in shaping the biological carbon pump (Mitra *et al.*, 2014b), stoichiometric implications for mixotrophy (Lundgren *et al.*, 2016), the functional classification of planktonic protists (Mitra *et al.*, 2016) and the biogeographies of the different types of mixotrophic protist plankton (Leles *et al.*, 2017, 2019). Many additional publications have also raised the profile of photo- and phago-mixotrophic plankton over the last two decades (e.g. Stickney *et al.*, 2000; Burkholder *et al.*, 2008; Zubkov and Tarran, 2008; Carvalho and Granéli, 2010; Brutemark and Granéli, 2011; Sanders, 2011; Hartmann *et al.*, 2013; Wilken *et al.*, 2013; Saad *et al.*, 2016; Selsos *et al.*, 2017; Stoecker *et al.*, 2017), and brought the subject to a wider audience (Mitra, 2016, 2018; Glibert *et al.*, 2019).

Why is this subject important? Because the combining of primary and secondary production in a single

organism radically changes biogeochemical and trophic dynamics involving those organisms (Hitchman and Jones, 2000; Mitra *et al.*, 2014b; Caron, 2016; Ward and Follows, 2016; Ghyoot *et al.*, 2017a, 2017b; Leles *et al.* 2018). It alters the flows of energy and materials in and out of organisms that form the base of the food chain, and changes the way that we understand and thence simulate processes from harmful algal blooms (HABs) and fisheries to global climate change.

In all areas of science, there comes a point when the weight of new evidence warrants a fundamental reassessment of the paradigm in which we express our understanding. Within marine ecology, while the components of the microbial loop had been known about for decades before the mid 1970s, the broad importance of the concept was only brought to wide attention with works such as those of Pomeroy (1974), Williams (1981) and Azam *et al.* (1983), with increasing acceptance following that (e.g. Pomeroy *et al.*, 2007; Fenchel, 2008). We suggest that a similar reassessment point has now been reached concerning the importance of photo- and phago-mixotrophic plankton, and its potential to radically reshape marine ecology. To consider such a reshaping represents a very significant, and most likely controversial task, cutting across all sectors of plankton research. Here we present some ideas on how this task could be undertaken. However, we start by considering naming conventions and some misconceptions that surround the topic of evolution and functional traits of planktonic mixotrophs. We then continue by exploring the implications for marine ecology, and then consider various specific challenges for plankton research.

To aid this discussion we present a glossary (Table I).

WHAT IS IN A NAME?

Traditional descriptions of plankton most obviously comprise phytoplankton and zooplankton, with increasing inclusion of bacterioplankton from the 1980s. Often these descriptors attract add-on terms denoting organism size (pico-, nano-, micro-, meso-; e.g. Sieburth *et al.*, 1978). Over a century of marine research has been built upon terminologies largely of terrestrial origin, dividing primary and secondary production neatly between plant-like phytoplankton and animal-like zooplankton. Thus, phytoplankton may be referred to as the “grass of the sea”,

Table I: Glossary to terms describing forms of nourishment and functional types of plankton

Term	Definition
Autotrophy	Nutrition involving the synthesis of complex organic substances using photosynthesis (<i>phototrophy</i>) or chemosynthesis. Typically associated with the use of inorganic nutrients.
Bacterioplankton	Bacterial (prokaryote) <i>plankton</i> acquiring nourishment via <i>osmo(hetero)trophy</i> , and some also via <i>chemo (auto)trophy</i>
Constitutive mixoplankton (CMs)	<i>Protist plankton</i> with an inherent capacity for <i>phototrophy</i> that can also exhibit <i>phagotrophy</i>
Cyanobacteria	<i>Bacterioplankton</i> (prokaryote) members of the <i>phytoplankton</i>
Generalists non-constitutive mixoplankton (GNCMs)	<i>NCMs</i> that acquire their capacity for <i>phototrophy</i> from general (i.e. non-specific) phototrophic prey (cf. <i>SNCM</i>)
Heterotrophy	Nutrition involving the consumption and interconversions of sources of organic carbon
Metazooplankton	Multicellular (i.e. non- <i>protist</i>) <i>zooplankton</i>
Mixoplankton	<i>Plankton protists</i> capable of obtaining nourishment via <i>photo(auto)trophy</i> and <i>phago(hetero)trophy</i> , as well as via <i>osmo(hetero)trophy</i> .
Mixotrophy	Nutrition involving both <i>autotrophy</i> and <i>heterotrophy</i>
Non-constitutive mixoplankton (NCMs)	<i>Protist plankton</i> that acquire the capability for <i>phototrophy</i> from consumption (via <i>phagotrophy</i>) of phototrophic prey
Osmotrophy	A mode of <i>heterotrophy</i> (i.e. <i>osmo(hetero)trophy</i>) involving the uptake and consumption of dissolved organic compounds
Phagotrophy	A mode of <i>heterotrophy</i> (i.e. <i>phago(hetero)trophy</i>) involving the engulfment of particles (often whole organisms) into a phagocytic vacuole in which digestion occurs
Phototrophy	A mode of <i>autotrophy</i> (i.e. <i>photo(auto)trophy</i>) involving the fixation of CO ₂ using energy derived from light.
Phytoplankton	<i>Plankton</i> obtaining nourishment via <i>photo(auto) trophy</i> and <i>osmo(hetero)trophy</i> . They are <u>incapable</u> of <i>phagotrophy</i> ; cf. <i>mixoplankton</i> .
Plankton	Organisms that cannot maintain a fixed location in the water column, and are thus moved by the tides and currents
Protist	Single-celled eukaryote organism
Protophytoplankton	<i>Protist phytoplankton</i>
Protozooplankton	<i>Protist zooplankton</i>
Specialist non-constitutive mixoplankton (SNCMs)	<i>NCMs</i> that acquire their capacity for <i>phototrophy</i> from specific phototrophic prey (cf. <i>GNCM</i>)
Zooplankton	<i>Plankton</i> obtaining nourishment via <i>heterotrophy</i> . They are <u>incapable</u> of <i>phototrophy</i> ; cf. <i>mixoplankton</i> .

Terms in *italics* are further defined within this glossary. To these terms may be applied (as appropriate), a prefix indicating organism size according to **pico-** (0.2–2 µm), **nano-** (2–20 µm), **micro-** (20–200 µm) and **meso-** (>200 µm). Thus, *nano-protazooplankton* include organisms often termed “heterotrophic nanoflagellates”, while *meso-metazooplankton* include copepods.

and their copepod predators as “herbivores” or “insects of the sea” (e.g. Cushing, 1975), with classic models (e.g. NPZ: Fasham *et al.*, 1990; Franks, 2002), their complex successors (e.g. DARWIN: Follows *et al.*, 2007; MEDUSA: Yool *et al.*, 2013; ERSEM: Baretta *et al.*, 1995; Butenschön *et al.*, 2016) and indeed climate change (Arora *et al.*, 2013) and fisheries management models (Plagányi, 2007), also deploying this classic dichotomy.

The growing realization that plankton communities often contain organisms that combine phototrophy and phagotrophy (i.e. primary and higher production), requires that we more readily and clearly discriminate between organisms that are actually or potentially photo- and phago-mixotrophic and organisms that are not. We need greater clarity during our conversations, in conferences, on posters, in scientific publications and critically also in teaching. The problem is that the terminology currently referencing these protist plankton is unwieldy and/or ambiguous. And the problem starts with the word “mixotroph” (the combining of autotrophy and heterotrophy in one organism; Table I).

For protist plankton, the mode of autotrophy is phototrophy, while heterotrophy may be enabled by osmotrophy and/or phagotrophy (Fig. 1; Table I). There is a profound difference between heterotrophy supported by osmotrophy versus that by phagotrophy, both for trophic dynamics and also for food web structuring. Phagotrophy typically involves the killing of other organisms through the process of capture, ingestion (either partially or totally) and digestion. In contrast, osmotrophy does not require the act of killing. Indiscriminate use of “mixotrophy” fails to draw attention to this important difference.

Osmotrophy, the ability to use dissolved organics such as amino acids and vitamins (auxotrophy; Droop, 2007), is common in plankton typically considered as phototrophs. In aquatic ecology the subject has a long and controversial history (see Flynn and Butler, 1984 for a review of older discussions on the ecological importance to plankton); for many species in nature the role of osmotrophy probably acts mainly to mitigate against metabolite leakage (Flynn and Berry, 1999).




Fig. 1. Overlaps in the trophic capacity of different microplankton. **(a)** Venn diagram of overlaps between photo-, osmo- and phagotrophy. “VOID” indicates that no organisms exist in this sector. Mixoplankton are, by definition, capable of all three modes. Phytoplankton are mixotrophic by virtue of photo- and osmotrophy; they are non-phagotrophic. Protozooplankton are incapable of phototrophy. **(b)** Overlaps between traditional allocations of chlorophyll (Chl) containing organisms as “phytoplankton” and microbial grazers as “protozooplankton”. Symbol sizes or area allocations do not appertain significance.

Nonetheless, phytoplankton are typically mixotrophic in this sense, and indeed various modelling explorations of mixotrophy also refer to photo- and osmo-mixotrophy (e.g. Våge *et al.*, 2013). This mode of mixotrophy (photo(auto)trophy + osmo (hetero)trophy) in protistan and cyanobacterial microalgae is also exploited for biotechnology (Chojnacka and Noworyta, 2004; Liang *et al.*, 2009; Bhatnagar *et al.*, 2011; Morales-Sánchez *et al.*, 2013). Confusingly the mode of mixotrophy being studied is neither always apparent from the title nor in the abstract of many scientific publications (e.g. Yelton *et al.*, 2016), and indeed both osmotrophy and phagotrophy may be implicated (Burkholder *et al.*, 2008). We need to be able to differentiate readily between generic “mixotrophy” and specifically that which involves photo-phagotrophy.

Because of the likely ubiquity of osmotrophy, use of this trait in the functional classification of protists is compromised (Flynn *et al.*, 2013). Accordingly, Mitra *et al.* (2016) presented a revised classification of the protist plankton based on functional types, centred on the potential for and the mode of expressing photo- and phago-mixotrophy. That publication differentiates between the “pure” photo(auto)trophs (phytoplankton), the mixotrophs that have a constitutive ability to photosynthesise (“constitutive mixotroph”, CM), those that need to acquire their phototrophic capability from their prey (“non-constitutive mixotrophs”, NCM) and the “pure” phago- and heterotrophs (protozooplankton).

NCMs are then themselves split between generalists (GNCM), plastidic specialists (pSNCM) and endosymbiotic specialists (eSNCM) according to their mode of acquiring phototrophic capabilities. All these groups are fundamentally different functional types (Fig. 2). While the terms CM and NCM are gradually gaining increased usage (e.g. Johnson and Moeller, 2018; Hansson *et al.*, 2019; Naselli-Flores and Barone, 2019), what we are still missing is a term akin to the traditional terms—phytoplankton and zooplankton—to use to collectively refer to all photo- and phago-mixotrophic planktonic protists.

As such a short descriptive term, we propose that the word “mixoplankton” be used to specifically reference photo- and phagotrophic protists. This term already appears in various online teaching resources as well as on Fishbase website (<https://www.fishbase.de/glossary/Glossary.php?q=mixoplankton>), where mixoplankton are defined as “Planktonic organisms that can be classified at several trophic levels. For example, some ciliates can be photosynthetic but also can ingest other plankton and are heterotrophic.” Such a description perhaps does not make it clear that one and the same organism is engaging simultaneously or alternatively in photo-*plus* phago-*trophic* activities. Nor does this description exclude reference to mixotrophs that are incapable of phagotrophy. We thus suggest that this definition is reworded to read:

“mixoplankton: planktonic protist organisms that express, or have potential to express, phototrophy and phagotrophy.”




Fig. 2. Schematics showing the distinct differences between different protist plankton physiologies. The protozooplankton are osmo–phagotrophic; they are incapable of phototrophy. The phytoplankton are photo–osmo mixotrophic; they are incapable of phagotrophy. The constitutive mixoplankton (CM) and non-constitutive mixoplankton (NCMs) are all photo-, osmo- and phago-mixotrophic. However, some mixoplankton have life stages that are not photo- and phagotrophic, with nutrition aligning with that of “phytoplankton” or “protozooplankton”. The generalist GNCCMs acquire phototrophy from many phototroph prey types; pSNCCMs are plastic specialists acquiring phototrophy from specialist prey type(s); eSNCCMs are endosymbiotic acquiring phototrophy by harbouring specific phototrophic prey. Note: illustrations are not to scale; in particular, eSNCCMs are in relative terms ca. 10 to 100 times larger than the others (see Fig. 3).

The word “potential” in the definition of mixoplankton is also important, because while some mixoplankton are obligatory photo- and phago-mixotrophs (or have life stages that are), other species (specifically many CMs) may not show such an obligatory requirement (Sanders *et al.*, 1990; Jones, 1997; Calbet *et al.*, 2011; Berge and Hansen, 2016; Stoecker *et al.*, 2017).

Usage of the overarching term “mixoplankton” has various advantages:

- It is short and not hyphenated, amenable to the addition of an allometric prefix (e.g. nano-mixoplankton).
- There is a clear linkage to plankton; mixotrophy as a mode of nutrition is distributed across many organism groups besides plankton (e.g. corals, the sea slug *Elysia viridis*, the Venus flytrap *Dionaea muscipula*).
- Importantly, “mixoplankton” discriminates between the mixotrophy expressed in protists unable to phagocytose (i.e. photo- and osmo-mixotrophy, as conducted by diatoms for example), versus that expressed by protists that can conduct phagotrophy (i.e. the mixotrophy of the CM and NCM functional groups described by

Mitra *et al.*, 2016). We can readily realign the original descriptions of Mitra *et al.*, 2016 by modifying the meaning of the abbreviations CM and NCM to “constitutive mixoplankton” and “non-constitutive mixoplankton”, respectively.

- Perhaps of equal importance to the above is that phytoplankton are then characterized as being incapable of phagotrophy, while protozooplankton are incapable of phototrophy (Table I; Fig. 2).

From here on, we shall use the term “mixoplankton” in reference to photo- and phago-mixotrophic planktonic protists, including where the original referenced paper termed such organisms as “mixotrophs”.

MISCONCEPTIONS ON THE EVOLUTION AND TRAITS OF MIXOPLANKTON

To help understand why different organisms occupy the niches that they do occupy requires a full understanding of their functional traits, and also of those organisms

with which they interact. A common conception is that mixoplankton are inferior to their phytoplankton or protozooplankton counterparts with respect to their expression of traits such as photosynthesis, nutrient uptake and feeding (Dolan and Pérez, 2000; Litchman *et al.*, 2007; cf. Calbet *et al.*, 2011). This assumes that an organism that specializes in two nutritional routes is a generalist in its expression of each of those routes. This argument could be considered as a trait-trade-off (TTO), a concept which has found considerable support in the plankton research community, especially with respect to plankton modelling (Litchman and Klausmeier, 2008; Litchman *et al.*, 2013). Such TTO arguments must only be applied to organisms from the same environment (Litchman *et al.*, 2007), and TTOs are in any case questionable from an evolutionary standpoint (Flynn *et al.*, 2015).

Clearly, we expect traits expressed by phytoplankton and protozooplankton to be advantaged in the environment in which they evolved, just as those species that combine photo- and phagotrophy must be advantaged by possessing both phototrophy and phagotrophy in other environments. Questions of which trait is better or not in a given environmental setting have been answered by evolution. To better clarify this matter requires a consideration of the evolution of the protists.

The genetic diversity of protists is vast, equalling that of all other eukaryotes combined (Burki 2014; Adl *et al.*, 2019). The evolution of eukaryotes (eukaryogenesis) was based on the phagotrophic acquisition of an alphaproteobacteria into a host, which subsequently lead to the formation (evolution) of mitochondria (Hampl *et al.*, 2019). The formation of mitochondria provided the cells with extra energy that allowed the evolution of eukaryotic cell structures (Martin *et al.*, 2017). One prerequisite for this event is that the host must have had an actin filament system that enabled the formation of protrusions and the subsequent engulfment of bacteria (Yutin *et al.*, 2009). Whether the last eukaryotic common ancestor was an Archaea or not is still not resolved (Martijn and Ettema, 2013), but under any scenario phagotrophy evolved very early in life on Earth. Moreover, secondary endosymbiotic events gave rise to photosynthesis within eukaryotes, which appears to have been lost and (re)gained several times in many protist groups (Archibald, 2009). Extant protozooplankton thus include organisms that gained and then lost their scope for phototrophy (i.e. that had ancestors that were mixoplankton).

Protist evolution saw many cycles of gaining and losing capabilities (Figuroa-Martinez *et al.*, 2015; Hampl *et al.*, 2019). Mixoplankton did not arise as a coming together of protist phytoplankton plus protozooplankton traits into one organism, and protist phytoplankton did not fail to acquire the capability for phagotrophy. On the contrary,

the most profound event in protist plankton evolution in some ways was the loss of the ancestral trait of phagotrophy. Not only did this remove an important nutritional pathway, but it also removed a major route through which protists have acquired genetic and physiological variety of importance for their evolution.

Ecologically the most important of those protist groups in contemporary aquatic environments that lost phagotrophy are the diatoms, which also evolved a non-cellulosic cell wall (frustule of silica) that was likely incompatible with phagotrophy from an early stage in their evolution. Planktonic diatoms are also effectively non-motile (they cannot swim) within the plankton. Motility is rather an important trait for a planktonic predator; the low motility of the mixoplanktonic Foraminifera and other Rhizaria are compensated for in this regard by the use of pseudopods for feeding. The environment in which diatoms flourish is also typically an immature one, of high turbulence, relatively high inorganic nutrient concentrations, and lacking in the prey that could otherwise provide nutrients. A similar set of arguments could be made for the often-assumed absence of phagotrophy in the ecologically dominant calcified form of coccolithophorids (cf. Rokitta *et al.*, 2011). In contrast, mixoplankton dominate in more mature systems containing competitors that are also potential prey, and in systems in which nutrients (which can include light) are supplied in unbalanced proportions, if not limiting amounts (Mitra *et al.*, 2014b). These are also the situations in which HABs occur (Granéli, 2006; Granéli *et al.*, 2008; Glibert *et al.*, 2018), where mixoplankton are often dominant species.

There is significant variation across different mixoplankton types with respect to their core physiology (Fig. 2), and allied variation in feeding types, and hence in the form of trophic interactions. Despite this, the traits of these organisms have often been referenced in the literature in a rather indiscriminate fashion, as if all mixotrophic plankton are effectively the same rather than being different (CM vs GNCM, vs SNCM etc.). This is especially so in the conceptual and modelling-orientated literature (e.g. Ward and Follows, 2016). The allometric “rule”-based analysis of Andersen *et al.* (2016), suggesting that mixoplankton are mechanistically aligned (optimized) at a position intermediate between smaller phototrophic phytoplankton and larger phagotrophic micro (proto) zooplankton, does not align with reality. In fact, mixoplankton span size ranges that encompass the whole breadth of protist phytoplankton and protozooplankton (Fig. 3); this range exceeds that similar to that of ant-to-cow. Equally important though, there appears to be important functional-allometric differences between the CMs, GNCMS and SNCMs, which is also seen in their biogeography (Leles *et al.*, 2017, 2019).




Fig. 3. Size ranges for different microbial plankton functional types. As labelled here, phytoplankton (prokaryotic cyanobacteria and allies, and eukaryotic protists) are photo- and osmo- mixotrophic; they are incapable of phagotrophy. Protozooplankton are osmo- and phagotrophic; they are incapable of phototrophy. For the mixoplankton, which all express or have potential to express photo-, osmo- and phagotrophy, the typical size ranges are as indicated by the boxes. Note that in total mixoplankton sizes span all other protist plankton types, though the CMs are the dominant examples in the sub 20 μm size categories. GNCM, generalist non-constitutive mixoplankton; pSNCM, plastidic specialist non-constitutive mixoplankton; eSNCM, endosymbiotic non-constitutive mixoplankton; CM, constitutive mixoplankton. The size ranges for mixoplankton have been taken from the data collected for *Leles et al.* (2017, 2019) and from personal observations made by authors. See also Figs 1 and 2.

In conclusion, mixoplankton should not be collectively viewed as inferior, nor indeed superior, to phytoplankton and protozooplankton. They are just different (and different within themselves—Fig. 2). They also, in consequence, have the potential to fulfil different functional roles within ecology, exploiting different niches. Before we appreciated the global abundance of mixoplankton we could perhaps justify ignoring their contribution. We can no longer afford to do so.

PLACING MIXOPLANKTON WITHIN MARINE ECOLOGY

Researching and understanding the role of mixoplankton within marine plankton systems has the potential to completely reform our understanding of marine ecology. However, the development of plankton science in this regard has been complicated by:

(i) the failure to appreciate the implications of different modes of mixoplankton functioning; the differing degrees with which mixotrophy is obligatory (*Sanders et al.*, 1990; *Jones*, 1997; *Stoecker et al.*, 2017; *Gomes et al.*, 2018) even within the same or closely related species (*Calbet et al.*, 2011; *Berge and Hansen*, 2016);

(ii) the afore-mentioned confusion in terminologies.

There is in consequence something of a legacy of published works that ignore or misrepresent mixoplankton and are in retrospect questionable as to whether they are best considered as incomplete, or explorations

in theoretical biology or ecology rather than of reality. The implications of the advance in our understanding and appreciation of mixoplankton within plankton ecology thus have similarities with the rise to prominence of the “microbial loop” (*Azam et al.*, 1983) and the microbial food web in the 1980s, and the inclusion of the “viral shunt” (*Wilhelm and Suttle*, 1999) in the 1990s. Before the 1980s, marine ecology saw at its base a system dominated by net-sized phytoplankton (i.e. cells retained in plankton nets (mesh size typically 20–35 μm) and meso-zooplankton (mesh size typically >200 μm ; *Cushing*, 1975). The pre-1980s paradigm can be seen to have been augmented by the addition of microbial food webs; the form of this revised trophic structure is indicated in Fig. 4a. The whole web contributes in various ways to the “biological carbon pump” (*Turner*, 2015) and to the “microbial carbon pump” (*Jiao et al.*, 2010; *Legendre et al.*, 2015).

The enhanced appreciation of the importance of the microbial food web was brought about in large measure by discovering and identifying the importance of certain types of biological entities (e.g. *Prochlorococcus*, viruses, etc.), as well as the overdue recognition of the importance of known groups (bacteria, cyanobacteria, heterotrophic nanoflagellates etc.). The recent expansion of interests in mixoplankton could be argued as just the overdue recognition of facets of ecology known for over a century (e.g. *Biecheler*, 1936; *Blackbourn et al.*, 1973; *Jones*, 1997; *Stoecker*, 1998; *Jones*, 2000). However, the inclusion of mixoplankton as a frequent and at times dominant




Fig. 4. Differences between the pre-2010 paradigm for plankton trophic dynamics, (a), and the post-2010 paradigm, (b), that acknowledges the significance and roles of mixoplankton. The microbial food web components are within the blue boxes; viruses impact all organism groups though (because of higher numeric host abundances) may impact smaller organisms more. Red dashed arrows indicate input and outputs of dissolved inorganic and organic nutrients, blue arrows are nutrient uptakes and black arrows indicate grazing routes. Non-motile phytoplankton include diatoms, and also non-motile life stages of others (notably organisms such as the nanoplanktonic coccolithorporid *Emiliania*, and the microplanktonic colonial *Phaeocystis*). In configuring (b), motile phytoplankton in (a) are considered to be potentially mixoplankton, as are ca. half of the protozooplankton indicated in (a). CM: constitutive mixoplankton. NCM: non-constitutive mixoplankton. HNF: heterotrophic nanoflagellates. p/n: pico/nano-sized organisms (i.e. 0.2–2.0 and 2.0–20 μm diameter, respectively). μ: micro-sized organisms (i.e. 20–200 μm diameter).

activity in plankton biogeography (Leles *et al.*, 2017, 2019; Faure *et al.*, 2019; Gutierrez-Rodriguez *et al.* in preparation) and trophic dynamics (e.g. Hartmann *et al.*, 2013) marks a radically different development in marine science compared to the inclusion of either the microbial loop or the viral shunt. It is different because it sees the bulk of the known members of the traditional trophic web description actually undertaking different (additional) trophic roles. Thus, most of the protist primary producers traditionally labelled as “phytoplankton”, other than those in immature ecosystems (notably diatoms), are actually also part-time grazers, while half those organisms traditionally labelled as “micro (proto)zooplankton” in the photic zone are also photosynthesising (Fig. 4b; see review by Stoecker *et al.*, 2009). This situation operates across the whole microbial plankton allometric scale, from picoflagellates (Zubkov and Tarran, 2008; Stoecker and Lavrentyev, 2018) to mm-dimension SNCMs (Decelle *et al.*, 2012; Biard *et al.*, 2017) (Fig. 3). The trophic linkages are redrawn and new links added.

One of the most frequent questions that arises in scientific discussions on mixoplankton is, how important is mixotrophy and when are these organisms functionally mixotrophic? Whether we need to know when they are eating, or whether it is sufficient just to know that they may do so if the need or opportunity arises, is a non-trivial question to answer. The ability to eat may supply small but vital amounts of nutrition to a mixoplanktonic organism that would be unavailable to non-mixoplankton

competitors at critical stages. At the same time competitors are removed by a combination of phagotrophy and production of compounds with allelopathic effects (John *et al.*, 2002, 2015; Skovgaard and Hansen, 2003; Tillmann, 2003); every difference is a difference. We suspect that many laboratory cultures of these organisms lose their ability to express different nutritional mechanisms over the years of them being held in constant light and high nutrient conditions (e.g. Blossom and Hansen in preparation). It thus follows that if organisms in nature can express a trait then that trait is, at least occasionally, exploited to advantage in nature.

Of the protist functional groups identified by Mitra *et al.* (2016), the NCMs must be regularly and significantly mixotrophic as many have to feed to acquire chloroplasts to support their obligatory need to undertake photosynthesis. The challenge is primarily with identifying when CM species are eating. Some CM species are perhaps only rare feeders in nature, while others (given the voracious behaviour of those brought into culture) are most likely regular feeders (reviewed by Jeong *et al.*, 2010; Hansen, 2011). The former group include species such as *Tripos furca* (= *Ceratium furca*; Smalley *et al.*, 2003, 2012), *Prorocentrum minimum* (Johnson, 2015) and *Heterocapsa triquetra* (Jeong *et al.*, 2005), while the latter include species such as *Prymnesium* (Skovgaard and Hansen, 2003; Carvalho and Granéli, 2010), *Karlodinium* (Berge *et al.*, 2008a and b; Calbet *et al.*, 2011) and *Fragilidium* spp (Jeong *et al.*, 1999; Hansen *et al.*, 2000; Skovgaard *et al.*, 2000).

As evidence of the importance of differentiating between the CM and NCM forms, it is perhaps noteworthy that the microbial food web interfaces directly with many CMs (Burkholder *et al.*, 2008; Jeong *et al.*, 2010; Hartmann *et al.*, 2013; Stoecker and Lavrentyev, 2018), while the NCMs generally tend to be larger (eSNCMs up to mm dimensions, e.g. green *Noctiluca*; Saito *et al.*, 2006; Stoecker *et al.*, 2009) (Figs 3 and 4b). That said, some NCMs are nano-sized, at least in the oligotrophic ocean (Pitta and Giannakourou, 2000; Pitta *et al.*, 2001).

Although we do not yet know the full significance of mixoplanktonic activity for community structure and dynamics, or indeed for the physiology and growth of most individual species, the structural difference between Fig. 4a and b appears so profound as to represent a paradigm shift. The mixoplankton paradigm is not a “loop”, or a “shunt”, appended to the pre-2010 trophic web (Fig. 4). Indeed, mixoplankton activity has potentially important implications for the functioning of the microbial food web (Fig. 4b). Is it perhaps a “multiplier”, improving assimilation efficiency and the allometrics of trophic transfer? We do not know. Perhaps until it becomes much clearer, we should just refer to it as simply the “*mixoplankton paradigm*”.

IMPLICATIONS OF THE MIXOPLANKTON PARADIGM FOR PLANKTON RESEARCH

The ramifications of the mixoplankton paradigm for plankton research are wide ranging. Some implications are perhaps more obvious than others, but taking a fresh look at everything may be the safer route forward. To really understand the implication of mixoplankton for marine ecology, future research will have to answer many questions such as: when and where are mixoplankton important? How and from where/who do mixoplankton acquire their energy and nutrients? What activity rates can be associated with mixoplankton? Which circumstances promote mixotrophy? Only a multidisciplinary approach can answer those questions; below we commence such a review, though it is readily apparent that the task will proceed for likely several decades.

Survey and fieldwork

Survey and fieldwork are ideally suited to answer questions such as when, where and how many organisms occur. These are still highly relevant questions regarding mixoplankton. However, survey methods, irrespective of

the approach and technology (remote sensing, autoboys, Continuous Plankton Recorder, etc.) have been designed and optimized essentially for organisms that are traditionally labelled as “phytoplankton” or “zooplankton” (both groups of which we now appreciate include mixoplankton), and typically for species that are nearest the surface, physically robust and relatively large. Many of these approaches are best suited to organisms with clear physiological traits (pigment signatures in cyanobacteria, reflective signatures in coccolithophorids, high structural integrity in diatoms). Many mixoplankton (flagellates and ciliates) are small, not structurally robust or easily sampled, nor amenable to identification. This means that long-term surveys (such as the CPR), focussed as they often are for specific groups or species, will unfortunately not provide historic data on most specific mixoplanktonic organisms. The work of Leles *et al.* (2017, 2019) shows clearly that there are serious gaps in our appreciation of the biogeography of, especially, the CMs stemming from such traditional methods. It may be possible, through suitable cross calibration with presence/absence of other plankton to glean some information. More profound, perhaps, is the need to detach a signal for chlorophyll from an automatic and exclusive association with primary production supported by inorganic nutrient acquisition, when in mixoplankton that same Chl signal is directly linked to primary production coupled to secondary production.

The paucity of suitably trained researchers who can readily identify mixoplankton not only hinders survey work directly, but will likely restrict the ground truthing of other technologies. Crucially this includes checking the veracity of molecular methodologies such as metabarcoding (who is there?) and metatranscriptomics (what might they be doing?); see discussion in Leles *et al.* (2019) and Faure *et al.* (2019). The challenge of exploiting molecular methods becomes even greater when one considers the presence of non-self genetic material within mixoplankton, as nucleic acids from their prey, in acquired plastids and/or in symbionts; more single-cell analyses are needed, including single-cell polymerase chain reaction (PCR) and genomic studies (Tai *et al.*, 2013; Kolisko *et al.*, 2014). Beisner *et al.* (2019) gives a review of the current states of the art in this subject arena, specifically for nano-sized mixoplankton. Molecular approaches have contributed much to elucidating patterns of mixotrophy in plankton (Burns *et al.*, 2015; McKie-Krisberg *et al.*, 2018; Faure *et al.*, 2019). Nonetheless, “old fashioned” techniques, such as optical and electron microscopy, also have important continuing roles and have been, and still are, frequently deployed in research on mixoplankton (Anderson, 1978; Swanberg and Caron, 1991; Mafra *et al.*, 2016; Kim *et al.*, 2017).

Experimental work

Knowing what is there, is one thing. Knowing what it is actually doing there and at what rate is more challenging, and arguably more important (Flynn *et al.*, 2018). Marine scientists have spent decades developing and deploying field and laboratory techniques for measuring primary production (photosynthesis, nutrient uptake) and, with markedly lesser success (Mitra *et al.*, 2014a), for grazing activities and secondary production by zooplankton. Researchers have studied laboratory cultures with emphasis on using axenic strains, and most frequently using strains that have been maintained in culture for decades in high inorganic nutrient, constant temperature/light environments (and thus have likely evolved, e.g. Martins *et al.* 2004). These approaches, although undertaken for perfectly sound reasons under the traditional paradigm of considering these phototrophic protists as phytoplankton, can now be seen as simplified too far. Our “laboratory rats”, as single strain isolates, have indeed not represented the true physiology of natural populations, which consists of a high heterogenic diversity, but for reasons far beyond just adapting to constant-temperature room conditions (e.g. Kremp *et al.*, 2012; Alpermann *et al.*, 2010; Brandenburg *et al.*, 2018).

Simply determining whether a mixoplankton is operating at that instant as a photo- and phago-mixotroph, and what its rate of grazing is upon what prey organisms, presents a profound challenge that few researchers and laboratories are well placed to even attempt to confront. We need to know those rates of photosynthesis, grazing, nutrient regeneration, respiration, etc.; without such data we cannot verify the performance of simulation models. To complicate things further, we need similar data for their prey species as well (which could also be mixoplankton), growing in the same water at the same time. There are very few studies of mixoplankton physiology that are adequately executed to provide data describing the dynamics of growth and trophic activity (e.g. Skovgaard *et al.*, 2000, 2003), and even fewer provide the necessary information on both the mixoplankton and its prey (Adolf *et al.*, 2003, Lin *et al.*, 2018); see discussion in Flynn and Mitra (2009). Simply catching these organisms in the act of eating is a problem (Anderson *et al.*, 2017). Consider a nano-mixoplankton eating bacteria; it may only need to eat one bacterium a day to acquire its P or Fe quota. Did we observe that event in our shipboard incubation? Do we know whether the time of day is important for the event? Is the prey presented for possible consumption in an experiment an appropriate species, and in the correct nutritional state?

Training in phytoplankton physiological ecology was perhaps at its zenith during 1970–1980. Since the emergence of molecular biology, however, the literature

has seen far fewer papers reporting all-engrossing experiments reporting time series incubations and cultures growth dynamics, etc. Molecular biology and genomics have played a pivotal role in elucidating the evolution of phagotrophy and eukaryogenesis, which is strongly linked to functionality of mixoplankton (Burns *et al.*, 2015). Those approaches were particularly useful in revealing features of the trophic interactions between NCMs (Takishita *et al.*, 2002). Emphasis has been placed upon who is there (at best only semi-quantitatively), and determining the scope for action (as implied from DNA and RNA data). Even meta-transcriptomic approaches can only indicate the potential contribution of genetic functionality at community level rather than the actual rate of action (Stewart *et al.*, 2012; Wohlrab *et al.*, 2018). The absence of genetic capability indicates an absence of functional ability; molecular biology can at best indicate potential capability for rate activity, but it (neither DNA sequences data, nor RNA data) cannot provide the actual rates. And will molecular biology be able to discriminate between a bacterial–mixoplankton interaction where the bacteria are external to the protist (as a contaminant or part of its associated microbiome) rather than internal having been eaten? There are many potential sources of conflicting signals even in cultured organisms. For example, membrane production and (auto)digestion processes occur in cells that are not phagocytic. So, to what extent molecular methods can provide help in establishing rate processes and relative contributions of photo- versus osmo- versus phagotrophy in mixoplankton is far from clear at present, though various studies have explored the topic (Santoferrara *et al.*, 2014; McKie-Krisberg *et al.*, 2018).

In short, there is much to do in the laboratory and in the field, and that applies whether these studies also involve reappraisals of species formally labelled as just phytoplankton or protozooplankton, or as prey, as competitors, or as predators. Critically, while phytoplankton are amenable to study in axenic conditions, or under conditions in which autecology dominates, studies of mixoplankton require a simultaneous study of their prey. This greatly complicates matters, generating problems comparable if not more complex, to those encountered in studies of zooplankton (Mitra *et al.*, 2014a).

Conceptual and simulation models

Survey and fieldwork, as well as experimental studies, are very important for providing information needed to support successful conceptual and numerical modelling of plankton ecology; they provide data and understanding to aid the construction and validation of the models. The added value of models, especially of numerical

simulation models, is the scope for their use to explore cause-and-effect questions, such as why and how are the pathways in trophic dynamics as they are. What is apparent is that modelled mixoplankton activity has clear scope to radically alter simulated trophic dynamics and biogeochemical flows (Flynn and Mitra, 2009; Mitra *et al.* 2014b, 2016; Ward and Follows, 2016; Ghyoot *et al.*, 2017b; Leles *et al.* 2018). The question is how closely those projections simulate reality, or whether they simply reflect the conceptual basis of the mixoplankton description. So, do we understand the physiology and functional roles of mixoplankton well enough to construct robust models to test hypotheses?

Mixotrophy in plankton has stimulated a curiosity-driven exploration of simulated ecology (Jost *et al.*, 2004; Hammer and Pitchford, 2005). Mixotrophy in such models has often been photo- and osmotrophic, rather than photo-, osmo- and phagotrophic. Modelled mixotrophs have often had traits that have been “traded” to prevent them from becoming superior and dominating over their pure phototrophic or heterotrophic counterparts (Jost *et al.*, 2004; Ward *et al.*, 2011). The rationale behind TTO models describing phytoplankton activity (Bruggeman and Kooijman, 2007; Follows *et al.*, 2007; Litchman and Klausmeier, 2008; Smith *et al.*, 2016) comes under a different spotlight if we consider that these organisms are also predators. Factors that appear of benefit for phytoplankton (e.g. small size favouring nutrient acquisition) or for consumers (e.g. larger size than the prey, and to minimize predation), and the relative importance of bottom-up and top-down factors become confused when we consider mixoplankton ecology and physiology. Then there are also the arguments against TTO approaches mentioned above. For modellers who wish to exploit TTO concepts for pragmatic computational reasons, at present we simply do not have sufficient empirical evidence from which to construct robust TTO arguments.

Other approaches (Flynn and Mitra, 2009; Ghyoot *et al.*, 2017a) have specifically sought to align model construction with the functional type descriptions that we observe (Mitra *et al.*, 2016). It is important that we recognize the importance of different functional forms of mixoplankton (Fig. 2). We need to do so at least for the forms that are most frequently dominant in terms of biomass (i.e. CM vs GNCM), though pSNCM and eSNCM forms are also important groups in terms of HABs and biogeochemistry. All mixoplankton types appear to be of significance across large areas of the oceans (Leles *et al.*, 2017, 2019). How does such knowledge shape a reappraisal of works considered as exemplars for global-scale plankton modelling, such as the DARWIN model approach of Follows *et al.* (2007), which sought to describe global “phytoplankton” productivities according

to standard phototrophic trait approaches, dominated by bottom-up controls? Equally problematic, is that plankton models that do not describe mixotrophic activity have had their skill considered with reference to data collected for natural populations that will most likely have included, on occasion, significant mixoplankton presence and activity.

Over the last decade, plankton simulation science has gone from the extreme of ignoring mixotrophy, to the opposite extreme of assuming that all plankton are mixotrophic (Ward and Follows, 2016). Ward and Follows (2016) deployed an allometric-linked sliding scale of mixoplanktonic capability. However, as noted above, far from being confined as an optimal strategy for mid-sized protists (Andersen *et al.*, 2016), significant levels of photo- and phago-mixotrophy operates in some of the smallest phototrophic flagellates (Hartmann *et al.*, 2013; Hansen and Hjorth, 2002; Anderson *et al.*, 2018), through 1 mm diameter green *Noctiluca* (Hansen *et al.*, 2004; Gomes *et al.*, 2018), to the very largest colonial Radiolaria (Collodaria; Swanberg, 1983; Biard *et al.*, 2016). There is a clear difference in the alignment of allometry and different mixoplankton functional types as well (Fig. 3). We should perhaps guard against developing and exploiting simple explanations that may lead to premature conclusions.

For sure, the modelling of protist plankton activities requires a reappraisal. We need computationally efficient models that describe these different organisms (Fig. 2), in a fashion that is acceptable in the eyes of those scientists who study the physiology and ecology of mixoplankton, before we place them in ecosystem simulators. To do otherwise, to describe theoretical mixoplankton forms that do not simulate the behaviour of real organisms, is surely at least as questionable as to continue to ignore the existence of mixoplankton.

Conceptual models of planktonic ecology also extend far beyond the form indicated in Fig. 4, to more overarching concepts, which also warrant a revisit under the mixoplankton paradigm. Thus, we can reconsider the underpinning of Margalef’s mandala (Margalef, 1978) and perhaps even revisit the conceptual basis of Longhurst (2007) provinces. When working on Leles *et al.* (2017, 2019) we sought to place biogeographies of CM and NCM protists within the Longhurst province descriptions aware that what we were doing was using a structure built in the era of the traditional paradigm, and allied concepts, to attempt to explain the biogeography of very different functional groups. It may come to pass that a merging of a mixoplanktonic centric revision of Margalef’s mandala (Glibert, 2016) and the Longhurst-style provinces will see a reforming of biogeographic interpretations to provide an enhanced holistic understanding of ocean life.

CONCLUSION

There is clearly much work to do across the entire gamut of plankton research in the context of mixoplankton. It is possible that our holistic view of how planet-scale processes operate may ultimately be little changed. Maybe, to borrow from Williams (1984) commentary upon how the microbial loop was not impacting the science of plankton ecology as much as one may have expected, the mixoplankton paradigm will prove to be just another of the “Emperor’s New Suit of Clothes” (i.e. *de facto* invisible). The microbial loop is still not included commonly in extant plankton models. While mixoplankton are ubiquitous, and we will eventually understand better how their food web operates, perhaps ultimately it is just a wheel spinning inside other wheels of greater consequence, with microbial primary, secondary and bacterial productions being so integrated that we would be better considering them as a whole (Flynn, 1988).

On the other hand, perhaps the mixoplankton paradigm will bring into sharper relief that the microbial loop and the allied bacterial food web, with its role in the microbial carbon pump generating refractory dissolved organic matter (DOM) (Jiao *et al.*, 2010; Lechtenfeld *et al.*, 2014), is indeed collectively worthy of more detailed inclusion in models. The important role of bacteria-mixoplankton interactions in oligotrophic systems is clear (Hartmann *et al.*, 2013; Mitra *et al.*, 2014b). By similar arguments, maybe our concepts for the management of HABs will need to be reappraised, recognizing the importance of alternative trophic interactions (including the needs of some species for certain prey) in addition to the primarily bottom-up light/inorganic-nutrient processes that are currently focussed upon (Glibert *et al.*, 2018; Shumway *et al.*, 2018).

If such events come to pass, then the mixoplankton paradigm will develop into more than just recognizing the importance of photo- and phago-mixotrophic protist plankton. We are only going to find out if we look, and we can only attain that goal by combining the different field of research in plankton ecology in a multidisciplinary approach. We will need a combination of laboratory and field rate determinations, biochemistry (stoichiometry) and molecular approaches, new methodologies for field and survey work and new models to test the mixoplankton paradigm. And, of course, we need to train the next generation of scientists to deliver this.

ACKNOWLEDGEMENTS

This work develops from the application documentation for MixITiN and from subsequent MixITiN workshops and interactions (this justifying the authors to this work).

FUNDING

This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 766327.

REFERENCES

- Adl, S. M., Bass, D., Lane, C. W., Lukeš, J., Schoch, C. L., Smirnov, A., Agatha, S., Berney, C. *et al.* (2019) Revisions to the classification, nomenclature, and diversity of eukaryotes. *J. Eukaryot. Microbiol.*, **66**, 4–119. doi:10.1111/jeu.12691
- Adolf, J. E., Stoecker, D. K. and Harding, L. W. (2003) Autotrophic growth and photoacclimation in *Karlodinium micrum* (Dinophyceae) and *Storeatula major* (Cryptophyceae). *J. Phycol.*, **39**, 1101–1108.
- Alpermann, T. J., Tillmann, U., Beszteri, B., Cembella, A. D. and John, U. (2010) Phenotypic variation and genotypic diversity in a planktonic population of the toxigenic marine dinoflagellate *Alexandrium tamarense*. *J. Phycol.*, **46**, 18–32. doi:10.1111/j.1529-8817.2009.00767.x
- Andersen, K. H., Berge, T., Gonçalves, R. J., Hartvig, M., Heuschele, J., Hylander, S., Jacobsen, N. S., Lindemann, C. *et al.* (2016) Characteristic sizes of life in the oceans, from bacteria to whales. *Annu. Rev. Mar. Sci.*, **8**, 217–241.
- Anderson, O. R. (1978) Fine-structure of a symbiont-bearing colonial radiolarian, *Collosphaera-globularis*, and ¹⁴C isotopic evidence for assimilation of organic-substances from its zooxanthellae. *J. Ultrastruct. Res.*, **62**, 181–189.
- Anderson, R., Charvet, S. and Hansen, P. J. (2018) Mixotrophy in chlorophytes and haptophytes—effect of irradiance, macronutrient, micronutrient and vitamin limitation. *Front. Microbiol.*, **9**, 1704. doi:10.3389/fmicb.2018.01704
- Anderson, R., Jürgens, K. and Hansen, P. J. (2017) Mixotrophic phytoflagellate bacterivory field measurements strongly biased by standard approaches: a case study. *Front. Microbiol.*, **8**, 1398.
- Archibald, J. M. (2009) The puzzle of plastid evolution. *Curr. Biol.*, **19**, R81–R88.
- Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G., Bopp, L. *et al.* (2013). Carbon-concentration and carbon-climate feedbacks in CMIP5 earth system models. *J. Climate*, **26**, 5289–5313. doi:10.1175/JCLI-D-12-00494.1
- Azam, F., Fenchel, T., Field, J. G., Gray, J. S., Meyer-Reil, L. A. and Thingstad, F. (1983) The ecological role of water-column microbes in the sea. *Mar. Ecol. Prog. Ser.*, **10**, 257–263.
- Baretta, J. W., Ebenhöh, W. and Ruurdij, P. (1995) The European Regional Seas Ecosystem Model (ERSEM), a complex marine ecosystem model. *Neth. J. Sea Res.*, **33**, 233–246.
- Beisner, B. E., Grossart, H-P. and Gasol, J. M. (2019) A guide to methods for estimating phago-mixotrophy in nanophytoplankton. *J. Plankt. Res.*, **41**, 77–89. doi:10.1093/plankt/fbz008
- Berge, T., Hansen, P. J. and Moestrup, Ø. (2008a) Feeding mechanism, prey specificity and growth of the plastidic dinoflagellate *Karlodinium armiger* in the light and dark. *Aquat. Microb. Ecol.*, **50**, 279–288.
- Berge, T., Hansen, P. J. and Moestrup, Ø. (2008b) Prey size spectrum and bioenergetics of the mixotrophic dinoflagellate *Karlodinium armiger*. *Aquat. Microb. Ecol.*, **50**, 289–299.
- Berge, T. and Hansen, P. J. (2016) Role of the tertiary plastid of the predatory dinoflagellate *Karlodinium armiger*. *Mar. Ecol. Prog. Ser.*, **549**, 41–54.

- Bhatnagar, A., Chinnasamy, S., Singh, M. and Das, K. C. (2011) Renewable biomass production by mixotrophic algae in the presence of various carbon sources and wastewaters. *Appl. Energy*, **88**, 3425–3431.
- Biard, T., Bigeard, E., Audic, S., Poulain, J., Gutierrez-Rodriguez, A., Pesant, S., Stemann, L. and Not, F. (2017) Biogeography and diversity of Collodaria (Radiolaria) in the global ocean. *ISME J.*, **11**, 1331–1344.
- Biard, T., Stemann, L., Picheral, M., Mayot, N., Vandromme, P., Hauss, H., Gorsky, G., Guidi, L. *et al.* (2016) In situ imaging reveals the biomass of giant protists in the global ocean. *Nat.*, **532**, 504–507.
- Biecheler, B. (1936) Observation de la capture et de la digestion des protées chez un peridinéen vert. *C.R. d la Soc. De boil.*, **121**, 1173–1175.
- Bird, D. F. and Kalf, J. (1986) Bacterial grazing by planktonic lake algae. *Science*, **231**, 493–495.
- Blackbourn, D. J., Taylor, F. J. R. and Blackbourn, J. (1973) Foreign organelle retention by ciliates. *J. Protozool.*, **20**, 286–288.
- Brandenburg, K. M., Wohlrab, S., John, U., Kremp, A., Jerney, J., Krock, B. and Van de Waal, D. B. (2018) Intraspecific trait variation and trade-offs within and across populations of a toxic dinoflagellate. *Ecol. Lett.*, **21**, 1561–1571.
- Bruggeman, J. and Kooijman, S. (2007) A biodiversity-inspired approach to aquatic ecosystem modeling. *Limnol. Oceanogr.*, **52**, 1533–1544.
- Brutemark, A. and Granéli, E. (2011) Role of mixotrophy and light for growth and survival of the toxic haptophyte *Prymnesium parvum*. *Harmful Algae*, **10**, 388–394.
- Burkholder, J. A. M., Glibert, P. M. and Skelton, H. M. (2008) Mixotrophy, a major mode of nutrition for harmful algal species in eutrophic waters. *Harmful Algae*, **8**, 77–93. doi:10.1016/j.hal.2008.08.010
- Burki, F. (2014) The eukaryotic tree of life from a global phylogenomic perspective. *Cold Spring Harb. Perspect. Biol.* **6**, a016147. doi:10.1101/cshperspect.a016147
- Burns, J. A., Paasch, A., Narechania, A. and Kim, E. (2015) Comparative genomics of a bacterivorous green alga reveals evolutionary causalities and consequences of phago-mixotrophic mode of nutrition. *Genome Biol. Evol.*, **7**, 3047–3061.
- Butenschön, M., Clark, J., Aldridge, J. N., Allen, J. I., Artioli, Y., Blackford, J., Bruggeman, J., Cazenave, P. *et al.* (2016) ERSEM 15.06: a generic model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels. *Geosci. Model Dev.*, **9**, 1293–1339 <https://doi.org/10.5194/gmd-9-1293-2016>.
- Calbet, A., Bertos, M., Fuentes-Grünwald, C., Alacid, E., Figueroa, R., Renom, B. and Garcés, E. (2011) Intraspecific variability in *Karlodinium venificum*: Growth rates, mixotrophy, and lipid composition. *Harmful Algae*, **10**, 654–667.
- Caron, D. A. (2016) Mixotrophy stirs up our understanding of marine food webs. *Proc. Natl. Acad. Sci.*, **113**, 2806–2808.
- Carvalho, W. F. and Granéli, E. (2010) Contribution of phagotrophy versus autotrophy to *Prymnesium parvum* growth under nitrogen and phosphorus sufficiency and deficiency. *Harmful Algae*, **9**, 105–115.
- Chojnacka, K. and Noworyta, A. (2004) Evaluation of *Spirulina sp.* growth in photoautotrophic, heterotrophic and mixotrophic cultures. *Enzyme Microb. Technol.*, **34**, 461–465. doi:10.1016/j.enzmictec.2003.12.002
- Cushing, D. H. (1975) *Marine Ecology and Fisheries*, Cambridge University Press, UK.
- Decelle, J., Suzuki, N., Mahé, F., de Vargas, C. and Not, F. (2012) Molecular phylogeny and morphological evolution of the Acantharia (Radiolaria). *Protist*, **163**, 435–450.
- Dolan, J. R. and Pérez, M. T. (2000) Costs, benefits and characteristics of mixotrophy in marine oligotrichs. *Freshwater Biol.*, **45**, 227–238.
- Droop, M. R. (2007) Vitamins, phytoplankton and bacteria: symbiosis or scavenging? *J. Plankton Res.*, **29**, 107–113 <https://doi.org/10.1093/plankt/fbm009>.
- Estep, K. W., Davis, P. G., Keller, M. D. and Sieburth, J. M. N. (1986) How important are oceanic algal nanoflagellates in bacterivory. *Limnol. Oceanogr.*, **31**, 646–650.
- Fasham, M. J. R., Ducklow, H. W. and McKelvie, S. M. (1990) A nitrogen-based model of plankton dynamics in the oceanic mixed layer. *J. Mar. Res.*, **48**, 591–639.
- Faure, E., Not, F., Benoiston, A.-S., Labadie, K., Bittner, L. and Ayata, S.-D. (2019) Mixotrophic protists display contrasted biogeographies in the global ocean. *ISME J.*, doi.org/10.1038/s41396-018-0340-5.
- Fenchel, T. (2008) The microbial loop—25 years later. *J. Exp. Mar. Biol. Ecol.*, **366**, 99–103.
- Figueroa-Martinez, F., Nedelcu, A. M., Smith, D. R. and Reyes-Prieto, A. (2015) When the lights go out: the evolutionary fate of free-living colorless green algae. *New Phytol.*, **206**, 972–982.
- Flynn, K. J. (1988) The concept of "primary production" in aquatic ecology. *Limnol. Oceanogr.*, **33**, 1215–1216.
- Flynn, K. J. and Berry, L. S. (1999) The loss of organic nitrogen during marine primary production may be overestimated significantly when estimated using ¹⁵N substrates. *Proc. Roy. Soc. Lond. B*, **266**, 641–647.
- Flynn, K. J. and Butler, I. (1986) Nitrogen sources for the growth of marine microalgae; role of dissolved free amino acids. *Mar. Ecol. Prog. Ser.*, **34**, 281–304.
- Flynn, K. J. and Hansen, P. J. (2013) Cutting the canopy to defeat the "selfish gene"; conflicting selection pressures for the integration of phototrophy in mixotrophic protists. *Protist*, **164**, 811–823. doi:10.1016/j.protis.2013.09.002
- Flynn, K. J. and Mitra, A. (2009) Building the "perfect beast": modelling mixotrophic plankton. *J. Plankton Res.*, **31**, 965–992. doi:10.1093/plankt/fbp044
- Flynn, K. J., Mitra, A., Glibert, P. M. and Burkholder, J.-A. M. (2018) Mixotrophy in harmful algal blooms: by whom, on whom, when, why, and what next. In Glibert, P. M., Berdalet, E., Burford, M., Pitcher, G. and Zhou, M. (eds), *Global Ecology and Oceanography of Harmful Algal Blooms*. Ecological Studies 232. Springer International Publishing AG. https://doi.org/10.1007/978-3-319-70069-4_7
- Flynn, K. J., St John, M., Raven, J. A., Skibinski, D. O. E., Allen, J. I., Mitra, A. and Hofmann, E. E. (2015) Acclimation, adaptation, traits and trade-offs in plankton functional type models: reconciling terminology for biology and modelling. *J. Plankton Res.*, **37**, 683–691. doi:10.1093/plankt/fbv036
- Flynn, K. J., Stoecker, D. K., Mitra, A., Raven, J. A., Glibert, P. M., Hansen, P. J., Granéli, E. and Burkholder, J. M. (2013) Misuse of the phytoplankton-zooplankton dichotomy: the need to assign organisms as mixotrophs within plankton functional types. *J. Plankton Res.*, **35**, 3–11. doi:10.1093/plankt/fbs062
- Follows, M. J., Dutkiewicz, S., Grant, S. and Chisholm, S. W. (2007) Emergent biogeography of microbial communities in a model ocean. *Science*, **315**, 1843–1846.

- Franks, P. J. S. (2002) NPZ models of plankton dynamics: their construction, coupling to physics, and application. *J. Oceanogr.*, **58**, 379–387.
- Ghyoot, C., Flynn, K. J., Mitra, A., Lancelot, C. and Gypens, N. (2017a) Modeling plankton mixotrophy: a mechanistic model consistent with the Shuter-type biochemical approach. *Front. Ecol. Evol.*, **5**, 78.
- Ghyoot, C., Lancelot, C., Flynn, K. K., Mitra, A. and Gypens, N. (2017b) Introducing mixotrophy into a biogeochemical model describing an eutrophied coastal ecosystem: The Southern North Sea. *Prog. Oceanogr.*, **157**, 1–11. doi:10.1016/j.pocean.2017.08.002
- Glibert, P. M. (2016) Margalef revisited: a new phytoplankton mandala incorporating twelve dimensions, including nutritional physiology. *Harmful Algae*, **55**, 25–30.
- Glibert, P. M., Berdalet E., Burford M., Pitcher G. and Zhou M. (eds) (2018) *Global Ecology and Oceanography of Harmful Algal Blooms*. Ecological Studies (Analysis and Synthesis) 232. Springer, Cham.
- Glibert P. M. Mitra, A., Flynn, K. J., Hansen, P. J., Jeong, H. J. and Stoecker, D. (2019) *Plants Are Not Animals and Animals Are Not Plants, Right? Wrong! Tiny Creatures in the Ocean Can Be Both at Once!* Frontiers for Young Minds <https://kids.frontiersin.org/article/10.3389/frym.2019.00048>
- Gomes, H. d. R., McKee, K., Mile, A., Thandapu, S., Al-Hashmi, K., Jiang, X. and Goes, J. I. (2018) Influence of light availability and prey type on the growth and photo-physiological rates of the mixotroph *Noctiluca scintillans*. *Front. Mar. Sci.*, **5**, 374.
- Granéli, E. (2006) In Turner, J. T. (ed.), *Ecology of Harmful Algae*, Vol. 189, Springer, Berlin, p. 406.
- Granéli, E., Weberg, M. and Salomon, P. S. (2008) Harmful algal blooms of allelopathic microalgal species: the role of eutrophication. *Harmful algae*, **8**, 94–102.
- Hammer, A. C. and Pitchford, J. W. (2005) The role of mixotrophy in plankton bloom dynamics, and the consequences for productivity. *ICES J. Mar. Sci.*, **62**, 833–840.
- Hampel, V., Čepička, I. and Eliáš, M. (2019) Was the mitochondrion necessary to start eukaryogenesis. *Trends Microbiol.*, **27**, 96–104.
- Hansen, P. J. (2011) The role of photosynthesis and food uptake for the growth of marine mixotrophic dinoflagellates. *J. Eukaryot. Microbiol.*, **58**, 203–214.
- Hansen, P. J. and Hjorth, M. (2002) Growth and grazing responses of the prymnesiophyte *Chrysochromulina ericina*: the role of irradiance, prey concentration and pH. *Mar. Biol.*, **141**, 975–983.
- Hansen, P. J., Miranda, L. and Azanza, R. (2004) Green *Noctiluca scintillans*: a dinoflagellate with its own green house. *Mar. Ecol. Prog. Ser.*, **275**, 79–87.
- Hansen, P. J., Skovgaard, A., Glud, R. N. and Stoecker, D. K. (2000) Physiology of the mixotrophic dinoflagellate *Fragilidium subglobosum*. II. Effects of time scale and prey concentration on photosynthetic performance. *Mar. Ecol. Prog. Ser.*, **201**, 137–146.
- Hansson, T. H., Grossart, H.-P., del Giorgio, P. A., St-Gelais, N. F. and Beisner, B. E. (2019) Environmental drivers of mixotrophs in boreal lakes. *Limnol. Oceanogr.*, doi:10.1002/lno.11144
- Hartmann, M., Zubkov, M. V., Scanlan, D. J. and Lepère, C. (2013) *In situ* interactions between photosynthetic picoeukaryotes and bacterioplankton in the Atlantic Ocean: evidence for mixotrophy. *Environ Microbiol. Rep.*, **5**, 835–40. doi:10.1111/1758-2229.12084
- Hitchman, R. B. and Jones, H. L. (2000) The role of mixotrophic protists in the population dynamics of the microbial food web in a small artificial pond. *Freshwater Biol.*, **43**, 231–241.
- Jiao, N., Herndl, G. J., Hansell, D. A., Benner, R., Kattner, G., Wilhelm, S. W., Kirchman, D. L., Weinbauer, S. W. et al. (2010) Microbial production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean. *Nature Rev. Microbiol.*, **8**, 593–599.
- Jeong, H. J., Shim, J. H., Kim, J. S., Park, J. Y., Lee, C. W. and Lee, Y. (1999) Feeding by the mixotrophic thecate dinoflagellate *Fragilidium* cf. *mexicanum* on red-tide and toxic dinoflagellates. *Mar. Ecol. Prog. Ser.*, **196**, 263–277.
- Jeong, H. J., Yeong, D. Y., Park, J. Y., Song, J. Y., Kim, S. T., Lee, S. H., Kim, K. Y. and Yih, W. H. (2005) Feeding by phototrophic red-tide dinoflagellates: five species newly revealed and six species previously known to be mixotrophic. *Aquat. Microb. Ecol.*, **40**, 133–150.
- Jeong, H. J., Yoo, Y. D., Kim, J. S., Seong, K. A., Kang, N. S. and Kim, T. H. (2010) Growth, feeding and ecological roles of the mixotrophic and heterotrophic dinoflagellates in marine planktonic food webs. *Ocean Sci.*, **45**, 65–91.
- John, U., Tillmann, U., Hülskötter, J., Alpermann, T. J., Wohlrab, S. and Van de Waal, D. B. (2015) Intraspecific facilitation by allelochemical mediated grazing protection within a toxigenic dinoflagellate population. *Proc. Roy. Soc. B.*, **282**, 20141268, <https://doi.org/10.1098/rspb.2014.1268>.
- John, U., Tillmann, U. and Medlin, L. K. (2002) A comparative approach to study grazer interference and lipid composition of a toxic and non-toxic clone of *Chrysochromulina polylepis* (Prymnesiophyceae). *Harmful Algae*, **1**, 45–57.
- Johnson, M. D. (2015) Inducible mixotrophy in the Dinoflagellate *Prorocentrum minimum*. *J. Eukaryot. Microbiol.*, **62**, 431–443.
- Johnson, M. D. and Moeller, H. V. (2018) Mixotrophy in protists: from model systems to mathematical models. *Front. Mar. Sci.*, **5**, 490.
- Jones, H. L. J. (1997) A classification of mixotrophic protists based on their behaviour. *Freshwater Biol.*, **37**, 35–43. doi:10.1046/j.1365-2427.1997.00138.x
- Jones, R. I. (1994) Mixotrophy in planktonic protists as a spectrum of nutritional strategies. *Mar. Microb. Food Webs*, **8**, 87–96.
- Jones, R. I. (2000) Mixotrophy in planktonic protists: an overview. *Freshwater Biol.*, **45**, 219–226.
- Jost, C., Lawrence, C. A., Campolongo, F., Van de Bund, W., Hill, S. and DeAngelis, D. L. (2004) The effects of mixotrophy on the stability and dynamics of a simple planktonic food web model. *Theor. Popul. Biol.*, **66**, 37–51.
- Kim, M., Drumm, K., Daugbjerg, N. and Hansen, P. J. (2017) Dynamics of sequestered cryptophyte nuclei in *Mesodinium rubrum* during starvation and refeeding. *Front. Microbiol.*, **8**, 423. doi:10.3389/fmicb.2017.00423
- Kolisko, M., Boscaro, V., Burki, F., Lynn, D. H. and Keeling, P. J. (2014) Single-cell transcriptomics for microbial eukaryotes. *Curr. Biol.*, **24**, R1081–R1082 <https://doi.org/10.1016/j.cub.2014.10.026>.
- Kremp, A., Godhe, A., Egardt, J., Dupont, S., Suikkanen, S., Casabianca, S. and Penna, A. (2012) Intraspecific variability in the response of bloom-forming marine microalgae to changed climate conditions. *Ecol. Evol.*, **2**, 1195–1207 <https://doi.org/10.1002/ece3.245>.
- Laval-Peuto, M. and Febvre, M. (1986) On plastid symbiosis in *Tontonia appendiculariformis* (Ciliophora, Oligotrichina). *BioSystems*, **19**, 137–158.

- Lechtenfeld, O. J., Kattner, G., Flerus, R., McCallister, S. L., Schmitt-Kopplin, P. and Koch, B. P. (2014) Molecular transformation and degradation of refractory dissolved organic matter in the Atlantic and Southern Ocean. *Geochim. Cosmochim. Acta*, **126**, 321–337 <https://doi.org/10.1016/j.gca.2013.11.009>.
- Legendre, L., Rivkin, R. B., Weinbauer, M. G., Guidi, L. and Uitz, J. (2015) The microbial carbon pump concept: potential biogeochemical significance in the globally changing ocean. *Prog. Oceanogr.*, **134**, 432–450.
- Leles, S. G., Luca Polimene L., Bruggeman, J., Blackford, J., Ciavatta, S., Mitra, A. and Flynn K. J. (2018) Modelling mixotrophic functional diversity and implications for ecosystem function. *J. Plankt. Res.*, **40**, 627–642. doi:10.1093/plankt/fby044
- Leles, S. G., Mitra A., Flynn, K. J., Stoecker, D. K., Hansen, P. J., Calbet, A., McManus, G. B., Sanders, R. W. *et al.* (2017) Oceanic protists with different forms of acquired phototrophy display contrasting biogeographies and abundance. *Proc. Roy. Soc. B*, **284**, doi:10.1098/rspb.2017.0664
- Leles, S. G., Mitra, A., Flynn, K. J., Tillmann, U., Stoecker, D. K., Jeong, H. J., Burkholder, J. A., Hansen, P. J. *et al.* (2019) Sampling bias misrepresents the biogeographic significance of constitutive mixotrophs across global oceans. *Glob. Ecol. Biogeogr.*, **28**, 418–428.
- Liang, Y., Sarkany, N. and Cui, Y. (2009) Biomass and lipid productivities of *Chlorella vulgaris* under autotrophic, heterotrophic and mixotrophic growth conditions. *Biotechnol. Lett.*, **31**, 1043–1049. doi:10.1007/s10529-009-9975-7
- Lin, C-H., Flynn, K. J., Mitra, A. and Glibert, P. M. (2018) Simulating effects of variable stoichiometry and temperature on mixotrophy in the harmful dinoflagellate *Karlodinium veneficum*. *Frontiers in Marine Science*, **5**, 320. doi:10.3389/fmars.2018.00320
- Litchman, E. and Klausmeier, C. A. (2008) Trait-based community ecology of phytoplankton. *Annu. Rev. Ecol. Evol. Syst.*, **39**, 615–639.
- Litchman, E., Klausmeier, C. A., Schofield, O. M. and Falkowski, P. G. (2007) The role of functional traits and trade-offs in structuring phytoplankton communities: scaling from cellular to ecosystem level. *Ecol. Lett.*, **10**, 1170–1181.
- Litchman, E., Ohman, M. D. and Kiørboe, T. (2013) Trait-based approaches to zooplankton communities. *J. Plankton Res.*, **35**, 473–484.
- Longhurst, A. (2007) *Ecological Geography of the Sea*, 2nd edn, Academic Press, San Diego, CA.
- Lundgren, V. M., Glibert, P. M., Granéli, E., Vidyarthna, N. K., Fiori, E., Ou, L., Flynn, K. J., Mitra, A. *et al.* (2016) Metabolic and physiological changes in *Prymnesium parvum* when grown under, and grazing on prey of, variable nitrogen:phosphorus stoichiometry. *Harmful Algae*, **55**, 1–12.
- Mafra, L. L. Jr., Nagai, S., Uchida, H., Tavares, C. P. S., Escobar, B. P. and Suzuki, T. (2016) Harmful effects of *Dinophysis* to the ciliate *Mesodinium rubrum*: Implications for prey capture. *Harmful Algae*, **59**, 82–90. doi:10.1016/j.hal.2016.09.009
- Margalef, R. (1978) Life-forms of phytoplankton as survival alternatives in an unstable environment. *Oceanol. Acta*, **1**, 493–509.
- Martijn, J. and Ettema, T. J. (2013) From archaeon to eukaryote: the evolutionary dark ages of the eukaryotic cell. *Biochem. Soc. Trans.*, **41**, 451–457. doi:10.1042/BST20120292
- Martin, W. F., Tielens, A. G. M., Mentel, M., Garg, S. G. and Gould, S. B. (2017) The physiology of phagocytosis in the context of mitochondrial origin. *Microbiol. Mol. Biol. Rev.*, **81**, e00008–17 doi:10.1128/MMBR.00008-17
- Martins, C. A., Kulis, D., Franca, S. and Anderson, D. M. (2004) The loss of PSP toxin production in a formerly toxic *Alexandrium lusitanicum* clone. *Toxicon*, **43**, 195–205.
- McKie-Krisberg, Z. M., Sanders, R. W. and Gast, R. J. (2018) Evaluation of mixotrophy-associated gene expression in two species of Polar marine algae. *Front. Mar. Sci.*, **5**, 273. doi:10.3389/fmars.2018.00273
- Mitra, A. (2016) *Uncovered: The Mysterious Killer Triffids that Dominate Life in Our Oceans*. <http://theconversation.com/uncovered-the-mysterious-killer-triffids-that-dominate-life-in-our-oceans-67387>
- Mitra, A. (2018) *Tiny Creatures, Part plant and Part Animal, May Control the Fate of the Planet*. Scientific American, April 2018. <https://www.scientificamerican.com/article/tiny-creatures-part-plant-and-part-animal-may-control-the-fate-of-the-planet/>
- Mitra, A., Castellani, C., Gentleman, W. C., Jónasdóttir, S. H., Flynn, K. J., Bode, A., Halsband, C., Kuhn, P. *et al.* (2014a) Bridging the gap between marine biogeochemical and fisheries sciences; configuring the zooplankton link. *Prog. Oceanogr.*, **129**, 176–199. doi:10.1016/j.pocean.2014.04.025
- Mitra, A., Flynn, K. J., Burkholder, J. M., Berge, T., Calbet, A., Raven, J. A., Granéli, E., Hansen, P. J. *et al.* (2014b) The role of mixotrophic protists in the biological carbon pump. *Biogeosciences*, **11**, 995–1005. doi:10.5194/bg-11-995-2014
- Mitra, A., Flynn, K. J., Tillmann, U., Raven, J. A., Caron, D., Stoecker, D. K., Not, F., Hansen, P. J. *et al.* (2016) Defining planktonic protist functional groups on mechanisms for energy and nutrient acquisition; incorporation of diverse mixotrophic strategies. *Protist*, **167**, 106–120 <http://dx.doi.org/10.1016/j.protis.2016.01.003>.
- Morales-Sánchez, D., Tinoco-Valencia, R., Kyndt, J. and Martínez, A. (2013) Heterotrophic growth of *Neochloris oleoabundans* using glucose as a carbon source. *Biotechnol. Biofuels*, **6**, 100.
- Naselli-Flores, L. and Barone, R. (2019) Mixotrophic phytoplankton dynamics in a shallow Mediterranean water body: how to make a virtue out of necessity. *Hydrobiologia*, **831**, 33–41. doi:10.1007/s10750-018-3507-1
- Pitta, P. and Giannakourou, A. (2000) Planktonic ciliates in the oligotrophic Eastern Mediterranean: vertical, spatial distribution and mixotrophy. *Mar. Ecol. Prog. Ser.*, **194**, 269–282.
- Pitta, P., Giannakourou, A. and Christaki, U. (2001) Planktonic ciliates in the oligotrophic Mediterranean Sea: Longitudinal trends of standing stocks, distributions and analysis of food vacuole contents. *Aquat. Microb. Ecol.*, **24**, 297–311.
- Plagányi, E. E. (2007) *Models for an Ecosystem Approach to Fisheries*. FAO (Food and Agriculture Organization of the United Nations) Fisheries Technical Paper 477, FAO, Rome, 108.
- Pomeroy, L. R. (1974) The ocean's food web, a changing paradigm. *BioScience*, **9**, 499–504.
- Pomeroy, L. R., Williams, P. J. LEB., Azam, F. and Hobbie, J. E. (2007) The microbial loop. *Oceanography*, **20**, 28–33.
- Rokitta, S. D., de Nooijer, L. J., Trimborn, S., de Vargas, C., Rost, B. and John, U. (2011) Transcriptome analyses reveal differential gene expression patterns between the life-cycle stages of *Emiliana huxleyi* (haptophyta) and reflect specialization to different ecological niches. *J. Phycol.*, **47**, 829–838.
- Saad, J. F., Unrein, F., Tribelli, P. M., López, N. and Izaguirre, I. (2016) Influence of lake trophic conditions on the dominant mixotrophic algal assemblages. *J. Plankton Res.*, **38**, 818–829.

- Saito, H., Furuya, K. and Lirdwitayaprasit, T. (2006) Photoautotrophic growth of *Noctiluca scintillans* with the endosymbiont *Pedinomonas noctilucae*. *Plankton Benthos Res.*, **1**, 97–101.
- Sanders, R. W. (1991) Mixotrophic protists in marine and freshwater ecosystems. *J. Protozool.*, **38**, 76–81.
- Sanders, R. W. (2011) Alternative nutritional strategies in protists: symposium introduction and a review of freshwater protists that combine photosynthesis and heterotrophy. *J. Eukaryotic Microbiol.*, **58**, 181–184.
- Sanders, R. W., Porter, K. G. and Caron, D. A. (1990) Relationship between phototrophy and phagotrophy in the mixotrophic chrysophyte *Poteroochromonas malhamensis*. *Microb. Ecol.*, **19**, 97–109.
- Santoferrara, L. F., Guida, S., Zhang, H. and McManus, G. B. (2014) De novo transcriptomes of a mixotrophic and a heterotrophic ciliate from marine plankton. *PLoS One*, **9**, e101418.
- Selosse, M. A., Charpin, M. and Not, F. (2017) Mixotrophy everywhere on land and in water: the grand écart hypothesis. *Ecol. Lett.*, **20**, 246–263.
- Shumway, S. E., Burkholder, J.-A. and Morton, S. L. (eds.) (2018) *Harmful Algal Blooms: A Compendium Desk Reference*. John Wiley & Sons, Hoboken, NJ.
- Sieburth, J. M. N., Smetacek, V. and Lenz, J. (1978) Pelagic ecosystem structure: Heterotrophic compartments of the plankton and their relationship to plankton size fractions. *Limnol. Oceanogr.*, **23**, 1256–1263.
- Skovgaard, A. and Hansen, P. J. (2003) Food uptake in the harmful alga *Prymnesium parvum* mediated by excreted toxins. *Limnol. Oceanogr.*, **48**, 1161–1166.
- Skovgaard, A., Hansen, P. J. and Stoecker, D. K. (2000) Physiology of the mixotrophic dinoflagellate *Fragilidium subglobosum*. I. Effects of phagotrophy and irradiance on photosynthesis and carbon content. *Mar. Ecol. Prog. Ser.*, **201**, 129–136.
- Skovgaard, A., Legrand, C., Hansen, P. J. and Granéli, E. (2003) Effects of nutrient limitation on food uptake in the toxic haptophyte *Prymnesium parvum*. *Aquat. Microb. Ecol.*, **31**, 259–265.
- Smalley, G. W., Coats, D. W. and Stoecker, D. K. (2003) Feeding in the mixotrophic dinoflagellate *Ceratium furca* is influenced by intracellular nutrient concentrations. *Mar. Ecol. Prog. Ser.*, **262**, 137–151.
- Smalley, G. W., Coats, D. W. and Stoecker, D. K. (2012) Influence of inorganic nutrients, irradiance, and time of day on food uptake by the mixotrophic dinoflagellate *Neoceratium furca*. *Aquat. Microb. Ecol.*, **68**, 29–41.
- Smith, S. L., Pahlow, M., Merico, A., Acevedo-Trejos, E., Sasai, Y., Yoshikawa, C., Sasaoka, K., Fujiki, T. et al. (2016) Flexible phytoplankton functional type (FlexPFT) model: size-scaling of traits and optimal growth. *J. Plankton Res.*, **38**, 977–992 <https://doi.org/10.1093/plankt/fbv038>.
- Stewart, F. J., Ulloa, O. and DeLong, E. F. (2012) Microbial metatranscriptomics in a permanent marine oxygen minimum zone. *Environ. Microbiol.*, **14**, 23–40. doi:10.1111/j.1462-2920.2010.02400.x
- Stickney, H. L., Hood, R. R. and Stoecker, D. K. (2000) The impact of mixotrophy on planktonic marine ecosystems. *Ecol. Modell.*, **125**, 203–230.
- Stoecker, D. K. (1998) Conceptual models of mixotrophy in planktonic protists and some ecological and evolutionary implications. *Eur. J. Protist.*, **34**, 281–290.
- Stoecker, D. K., Hansen, P. J., Caron, D. A. and Mitra, A. (2017) Mixotrophy in the Marine Plankton. *Ann. Rev. Mar. Sci.*, **9**, 311–335.
- Stoecker, D. K., Johnson, M. D., de Vargas, C. and Not, F. (2009) Acquired phototrophy in aquatic protists. *Aquat. Microb. Ecol.*, **37**, 279–310.
- Stoecker, D. K. and Lavrentyev, P. J. (2018) Mixotrophic plankton in the Polar Seas: a Pan-Arctic review. *Front. Mar. Sci.*, **5**, 292 <https://www.frontiersin.org/article/10.3389/fmars.2018.00292>.
- Stoecker, D. K., Michaels, A. E. and Davis, L. H. (1987) Large proportion of marine planktonic ciliates found to contain functional chloroplasts. *Nature*, **326**, 790.
- Swanberg, N. R. (1983) The trophic role of colonial Radiolaria in oligotrophic oceanic environments. *Limnol. Oceanogr.*, **28**, 655–666.
- Swanberg, N. R. and Caron, D. A. (1991) Patterns of sarcodine feeding in epipelagic oceanic plankton. *J. Plankton Res.*, **13**, 287–312.
- Tai, V., James, E. R., Perlman, S. J. and Keeling, P. J. (2013) Single-cell DNA barcoding using sequences from the small subunit rRNA and internal transcribed spacer region identifies new species of *Trichonympha* and *Trichomitopsis* from the hindgut of the termite *Zootermopsis angusticollis*. *PLoS One*, **8**, e58728, <https://doi.org/10.1371/journal.pone.0058728>.
- Takishita, K., Kolke, K., Maruyama, T. and Ogata, T. (2002) Molecular evidence for plastid robbery (kleptoplastidy) in *Dinophysis*, a dinoflagellate causing diarrhetic shellfish poisoning. *Protist*, **153**, 293–302. doi:10.1078/1434-4610-00106
- Tillmann, U. (2003) Kill and eat your predator: a winning strategy of the planktonic flagellate *Prymnesium parvum*. *Aquat. Microb. Ecol.*, **32**, 73–84.
- Turner, J. T. (2015) Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's biological pump. *Prog. Oceanogr.*, **130**, 205–248. doi:10.1016/j.pocean.2014.08.005
- Våge, S., Castellani, M., Giske, J. and Thingstad, T. F. (2013) Successful strategies in size structured mixotrophic food webs. *Aquat. Ecol.*, **47**, 329–347 <https://doi.org/10.1007/s10452-013-9447-y>.
- Ward, B. A., Dutkiewicz, S., Barton, A. D. and Follows, M. J. (2011) Biophysical aspects of resource acquisition and competition in algal mixotrophs. *Am. Nat.*, **178**, 98–112.
- Ward, B. A. and Follows, M. J. (2016) Marine mixotrophy increases trophic transfer efficiency, mean organism size, and vertical carbon flux. *Proc. Natl. Acad. Sci.*, **113**, 2958–2963. doi:10.1073/pnas.151711.8113
- Wilhelm, S. W. and Suttle, C. A. (1999) Viruses and nutrient cycles in the sea: viruses play critical roles in the structure and function of aquatic food webs. *BioScience*, **49**, 781–788.
- Wilken, S., Huisman, J., Naus-Wiezer, S. and Donk, E. (2013) Mixotrophic organisms become more heterotrophic with rising temperature. *Ecol. Lett.*, **16**, 225–233.
- Williams, P. J. LeB. (1981) Incorporation of microheterotrophic processes into the classical paradigm of the planktonic food web. *Kieler Meeresforsch. Sondh.*, **5**, 1–28.
- Williams, P. J. LeB. (1984) Bacterial production in the marine food chain: The emperor's new suit of clothes? In Fasham, M. J. R. (eds) *Flows of Energy and Materials in Marine Ecosystems*. NATO Conference Series 13. Springer, Boston, MA.
- Wohlrab, S., Falcke, J., Lin, S., Zhang, H., Elferink, S., Neuhaus, S., Voss, D., Zielinski, O. et al. (2018) Metatranscriptome profiling indicates size-dependent differentiation in plastic and conserved community traits and functional diversification in dinoflagellate communities. *Front. Mar. Sci.*, **5**, 358. doi:10.3389/fmars.2018.00358.

- Wyndham, J. (1951) *The Day of the Triffids*, Michael Joseph, 0-7181-0093-X.
- Yelton, A. P., Acinas, S. G., Sunagawa, S., Bork, P., Pedrós-Alió, C. and Chisholm, S. W. (2016) Global genetic capacity for mixotrophy in marine picocyanobacteria. *ISME J.*, **10**, 2946.
- Yool, A., Popova, E. E. and Anderson, T. R. (2013) MEDUSA-2.0: an intermediate complexity biogeochemical model of the marine carbon cycle for climate change and ocean acidification studies. *Geosci. Model Dev.*, **6**, 1767–1811.
- Yutin, N., Wolf, M. Y., Wolf, Y. I. and Koonin, E. V. (2009) The origins of phagocytosis and eukaryogenesis. *Biol. Direct*, **26**, **4**, 9.
- Zubkov, M. V. and Tarran, G. A. (2008) High bacterivory by the smallest phytoplankton in the North Atlantic Ocean. *Nature Letters*, **455**, 224–227.