

Raman characterization of the UV-protective pigment gloeocapsinpotential biosignatures of early Earth records

Jean-Yves STORME^{1,*}, Stjepko GOLUBIC^{1,2,3}, Annick WILMOTTE³, Julia KLEINTEICH³, David VELAZQUEZ³, Emmanuelle J. JAVAUX¹

Keywords: Raman microspectroscopy, Biosignatures, Gloeocapsin, Parietin, Scytonemin

Abstract

Extracellular UV-screening pigments gloeocapsin and scytonemin present in the EPS envelopes of freshwater and marine cyanobacteria were studied by different spectroscopy methods. The Raman spectral analysis and the UV-VIS light absorbance spectra of the extracellular pigment gloeocapsin showed that it shared Raman and UV spectral signatures with parietin, a radiation-protective pigment present in lichens. Gloeocapsin occurs in several cyanobacterial species, mostly with exclusion of scytonemin, indicating that these pigments have evolved in cyanobacteria as separate protective strategies. Both gloeocapsin and scytonemin are widely and species-specifically distributed across different cyanobacterial taxa. The widespread occurrence of these pigments may suggest an early origin, while their detection by Raman Spectroscopy makes them potential biosignatures for cyanobacteria in the early Earth record and demonstrates the usefulness of non-destructive Raman spectroscopy analyses for the search of complex organics, including possible photosynthetic pigments, in early Earth and extraterrestrial samples.

¹ University of Liège, Palaeobiogeology, Palaeobotany, Palaeopalynology, Department of Geology, Quartier Agora (bat. B18) – Allée du 6 Août, 14, 4000 Liège (Sart-Tilman), BELGIUM

² Boston University, Biological Science Center, Cummington Street 5, 02215 Boston, USA

University of Liège, Center for Protein Engineering, Department of Life Sciences, Quartier Agora (bat. B6) – Allée du 6 Août, 11/, 4000 Liège (Sart-Tilman), BELGIUM

^{*}Corresponding author: jystorme@ulg.ac.be, +32 (0) 4 3665122