MCZ

UNIVERSETY

# New Muricidae (Mollusca: Neogastropoda) from the Lower Miocene Cantaure Formation of Venezuela

Bernard LANDAU \* Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, Netherlands

Centro de Geologia da Universidade de Lisboa. Campo Grande, 1749-016 Lisboa, Portugal

and

International Health Centres, Av. Infante de Henrique 7, Areias São João, 8200 Albufeira, Portugal bernielandau@sapo.pt

> Roland HOUART Research Associate Institut royal des Sciences naturelles de Belgique Rue Vautier, 29, 1000 Bruxelles, Belgium. roland.houart@skynet.be

> > \* Corresponding author

**ABSTRACT.** Seven new muricine records are added to that for the Lower Miocene Cantaure Formation of Venezuela, four of which are new: *Siratus harzhauseri* nov. sp. *Ocenebra etteri* nov. sp., *Typhina canaliculata* nov. sp. and *Laevityphis jungi* nov. sp., and three are identified to genus level *Phyllonotus* sp. cf. *P. infrequens* (Vokes, 1963), *Purpurellus* sp. cf. *P. repetiti* (Vokes, 1970) and *Vitularia* sp. cf. *V. salebrosa* (King & Broderip, 1832). The genus *Ocenebra* is recorded in the western Atlantic for the first time.

**KEYWORDS.** Muricidae, Mollusca, Miocene, Cantaure Formation, Venezuela, new records, new taxa.

#### **INTRODUCTION**

In his landmark monograph on the molluscan assemblage of the Cantaure Formation, Jung (1965) recorded and described ten muricine species from the deposits. Further new species were added in the 1990's by Vokes (1992, 1994, 1995), Vermeij & Vokes (1997), and later by Vermeij (2001) and Landau & Vermeij (2010). A full list of all species described or recorded from the Cantaure Formation subsequent to Jung's (1965) monograph were listed by Landau & Vermeij (2010, p. 105, appendix 1).

A short account of the geographical and stratigraphical range was given by Landau & Vermeij (2010, p. 99). This will not be repeated here, except to say that the assemblage is now considered to be late early Miocene in age (Landau *et al.*, 2012, p. 258, chart 18). In this contribution to the knowledge of the gastropod assemblage found in the late early Cantaure Formation

of Venezuela, we focus on new species, or taxa not previously recorded from the assemblage. This increases the number of muricine species known from the formation to 29.

#### Material and methods

The material described here is from the Gibson-Smith collection housed in the Naturhistorisches Museum Basel (NHMB coll.), Switzerland and the Bernard

Landau collection (BL coll.), now deposited in the Naturhistorisches Museum Wien (NHMW coll.), Vienna.

The classification adopted here is according to WoRMS (http://www.marinespecies.org/index.php). The descriptions adopt the terminology suggested by Merle (1999, 2001) (Figs 1, 8), see next page:

Р	Primary cord
S	secondary cord
t	tertiary cord
Ad	adapical (or adapertural)
Ab	abapical (or abapertural)
SP	Subsutural cord
IP	Infrasutural primary cord (primary cord on shoulder)
adis	adapical infrasutural secondary cord (shoulder)
abis	abapical infrasutural secondary cord (shoulder)
P1	Shoulder cord
P2-P6	Primary cords of the convex part of the teleoconch whorl
s1-s6	secondary cords of the convex part of the teleoconch whorl
example: s1 = secondary cord between P1 and P2; s2 = secondary cord between P2 and P3, etc.	
ADP	adapertural primary cord on the siphonal canal
MP	median primary cord on the siphonal canal
ABP	abapertural primary cord on the siphonal canal
ads :	adapertural secondary cord on the siphonal canal
ms :	median secondary cord on the siphonal canal
abs :	abapertural secondary cord on the siphonal canal

#### SYSTEMATIC PALAEONTOLOGY

Superfamily **MURICOIDEA** Rafinesque, 1815 Family **MURICIDAE** Rafinesque, 1815 Subfamily **MURICINAE** Rafinesque, 1815 Genus *Siratus* Jousseaume, 1880

Type species. – *Murex senegalensis* Gmelin, 1791, by original designation. Recent, Brazil.

Discussion. As discussed by Barco et al. (2010), Siratus Jousseaume, 1880 is closer to Vokesimurex Petuch, 1994 than it is to Chicoreus de Montfort, 1810, and all three should be used at full generic rank. Merle et al. (2011, p. 72) discussed the difficulty in separating some species of Siratus from Vokesimurex. They suggested that well-developed secondary internal denticles and numerous columellar denticles distinguished many Siratus from Vokesinurex, but that these characters are variable. A more reliable character is the shape of the siphonal canal, which is bent in Siratus and straight in Vokesimurex. The new species from Cantaure has weak apertural dentition on both lips, which can be present in either genus, but the siphonal canal, although incomplete, is clearly bent, placing it in the genus Siratus.

### Siratus harzhauseri nov. sp. Figures 1, 9-11

**Dimensions and type material.** Holotype; NHMW 2013/0476/0010 (ex BL coll.) (Figs 1, 9-11), height 48.6 mm.

**Etymology.** Named for Mathias Harzhauser of the Naturhistorisches Museum Wicn, friend of the senior author, in recognition of his contributions to palacomalacology.

**Type locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela (= locality GS1PGNA of Gibson-Smith & Gibson-Smith, 1979).

**Type stratum.** Cantaure Formation (late early Miocene: Burdigalian).

**Diagnosis.** A medium sized *Siratus* species, with a relatively elongated shell, a narrowly canaliculated suture, three varices per whorl bearing short, stout spines placed mid-whorl, with three intervarical axial ridges on early whorls, four with prominent tubercles mid-whorl on last 1½ whorls.

Description. Shell medium-sized, fusiform, spire tall. Protoconch not preserved. Teleoconch of six high, weakly-shouldered whorls, with a narrow, concave, sub-horizontal subsutural area; whorls convex below. Suture impressed, shallowly canaliculated. Three elevated varices per whorl, placed at 120°, aligned almost vertically, bearing a short, stout spine placed just above mid-whorl corresponding to P1, giving whorls a somewhat angular appearance. Intervarical axial sculpture consisting of three nodulose ridges on early whorls, increasing to four on abapical half of penultimate whorl, where the nodules mid-whorl become more prominent. Last whorl fusoid; subsutural area with IP, adis and abis of almost equal strength. Below shoulder P2-P6 narrow, s2-s6 developed in the interspaces between the primary cords. ADP and MP developed on base, ads and ms weaker. Tertiary cords developed throughout between primary and secondary elements. Siphonal canal below ms missing. Aperture of moderate size, ovate. Outer lip convex, margin erect bearing 16 denticles at the inner edge. Labral varix moderately expanded, rounded, bearing straight spine at shoulder. Anal sinus broad, shallow; siphonal canal incomplete, but long, narrow, open, pointing slightly to left. Columellar lip well delimited, narrowly expanded, adherent adapically, erect abapically, bearing clongated denticles along its entire length.



**Figure 1.** *Siratus harzhauseri* nov. sp., holotype NHMW 2013/0476/0010 (ex BL coll.), height 48.6 mm, 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela, Cantaure Formation, Burdigalian, late early Miocene.

Discussion. Siratus harzhauseri nov. sp. is most similar to the Caribbean Pliocene-Recent S. formosus (Sowerby, 1841), but differs in having a more elongated fusiform shell shape, in having a shallowly canaliculated suture, in having P1 and hence the shoulder spine placed lower, closer to mid-whorl and shorter, and in having the intervarical axial ridges more strongly nodulose mid-whorl on the last 11/2 whorls. Siratus domingensis (Sowerby, G. B. I, 1850) from the early Pliocene Gurabo Formation of the Dominican Republic is similar to S. formosus and was considered to be ancestral to it by Vokes (1989). Apart for a few exceptionally large specimens, S. domingensis is smaller than either S. formosus or S. harzhauseri, it has smaller spines and a more recurved siphonal canal. Siratus domingensis has three-five intervarical axial ridges, which are narrower and the suture is not canaliculated. Siratus articulatus (Reeve, 1845) from the Pliocene-Recent Caribbean can easily be separated as P1, P3, P5 and P6 form more or less long spines at the apertural varix and the suture is not canaliculated. Another closely similar species is the early Pleistocene to Recent Caribbean Siratus springeri (Bullis, 1964), but this species differs from S. harzhauseri in having convex rather than angular whorl profile, in having only two or three axial intervarical ridges, in having the labral varix dceply excavated on the abapertural side, and in most specimens P1, P5, P6 are strongly developed.

Two further *Siratus* species occur in the Cantaure assemblage: *S. quirosensis* (F. Hodson, 1931), which is much smaller than *S. harzhanseri*, with only two axial intervarical ridges that are more prominent than

in *S. harzhauseri*, and P1, P3 and P5 are strongly developed. The second, *Siratus denegatus* (Jung, 1966), is quite different from all its congeners in having a rather triangular shell, a sharply carinate last whorl, only one axial intervarical ridge, very fine spiral sculpture, and P1 developed into a short, sharp spine at the shoulder.

#### Genus Phyllonotus Swainson, 1833

Type species. – *Murex imperialis* (var. a) Swainson, 1833, by subsequent designation, Swainson, 1833, pl. 109 (= *Murex imperialis* Swainson, 1831) (not Fischer de Waldheim, 1807) (= *Murex margaritensis* Abbott, 1958). Recent, West Atlantic.

#### *Phyllonotus* sp. cf. *P. infrequens* (Vokes, 1963) Figures 12-13

cf. 1963. *Murex (Phyllonotus) infrequens* Vokes, p. 156, pl. 1, fig. 4.

**Material.** 1 specimen NHMW 2013/0476/0011 (ex BL coll.), height 35.3 mm (Figs 12-13).

**Locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela, Cantaure Formation, Burdigalian, late early Miocene.

**Discussion.** This specimen resembles the species illustrated by Vokes (1989, 1990) as *Phyllonotus infrequens* from the Early Miocene Baitoa Formation of the Dominican Republic. However *Phyllonotus* 

*infrequens* from the Early Miocene Chipola Formation of Florida has a more globose shell with a less impressed suture, a more rounded aperture and less conspieuous secondary and tertiary spiral cords. The specimen from the Baitoa Formation and our specimen from the Cantaure Formation are somewhat related, although the Cantaure one is more globose, has a more impressed suture and a narrower siphonal canal.

### Genus Purpurellus Jousseaume, 1880

Type species. – *Murex gambiensis* Reeve, 1845, by original designation. Recent, West Africa.

Emerson & D'Attilio (1969) demonstrated that, on the basis of radular morphology, *Purpurellus* Jousseaume, 1880 was closely related to *Pterynotus*. *Purpurellus* differs from *Pterynotus* in having a sealed siphonal canal and an operculum with a eentral nucleus, instead of apieal in *Pterynotus s.s.* The use of the taxon at full generic rank is now generally accepted (Landau *et al.*, 2007, Merle *et al.*, 2011, Bouehet & Houart, 2013).

#### *Purpurellus* sp. cf. *P. repetiti* (Vokes, 1970) Figures 14-17

cf. 1970 Pterynotus (Purpurellus) repetiti Vokes, p. 16, pl. 3, fig. 4.

Material. 1 incomplete specimen, NHMW 2013/0476/0008 (ex BL coll.), height 27.2 mm (Figs 14-15); 1 incomplete specimen, NHMW 2013/0476/0009 (ex BL eoll.), height 32.4 mm (Figs 16-17).

**Locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela (=locality GS1PGNA of Gibson-Smith & Gibson-Smith, 1979), Cantaure Formation (early Miocene: Burdigalian).

Discussion. Vokes (1989, p. 55) commented that Purphrellus species could be difficult to separate. The two incomplete speeimens illustrated here resemble the specimen described as Purphrellus repetiti (Vokes, 1970) from the "Silverdale Beds" of North Carolina, originally considered Early Miocene, but placed in the Late Oligocene Chiekasawhayan Stage by Gibson (1977, p. 202; 1983, p. 38). This species is characterised by its relatively tall spire, moderately expanded varices, with short straight spines at the shoulder, single sharp intervarical node, and very weak spiral sculpture. It is most like the European Miocene-Pliocene Purpurellus veranyi (Paulucci, 1866) (see Landau et al., 2007; Merle et al., 2011), but this species has broader intervarical nodes and a relatively smaller, more circular aperture. Apart from these constant differences, most specimens of P. veranyi arc broader, have stronger, recurved spines at the shoulder, slightly more prominent spiral sculpture,

and more expanded labral varix. However, given the badly preserved material from the Cantaure Formation and the few specimens involved, the identification remains doubtful.

The only other western Atlantie eongener is *Purpurellus mirificus* Vokes, 1989 from the early Pliocene Gurabo Formation of the Dominican Republic, which is quite different from *P. repetiti* and *P. verauyi* in having greatly expanded wing-like varices, similar to the Reeent West African *P. gambieusis* (Reeve, 1845).

In Europe the fossil record also starts in the Late Oligocene with an undescribed speeies listed by Lozouet (1986, p. 377) from the Atlantic Late Oligocene of France. Today *Purpurellus* has a disjunct distribution, P. gambiensis (Reeve, 1845) from West Africa and two species, P. pinniger (Broderip, 1833) and P. macleani (Emerson & D'Attilio, 1969) from the Paeific coast of Tropical America. In the fossil record, specimens of Purpurellus are always rare, and although the first record is Late Oligocene in both the eastern and western Atlantic, it is likely that the genus is a Palaeogene offshoot of Pterynotus, which originated in the Tethys, immigrated to the western Atlantic and colonised the Tropical American Pacific before the closure of the central American Seaway. subsequently dying out in the western Atlantic.

#### Subfamily **OCENEBRINAE** Cossmann, 1903 Genus *Ocenebra* Gray, 1847

Type species. – *Murex erinaceus* Linnaeus, 1758, by monotypy. Recent, northeastern Atlantic, Mediterranean.

# *Ocenebra etteri* nov. sp. Figures 18-23

**Dimensions and type material.** Holotype, NHMW 2013/0476/0018 (Figs 20-21), height 22.3 mm; paratype 1, NHMW 2013/0476/0017 (Figs 18-19), height 17.5 mm; paratype 2, 2013/0476/0019, height 22.4 mm; paratype 3, NHMW 2013/0476/0021, height 12.1 mm.

**Other material.** NHMW 2013/0476/0020 (1), height 14.0 mm, juvenile (Figs 22-23).

**Etymology.** Named after Dr. Walter Etter of the Naturhistorisches Museum Basel, Switzerland, in recognition of his help and support.

**Type locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela (= locality GS1PGNA of Gibson-Smith & Gibson-Smith, 1979).

**Type stratum.** Cantaure Formation (late early Miocene: Burdigalian).

**Diagnosis.** A medium sized *Ocenebra* species with a strongly nodulose shoulder, three varices per whorl on the last three whorls, with one intervarical node, spiral cords P1-P5 equal in strength, P6 slightly narrower, with a single intercalated secondary cord, aperture small with very shallow ID and strong D1-D5 within and a short, sealed siphonal canal.

**Description.** Shell medium sized, heavy, nodose. Subsutural ramp broad, weakly sloping, slightly concave. Spire high with teleoconch up to five or six strongly shouldered, nodose whorls. Suture adpressed. Protoconch unknown. Axial sculpture of teleoconch whorls consisting of high ribs and high, broad, varices. Each varix with short, blunt nodes extending from primary spiral cords. Antepenultimate, penultimate and last whorls with three varices and a single, strong, prominent, intervarical node. Other spire whorls with narrow axial ribs. Last whorl broad. Subsutural area with adis, IP, abis, followed on convex part of whorl by P1, s1, P2, s2, P3, s3, P4, s4, P5, s5, P6, s6; ADP and MP on siphonal canal. P1-P6 almost similar in size and strength, P6 weakly narrower. Secondary cords obviously narrower than primary cords. Siphonal canal below MP missing or damaged. moderately Aperture small, ovate. Columellar lip narrow, smooth, adherent. Anal notch shallow, broad. Outer apertural lip with very shallow ID and strong D1-D5 within. Labral varix rounded, strong, broad, with small nodes. Siphonal canal short, narrow, ventrally sealed, partly broken.

**Discussion.** Vermeij & Vokes (1997, p. 72) restrict the genus *Ocenebra* to "a relatively small number of Miocene to Recent species from western Europe, the Mediterranean region and Tropical West Africa. These are characterized by the tendency to form three varices on the last whorl, by the presence of six to eight primary spiral cords on the last whorl, a crenulated outer lip without a labral tooth, an adherent or very lightly erect inner lip, and six weak to strong denticles on the inner side of the outer lip. In species with varices, the latter are separated from each other by a single intervarical node".

The Tropical West African species are now classified in other genera. However, Houart & Sirenko (2003) also assigned three Recent and fossil species from the northwestern Pacific to *Oceuebra*.

Two more or less related species were assigned to *Miocenebra: M. silverdalensis* (Vokes, 1963) and to *Fenolignum: F. umbilicatum* Vermeij & Vokes, 1997, by Vermeij & Vokes (1997), both species from the Belgrade Formation, North Carolina.

This is the first occurrence of the genus *Ocenebra* in the tropical western Atlantic and therefore this new species cannot be compared to any Caribbean fossil or Recent taxon. *Ocenebra etteri* nov. sp. is far more reminiscent of some of the Early Miocene Aquitanian and Burdigalian species from the French Aquitaine Basin (see Lozouet *et al.*, 2001, pl 22, 23). Genus Vitularia Swainson, 1840

Type species. – *Murex miliaris* Gmelin, 1791, by monotypy. Recent, western Indo-Pacific.

Note. – The placement of this genus in the Ocenebrinae Cossmann, 1903 is provisional, as molecular data presented by Barco *et al.* (2010) showed the subfamily to be polyphyletic.

Vitularia sp. cf. V. salebrosa (King & Broderip, 1832) Figures 24-27

cf. 1832 Murex salebrosa King & Broderip, p. 347.

Material. 1 incomplete specimen, NHMW 2012/0197/0008 (ex BL coll.), height 29.1 mm (Figs 24-25); 1 incomplete specimen, NMB H20169, NMB locality 12842, height 72.2 mm (Figs 26-27).

**Locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela (= locality GS1PGNA of Gibson-Smith & Gibson-Smith, 1979), Cantaure Formation (early Miocene: Burdigalian).

**Discussion.** A *Vitularia* species occurs in the Cantaure formation, which is difficult to characterise as only two incomplete specimens are available to us, and they are somewhat different from each other. The first of these is smaller and probably represents a subadult stage (Figs 24-25). The shell is slender and the spire high. The carina on the spire whorls is sharp and runs mid-whorl. The subsutural ramp is broad and weakly concave, below the carina the whorl is straight-sided and tapers inwards to the suture. The axial sculpture consists of eight prosocline ribs developed into prominent rounded tubercles at the shoulder. The ribs cross the subsutural ramp as narrow lamellae. The surface is pustulose, a feature of the genus. Unfortunately the adapical portion of the last whorl and aperture are missing. This is the specimen discussed by Landau et al. (2013). The second specimen (Figs 26-27) is larger, but strongly abraded and incomplete. The spire is also high, and the shoulder bears rounded tubercles. However, it differs in the position of the carina on the spire whorls, which is closer to the adapical suture, giving the spire a conical rather than scalate shape. The two Vitularia specimens from Cantaure probably represent a single species, as these differences in the position of the suture and spire shape can also be seen in the European Miocene Vitularia linguabovis (Basterot, 1825) (Landau et al., 2013).

The genus is not speciose. In the Caribbean faunas only two species are known. *Vitularia dominicana* Vokes, 1989 from the early Pliocene Gurabo Formation of the Dominican Republic is low spired, and according the illustrated holotype, has large round

#### B. LANDAU & R. HOUART

tubercles at the shoulder, which is placed high on the last whorl. At the time of description only three specimens were available to Emily Vokes, with a maximum height of 28.2 mm. Subsequent collections made by the senior author have brought further specimens to light, which attain a greater maximum height of 53.9 mm. As the shell increases in size the penultimate and last whorls become rounded and lose the tubercles at the shoulder. In the largest specimens from the Dominican Republic the shoulder of the last whorl is rounded and quite smooth. In view of these important and, so far undocumented ontogenic changes, we take the opportunity to illustrate further specimens from the Dominican Republic (Figures 2-7).



**Figures 2-7.** *Vitularia dominicana* Vokes, 1989, Tulane locality 1215 (see Vokes, 1989), Gurabo River, Gurabo Formation, Lower Pliocene, Dominican Republic. Figs 2-3 NHMW 2013/0476/0005, height 40.8 mm; figs 4-5 NHMW 2013/0476/0006, height 53.7 mm; figs 6-7 NHMW 2013/0476/0007, height 49.5 mm.

Vokes (1989) compared the Dominican shell to the Recent Indo-Pacific species *Vitularia miliaris* (Gmelin, 1791), but large specimens of this species always have large tubercles at the shoulder, which remains angular and placed high on the last whorl.

The second species is Vitularia salebrosa (King & Broderip, 1832) (= V. ecuadoriana Marks, 1951 = V.linguabison Vokes, 1967). This species occurs in the Neogene in the late Pliocene Caribbean Agueguexquite Formation of Mexico and Pinecrest Formation of Florida, in the Panamic faunas from the early Pliocene Esmeraldas Group of Ecuador, through the Pleistocene of Baja California, Ecuador and the Galapagos, and is found today from Baja California to Peru. It is one of the paciphile species discussed by Landau et al. (2009). Vitularia salebrosa is tall-spired, like the Cantaure specimens, but seems to have smaller and more numerous tubercles at the shoulder. Vitularia linguabovis (Basterot, 1825), which occurs in the early-late European Miocene, is broader, usually with a less elevated spire, and more numerous tubercles at the shoulder. We wait in the hope of finding better specimens from Cantaure, which is the earliest occurrence of the genus in the tropical American Neogene assemblages.

Subfamily **TYPHINAE** Cossmann, 1903 Genus *Typhina* Jousseaume, 1880

Type species. – *Typhis belcheri* Broderip, 1833 (= *Murex cleryi* Petit, 1840) by original designation, Recent, West Africa and Brazil. = *Talityphis* Jousseaume, 1882

*Talityphis* Jousseaume, 1882 and *Typhina* Jousseaume, 1880 were considered congeneric by Houart (2002).

*Typhina canaliculata* nov. sp. Figures 8, 28-34

**Dimensions and type material.** Holotype; NHMW 2013/0476/0001 (ex BL coll.) (Figs 28-31), height 28.6 mm; paratype 1; NHMW 2013/0476/0002 (ex BL coll.) (Figs 8, 32-34), height 23.8 mm; paratype 2; NHMW 2013/0476/0003 (ex BL coll.), height 28.9 mm. paratype 3 NMB H20171 (NMB locality 17520), height 19.9 mm; paratype 4 NMB H20170 (NMB locality 17516), height 27.9 mm; paratype 5 NMB H19068 (NMB locality 17516), height 27.7 mm.

**Other material.** Maximum height 30.3 mm; 10 specimens NHMW 2013/0476/0004 (ex BL coll.), NMB locality 17516 NMB coll. (6), same locality.

**Etymology.** Name reflecting the deeply canaliculated suture seen in this species.

**Type locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela (= locality GS1PGNA of Gibson-Smith & Gibson-Smith, 1979).

**Type stratum.** Cantaure Formation (late early Miocene: Burdigalian).

**Diagnosis.** A medium-sized *Typlina* species, with a relatively high spire, strongly concave subsutural area resulting in a deeply canaliculated suture, four varices per whorl, short tubes, five weakly developed spiral cords forming weakly recurved spines on the varices, and a moderately expanded labral varix.

Description. Shell medium-sized, stout; spire

relatively high. Protoconch not preserved. Six or seven sharply-angular, straight-sided teleoconch whorls, with narrow, deeply concave subsutural area. Suture impressed, deeply canaliculated. Four varices per whorl. Tubes (P1) of moderate length, pointing laterally and abaperturally, placed closer to preceding than succeeding varices. A laminar partition above the aperture crossing the shoulder joins the varix to the varix of the preceding whorl. Below tube shell surface very weakly swollen and two lines pass anteriorly, one being the margin of the old mouth, the other, in advance of the tube, being similar in character and indicating another arrest in growth. P2-P6 weakly developed, forming small, posteriorly-reflected spines over the varices. Surface bearing faint growth lines. Aperture small, ovate, bordered by raised rim; labral varix constant in width, moderately expanded, consisting of inner band bearing spiral cords and outer flange with a scalloped edge formed by spines at termination of P2-P6. Siphonal canal closed, broad, short, pointing to the right and relatively abaperturally.



**Figure 8.** *Typhina canaliculata* nov. sp., paratype 1 NHMW 2013/0476/0002 (ex BL coll.), height 23.8 mm, 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela, Cantaure Formation, Burdigalian, late early Miocene.

**Discussion.** The four varices and tubes per whorl, the tubes being placed closer to the preceding than succeeding varix, the apertural varix greatly expanded and the partition above the aperture place this new species in the genus *Typluina* Jousscaume, 1880.

*Typhina canaliculata* nov. sp. resembles the Recent and type species *T. belcheri* (= *T. cleryi*) but it differs in having a comparatively larger shell with a higher spire, a straighter, less upward bent and shorter P2 spine instead of long and strongly adapically recurved in *T. belcheri*, a more sculptured, more strongly folded apertural varix and a comparatively broader siphonal canal.

*Typhina canaliculata* nov. sp. is also closely similar to *Typhina obesa* (Gabb, 1873), which is widespread in the Caribbean early Miocene (Vokes, 1989), but differs in being less broad and in having a taller spire, whereas *T. obesa* has a very short, depressed spire. The subsutural area in *T. obesa* is broad and weakly concave, whereas in *T. canaliculata* it is narrower and

strongly concave, resulting in a deeply canaliculated suture, bordered by a high-ridged shoulder. In Typhina canaliculata the spiral cords are more strongly developed than in T. obesa, forming short recurved spines at the varices on the last whorl. In T. obesa the spiral sculpture is very weakly developed, hardly marked over the sharp, straight varices. Lastly, the aperture is relatively smaller and the labral varix narrower than in T. obesa. Typhina expansa (Sowerby, 1874) from the early Pliocene to Recent Caribbean, is very closely similar to T. obesa (differences between the two were discussed by Vokes, 1989, p. 332), but this species can also be distinguished from T. canaliculata in having a broader shell shape and in having a wider apertural varix, but above all by the less concave subsutural area. Typhina alata (Sowerby, 1850) from the early Pliocene Gurabo Formation of the Dominican Republic and Pliocene formations of Ecuador differs in having a more fusiform shell shape, a less depressed subsutural area, a wider labral varix and in forming an intervarical node below the tubes, a character not seen in T. obesa, T. expansa or T. canaliculata.

The earliest member of the genus in the Caribbean was thought to be *Typhina precursor* (Keen & Campbell, 1964) from Las Perdices Shale of Colombia, which, according to Gertman (1969), was dated as Aquitanian early Miocene. This species also occurs in the Manzanilla Formation of Trinidad, dated by Gertman as early Miocene. The age of the Las Pedices Group is now considered middle-late Miocene (Landau *et al.*, 2012, chart 2), and the Manzanilla Formation is considered to be late Miocene (Landau *et al.*, 2012, chart 15). *Typhina precursor* is a very large solid species, 47-49 mm in height, which also seems to have a narrow shoulder, but differs importantly from *T. canaliculata* and other *Typhina* species in having no spiral sculpture developed.

Other Caribbean congeners are: *Typhina pterina* (Gardner, 1947) from the middle Miocene Shoal River Formation of Florida and *T. eucteana* (Woodring, 1970) from the late Miocene Gatun formation of Panama, which are both much smaller, with a more fusiform shell shape. The subsutural area in both of

these is concave, but not as deep as in *T. canaliculata*. Typhina eucteana is further distinguished by its long, narrow siphonal canal. Typhina siphon (Woodring, 1928) from the Pliocene Bowden Formation of Jamaica is a more slender, fusiform species, with recurved spines at the shoulder, and a longer siphonal canal. Typhina acuticosta (Conrad, 1830) from the middle Miocene St. Mary Formation of Maryland and late Miocene Choctawhatchee Formation of Florida is quite different, with a slender shell shape and acutely angled, straight-sided whorls. Typhina carmenae (Gertman, 1969) Pliocene from the late Agueguexquite Formation of Mexico has a broader shell shape, more prominent spiral sculpture and a far more expanded apertural varix than T. canaliculata. The Recent Panamic Typhina latipennis (Dall, 1919) has weaker spiral sculpture than the fossil Cantaure species, and a greatly expanded flange on the apertural varix.

#### Genus Laevityphis Cossmann, 1903

Type species. – *Typhis coronarins* Deshayes, 1865 [*Laevityphis muticus* (Sowerby, 1835)], by original designation. Eocene, France.

# *Laevityphis jungi* nov. sp. Figures 35-38

*Typhis (Laevityphis) sawkinsi.* Jung, 1965, p. 525, pl. 70, figs 7, 8 [not *Laevityphis sawkinsi* (Mansfield 1925)].

**Dimensions and type material.** Holotype; NHMW 2013/0476/0012 (ex BL coll.) (Figs 35-36), height 28.3 mm; paratype 1; NHMW 2013/0476/0013 (ex BL coll.) (Figs 37-38), height 27.0 mm; paratype 2; NHMW 2013/0476/0014 (ex BL coll.), height 27.1 mm; paratype 3; NHMW 2013/0476/0015 (ex BL coll.), height 23.8 mm; paratype 4 NMB H18542, height 23.0 mm; paratype 5 NMB H18543, height 20.7 mm; paratype 6 NMB H13719, height 23.0 mm [specimen figured by Jung, 1965, pl. 70, figs 7, 8 as *Typhis (Laevityphis) sawkinsi* Mansfield].

### Figures 9-23

**12-13.** *Phyllonotus* sp. cf. *P. infrequens* (Vokes, 1963), NHMW 2013/0476/0011 (ex BL coll.), height 35.3 mm. **14-17.** *Purpurellus* sp. cf. *P. repetiti* (Vokes, 1970)

14-15. NHMW 2013/0476/0008 (ex BL coll.), height 27.2 mm; 16-17. NHMW 2013/0476/0009 (ex BL coll.), height 32.4 mm.

18-23. Ocenebra etteri nov. sp.

**18-19.** Paratype 1, NHMW 2013/0476/0017, height 17.5 mm; **20-21.** Holotype, NHMW 2013/0476/0018, height 22.3 mm; **22-23.** Juvenile, NHMW 2013/0476/0020, height 14.0 mm.

All: 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela, Cantaure Formation, Burdigalian, late early Miocene.

<sup>9-11.</sup> Siratus harzhauseri nov. sp., holotype NHMW 2013/0476/0010 (ex BL coll.), height 48.6 mm.



**Other material.** Maximum height 31.7 mm; 20 specimens NHMW 2013/0476/0016 (ex BL coll.), NMB locality 17516 NMB coll. (66), NMB locality 17520 NMB (100+), same locality.

**Etymology.** Named in honour or Peter Jung, in recognition of his work on the Cantaure assemblage.

**Type locality.** 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela (= locality GS1PGNA of Gibson-Smith & Gibson-Smith, 1979).

**Type stratum.** Cantaure Formation (late early Miocene: Burdigalian).

**Diagnosis.** A medium-sized *Laevityplus* species, with a high spire, a narrow, concave subsutural area, four varices per whorl placed in the middle of the intervarical area, bearing spines pointing adapically and inwards, short tubes in the intervarical spaces, an almost smooth surface, a small aperture, a narrowly expanded labral varix, and a long, narrow siphonal canal.

Description. Shell medium-sized, elongate; spire high. Protoconch paucispiral, consisting of two elevated, bulbous whorls. Six or seven sharply-angular teleoconch whorls, with narrow, concave, subhorizontal subsutural area, weakly convex below shoulder to suture. Suture impressed, undulating. Four simple, convex varices per whorl, with a large, sharp spine at the shoulder of each varix pointing adapically and inwards; end usually broken away and appearing as small tube-like structures, but not connecting to interior of shell. Tubes (P1) of moderate length, separated from varices, pointing laterally and abaperturally, placed in the middle of the intervarical area. A ridge running abapically from the tube dividing the intervarical area into two almost equal halves. Last whorl relatively short, convex, strongly constricted at the base. P2-P6 not clearly developed, very weak spiral sculpture just visible on penultimate and last whorls. Aperture small, ovate, bordered by raised rim; labral varix constant in width, moderately expanded. Siphonal canal closed, narrow, long, pointing to the right and abaperturally at distal end.

**Discussion.** The four varices and tubes per whorl, the tubes not being attached to the varix, the varix bearing a spine pointing apically and the narrowly expanded apertural varix place this new species in the genus *Laevityphis* Cossmann, 1903 (Gertman, 1969). In this genus the tube is usually placed closer to the succeeding than the preceding varix, although in *Laevityphis jungi* nov. sp. the tube is placed almost midway between the varices.

This species was first identified by Jung (1965) as *Typhis (Laevityphis) sawkinsi* (Mansfield, 1925) from the early-middle Miocene Brasso Formation of Trinidad, although the author noted that the Cantaure shells were not identical with those from Trinidad. When compared to the holotype figured by Gertman (1969, pl. 7, fig. 1), the Trinidadian species is much stockier, with a far shorter spire, the varices are far wider and more prominent than in *L. jungi*, the base is less strongly constricted and the siphonal canal is broader and shorter than in the Venezuelan species.

The fact that the Venezuelan and Trinidadian species were not conspecific was noted by Vokes (1989, p. 81), who considered the Cantaure shells to be a rather large variant of Laevityphis linguiferus (Dall, 1890) a species common in the early Miocene Chipola Formation of Florida. Indeed, Vokes (1989) considered the two identical, apart from their size. We have compared the Cantaure material with numerous specimens of L. linguiferus from the Chipola Formation and cannot agree with this conclusion. Laevityphis jungi is indeed a much larger shelled species, adults ranging from 22.7 mm - 31.7 mm in height, whereas the largest specimens of L. linguiferus do not attain 15 mm in height. Further differences can be seen in the spire, which is much taller in the Venezuelan species and the last whorl, which is more

# Figures 24-38

24-27. *Vitularia* sp. cf. *V. salebrosa* (King & Broderip, 1832)
24-25. NHMW 2012/0197/0008 (ex BL coll.), height 29.1 mm; 26-27. NMB H12842, height 72.2 mm.
28-34. *Typhina canaliculata* nov. sp.
28-31. Holotype; NHMW 2013/0476/0001 (cx BL coll.), height 28.6 mm; 32-34. Paratype 2; NHMW 2013/0476/0003 (ex BL coll.), height 28.9 mm.

**35-38.** *Laevityphis jungi* nov. sp.

**35-36.** Holotype; NHMW 2013/0476/0012 (ex BL coll.), height 28.3 mm; **37-38.** Paratype 1; NHMW 2013/0476/0013, (ex BL coll.), height 27.0 mm.

All: 1 km southwest of Casa Cantaure, about 10 km west of Pueblo Nuevo, Falcón, Venezuela, Cantaure Formation, Burdigalian, late early Miocene.



constricted at the base. The proximal end of the siphonal canal is proportionally broader in *L. linguiferus* than in *L. jungi*, but then narrows rapidly at the distal end to form a very small siphonal exit, whereas in *L. jungi* the narrowing in gradual and the siphonal exit proportionally wider. The sculpture, or lack of it, is similar in both species.

Several congeners occur in the tropical American fossil record. The oldest member of the group, *L. gracilis* (Conrad, 1833) from the late middle Eocene Gosport Formation of Alabama, USA also has a slender shell and tall spire, but is distinguished by its crenulated varices. The other Miocene and Pliocene congeners, such as *L. costaricensis* (Olsson, 1922), *L. bullisi* (German, 1969), *L. apheles* (Vokes, 1989) and *L. spinirectus* (Vokes, 1989) all differ in having smaller, less slender shells with lower spires.

## ACKNOWLEDGEMENTS

We would like to thank Emily Vokes for her comments and for acting as referee. Also to Walter Etter and Olivier Schmidt of the Naturhistorisches Museum Basel, Switzerland, for access to the PPP collection.

# REFERENCES

Barco, A., Claremont, M., Reid, D.G., Houart, R.,
Bouchet, P., Williams, S.T., Cruaud, C., Couloux,
A., Oliverio, M. 2010. A molecular phylogenetic
framework for the Muricidae, a diverse family of
carnivorous gastropods. *Molecular Phylogenetics* and Evolution 56: 1025-1039.

Bouchet, P. & Houart, R. 2013. World Register of Marine Species. <u>http://www.marinespecies.org/aphia.php?p=taxdet</u> <u>ails&id=225362</u>. Accessed on 19 November 2013.

Emerson, W.K., D'Attilio, A. 1969. Remarks on the taxonomic placement of *Purpurellus* Jousseaume, 1880, with the description of a new species. *The Veliger* 12: 145-148.

Gertman, R.L. 1969. Cenozoic Typhinae (Mollusca: Gastropoda) of the western Atlantic region. *Tulane Studies in Geology* 7: 143-191.

Gibson-Smith, J. & Gibson-Smith, W. (1979). The genus *Arcinella* (Mollusca: Bivalvia) in Venezuela and some associated faunas. *Geos* 24: 11-32.

Gibson, T.G. 1977. *In* Geologic and hydrologic principles, processes, and techniques; Geologic Survey Research 1977. *United States Geological Survey Professional paper* 1050: 153-221.

Gibson, T.G. 1983. Stratigraphy of Miocene through lower Pleistocene strata of the United States central Atlantic Coastal Plain. *In* Ray, C:E., ed., Geology and paleontology of the Lee Creek mine, North Carolina, I. *Swithsonian Cotributions to Paleobiology* 53: 35-80.

Houart, R. 2002. Description of a new typhine (Gastropoda: Muricidae) from New Caledonia with comments on some generic classifications within the subfamily. *Venus* 61 (3-4): 147-159.

- Houart, R. & Sirenko, B. 2003. Review of the Recent species of *Ocenebra* Gray, 1847 and *Ocinebrellus* Jousseaume, 1880 in the Northwestern Pacific. *Ruthenica* 13 (1): 53-74.
- Jung, P. 1965. Miocene Mollusca from the Paraguana Peninsula, Venezuela. *Bulletins of American Paleontology* 49 (223): 387-644.
- King, P.P. and Broderip, W.J. 1832. Description of the Cirrhipeda, Conchifera and Mollusca, in a collection formed by the Officers of H.M.S.
  Adventure and Beagle employed between the years 1826 and 1830 in surveying the Southern Coasts of South America, including the Straits of Magalhaens and the Coast of Tierra del Fuego. *Zoological Journal* 5: 332-349.

Landau, B.M., Harzhauser M., İslamoğlu, Y., Silva, C.M. da 2013. Systematics and palaeobiogeography of the gastropods of the middle Miocene (Serravallian) Karaman Basin, Turkey. *Cainozoic Research* 11-13: 3-584.

Landau, B.M., Houart, R., Silva, C.M. da 2007. The early Pliocene Gastropoda (Mollusca) of Estepona, southern Spain, 7. Muricidae. *Palaeontos* 11: 1-87.

- Landau, B., Petit, R.E., Etter, W., Silva, C.M. da 2012. New species and records of Cancellariinae (Caenogastropoda) from tropical America, together with a catalogue of Neogene to Recent species from this region. *Cainozoic Research* 9: 193-279.
- Landau B.M. & Vermeij, G.J. 2010. A new species of *Plicopurpura* (Mollusca: Rapaninae) from the Lower Miocene Cantaure Formation of Venezuela. *Novapex* 11: 99-106.
- Landau B.M., Vermeij, G.J., Silva, C.M. 2009. Pacific elements in the Caribbean Neogene gastropod fauna: the source-sink model, larval development, disappearance, and faunal units. *Bulletin de la Société Géologique de France* 180(4): 249-258.

Lozouet, P. 1986. Les Gastéropodes prosobranches de l'Oligocène supérieur du Bassin de l'Adour (systématique, paléoenvironnements, paléoclimatologie, paléobiogéographie). Diplôme de l'E.P.H.E., 475pp.

Lozouet, P., Lesport, J.F., & Renard, P. 2001. Révision des Gastropoda (Mollusca) du stratotype de l'Aquitanien (Miocène inf.): site de Saucats 'Lariey', Gironde, France. *Cossuanuiana* (hors série 3): 1-189.

- Merle, D. 1999. La radiation des Muricidae (Gastropoda : Neogastropoda) au Paléogène: approche phylogénétique et évolutive. Paris. Unpublished thesis, Muséum national d'Histoire naturellc : i-vi, 499 pp.
- Merle, D. 2001. The spiral cords and the internal denticles of the outer lip in the Muricidae: terminology and methodological comments. *Novapex*, **2** (**3**), 69-71.

- Merle, D., Garrigues, B., Pointier, J.P. 2011. Fossil and Recent Muricidae of the World. Part Muricinae. Hackenheim (ConchBooks): 648 pp.
- Swainson, W. 1820-33. The Zoological illustrations, or original figures and descriptions of new, rare, or interesting animals, selected chiefly from the classes of ornithology, entomology, and conchology. London, ser. 1, vols 1-3, pls 1-182 (1820-23); ser. 2, vols 1-3, pls 1-136 (1829-33).
- Vermeij G.J. 2001. Distribution, history, and taxonomy of the *Thais* clade (Gastropoda: Muricidae) in the Neogene of tropical America. *Journal of Paleontology* 75: 697-705.
- Vermeij G.J. & Vokes E.H. 1997. Cenozoic Muricidae of the western Atlantic region. Part 12. The subfamily Ocenebrinae (in part). *Tulane Studies in Geology and Paleontology* 29: 69-118.
- Vokes, E.H. 1963. Cenozoic Muricidae of the western Atlantic region. Part 1 – *Murex sensu stricto*. *Tulane Studies in Geology* 1: 93-123.
- Vokes, E.H. 1970. Cenozoic Muricidae of the western Atlantic region. Part V. *Pterynotns* and *Poirieria*. *Tnlane Studies in Geology* 8: 1-50.

- Vokes, E.H. 1989. Neogene paleontology in the northern Dominican Republic, 8. The family Muricidae (Mollusca: Gastropoda). *Bulletins of American Paleoutology* 97 (332): 5-94.
- Vokes, E.H. 1990. Cenozoic Muricidae of the Western Atlantic region. Part VIII – *Murex s.s.*, *Haustellum*, *Chicoreus*, and *Hexaplex*; additions and corrections. *Tulane Studies in Geology and Paleontology* 23: 1-96.
- Vokes E.H. 1992. Cenozoic Muricidae of the western Atlantic region. Part 9. *Pterynotns, Poirieria, Aspella, Dermonmrex, Calotrophon, Acantholabia,* and *Attiliosa*; additions and corrections. *Tulane Studies in Geology and Paleontology* 25: 1-108.
- Vokes E.H. 1994. Cenozoic Muricidae of the western Atlantic region. Part 10. The subfamily Muricopsinae. *Tulane Studies in Geology and Paleontology* 26: 49-160.
- Vokes E.H. 1995. Two new Cenozoic Muricinae (Gastropoda: Muricidae) of the western Atlantic region. *Thlane Studies in Geology and Paleontology* 28: 119-122.