

## **Alien species on the coasts of Turkey**

**M. E. ÇINAR<sup>1</sup>, M. BILECENOĞLU<sup>2</sup>, B. ÖZTÜRK<sup>1</sup>, T. KATAGAN<sup>1</sup> and V. AYSEL<sup>3</sup>**

<sup>1</sup>Ege University, Faculty of Fisheries, Department of Hydrobiology, 35100 Bornova, Izmir, Turkey

<sup>2</sup>Adnan Menderes University, Faculty of Arts & Sciences, Department of Biology, 09010 Aydin, Turkey

<sup>3</sup>Onsekiz Mart University, Faculty of Arts & Sciences, Department of Biology, Çanakkale, Turkey

e-mail: melih.cinar@ege.edu.tr

---

### **Abstract**

*The compilation of data on alien species reported from the Turkish coasts yielded a total of 263 species belonging to 11 systematic groups, of which Mollusca had the highest number of species (85 species), followed by Crustacea (51), fishes (43) and phytobenthos (39). The Black Sea is represented by a total of 20 alien species, the Sea of Marmara by 48 species, the Aegean Sea by 98 species and the Levantine Sea by 202 species. The majority of aliens found in the Black Sea and the Sea of Marmara were transported via shipping, whereas the Levantine coast is extensively subjected to Lessepsian migration. Benthic habitats (soft and hard substrata) comprise 76% of the total alien species and the pelagic environment is inhabited by thirty-nine species. Almost 50% of aliens collected from the Turkish coasts were found only at 0-10 m depth. Eight species occur at depths deeper than 100 m. The impacts of aliens on the benthic and pelagic ecosystems are presented.*

**Keywords:** Alien species; Species list; Impact; Black Sea; Sea of Marmara; Aegean Sea; Levantine Sea; Turkey.

---

### **Introduction**

Species introduction is one of the major factors adversely affecting biological diversity (ELTON, 1958). The impacts of alien species on their new environment include restructuring established food webs, importing new diseases and competition with native organisms for food and space. Other significant ecological changes may occur when the invading organisms reproduce with native species, altering the gene pool (OCCHIPINTI AMBROGI, 2001). Invaders may belong to any taxonomical group, and are

capable of colonizing every ecosystem on earth, changing the ecological relations within communities, altering evolutionary processes and causing dramatic changes in native populations, including extinctions (MACK *et al.*, 2000).

The negative effects of aliens experienced world-wide have triggered the social and scientific media to take some precautions against the dispersal of aliens among regions or among localities within a particular region. Thus, compiled data regarding all aliens or a target organism (i.e. *Caulerpa taxifolia* in the Mediterranean Sea) are urgently required by scientists

and decision makers. In addition, monitoring programmes on spatio-temporal structures of communities particularly in the hot spot areas for aliens such as harbours, brackish and polluted waters should be undertaken both on local and global scales.

Turkey is surrounded by four seas with different hydrographical characteristics. The maritime traffic taking place through the Dardanel and Bosphorus straits and among commercial harbours makes the Turkish coasts more susceptible to invasions by aliens. The proximity of Turkey to the Suez Canal has resulted in dense settlements of Lessepsian migrants, especially in habitats along the Levantine coast of Turkey. Alien species and their roles in the benthic and pelagic ecosystems is increasingly becoming a subject of study in the country.

This paper reviews the alien species reported from the Turkish coasts and constitutes the first comprehensive database for future studies.

## Methods

### *Data on alien species along the Turkish coasts*

In this study, only reported data on aliens along the Turkish coasts have been taken into account. All calculations are based on species' records up to the end of 2005. Cryptogenic species have not been included in our compiled list. Some of the species in Table 1, which represent uncertainty in their real taxonomic position and distributional pattern, are classified as questionable or excluded. The questionable species will turn into aliens if their real taxonomic and distributional identities are clarified. In all calculations, including number of species per sea, only the established and questionable species have been taken into account. The importance of aliens in the total biota was estimated, based on the authors' databases.

## Results and Discussion

### *List of alien species from the Turkish coasts*

Table 1 includes all alien species reported from the Turkish coasts. The species are categorized by their origins, the mode of introduction and their habitat and depth preferences. The establishment success is assigned for each species. The first reported years of the species together with the relevant publications are also given for each sea.

The present data show that a total of 263 alien species, which belong to 11 systematic groups, occur along the Turkish coasts. Among the groups, Mollusca had the highest number of species (85 species), followed by Crustacea (51), fishes (43) and phytobenthos (39). Cnidaria is represented by only two species (*Rhopilema nomadica* and *Cassiopea andromeda*), Ctenophora by 2 species (*Mnemiopsis leidyi* and *Beroe ovata*), Pantopoda by one species (*Anoplodactylus californicus*), Bryozoa by one species (*Rhynchozoon larreyi*) and Echinodermata by 2 species (*Ophiactis savignyi* and *Synaptula reciprocans*).

Thirty-two species are considered as questionable or excluded in Table 1, mainly due to the uncertainty in their real taxonomical status and distributional patterns. The polychaetes, *Branchiosyllis exilis*, *Opisthosyllis brunnea*, *Rhodine loveni* and *Monticellina dorsobranchialis*, which were previously considered as Lessepsian migrants, indeed occur also in the western Mediterranean and the Atlantic Ocean. These species were also previously proposed to be excluded from the list of Lessepsian species (ERGEN *et al.*, 2002; ÇINAR, 2003; 2005). RULLIER (1963) found 11 polychaete species with Indo-Pacific affinity from the Sea of Marmara (Table 1). Of the species, *Nereis zonata persica* and *Timarete ancylochaeta* were also reported along the Levantine coast and considered as Lessepsian migrants (BEN-ELIAHU, 1995). The other species have not been subsequently reported from elsewhere in the Mediterranean Sea.

A total of 176 Lessepsian and 48 ship-transferred species were reported along the Turkish

Table 1.

The list of alien species and their first year of reports from the Turkish coasts. The habitat and depth preferences of aliens along the coasts together with their origins and establishment success are also given. The marked species in the list are questionable or excluded ones. BS: Black Sea, SM: Sea of Marmara, AS: Aegean Sea, LS: Levantine Sea, O: Origin (IP: Indopacific, RS: Red Sea, AT: Atlantic, WA: Western Atlantic, ST: Subtropical, IO: Indian Ocean, PG: Persian Gulf, PO: Pacific Ocean, TA: Tropical Atlantic, CT: Circumtropical, BA: Boreal Atlantic) Es: Establishment success (E: Established, Q: Questionable, Ex: Excluded), MI= Mode of Introduction (L: Lessepsian, S: Shipping, G: Gibraltar, Aq: Aquaculture), H: Habitat [Hs: hard substrata (including algae, sponge), Ss: soft substrata (including phanerogams), P: pelagic, Pz: parasite], D: Depth range (I: 0-10 m, II: 11-50m, III: 51-100 m, IV: 101-200 m, V: 201-400, VI: 401-500 m).

	BS	SM	AS	LS	Es	O	MI	H	D
<b>PHYTOPLANKTON</b>									
<i>Alexandrium tamarens</i> (Lebour) Balech 1992	-	-	1984 <sup>124</sup>	-	Q	WA	S	P	I
<i>Chaetoceros coarctatus</i> Lauder 1864	-	-	1987 <sup>125</sup>	-	E	AT,PO	S	P	I
<i>Gymnodinium</i> cf. <i>mikimotoi</i> Miyake et Kominami ex Oda 1935	-	-	2002 <sup>44</sup>	-	Q	?	S	P	I
<i>Heterosigma</i> cf. <i>akashiwo</i> (Hada) Hada, 1968	-	-	2002 <sup>44</sup>	-	Q	?	S	P	I
<i>Rhizosolenia calcar-avis</i> M. Schultze, 1858	1999 <sup>71</sup>	1993 <sup>123</sup>	-	-	E	AT	S	P	I
<i>Scrippsiella trochoidea</i> (Stein) Loeblich III, 1976	1999 <sup>71</sup>	-	-	-	E	AT	S	P	I
<i>Thalassiosira nordenskiöldii</i> Cleve 1873	2002 <sup>160</sup>	-	-	-	E	AT	S	P	I
<b>PHYTOBENTHOS</b>									
<b>Cyanophyta (=Cyanobacteria)</b>									
<i>Trichodesmium erythraeum</i> Ehrenberg, 1830	-	-	1996 <sup>69</sup>	-	E	IP,RS	L	Hs,Ss	I
<b>Rhodophyta</b>									
<i>Acanthophora nayadiformis</i> (Delile) Papenfuss, 1968	-	1973 <sup>177</sup>	1970 <sup>103</sup>	1997 <sup>20</sup>	E	RS	L	Hs	I,II
<i>Acanthophora muscoides</i> Linnaeus, 1753	-	-	1986 <sup>178</sup>	-	E	AT	S	Hs	I,II
<i>Acrochaetium codicolum</i> Børgesen, 1927	1996 <sup>28</sup>	1986 <sup>178</sup>	1990 <sup>55</sup>	1997 <sup>19</sup>	E	IP,AT	S	Hs,Ss	I
<i>Asparagopsis armata</i> Harvey, 1855	1973 <sup>176</sup>	1986 <sup>178</sup>	1973 <sup>176</sup>	1969 <sup>174</sup>	E	IP,AT	S	Hs	I
<i>Asparagopsis taxiformis</i> (Delile) Trevisan de Saint-Léon, 1845	-	-	2001 <sup>27</sup>	-	E	CT	L	Hs	I
<i>Bonnemaisonia hamifera</i> Hariot, 1891	-	1986 <sup>178</sup>	-	1997 <sup>20</sup>	E	PO,AT	?S	Hs	I
<i>Botryocladia madagascariensis</i> G.Feldmann, 1945	-	-	-	2000 <sup>161</sup>	E	IO	?S	Hs	I
<i>Chondria collinsiana</i> Howe, 1920	-	1986 <sup>178</sup>	-	-	E	AT, IO	?L	Hs	II
<i>Chondrophycus papillosus</i> (C. Agardh) Garbary & Harper 1998	1973 <sup>175</sup>	1957 <sup>143</sup>	1969 <sup>174</sup>	1969 <sup>174</sup>	E	RS	?L	Hs	I
<i>Ganonema farinosum</i> (Lamouroux) Fan & Wang, 1974	1995 <sup>21</sup>	1899 <sup>91</sup>	1986 <sup>26</sup>	1997 <sup>19</sup>	Q	IP,RS	?L	Hs,Ss	I

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Gracilaria arcuata</i> Zanardini 1858	-	1986 <sup>178</sup>	-	-	E	RS,IP	L	Hs	I
<i>Griffithsia corallinoides</i> (Linnaeus) Trevisan, 1845	-	1993 <sup>23</sup>	-	-	E	AT, IP	G	Hs	I
<i>Hypnea spinella</i> (C. Agardh) Kützing, 1847	-	-	1987 <sup>18</sup>	-	E	AT,IP	S	Hs	I
<i>Hypnea variabilis</i> Okamura, 1909	-	1986 <sup>178</sup>	1986 <sup>178</sup>	-	E	PO	S	Hs	I
<i>Laurencia intermedia</i> Yamada, 1931	1986 <sup>26</sup>	-	-	-	E	PO,RS	S	Hs	I,II
<i>Lophocladia lallemandii</i> (Montagne) Schmitz, 1893	-	-	1970 <sup>103</sup>	1986 <sup>53</sup>	E	IP	L	Hs	I,II
<i>Polysiphonia fucoides</i> (Hudson) Greville, 1824	1973 <sup>176</sup>	-	1973 <sup>176</sup>	-	E	AT, IO	S	Hs	I,II
<i>Polysiphonia paniculata</i> Montagne, 1842	1986 <sup>26</sup>	-	-	-	E	AT	S	Hs	I
<i>Polysiphonia kampsaxiii</i> Boergesen, 1939	-	-	1984 <sup>17</sup>	1997 <sup>20</sup>	E	IP	S	Hs	I
<i>Radicilingua thysanorhizans</i> (Holmes) Papenfuss, 1956	-	1986 <sup>178</sup>	1986 <sup>26</sup>	1986 <sup>53</sup>	E	AT	S	Hs	I
<i>Rhodophysema georgii</i> Batters, 1900	-	1986 <sup>178</sup>	-	-	E	AT,PO	?S	Hs	I
<i>Spyridia hypnoides</i> (Bory de Saint-Vincent) Papenfuss, 1968	-	-	1987 <sup>18</sup>	1997 <sup>19</sup>	E	IO	S	Hs	I
<b>Heterokontophyta</b>									
<i>Chorda filum</i> (Linnaeus) Stackhouse 1797	-	1986 <sup>178</sup>	-	-	?	AT,PO	?S	Hs	I,II
<i>Ectocarpus siliculosus</i> (Dillwyn) Lyngbye, 1819	1973 <sup>175</sup>	1899 <sup>91</sup>	1899 <sup>91</sup>	1986 <sup>146</sup>	E	AT	S	Hs	I
<i>Halothrix lumbricalis</i> (Kützing) Reinke, 1888	2004 <sup>29</sup>	1993 <sup>23</sup>	1986 <sup>178</sup>	1993 <sup>145</sup>	E	AT,PO	S	Hs	I
<i>Pilayella littoralis</i> (Linnaeus) Kjellman, 1872	1998 <sup>24</sup>	1993 <sup>23</sup>	1985 <sup>105</sup>	1985 <sup>105</sup>	E	AT	S	Hs	I
<i>Protectocarpus speciosus</i> Boergesen, 1902	-	1993 <sup>23</sup>	-	-	E	AT	S	Hs	I
<i>Sargassum latifolium</i> (Turner) C. Agardh, 1820	-	1986 <sup>178</sup>	1986 <sup>178</sup>	-	E	RS	?L	Hs	I,II
<i>Sphaerotrichia divaricata</i> (Agardh) Kylin, 1940	-	1986 <sup>178</sup>	1970 <sup>103</sup>	1986 <sup>178</sup>	E	AT	S	Hs	I
<i>Styopodium schimperi</i> (Buchinger ex Kützing) Verlaque & Boudouresque, 1991	-	-	-	1997 <sup>20</sup>	E	RS,AT	?L	Hs	I
<b>Chlorophyta</b>									
<i>Bryopsis pennata</i> Lamouroux, 1809	-	1986 <sup>26</sup>	1978 <sup>52</sup>	1978 <sup>52</sup>	E	AT	S	Hs	I
<i>Caulerpa racemosa</i> (Forsskål) J. Agardh, 1873	-	-	1991 <sup>54</sup>	1986 <sup>53</sup>	E	CT	?L	Hs,Ss	I,II
<i>Caulerpa scalpelliformis</i> (R.Brown ex Turner) C. Agardh, 1817	-	-	-	1998 <sup>86</sup>	E	AT,IP	?L	Ss	I,II
<i>Cladophora boodleoides</i> Børgesen 1925	-	-	1996 <sup>70</sup>	-	E	AT	G	Hs	I

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Codium fragile</i> (Suringar) Hariot ssp. <i>tomentosoides</i> (Goor) P.C. Silva, 1955	-	-	1986 <sup>178</sup>	-	E	AT,PO	?S	Hs	I
<i>Ulva fasciata</i> Delile, 1813	1990 <sup>25</sup>	1986 <sup>26</sup>	1977 <sup>104</sup>	1996 <sup>22</sup>	E	?RS	?L	Hs	I
<b>Phanerogame</b>									
<i>Halophila stipulacea</i> (Forsskål) Ascherson	-	-	1985 <sup>105</sup>	1985 <sup>105</sup>	E	IO,RS	L	Hs	I
<b>FORAMINIFERA (Benthic)</b>									
<i>Astaculus insolitus</i> (Schwager, 1866)	-	-	2004 <sup>132</sup>	-	E	IP,RS	L	Ss	III
<i>Astaculus sublegumen</i> (Parr, 1950)	-	-	2004 <sup>132</sup>	-	E	IP,RS	L	Ss	III
<i>Planogypsina acervalis</i> (Brady, 1884)	-	-	2004 <sup>132</sup>	-	E	IP,RS	L	Ss	II
<i>Planogypsina squamiformis</i> (Chapman, 1901)	-	-	2004 <sup>132</sup>	-	E	IP,RS	L	Ss	II,III
<i>Amphistegina lobifera</i> Larsen, 1976	-	-	2004 <sup>132</sup>	2002 <sup>133</sup>	E	IP,RS	L	Ss	I,II
<b>CNIDARIA</b>									
<i>Cassiopea andromeda</i> (Forsskål, 1775)	-	-	-	2002 <sup>39</sup>	E	IP,RS	L	Ss	I
<i>Rhopilema nomadica</i> Galil, Spanier & Ferguson, 1990	-	-	-	1995 <sup>113</sup>	E	IP,RS	L	P	?
<b>CTENOPHORA</b>									
<i>Mnemiopsis leidyi</i> (Agassiz, 1865)	1994 <sup>137</sup>	1994 <sup>114</sup>	1994 <sup>114</sup>	1993 <sup>167</sup>	E	WA	S	P	I,II
<i>Beroe ovata</i> Mayer 1912	2001 <sup>115</sup>	2004 <sup>108</sup>	-	-	E	WA	S	P	I,II
<b>POLYCHAETA</b>									
<i>Lepidonotus carinulatus</i> (Grube, 1870)	-	1963 <sup>149</sup>	-	-	Q	IP,RS	?S	?	?
<i>Harmothoe boholensis</i> (Grube, 1878)	-	1963 <sup>149</sup>	-	-	Q	IP,RS	?S	?	?
<i>Harmothoe minuta</i> (Potts, 1910)	-	1963 <sup>149</sup>	-	-	Q	IP,RS	?S	?	?
<i>Ancistrosyllis rigida</i> Fauvel, 1919	-	1963 <sup>149</sup>	-	-	Q	IP,RS	?S	?	?
<i>Sigambra constricta</i> (Southern, 1921)	-	1963 <sup>149</sup>	-	-	Q	IP,RS	?S	?	?
<i>Opisthosyllis brunnea</i> Langerhans, 1879	-	-	2002 <sup>63</sup>	-	Ex	IP,RS	L	Hs	I
<i>Branchiosyllis exilis</i> (Gravier, 1900)	-	-	2002 <sup>63</sup>	-	Ex	IP,RS	L	Hs	I
<i>Leonnates persicus</i> Wesenberg-Lund, 1949	-	-	2002 <sup>65</sup>	2003 <sup>85</sup>	E	IP,RS	L	Ss	II-IV
<i>Nereis zonata persica</i> Fauvel, 1911	-	1963 <sup>149</sup>	-	-	Q	IP,RS	?L	?	?
<i>Pseudonereis anomala</i> Gravier, 1900	-	-	2005 <sup>64</sup>	1989 <sup>33</sup>	E	IP,RS	L	Hs	I
<i>Glycera alba adpersa</i> Fauvel, 1939	-	1963 <sup>149</sup>	-	-	Ex	IP	?S	?	?
<i>Eurythoe complanata</i> (Pallas, 1766)	-	-	-	1997 <sup>81</sup>	Q	?IP	L	Hs	I
<i>Linopherus acarunculata</i> Monro, 1937	-	-	-	1997 <sup>81</sup>	E	RS	L	Ss	I
<i>Lumbrineris debilis</i> Grube, 1878	-	1963 <sup>149</sup>	-	-	Q	IP	?S	?	?
<i>Lysidice collaris</i> Grube, 1870	-	-	1998 <sup>61</sup>	1997 <sup>81</sup>	Q	IP,RS	L	Hs,Ss	I,II
<i>Prionospio saccifera</i> Mackie & Hartley, 1990	-	-		1999 <sup>62</sup>	E	IP,RS	L	Ss	III
<i>Polydora cornuta</i> Bosc, 1802	-	-	2005 <sup>66</sup>	-	E	?	S	Ss	I,II
<i>Streblospio gynobranchiata</i> Rice & Levin, 1998	-	-	2005 <sup>66</sup>	-	E	WA	S	Ss	I,II
<i>Dasybranchus carneus</i> Grube, 1870	-	1963 <sup>149</sup>	-	-	Q	RS	?S	?	?
<i>Notomastus aberans</i> Day, 1957	-	-	1983 <sup>141</sup>	-	E	IP	L	Ss	I

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Monticellina dorsobranchialis</i> (Kirkegaard, 1959)	-	-	2006 <sup>84</sup>	-	Ex	?	L	Ss	?
<i>Timarete dasylophius</i> (Marenzeller, 1879)	-	1963 <sup>149</sup>	-	-	Q	IP	?S	?	?
<i>Timarete anchylochaeta</i> (Schmarda, 1861)	-	1963 <sup>149</sup>	-	-	Q	IP	?S	?	?
<i>Metasychis gotoi</i> (Izuka, 1902)	-	-	2002 <sup>83</sup>	-	E	IP,RS	?L	Ss	?
<i>Rhodine loveni</i> Malmgren, 1865	-	-	1992 <sup>80</sup>	1998 <sup>82</sup>	Ex	IP,AT	L	Ss	II
<i>Pista unibranchia</i> Day, 1963	-	-	1998 <sup>142</sup>	1997 <sup>81</sup>	E	IP,RS	L	Ss	I,II
<i>Ficopomatus enigmaticus</i> (Fauvel, 1923)	-	1952 <sup>67</sup>	1976 <sup>79</sup>	-	E	ST	S	Hs	I
<i>Hydroides dianthus</i> (Verrill, 1873)	-	-	1865 <sup>148</sup>	-	E	WA	S	Hs	I
<i>Hydroides elegans</i> (Haswell, 1883)	-	-	1976 <sup>79</sup>	1991 <sup>34</sup>	E	A,P	S	Hs	I
<i>Spirorbis marioni</i> Caullery & Mesnil, 1897	-	-	1991 <sup>117</sup>	-	E	P	S	Hs	I
<b>PANTOPODA</b>									
<i>Anoplodactylus californicus</i> Hall, 1912	-	-	-	1962 <sup>153</sup>	E	CT	L	Hs	I
<b>CRUSTACEA</b>									
<b>Copepoda</b>									
<i>Centropages furcatus</i> (Dana, 1846)	-	2000 <sup>164</sup>	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Calanopia elliptica</i> (Dana, 1846)	-	-	-	2003 <sup>157</sup>	E	IP,RS	L	P	I
<i>Calanopia biloba</i> Bowman, 1957	-	-	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Calanopia minor</i> Scott, 1902	-	-	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Labidocera pavo</i> Giesbrecht, 1889	-	-	-	2003 <sup>157</sup>	E	IP,RS	L	P	I
<i>Euchaeta concinna</i> Dana, 1846	-	-	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Eucalanus crassus</i> Giesbrecht, 1888	-	-	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Eucalanus subcrassus</i> Giesbrecht, 1888	-	-	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Parvocalanus latus</i> Andronov, 1972	-	2000 <sup>164</sup>	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Parvocalanus elegans</i> Andronov, 1972	-	2000 <sup>164</sup>	-	2002 <sup>168</sup>	E	IP,RS	L	P	I-III
<i>Acartia tonsa</i> Dana, 1848	-	2000 <sup>164</sup>	2001 <sup>155</sup>	-	E	WA	S	P	?
<i>Acartia hasanii</i> Unal, Shmeleva & Kideys, 2002	-	-	-	2002 <sup>163</sup>	Q	?RS	?L	P	I
<i>Paracartia ioannae</i> Unal, Shmeleva & Kideys, 2002	-	-	-	2002 <sup>163</sup>	Q	?RS	?L	P	I
<i>Paracartia janetae</i> Unal, Shmeleva & Kideys, 2002	-	-	-	2002 <sup>163</sup>	Q	?RS	?L	P	I
<b>Cirripedia</b>									
<i>Heterosaccus dollfusi</i> Boschma, 1960	-	-	-	1997 <sup>140</sup>	E	RS	L	Pz	I,II
<b>Amphipoda</b>									
<i>Maera hamigera</i> Haswell, 1879	-	-	1978 <sup>121</sup>	1978 <sup>121</sup>	E	IP	L	Ss	I,II
<i>Stenothoe gallensis</i> Walker, 1904	-	-	1978 <sup>121</sup>	1978 <sup>121</sup>	E	IP	L	Ss	I,II
<b>Isopoda</b>									
<i>Sphaeroma walkeri</i> (Stebbing, 1905)	-	-	1976 <sup>118</sup>	-	E	IP	L	Hs	I,II

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<b>Tanaidacea</b>									
<i>Aapseudes intermedius</i> Hansen, 1985	-	-	1978 <sup>119</sup>	-	E	AT	?S	Hs	I,II
<b>Cumacea</b>									
<i>Eocuma sarsii</i> (Kossmann, 1880)	-	-	1983 <sup>109</sup>	-	E	IP	L	Ss	I,II
<b>Decapoda</b>									
<i>Marsupenaeus japonicus</i> (Bate, 1888)	-	2001 <sup>173</sup>	2001 <sup>173</sup>	1930 <sup>136</sup>	E	IP	L	Ss	I-III
<i>Melicertus hathor</i> (Burkenroad, 1959)	-	-	-	2002 <sup>128</sup>	E	RS,IO	L	Ss	I-III
<i>Metapenaeopsis aegyptia</i> Galil & Golani, 1990	-	-	-	2004 <sup>170</sup>	E	IP,RS	L	Ss	I-III
<i>Metapenaeopsis mogiensis consobrina</i> (Nobili, 1904)	-	-	-	2004 <sup>170</sup>	E	IP,RS	L	Ss	I-III
<i>Metapenaeus monoceros</i> (Fabricius, 1798)	-	-	-	1961 <sup>106</sup>	E	IP,RS	L	Ss	I-III
<i>Penaeus semisulcatus</i> de Haan, 1844	-	-	-	1930 <sup>101</sup>	E	IP,RS	L	Ss	I-IV
<i>Metapenaeus stebbingi</i> (Nobili, 1904)	-	-	-	1981 <sup>120</sup>	E	RS,IO	L	Ss	I-III
<i>Trachysalambria palaestinensis</i> Steinitz, 1932	-	-	-	1968 <sup>95</sup>	E	RS	L	Ss	II,III
<i>Leptochela pugnax</i> de Man, 1916	-	-	-	1981 <sup>120</sup>	E	IP,RS	L	Ss	I-IV
<i>Palaemonella rotumana</i> (Borradaile, 1898)	-	-	-	2002 <sup>92</sup>	E	IP,RS	L	Ss	I-IV
<i>Alpheus audouini</i> (Coutière, 1905)	-	-	-	2002 <sup>92</sup>	E	IP,RS	L	Hs	I,II
<i>Alpheus inopinatus</i> Holthuis & Gottlieb, 1958	-	-	-	1969 <sup>93</sup>	E	IO,RS	L	Hs,Ss	I,II
<i>Alpheus migrans</i> Lewinsohn & Holthuis, 1978	-	-	-	1994 <sup>122</sup>	E	RS	L	Ss	II,III
<i>Alpheus rapacida</i> de Man, 1908	-	-	-	1981 <sup>120</sup>	E	IP	L	Ss	I-III
<i>Urocaridella antonbrunii</i> (Bruce, 1967)	-	-	-	2004 <sup>170</sup>	Ex	IP	L	Ss	I,II
<i>Ixa monodi</i> Holthuis & Gottlieb, 1956	-	-	-	1956 <sup>107</sup>	E	RS	L	Ss	II,III
<i>Leucosia signata</i> Paulson, 1875	-	-	-	1982 <sup>99</sup>	E	IP	L	Ss	I,II
<i>Myra subgranulata</i> Kossmann, 1877	-	-	-	1930 <sup>136</sup>	E	IP,RS	L	Ss	I-IV
<i>Micippa thalia</i> (Herbst, 1803)	-	-	-	1995 <sup>74</sup>	E	IP	L	Hs,Ss	I-III
<i>Callinectes sapidus</i> Rathbun, 1896	-	-	2001 <sup>173</sup>	1961 <sup>106</sup>	E	WA	S	Ss	I-III
<i>Charybdis hellerii</i> (Milne Edwards, 1867)	-	-	-	1981 <sup>120</sup>	E	IP,RS	L	Hs	I,II
<i>Charybdis longicollis</i> Leene, 1938	-	-	-	1961 <sup>106</sup>	E	IP,RS	L	Ss	I-III
<i>Portunus pelagicus</i> (Linnaeus, 1758)	-	-	-	1928 <sup>100</sup>	E	IP,RS	L	Ss	I-III
<i>Thalamita poissonii</i> (Audouin, 1826)	-	-	-	1961 <sup>106</sup>	E	IP,RS	L	Hs,Ss	I-III
<i>Carupa tenuipes</i> Dana, 1851	-	-	-	2004 <sup>170</sup>	?	IP,RS	L	Hs	I-III
<i>Pilumnopus vauquelini</i> (Audouin, 1826)	-	-	-	1981 <sup>120</sup>	E	IP,RS	L	Hs	I
<i>Calappa hepatica</i> (Linnaeus, 1758)	-	-	-	2003 <sup>30</sup>	E	IP,RS	L	Ss	I-III
<i>Atergatis roseus</i> (Rüppell, 1830)	-	-	-	1990 <sup>76</sup>	E	IP,RS	L	Hs	I,II

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Daira perlata</i> (Herbst, 1790)	-	-	-	1995 <sup>74</sup>	?	IP	L	Hs,Ss	I
<i>Eucrate crenata</i> de Haan, 1835	-	-	-	1992 <sup>75</sup>	E	IP,RS	L	Ss	I-III
<i>Macrophthalmus graeffei</i> Milne Edwards, 1873	-	-	-	1995 <sup>74</sup>	?	IP,RS	L	Ss	I-III
<b>Stomatopoda</b>									
<i>Erugosquilla massavensis</i> (Kossmann, 1880)	-	2004 <sup>110</sup>	-	1961 <sup>106</sup>	E	IP,RS	L	Ss	I-III
<b>MOLLUSCA</b>									
<b>Gastropoda</b>									
<i>Diodora ruppellii</i> (Sowerby, 1834)	-	-	-	1995 <sup>72</sup>	E	IP,RS	L	Hs	I
<i>Smaragdia souverbiana</i> (Montrouzier, 1863)	-	-	-	1994 <sup>48</sup>	E	IP,RS	?	Ss	I
<i>Trochus erythreus</i> Brocchi, 1821	-	-	-	1995 <sup>72</sup>	E	RS	L	Hs	I
<i>Pseudominolia nedyma</i> (Melville, 1897)	-	-	-	1995 <sup>72</sup>	E	RS,IO	L	Ss	I,II
<i>Stomatella impertusa</i> (Burrow, 1815)	-	-	-	2000 <sup>152</sup>	?	IP,RS	?	Hs	I
<i>Cerithium scabridum</i> Philippi, 1848	-	-	1990 <sup>3</sup>	1987 <sup>129</sup>	E	RS,IO	L	Hs,Ss	I
<i>Rhinoclavis kochi</i> (Philippi, 1848)	-	-	-	1987 <sup>129</sup>	E	IP,RS	?L	Ss	I
<i>Gibborissoa virgata</i> (Philippi, 1849)	-	-	-	2002 <sup>1</sup>	E	IP	L	?	?
<i>Finella pupoides</i> Adams, 1860	-	-	-	1990 <sup>159</sup>	E	IP	L	Ss	I
<i>Clathrofenella ferruginea</i> (Adams, 1860)	-	-	-	1995 <sup>72</sup>	E	IP,RS	L	Ss	I,II
<i>Cerithiopsis pulvis</i> (Issel, 1869)	-	-	-	1990 <sup>159</sup>	E	RS	?L	Ss	I
<i>Cerithiopsis tenthrenois</i> (Melville, 1896)	-	-	-	1990 <sup>159</sup>	E	IO	?L	Ss	I
<i>Metaxia bacillum</i> (Issel, 1869)	-	-	-	1995 <sup>72</sup>	E	RS	?L	?	?
<i>Rissoina ambigua</i> (Gould, 1849)	-	-	-	2004 <sup>135</sup>	?	IP	?	?	II
<i>Rissoina bertholleti</i> Issel, 1869	-	-	-	1991 <sup>139</sup>	E	RS,IO	?L	Ss	I
<i>Strombus persicus</i> (Swainson, 1821)	-	-	1991 <sup>139</sup>	1983 <sup>138</sup>	E	PG	?	Hs,Ss	I,II
<i>Purpuradusta gracilis notata</i> (Gill, 1858)	-	-	-	1983 <sup>45</sup>	E	RS,IO	?L	Hs	I
<i>Cycloscala hyalina</i> (Sowerby, 1844)	-	-	-	1999 <sup>96</sup>	E	IP,RS	?	?	?
<i>Sticteulima cf. lentiginosa</i> (Adams, 1861)	-	-	-	1994 <sup>158</sup>	?	IP	?	?	?
<i>Ergalatax obscura</i> Houart, 1996	-	-	-	1995 <sup>50</sup>	E	RS	?	Hs	I
<i>Thais lacera</i> (Born, 1778)	-	-	-	1991 <sup>139</sup>	E	PG,IO	S	Hs,Ss	I
<i>Rapana venosa</i> (Valenciennes, 1846)	1960 <sup>90</sup>	1996 <sup>12</sup>	1995 <sup>72</sup>	-	E	PO	S	Ss	I
<i>Zafra savignyi</i> (Moazzo, 1939)	-	-	-	1991 <sup>139</sup>	E	RS	?L	Ss	I,II
<i>Zafra selasphora</i> (Melville & Standen, 1901)	-	-	-	1993 <sup>147</sup>	E	RS,IO	?L	?	?
<i>Lienardia mighelsi</i> Iredale & Tomlin, 1917	-	-	2003 <sup>135</sup>	-	?	IP	?	?	III
<i>Murchisonella columna</i> (Hedley, 1907)	-	-	-	1995 <sup>46</sup>	?	IP	?	Hs	I
<i>Chrysallida fischeri</i> (Hornung & Mermod, 1925)	-	-	-	1992 <sup>134</sup>	E	RS	?L	?	I

(continued)



Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Chrysalida maiae</i> (Hornung & Mermod, 1924)	-	-	-	1990 <sup>159</sup>	E	RS	L	Hs,Ss	I
<i>Chrysalida pirinthella</i> (Melvill, 1910)	-	-	-	1992 <sup>134</sup>	E	RS	?L	Ss	I
<i>Adelactaeon amoenus</i> (Adams, 1851)	-	-	-	1992 <sup>134</sup>	E	IP	?L	Ss	I
<i>Adelactaeon fulvus</i> (Adams, A., 1851)	-	-	-	1992 <sup>134</sup>	E	IP	?L	Ss	I,II
<i>Styloptygma beatrix</i> Melvill, 1896	-	-	-	1992 <sup>134</sup>	E	PG	?L	?	I
<i>Cingulina isseli</i> (Tryon, 1886)	-	-	-	1989 <sup>5</sup>	E	RS	?L	?	?
<i>Turbonilla edgarii</i> (Melvill, 1896)	-	-	-	1992 <sup>134</sup>	E	IP	?L	?	?
<i>Syrnola cincitella</i> Adams, A., 1860	-	-	-	1998 <sup>4</sup>	?	IP,RS	?	?	I
<i>Syrnola fasciata</i> Jickeli, 1882	-	-	-	1989 <sup>5</sup>	E	IP	?L	Ss	I
<i>Iolaea neofelixoides</i> (Nomura, 1936)	-	-	-	1998 <sup>4</sup>	?	PO	?	Ss	I
<i>Hinemoa cylindrica</i> (de Folin, 1879)	-	-	-	2001 <sup>51</sup>	?	IP	?	?	?
<i>Leucotina</i> cf. <i>eva</i> Thiele, 1925	-	-	-	2001 <sup>96</sup>	?	IP	?	?	?
<i>Acteocina crithodes</i> Melvill & Standen, 1907	-	-	-	2004 <sup>135</sup>	?	IP	?	Ss	II
<i>Acteocina mucronata</i> (Philippi, 1849)	-	-	1990 <sup>6</sup>	1990 <sup>6</sup>	E	RS	?L	Ss	?
<i>Cylichnina girardi</i> (Audouin, 1826)	-	-	1996 <sup>49</sup>	1990 <sup>159</sup>	E	IP	L	Ss	I
<i>Pyrunculus fourierii</i> (Audouin, 1826)	-	-	-	1989 <sup>5</sup>	E	IP,RS	L	Ss	II,III
<i>Bulla ampulla</i> Linnaeus, 1758	-	-	-	2004 <sup>171</sup>	E	IP	?L	Ss	II
<i>Haminoea cyanomarginata</i> Heller & Thomson, 1983	-	-	2004 <sup>171</sup>	2004 <sup>171</sup>	?	RS	?L	Hs	II
<i>Chelidonura fulvipunctata</i> Baba, 1938	-	-	-	1961 <sup>154</sup>	E	IP	?L	Ss	I,II
<i>Oxynoe viridis</i> (Pease, 1861)	-	-	-	2004 <sup>171</sup>	?	IP	?L	Ss	I
<i>Elysia grandifolia</i> Kelaart, 1858	-	-	-	2004 <sup>171</sup>	Q	?IO	?L	Hs	I
<i>Elysia tomentosa</i> Jensen, 1997	-	-	-	2004 <sup>171</sup>	?	?IP	?	Ss	I,II
<i>Bursatella leachii</i> Blainville, 1817	-	-	1961 <sup>154</sup>	1961 <sup>154</sup>	E	CT	?L	Ss	I
<i>Syphonota geographica</i> (Adams & Reeve, 1850)	-	-	-	2004 <sup>171</sup>	?	IP	L	Ss	I,II
<i>Plocamopherus ocellatus</i> Rüppell & Leuckart, 1830	-	-	-	2004 <sup>171</sup>	E	RS	L	Hs	I
<i>Hypselodoris infucata</i> Rüppell & Leuckart, 1828	-	-	-	2001 <sup>56</sup>	E	IP,RS	?L	Hs	I
<i>Melibe viridis</i> Kelaart, 1858	-	-	-	2004 <sup>171</sup>	?	IP	?	Ss	I
<i>Flabellina rubrolineata</i> (O'Donoghue, 1929)	-	-	2004 <sup>171</sup>	2004 <sup>171</sup>	E	IP,RS	?L	Hs	I,II
<i>Siphonaria belcheri</i> Hanley, 1858	-	-	-	2001 <sup>13</sup>	E	IP,RS	L	Hs	I
<b>Bivalvia</b>									
<i>Anadara demiri</i> (Piani, 1981)	-	-	1977 <sup>68</sup>	-	E	IO	?S	Ss	I,II
<i>Anadara inflata</i> (Reeve, 1846)	-	-	-	2002 <sup>60</sup>	?	IO	?S	Ss	?
<i>Anadara inaequivalvis</i> (Bruguière, 1789)	2003 <sup>10</sup>	1996 <sup>11</sup>	1995 <sup>72</sup>	-	E	IP	?S	Ss	I,II
<i>Anadara natalensis</i> (Krauss, 1848)	-	-	-	1991 <sup>139</sup>	E	IO,RS	L	Ss	I,II
<i>Brachidontes pharaonis</i> (Fischer, 1870)	-	-	1990 <sup>3</sup>	1985 <sup>116</sup>	E	IO,RS	L	Hs	I
<i>Septifer forskali</i> Dunker, 1855	-	-	-	2001 <sup>13</sup>	?	RS	?	Hs	II

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Crassostrea gigas</i> (Thunberg, 1793)	-	2004 <sup>15</sup>	-	2001 <sup>57</sup>	E	PO	Aq	Hs	I
<i>Saccostrea cucullata</i> (Born, 1778)	-	-	-	2001 <sup>57</sup>	E	IP,RS	?S	Hs	I,II
<i>Dendrostroma frons</i> (Linnaeus, 1758)	-	-	-	2001 <sup>59</sup>	E	IP	?	Hs	?
<i>Pinctada radiata</i> (Leach, 1814)	-	-	1990 <sup>159</sup>	1985 <sup>116</sup>	E	IP,RS	?L	Hs	I
<i>Electroma vexillum</i> (Reeve, 1857)	-	-	-	2005 <sup>58</sup>	?	IP,RS	?L	Hs	I
<i>Malvufundus regulus</i> (Forsskål, 1775)	-	-	-	1974 <sup>88</sup>	E	IP,RS	L	Hs	I
<i>Spondylus cf. multisetosus</i> Reeve, 1856	-	-	-	2001 <sup>59</sup>	Q	IP	?	Hs	?
<i>Spondylus spinosus</i> Schreibers, 1793	-	-	-	1999 <sup>73</sup>	E	IP,RS	L	Hs	I,II
<i>Chama pacifica</i> Broderip, 1834	-	-	-	2001 <sup>59</sup>	E	IP,RS	L	Hs	I,II
<i>Fulvia fragilis</i> (Forsskål, 1775)	-	-	2005 <sup>144</sup>	1987 <sup>129</sup>	E	IO,RS	?L	Ss	I
<i>Afrocardium richardi</i> (Audouin, 1826)	-	-	-	2000 <sup>2</sup>	E	RS	L	Ss	?
<i>Tellina valtonis</i> Hanley, 1844	-	-	-	2001 <sup>96</sup>	E	IO,RS	L	Ss	I
<i>Psammotreta praerupta</i> (Salisbury, 1934)	-	-	-	1999 <sup>73</sup>	?	WA	?	Ss	IV
<i>Gafrarium pectinatum</i> (Linnaeus, 1758)	-	-	-	1987 <sup>129</sup>	E	IP,RS	L	Ss	I
<i>Clementia papyracea</i> (Gray, 1825)	-	-	-	1995 <sup>72</sup>	E	IP,RS	L	Ss	III
<i>Paphia textile</i> (Gmelin, 1791)	-	-	-	1991 <sup>139</sup>	E	IP,RS	L	Ss	?
<i>Ruditapes philippinarum</i> (Adams & Reeve, 1850)	-	-	2001 <sup>14</sup>	-	E	PO	Aq	Ss	I
<i>Antigona lamellaris</i> Schumacher, 1817	-	-	-	1999 <sup>73</sup>	?	IP,RS	?	Ss	?
<i>Mya arenaria</i> Linnaeus, 1758	-	1996 <sup>11</sup>	-	-	E	WA	?S	Ss	II
<i>Gastrochaena cymbium</i> Spengler, 1783	-	-	-	1991 <sup>139</sup>	E	IP,RS	L	Ss	I-III
<i>Laternula anatina</i> (Linnaeus, 1758)	-	-	-	1995 <sup>72</sup>	E	IP,RS	L	Ss	I-III
<b>Cephalopoda</b>									
<i>Octopus aegina</i> Gray, 1849	-	-	-	1999 <sup>151</sup>	E	IP	?L	Ss	III
<i>Sepioteuthis lessoniana</i> Lesson, 1830	-	-	-	2002 <sup>150</sup>	E	IP	L	P	I-III
<b>BRYOZOA</b>									
<i>Rhynchozoon larreyi</i> (Audouin, 1826)	-	-	1979 <sup>165</sup>	-	E	IP,RS	L	Ss	I
<b>ECHINODERMATA</b>									
<i>Ophiactis savignyi</i> (Müller & Troschel, 1842)	-	-	1998 <sup>61</sup>	-	E	IP,RS	L	Hs	I
<i>Synaptula reciprocans</i> (Forsskål, 1775)	-	-	2001 <sup>173</sup>	-	E	IP,RS	L	Hs,Ss	?
<b>PISCES</b>						IP,RS			
<b>Chondrichthyes</b>									
<i>Carcharhinus altimus</i> (Springer, 1950)	-	-	-	2000 <sup>31</sup>	Q	TA	G	Ss	I-VI
<i>Himantura uarnak</i> (Forsskål, 1775)	-	-	-	1966 <sup>37</sup>	C	IP,RS	L	Ss	I,II
<b>Osteichthyes</b>									
<i>Dussumieria elopsoides</i> Bleeker, 1849	-	-	-	1953 <sup>36</sup>	E	IP,RS	L	P	I,II
<i>Etrumeus teres</i> (DeKay, 1848)	-	-	-	1997 <sup>32</sup>	E	IP,RS	L	P	I,II
<i>Herklotsichthys punctatus</i> (Rüppell, 1837)	-	-	-	1984 <sup>169</sup>	E	RS	L	P	I,II
<i>Enchelycore anatina</i> (Lowe, 1839)	-	-	-	2002 <sup>172</sup>	E	TA	G	Hs	I,II

(continued)

Table 1 (continued)

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Saurida undosquamis</i> (Richardson, 1848)	-	-	1973 <sup>38</sup>	1966 <sup>37</sup>	E	IP,RS	L	Ss	I-III
<i>Parexocoetus mento</i> (Valenciennes, 1846)	-	-	1966 <sup>37</sup>	1966 <sup>37</sup>	E	IP,RS	L	P	I
<i>Hemiramphus far</i> (Forsskål, 1775)	-	-	1950 <sup>126</sup>	1950 <sup>126</sup>	E	IP,RS	L	P	I
<i>Hyporhamphus affinis</i> (Günther, 1866)	-	-	-	1954 <sup>7</sup>	Q	IP,RS	L	P	I
<i>Fistularia commersonii</i> (Rüppell, 1835)	-	-	2002 <sup>43</sup>	2002 <sup>42</sup>	E	IP,IO	L	Hs,Ss	I,II
<i>Hippocampus fuscus</i> Rüppell 1838	-	-	-	2004 <sup>98</sup>	?	IO,RS	L	Ss	I
<i>Syngnathus rostellatus</i> Nilsson, 1855	-	-	-	2004 <sup>98</sup>	Q	BA	G	Hs,Ss	I
<i>Atherinomorus lacunosus</i> (Forster, 1801)	-	-	1969 <sup>94</sup>	1950 <sup>126</sup>	E	IP,RS	L	P	I
<i>Sargocentron rubrum</i> (Forsskål, 1775)	-	-	1950 <sup>126</sup>	1950 <sup>126</sup>	E	IP,RS	L	Hs	I-III
<i>Pelates quadrilineatus</i> (Bloch, 1790)	-	-	-	1987 <sup>131</sup>	E	IP,RS	L	Ss	I,II
<i>Apogon pharaonis</i> Bellotti, 1874	-	-	-	1987 <sup>131</sup>	E	IP,RS	L	Hs,Ss	I,II
<i>Sillago sihama</i> (Forsskål, 1775)	-	-	2004 <sup>41</sup>	1994 <sup>102</sup>	E	IP,RS	L	Ss	I,II
<i>Alepes djedaba</i> (Forsskål, 1775)	-	-	1969 <sup>94</sup>	1957 <sup>9</sup>	E	IP,RS	L	P	I-III
<i>Leiognathus klunzingeri</i> (Steindachner, 1898)	-	-	1966 <sup>37</sup>	1943 <sup>78</sup>	E	RS	L	Ss	I-III
<i>Upeneus moluccensis</i> (Bleeker, 1855)	-	-	1956 <sup>127</sup>	1950 <sup>126</sup>	E	IP	L	Ss	I-IV
<i>Upeneus pori</i> Ben-Tuvia & Golani, 1989	-	-	-	1950 <sup>126</sup>	E	IP,RS	L	Ss	I,II
<i>Pempheris vanicolensis</i> Cuvier, 1831	-	-	1999 <sup>130</sup>	1994 <sup>102</sup>	E	IP,RS	L	Hs	I,II
<i>Heniochus intermedius</i> Steindachner, 1893	-	-	-	2003 <sup>97</sup>	Q	IP,RS	L	Hs,Ss	I
<i>Chelon carinata</i> (Valenciennes, 1836) [= <i>Liza carinata</i> ]	-	-	-	1956 <sup>127</sup>	E	RS,IO	L	P	I
<i>Liza haematocheila</i> (Temminck & Schlegel, 1845) [= <i>Mugil soiuy</i> Basilevsky, 1855]	1992 <sup>166</sup>	1998 <sup>112</sup>	1998 <sup>112</sup>	-	E	PO	Aq	P	I
<i>Sphyræna pinguis</i> Doiuchi & Nakabo, 2005 [= <i>S. chrysotaenia</i> Klunzinger, 1884]	-	-	1969 <sup>94</sup>	1957 <sup>9</sup>	E	IP,RS	L	P	I,II
<i>Sphyræna obtusata</i> Cuvier, 1829 [= <i>S. flavicauda</i> Rüppell, 1838]	-	-	-	2002 <sup>42</sup>	E	IP,RS	L	P	I,II
<i>Pteragogus pelycus</i> Randall, 1981	-	-	-	2000 <sup>156</sup>	E	IP,RS	L	Hs,Ss	I,II
<i>Petroscirtes ancyllodon</i> Rüppell, 1838	-	-	-	2000 <sup>156</sup>	E	IP,RS	L	Hs,Ss	I
<i>Oxyurichthys petersi</i> (Klunzinger, 1871)	-	-	1999 <sup>35</sup>	1992 <sup>111</sup>	E	RS	L	Ss	I-III
<i>Callionymus filamentosus</i> Valenciennes, 1837	-	-	-	1994 <sup>102</sup>	E	IP,RS	L	Ss	I-III
<i>Siganus luridus</i> (Rüppell, 1829)	-	-	1973 <sup>38</sup>	1973 <sup>38</sup>	E	IP,RS	L	Hs	I,II
<i>Siganus rivulatus</i> Forsskål, 1775	-	-	1950 <sup>126</sup>	1950 <sup>126</sup>	E	IP,RS	L	Hs	I,II
<i>Scomberomorus commerson</i> Lacepède, 1800	-	-	1997 <sup>47</sup>	1987 <sup>89</sup>	E	IP,RS	L	P	I,II
<i>Solea senegalensis</i> Kaup, 1858	-	1942 <sup>77</sup>	1942 <sup>77</sup>	1942 <sup>77</sup>	Q	TA	G	Hs	I-III
<i>Cynoglossus sinusarabici</i> (Chabanaud, 1931)	-	-	-	1957 <sup>9</sup>	E	RS	L	Hs	I,II

(continued)

**Table 1 (continued)**

	BS	SM	AS	LS	Es	O	MI	H	D
<i>Stephanolepis diaspros</i> Fraser-Brunner, 1940	-	-	1950 <sup>126</sup>	1950 <sup>126</sup>	E	IP,RS	L	Hs,Ss	I,II
<i>Lagocephalus sceleratus</i> (Gmelin, 1789)	-	-	2005 <sup>8</sup>	-	E	IP,RS	L	Ss	I,II
<i>Lagocephalus spadiceus</i> (Richardson, 1844)	-	-	1966 <sup>37</sup>	1950 <sup>126</sup>	E	IP,RS	L	Ss	I,II
<i>Lagocephalus suezensis</i> Clark & Gohar, 1953	-	-	2002 <sup>42</sup>	1999 <sup>16</sup>	E	RS	L	Ss	I,II
<i>Sphoeroides pachygaster</i> (Müller & Troschel, 1848)	-	-	2003 <sup>87</sup>	1999 <sup>130</sup>	E	TA	G	Hs,Ss	I-V
<i>Torquigener flavimaculosus</i> Hardy & Randall, 1983	-	-	-	2003 <sup>40</sup>	E	IP,RS	L	Ss	I,II

coast. Forty-eight alien Crustaceans (94% of the total aliens) belong to Lessepsian migrants. Most of the plants (29 species) were introduced via ships. The other modes of introduction of species were migration through the Gibraltar straits and Aquaculture.

#### ***Distribution of alien species on the Turkish coasts***

The number of alien species found along the Turkish coasts and their modes of introduction are presented in Figure 1. The Turkish Black Sea coast possessed the lowest number of species (20 species), whereas the Turkish Levantine coast had the highest (202 species). The importance of shipping as a vector of introduction of species gradually diminishes from the Black Sea to the Levantine Sea. Lessepsian migrants exclusively dominate benthic habitats of the Levantine Sea. The Aegean Sea appears to be more influenced by Lessepsian migration than ship transportation.

The majority (70% of the species) of the marine animals introduced by ships in the Aegean Sea were reported from Izmir Bay; *Polydora cornuta*, *Streblospio gynobranchiata*, *Ficopomatus enigmaticus*, *Hydroides dianthus*, *H. elegans*, *Spirorbis marioni*, *Anadara demiri* and *A. inequivalvis*. This is mainly attributed to two facts; 1) Alsancak Harbour, which is one of the most important commercial harbours in Turkey, is located in the polluted part of Izmir Bay, 2) This bay has been relatively well studied and

continuously monitored since the 1970's. Species-poor communities such as polluted or physically degraded environments are known to be more vulnerable to invasion than are other communities (ZIBROWIUS, 1992). Izmir Bay has been subjected to various pollution discharges, intense marine transportation and increasing human populations. Ocean-going ships approaching Alsancak Harbour for loading or unloading processes empty their ballast tanks just before entering the harbour and the weakening in the ecosystem due to pollution greatly facilitates the settlement of alien species in the area.

#### ***Rate of invasion***

The new additions to the inventory of the marine alien species on the Turkish coasts greatly increased after 1980 and the last five years alone a total of 92 new species were recorded (Fig. 2). A decreasing pattern is obvious in the yearly rate of introduction over the past years (Table 2). While one new alien species was introduced every 15.3 weeks over the period 1981-2000 in the Aegean Sea, the time span has decreased to 8.7 between 2001 and 2005. The special interest in aliens by Turkish and foreign scientists in the last decade has increased and thereby, the real diversity of aliens within the groups of phyto-benthos, fishes and crustaceans inhabiting the Turkish coasts seems to be well documented. Nowadays, a number of projects regarding the actual status of aliens along the Turkish coasts




Fig. 1: The number of alien species along the Turkish coasts and their mode of introduction. PLT: All plants, PP: Phytoplankton, PhB: Phytobenthos, CTN: Ctenophora, POL: Polychaeta, CRU: Crustacea, MOL: Mollusca, OTH: Others.  $\Sigma S$  indicates the total number of species reported from the sea.




Fig. 2: Rate of introduction of alien species along the Turkish coasts.

**Table 2.**  
Time span (weeks) for alien species introduction along the Turkish coasts.

	1961-1980	1981-2000	2001-2005
<b>Weeks per 1 alien species</b>			
Black Sea	208	69.3	65
Sea of Marmara	80	31.5	65
Aegean Sea	40	15.3	8.7
Levantine Sea	52	6.4	4.7
Total	16.3	3.7	2.8


and their probable impacts on the native biota are undertaken.

### Depth and habitat preferences of alien species

The distribution of alien species by depth and habitat is indicated in Figure 3. The majority of aliens collected from the Turkish coasts were found in shallow waters. A total of 106 species (41% of the total number of aliens) solely occurred at depths ranging from 0 to 10 m, and 180 species (68%) at the depth interval 0-50 m. Species with a wide distributional range in the area are *Leonnates persicus* (10-200 m), *Penaeus semisulcatus* (0-200 m), *Leptochela pugnax* (0-200 m), *Palaemonella rotumana* (0-200 m), *Myra subgranulata* (0-200 m), *Carcharhinus altimus* (0-500 m), *Upeneus moluccensis* (0-200 m) and *Spherooides pachygaster* (0-400 m).

Benthic habitats (soft and hard substrata) harboured 76% of the total number of alien species (Figure 3). Thirty-nine species inhabited the pelagic environment. In soft substrata, all major groups, except for algae, are represented by a higher number of species. The majority of hard bottom organisms were reported among algae.

The echinoderm *Ophiactis savignyi* was only reported in canals of the sponge *Sarcotragus muscarum* from the southern part of the Aegean Sea [ÇINAR & ERGEN, 1998 (as *Ophiactis virens* (M. Sars, 1857)); ÇINAR *et al.*, 2002). The sweeper, *Pempheris vanicolensis*, prefers cavern habitats (BILECENOĞLU & TASKAVAK, 1999). The shrimp scad, *Alepes djedaba*, is generally associated with the venomous jellyfish *Rhopilema nomadica* (KIDEYS & GÜCÜ, 1995). ØKSNEBJERG *et al.* (1997)



**Fig. 3:** The habitat (upper graphic) and depth (lower graphic) preferences of alien species along the Turkish coasts.

reported the parasitic cirriped *Heterosaccus dollfusi* on the Lessepsian crab *Charybdis longicollis*. The nudibranch *Flabellina rubrolineata* was found on hydroids (YOKES & RUDMAN, 2004). ALBAYRAK & ÇEVİKER (2001) reported *Septifer forskali* on the bivalve *Spondylus spinosus*. *Electroma vexillum* was encountered in discharge canals of the Iskenderun Iron and Steel Factory, where hot water is discharged into the sea (ÇEVİK *et al.*, 2005).

## Impacts of alien species

The phytoplankton species, *Alexandrium tamarense*, *Gymnodinium* cf. *mikimotoi* and *Heterosigma* cf. *akashii*, which were reported from Izmir Bay and classified as questionable in Table 1, have a potential to cause a toxic and/or harmful bloom along the Turkish coasts (KORAY, 1984; BIZSEL & BIZSEL, 2002).

The green alga *Caulerpa racemosa* invaded soft and hard bottom habitats of the Levantine and Aegean coasts but its impact on the native biota has not yet been documented in the area. However, its occurrence off the Dardanel (salinity: 27 psu) indicates its highly adaptive and invasive character (OKUDAN *et al.*, 2002).

The negative effect of *Mnemiopsis leidyi* on the Black Sea ecosystem was summarized by KIDEYS (2002). This zooplanktonic predator reached enormous biomass levels in the summer of 1989, devastating the food chain of the entire Black Sea basin, and subsequently led to a sharp decrease in the anchovy production along the Turkish coast. The other ship-mediated ctenophore, *Beroe ovata*, which largely feeds on ctenophores, appeared in 1997 in the Black Sea and has resulted in the decrease in the population of *M. leidyi*.

The outburst of the population of *Rhopilema nomadica* off the Levantine coast of Turkey was reported to have negative consequences on human health, tourism and fisheries (KIDEYS & GÜCÜ, 1995). Many swimmers were stung and sought medical treatment. The blockage of nets of fishermen by individuals of *R. nomadica* also created major economical losses.

*Rapana venosa*, which feeds exclusively on bivalves, is responsible for the decrease in stocks of the bivalves *Ostrea edulis* Linné, 1758 and *Mytilus galloprovincialis* (Lamarck, 1819) in the Black Sea (BILECIK, 1990). The fishery of *R. venosa* was intense along the Turkish Black Sea coast, reaching up to 1166 tons in 1986, yielding 3.415.884 US \$ (BILECIK, 1990)

Low faunal diversity found on sponge samples collected from the southern Aegean Sea was due to the invasion of the Lessepsian species *Ophiactis savignyi* (ÇINAR *et al.*, 2002).

The species number and density of other groups within the sponge samples were sharply declined as compared to those collected from the northern sites where this echinoderm was not present. A negative correlation was estimated between number of specimens of the ophiuroid and the total number of specimens of other taxa.

The soft bottom near Alsancak Harbour in Izmir Bay was exclusively dominated by the spionid polychaete *Streblospio gynobranchiata*, which originated from the western Atlantic (ÇINAR *et al.*, 2005). Its density reached almost 34 300 ind.m<sup>-2</sup> in the area and accounted for almost 100% of faunal populations at some stations. The soft bottom benthic community around Alsancak Harbour seems to be restructured by this species.

KOÇAK *et al.* (1999) found dense populations of two alien serpulid species, *Hydroides elegans* and *H. dianthus*, on panels submerged in a polluted marina in Izmir Bay. The species density of *H. elegans* reached up to 98000 ind.m<sup>-2</sup> and that of *H. dianthus* up to 2 000 ind.m<sup>-2</sup>. The above authors also reported that an alien serpulid species of eastern Pacific origin, *Spirorbis marioni*, inhabited port environments subjected to relatively less pollution.

The density of *Anadara demiri* was found to be 300 ind.m<sup>-2</sup> on grey mud and 30 ind.m<sup>-2</sup> on black mud in Izmir Bay (DEMİR, 1977).

Aliens might bring their own parasites with them. ØKSNEBJERG *et al.* (1997) reported that 90% of the specimens of *Charybdis longicollis* collected between Karatas and Fener Burnu (Levantine coast of Turkey) were infected by the rhizocephalan *Heterosaccus dollfusi*. They also stated that the prevalence of this parasite was estimated at 50-60% in the east of Karatas, with its prevalence gradually declining towards the east into Iskenderun Bay. GALİL & LÜTZEN (1995) found that 55% of the males and 43% of the females of *C. longicollis* were visibly infected by this parasite along the Israeli coast in May 1994.

From a large number of aliens inhabiting the Turkish coasts, only a few possess commercial value, mostly belonging to fishes and crustaceans. Little is known concerning the impact of




Fig. 4: Percentages of alien species in total fauna along the Turkish coasts.

commercially important alien species on local fisheries, since there is no available catch statistic that distinguishes native and alien species' landings. Due to the great number of species caught during various fisheries activities, governmental statistics tend to present relevant data on a family basis. For example, the "mullids" category includes four species (native *Mullus spp.* and Lessepsian *Upeneus spp.*); a similar situation is valid for sphyraenids (two native species vs. two Lessepsian species). Lessepsian clupeids, especially *Etrumeus teres*, form a remarkable proportion of the pelagic catch along the Levantine coast. The siganids are generally sold together with the native *Sarpa salpa* due to their similar morphologies (BILECENOĞLU & KAYA, 2002). At least seven fish species (*Upeneus moluccensis*, *Saurida undosquamis*, *Scomberomorus commerson*, *E. teres*, *Sphyraena pinguis*, *Siganus spp.*) are captured in large amounts (mainly by bottom trawls), and consumed throughout the Levantine and southern Aegean Sea coasts. In terms of biomass, *S. undosquamis* is the most prominent species in the bottom trawl catch composition (31.9% of total CPUE in Iskenderun Bay), followed by *U. moluccensis* and *Siganus rivulatus* (GÜCÜ *et al.*, 1994).

Some species have relatively smaller catches and thus mostly consumed locally, i.e.

*Alepes djedaba*, *Sillago sihama*, *Dussumieria elopsoides*, *Atherinomorus lacunosus*. Among alien crustaceans, the highest annual production belongs to *Callinectes sapidus* (ca. 200 tonnes/year), which is captured especially in lagoons in amounts as much as 2 tonnes/day during summer periods (ÖZCAN *et al.*, 2003). *Marsupenaeus japonicus* is one the most valuable shrimp species along the Levantine coast, which is also imported to various cities in Turkey. Since they are not captured in large quantities, other commercial shrimps (*Metapenaeus spp.*, *Penaeus semisulcatus*) and brachyurans (*Portunus pelagicus*) are, in general, consumed locally.

#### Importance of alien species in total fauna

The relative percentages of aliens in total fauna are shown in Figure 4. Mollusca had the highest score in the Black (3.1%) and Levantine (16.1%) Seas, Polychaeta in the Sea of Marmara (4.4%) and fishes in the Aegean Sea (5.9%). The score for Polychaeta in the Sea of Marmara does not reflect its true number as 11 of 12 aliens are questionable. The score seems to be accurate for the better-studied group, fishes, which has more aliens on the southern coasts than the northern coasts due to the influence of Lessepsian migrants. The highest alien percentages for all groups examined were found on the Levantine



coast. This partly shows the impact of Lessepsian migration on the Levantine fauna and partly the lack of detailed works on some groups on the coast. The total number of species given for Mollusca and Polychaeta from the Levantine coast of Turkey is highly underestimated and undoubtedly would increase as further studies are carried out. It is a common assumption that Lessepsian migrants account for at least 10% of the species inventory of the Levantine Sea (POR, 1978). BEN-ELIAHU (1995) estimated that the proportion of migrant polychaetes is almost 9%; for fishes 13% (GOLANI, 1998) and for decapod crustaceans 20% (GALIL, 1986).

### Similarity among seas in terms of alien species

The dendrogram and MDS configuration in Figure 5 show that there are two groups in terms of the presence and absence of alien species in the Turkish seas. The Black Sea and the Sea of Marmara constitute one group, in which 16 species (mainly algae) are common. The other group with a similarity of 40% is composed from the Aegean and Levantine Seas, where 63 species (mainly lessepsian fishes) are common. There is a very weak similarity (20%) between the groups, which are placed at different corners of the MDS plot. As pointed out earlier, the southern seas seem to be more affected by Lessepsian migrants than ship-borne species, whereas shipping is the main vector in the transportation of aliens to the northern seas. The difference in the way of introduction seems to be the main reason for similarity or dissimilarity among the seas.

### Conclusions

The compiled data show that a total of 263 established or questionable alien species inhabit the coast of Turkey, with the Levantine Sea having the majority of them (202 species). Detecting new aliens depends totally on accurate taxonomic identifications and the knowledge of local biodiversity. This requires the availability of systematic experts on biotal components in the area and a close cooperation among scientists both on local and global scales. The wealth




Fig. 5: Dendrogram and MDS plot showing affinities of the seas in terms of alien species they included.

of data accumulated on aliens depicts the importance of monitoring studies on the understanding of ecological and functional roles of aliens in the ecosystems. Finally, results of studies and projects should not be concealed among pages of reports and papers. They should be presented to governmental establishments and politicians to force them to take immediate precautions on the dispersal of aliens, particularly via ballast water discharges.

### Addendum

Before the publication of this contribution, a total of 14 new alien species were reported from the Levantine (13 species) and Aegean [1 species (*Platax teira*)] coasts of Turkey (BILECENOĞLU & KAYA, 2006; ÇEVİKER & ALBAYRAK, 2006; ÇINAR *et al.*, 2006; ÖZTÜRK & CAN, 2006). These species belong to Hydrozoa (*Macrorhynchia philippina* (Kirch-enpauer, 1872)), Anthozoa (*Oculina patagonica*

De Angelis, 1908), Polychaeta (*Branchiomma luctuosum* Grube, 1869), Gastropoda (*Amathina tricarinata* (Linnaeus, 1767), *Aplysia dactylomela* Rang, 1828, *Chromodoris quadricolor* (Rüppell & Leuckart 1830)], Bivalvia [*Cardites akabana* (Sturany, 1899) and *Petricola hempri* Issel, 1869], Echinodermata [*Synaptula reciprocans* (Forsskål, 1775)], Tunicata [*Phallusia nigra* Savigny, 1816, *Pyura* (= *Herdmania*) *momus* (Savigny, 1816) and *Symplegma brakenhielmi* (Michaelsen, 1904)] and Osteichthyes [*Parupeneus forsskali* (Fourmanoir and Guézé, 1976) and *Platax teira* (Forsskål, 1775)]. These findings increased the number of alien species known from the coast of Turkey from 263 to 277, and from 202 to 216 on the Levantine coast of Turkey.

## References

- <sup>1</sup>AARTSEN, J. J. van, 2002. Indo-Pacific migrants into the Mediterranean. 1. *Gibborissoa virgata* (Philippi, 1849). *La Conchiglia*, 303: 56-58.
- <sup>2</sup>AARTSEN, J. J. van & GOUD, J., 2000. European marine Mollusca: Notes on less well-known species. XV. Notes on Lusitanian species of *Parvicardium* Monterosato, 1884, and *Afrocardium richardi* (Audouin, 1826) (Bivalvia:Heterodonta: Cardiidae). *Basteria*, 64: 171-186.
- <sup>3</sup>AARTSEN, J. J. van & KINZELBACH, R., 1990. Marine molluscs from the Iztuzu beach near Dalyan (Mediterranean coast of Turkey). *Zoology in the Middle East*, 4: 103-112.
- <sup>4</sup>AARTSEN, J. J. van & RECEVIK, M., 1998. Two more Indo-Pacific species found in the Mediterranean. *La Conchiglia*, 286: 13-15.
- <sup>5</sup>AARTSEN, J. J. van, BARASH, A. & CARROZZA, F., 1989. Addition to the knowledge of the Mediterranean mollusca of Israel and Sinai. *Bollettino Malacologico*, 25: 63-76.
- <sup>6</sup>AARTSEN, J. J. van, CARROZZA, F. & LINDNER, G., 1990. *Acteocina mucronata* (Philippi, 1849), a recent Red Sea immigrant species in the eastern Mediterranean. *Bollettino Malacologico*, 25: 285-298.
- <sup>7</sup>AKSIRAY, F., 1954. Türkiye deniz balıkları tayin anahtarı. *Istanbul Üniversitesi Fen Fakültesi Hidrobiyoloji Araştırma Enstitüsü Yayınları*, İstanbul, 277 p.
- <sup>8</sup>AKYOL, O., ÜNAL, V., CEYHAN, T. & BILECENOĞLU, M., 2005: First record of the silverside blaasop, *Lagocephalus sceleratus* (Gmelin, 1789), in the Mediterranean Sea. *Journal of Fish Biology*, 66: 1183-1186.
- <sup>9</sup>AKYUZ, E., 1957. Observations on the Iskenderun red mullet (*Mullus barbatus*) and its environment. *GFCM Proceedings and Technical Papers*, 4: 305-326.
- <sup>10</sup>ALBAYRAK, S., 2003. On the Mollusca Fauna of the Black Sea near Istanbul. *Zoology in the Middle East*, 30: 69-75.
- <sup>11</sup>ALBAYRAK, S. & BALKIS, N., 1996a. Bivalve Fauna of the Bosphorus. *Istanbul Üniversitesi Fen Fakültesi Biyoloji Dergisi*, 59: 1-15.
- <sup>12</sup>ALBAYRAK, S. & BALKIS, N., 1996b. Benthic Prosobranch Gastropods of the Bosphorus. *Istanbul Üniversitesi Fen Fakültesi Biyoloji Dergisi*, 59: 17-26.
- <sup>13</sup>ALBAYRAK, S & ÇEVİKER, D., 2001. Two extra-Mediterranean molluscs from southeast Turkey: *Siphonaria belcheri* Hanley, 1858 (Gastropoda: Siphonariidae) and *Septifer bilocularis* (Linnaeus, 1758) (Bivalvia:Veneridae). *Israel Journal of Zoology*, 47: 297-298.
- <sup>14</sup>ALBAYRAK, S., ASLAN, H. & BALKIS, H., 2001. A contribution to the Aegean Sea fauna: *Ruditapes philippinarum* (Adam & Reeve, 1850) [Bivalvia : Veneridae]. *Israel Journal of Zoology*, 47: 299-300.
- <sup>15</sup>ALBAYRAK, S., BALKIS, H. & BALKIS, N., 2004. Bivalvia (Mollusca) fauna of the Sea of Marmara. *Acta Adriatica*, 45: 9-26.
- <sup>16</sup>AVSAR, D. & ÇIÇEK, E., 1999. A new species record for the central and eastern Mediterranean; *Sphoeroides cutaneus* (Günther, 1870) (Pisces: Tetraodontidae). *Oebalia*, 25: 17-21.

- <sup>17</sup>AYSEL, V., 1984. Türkiye'nin Ege Denizi'ndeki *Polysiphonia* Greville (*Rhodomelaceae*, *Ceramiales*) türleri 1-Bölüm *Oligosiphonia*. *Doga Bilim Dergisi*, 8: 29-42.
- <sup>18</sup>AYSEL, V., 1987. Türkiye'nin Ege Denizi Florası II.Kırmızı Algler (*Rhodophyta*). *Doga Türk Botanik Dergisi*, 11: 1-21.
- <sup>19</sup>AYSEL, V., 1997a. Marine flora of Turkish Mediterranean coast 1. Red algae (*Rhodophyta*). *Turkish Journal of Botany*, 21: 155-163.
- <sup>20</sup>AYSEL, V., 1997b. Marine flora of Turkish Mediterranean coast 2. Brown algae (*Fucophyceae* = *Phaeophyceae*). *Turkish Journal of Botany*, 21: 329-334.
- <sup>21</sup>AYSEL, V. & ERDUGAN, H., 1995. Check-list of Blacksea seaweeds, Türkiye (1823-1994). *Turkish Journal of Botany*, 19: 545-554.
- <sup>22</sup>AYSEL, V. & GEZERLER-SIPAL, U., 1996. Türkiye'nin Akdeniz kıyılarının deniz florası, 3. Cyanophyceae, Chlorophyceae, Charophyceae ve Angiospermae. *Ege Üniversitesi Su Ürünleri Fakültesi Dergisi*, 13: 247-257.
- <sup>23</sup>AYSEL, GÜNER, H. & DURAL, B., 1993. Türkiye Marmara Denizi florası II. *Phaeophyta* ve *Rhodophyta*. *Ege Üniversitesi Su Ürünleri Fakültesi Dergisi*, 10: 115-168.
- <sup>24</sup>AYSEL, V., DURAL, B., GÖNÜZ, A. & OKUDAN, E. S., 1998. Kırklareli (Karadeniz, Trakya, Türkiye) deniz florası. *XIV Ulusal Biyoloji Kongresi*, Samsun, 2: 333-342.
- <sup>25</sup>AYSEL, V., KESERCIOGLU, T., GÜNER, H. & AKÇAY, H., 1990. Trabzon deniz algleri. *X Ulusal Biyoloji Kongresi*, Erzurum, 2: 183-192.
- <sup>26</sup>AYSEL, V., ZEYBEK, N., GÜNER, H. & SUKATAR, A., 1986. Türkiye'nin bazı derin deniz algleri III. *Rhodophyta* (Kırmızı Algler). *Doga Türk Biyoloji Dergisi*, 10: 8-29.
- <sup>27</sup>AYSEL, V., DURAL, B., OKUDAN, E. S., ALPASLAN, M. & UYSAL, I., 2001. Gökçeada (Ege Denizi, Çanakkale) Deniz Florası. *Ulusal Ege Adaları 2001 Toplantısı*, GökÇeada: 125-142.
- <sup>28</sup>AYSEL, V., DURAL, B., SUKATAR, A., GÜNER, H. & ERDUGAN, H., 1996a. Zonguldak Deniz Algleri, Karadeniz, Türkiye. *XIII Ulusal Biyoloji Kongresi*, İstanbul, 5: 311-321.
- <sup>29</sup>AYSEL, V., ERDUGAN, H., DURAL-TARAKÇI, B., OKUDAN, E. S., SENKARDESLER, A. & AYSEL, F. 2004. Marine Flora of Sinop (Black Sea, Turkey), *Ege University Journal of Fisheries & Aquatic Sciences*, 21: 59-68.
- <sup>30</sup>BALKIS, H. & ÇEVİKER, D., 2003. A new exotic crab species [*Calappa hepatica* (Linnaeus, 1758)] for the Mediterranean fauna. *Israel Journal of Zoology*, 49: 320-321.
- <sup>31</sup>BASUSTA, N. & ERDEM, Ü., 2000. Iskenderun Körfezi balıkları üzerine bir araştırma. *Turkish Journal of Zoology*, 24:1-19.
- <sup>32</sup>BASUSTA, N., ERDEM, Ü. & MATER, S., 1997. Iskenderun körfezi'nde yeni bir Lesepsiyen göçmen balık türü; Kizilgözlü Sardalya, *Etrumeus teres* (DeKay, 1842). *Mediterranean Fisheries Congress*, İzmir: 921-924.
- <sup>33</sup>BEN-ELIAHU, M. N., 1989. Lessepsian migration in Nereidae (Annelida: Polychaeta). Some case histories. p. 125-134. In: *Proceedings of Environmental Quality and Ecosystem Stability*, edited by E. Spanier, Y. Steinberger & M. Luria.
- <sup>34</sup>BEN-ELIAHU, M. N., 1991. Red Sea serpulids (Polychaeta) in the eastern Mediterranean. p. 515-528. In: *Proceedings of 2nd International Polychaeta Conference, Copenhagen, 1986. Systematics, Biology and Morphology of Word Polychaeta*, edited by M. E. Petersen & J. B. Kirkegaard, Ophelia Supplement, 5.
- BEN-ELIAHU, M. N., 1995. A list of Polychaeta along the Levant coast. *Haasiana*, 1: 78-93.
- <sup>35</sup>BENLİ, H. A., CIHANGİR, B. & BİZSEL, K. C., 1999. Ege Denizi'nde bazı demersal balıkçılık kaynakları üzerine

- arastirmalar. *Istanbul Üniversitesi Su Ürünleri Dergisi*, özel sayı: 301-369.
- <sup>36</sup>BEN-TUVIA, A., 1953. Mediterranean fishes of Israel. *Bulletin of the Sea Fisheries Research Station*, 8:1-40.
- <sup>37</sup>BEN-TUVIA, A., 1966. Red Sea fishes recently found in the Mediterranean. *Copeia*, 2: 254-275.
- <sup>38</sup>BEN-TUVIA, A., 1973. Man-made changes in the eastern Mediterranean Sea and their effect on the fishery resources. *Marine Biology*, 19: 197-203.
- <sup>39</sup>BILECENOĞLU, M., 2002. Türkiye kıyılarında ilk kez rastlanan bir denizanası türü: *Cassiopa andromeda* (Forsskal, 1775) (Cassiopidae, Scyphozoa). *Sualtı Dünyası Dergisi*, 72: 42-43.
- <sup>40</sup>BILECENOĞLU, M., 2003. Kizildeniz Göçmeni Balon Balığı (*Torquigener flavimaculosus* Hardy & Randall, 1983), Türkiye kıyılarında ilk gözlemler. *Sualtı Dünyası Dergisi*, 74: 38-39.
- <sup>41</sup>BILECENOĞLU, M., 2004. Occurrence of the Lessepsian migrant fish, *Sillago sihama* (Forsskal, 1775) (Osteichthyes: Sillaginidae), from the Aegean Sea. *Israel Journal of Zoology*, 50: 420-421.
- BILECENOĞLU, M. & KAYA, M., 2002. Growth of marbled spinefoot *Siganus rivulatus* Forsskal, 1775 (Teleostei: Siganidae) introduced to Antalya Bay, eastern Mediterranean Sea (Turkey). *Fisheries Research*, 54: 279-285.
- BILECENOĞLU, M. & TASKAVAK, E., 1999. Some observations on the habitat of Red Sea immigrant sweeper, *Pempheris vanicolensis*, at the Mediterranean coast of Turkey. *Zoology in the Middle East*, 17: 67- 70.
- <sup>42</sup>BILECENOĞLU, M., TASKAVAK, E. & KUNT, K. B., 2002. Range extension of three Lessepsian migrant fish (*Fistularia commersoni*, *Sphyræna flavicauda*, *Lagocephalus suezensis*) in the Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom*, 82: 525-526.
- <sup>43</sup>BILECENOĞLU, M., TASKAVAK, E., MATER, S. & KAYA, M., 2002. Checklist of the marine fishes of Turkey. *Zootaxa*, 113: 1-194.
- BILECENOĞLU, M. & KAYA, M., 2006. A new alien fish in the Mediterranean Sea – *Platax teira* (Forsskal, 1775) (Osteichthyes: Ephippidae). *Aquatic Invasions*, 1: 80-83.
- BILECIK, N., 1990. Distribution of sea snail “*Rapana venosa* (V.)” in the Turkish coasts of Black Sea and its impact on Black Sea fisheries. *Tarım Orman ve Koyisleri Bakanlığı Su Ürünleri Arastırma Enstitüsü Müdürlüğü Yayinlari*, seri B, 1, 34 pp.
- <sup>44</sup>BIZSEL, N. & BIZSEL, C., 2002. New records of toxic algae *Heterosigma* cf. *akashwo* and *Gymnodinium* cf. *mikimotoi* in the hypereutrophic Izmir Bay (Aegean Sea): Coupling between organisms and water quality parameters. *Israel Journal of Plant Sciences*, 50: 33-44.
- <sup>45</sup>BLÖCHER, M., 1983. A dead specimen of *Palmadusta gracilis* (Gaskoin, 1849) found in the Mediterranean. *La Conchiglia*, 10: 166-167.
- <sup>46</sup>BOGI, C., BUZZURRO, G. & GREPPI, E., 1995. Presenza di *Murchisonella columna* (Hedley, 1907) Nel Mediterraneo Orientale. *Bollettino Malacologico*, 31: 69-76.
- <sup>47</sup>BUHAN, E., YILMAZ, H., MORKAN, Y., BÜKE, E. & YÜKSEK, A., 1997. Güllük ve Gökova Körfezleri için yeni bir av potansiyeli: *Scomberomorus commerson* (Lacepède, 1800) (Pisces-Teleostei). *Mediterranean Fisheries Congress*, İzmir: 937-944.
- <sup>48</sup>BUZZURRO, G. & GREPPI, E., 1994. Presenza di *Smaragdia (Smaragdella) souverbiana* (Montrouzier, 1836) nel Mediterraneo Orientale. *Bollettino Malacologico*, 29: 319-321.
- <sup>49</sup>BUZZURRO, G. & GREPPI, E., 1996. The lessepsian molluscs of Tasucu (South-East Turkey). *La Conchiglia*, 279 (supplement): 3-22.
- <sup>50</sup>BUZZURRO, G., ENGL, W. & TÜMTÜRK, I., 1995. Bivalven und Gastropoden

- der Europäischen Meer (4): *Ergalatax martensi* (Dall, 1923) (Muricidae), ein neuer lessepsischer einwanderer von der Türkischen Südküste. *Club Conchylia Informationen*, 27: 17-18.
- <sup>51</sup>BUZZURRO, G., HOARAU, A., GREPPI, E. & PELORCE, J., 2001. Prima segnalazione di *Hinemoa cylindrica* (de Folin, 1879) per il Mediterraneo. *Bollettino Malacologico*, 37: 23-26.
- <sup>52</sup>CIRIK, S., 1978. Recherches sur la végétation marine des côtes Turques de la mer Egée. Etude particuliere des Peyssonneliacées de Turquie. *Thèse Doctora III, Cycle Biologie Végétale, Université Pierre et Marie Curie*, Paris: 172 pp.
- <sup>53</sup>CIRIK, S., 1986. A propos de la végétation marine de la baie d'Akkuyu, Mersin, Turquie. *5th OPTIMA Meeting*, Istanbul: 69-81.
- <sup>54</sup>CIRIK, S. & ÖZTÜRK, B., 1991. Notes sur la présence d'une forme rare du *Caulerpa racemosa*, en Méditerranée orientale. *Flora Mediterranea*, 1: 217-219.
- <sup>55</sup>CIRIK, S., ZEYBEK, N., AYSEL, V. & CIRIK, S., 1990. Note préliminaire sur la végétation marine de l'île Gökçeada (Mer Egée-Nord, Turquie). *Thalassographica*, 13: 33-37.
- <sup>56</sup>ÇEVİK, C. & ÖZTÜRK, B., 2001. A new lessepsian mollusc *Hypselodoris infucata* (Rüppell & Leuckart, 1828) (Gastropoda: Nudibranchia) for the coasts of Turkey. *Turkish Journal of Zoology*, 25: 27-30.
- <sup>57</sup>ÇEVİK, C., ÖZTÜRK, B. & BUZZURRO, G., 2001. The presence of *Crassostrea virginica* (Gmelin, 1791) and *Saccostrea commercialis* (Iredale & Roughley, 1933) in the eastern Mediterranean Sea. *La Conchiglia*, 298: 25-28.
- <sup>58</sup>ÇEVİK, C., DOĞAN, A., ÖNEN, M. & ZENETOS, A., 2005. First record of the Indo-Pacific species *Electroma vexillum* (Mollusca: Bivalvia: Pterioidea) in the eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom*, 2, *Biodiversity Records*, 4996.
- <sup>59</sup>ÇEVİK, D., 2001. Recent immigrant Bivalves in the northeastern Mediterranean off Iskenderun. *La Conchiglia*, 298: 39-46.
- <sup>60</sup>ÇEVİK, D. & ALBAYRAK, S., 2002. New record of an exotic Bivalve from the southeastern coast of Turkey: *Scapharca inflata* (Reeve, 1844). *La Conchiglia*, 305: 56-60.
- ÇEVİK, D. & ALBAYRAK, S., 2006. Three alien molluscs from Iskenderun Bay (SE Turkey). *Aquatic Invasions*, 1: 76-79.
- ÇINAR, M. E., 2003. Ecology of Syllidae (Annelida: Polychaeta) from northern Cyprus (Eastern Mediterranean Sea). *Bulletin of Marine Science*, 72: 795-811.
- ÇINAR, M. E., 2005. Polychaetes from the coast of northern Cyprus (eastern Mediterranean Sea), with two new records for the Mediterranean Sea. *Cahiers de Biologie Marine*, 46: 143-161.
- <sup>61</sup>ÇINAR, M. E. & ERGEN, Z., 1998. Polychaetes associated with the sponge *Sarcotragus muscarum* Schmidt, 1864 from the Turkish Aegean Coast. *Ophelia*, 48: 167-183.
- <sup>62</sup>ÇINAR, M. E. & ERGEN, Z., 1999. Occurrence of *Prionospio saccifera* (Spirochaeta: Polychaeta) in the Mediterranean Sea. *Cahiers de Biologie Marine*, 40: 105-112.
- <sup>63</sup>ÇINAR, M. E. & ERGEN, Z., 2002. Faunistic analysis of Syllidae (Annelida: Polychaeta) from the Aegean Sea. *Cahiers de Biologie Marine*, 43: 171-178.
- <sup>64</sup>ÇINAR, M. E. & ERGEN, Z., 2005. Lessepsian migrants expanding their distributional ranges; *Pseudonereis anomala* (Polychaeta: Nereididae) in Izmir Bay (Aegean Sea). *Journal of the Marine Biological Associations of the United Kingdom*, 85: 313-321.
- <sup>65</sup>ÇINAR, M. E., ERGEN, Z. & DAGLI, E., 2002. Occurrence of the lessepsian species *Leonnates persicus* (Polychaeta: Nereididae) in Izmir Bay, Aegean Sea. *Journal of the Marine Biological Associations of the United Kingdom*, 82: 811-815.

- ÇINAR, M. E., KATAGAN, T., ERGEN, Z. & SEZGIN, M., 2002. Zoobenthos-inhabiting *Sarcotragus muscarum* (Porifera: Demospongia) from the Aegean Sea. *Hydrobiologia*, 482: 107-117.
- <sup>66</sup>ÇINAR, M. E., ERGEN, Z., DAGLI, E. & PETERSEN, M. E., 2005. Alien species of spionid polychaetes (*Streblospio gynobranchiata* and *Polydora cornuta*) in Izmir Bay, eastern Mediterranean. *Journal of the Marine Biological Associations of the United Kingdom*, 85: 821-827.
- ÇINAR, M. E., BILECENOĞLU, M., ÖZTÜRK, B. & CAN, A., 2006. New records of alien species on the Levantine coast of Turkey. *Aquatic Invasions*, 1: 84-90.
- <sup>67</sup>DEMİR, M., 1952-1954. Bogazlar ve adalar sahillerinin omurgasız dip hayvanları. *Istanbul Üniversitesi Fen Fakültesi Hidrobiyoloji Araştırma Enstitüsü Yayınlarından*, 3: 615 pp.
- <sup>68</sup>DEMİR, M., 1977. On the presence of *Arca (Scapharca) amygdalum* Philippi, 1847 in the harbour of Izmir, Turkey. *Istanbul Üniversitesi Fen Fakültesi Mecmuası*, Serie B, 42: 197-202.
- <sup>69</sup>DURAL, B., 1996. Ege Denizi Cyanophyceae türleri. *Ege Üniversitesi Su Ürünleri Fakültesi Dergisi*, 12: 267-292.
- <sup>70</sup>DURAL, B. & AYSEL, V., 1996. A new record for the algal flora of Mediterranean 1. *Cladophora boodleoides* Börgesen. *Turkish Journal of Botany*, 20: 465-468.
- <sup>71</sup>EKER, E., GEORGIEVA, L., SENICHKINA, L. & KIDEYS, A. E., 1999. Phytoplankton distribution in the western and eastern Black Sea in spring and autumn 1995. *ICES Journal of Marine Science*, 56 (Supplement): 15-22.
- ELTON, C. S., 1958. The ecology of invasions by animals and plants. The University of Chicago Press Chicago, 181 pp.
- <sup>72</sup>ENGL, W., 1995. Specie prevalentemente lessepsiane attestate lungo le coste Turche. *Bollettino Malacologico*, 31: 43-50.
- <sup>73</sup>ENGL, W. & ÇEVİKER, D., 1999. New migrant species from southeast Turkey *Psammotreta praerupta* (Salisbury, 1934) and *Antigona lamellaris* Schumacher, 1817. *La Conchiglia*, 290: 17-20.
- <sup>74</sup>ENZENROSS, R. & ENZENROSS, L., 1995. Erstnachweise indopazifischer Brachyura (Crustacea: Decapoda) von detürkischen Mittelmeerküste. *Stuttgarter Beiträge zur Naturkunde*, 521: 1-4.
- <sup>75</sup>ENZENROSS, R., ENZENROSS, L. & KOCATAŞ, A., 1992. Lessepsian migration: two Indo-Pacific crabs species (Crustacea, Decapoda) found on the Turkish Mediterranean coast. *Journal of the Faculty of Science Ege University*, 14: 3-10.
- <sup>76</sup>ENZENROSS, L., ENZENROSS, R. & NIEDERHÖFER, H. J., 1990. Wissenschaftlich interessante Funde aus der Sammlung Enzenross (Marine Invertebraten). *Jahreshefte der Gesellschaften fuer Naturkunde in Wuerttemberg*, 145: 283-294.
- <sup>77</sup>ERAZI, R. A. R., 1942. The heterosomata of the Bosphorus, the Golden Horn and the Sea of Marmara. *Revue de la Faculte des Sciences de l'universite d'Istanbul*, 7: 235-262.
- <sup>78</sup>ERAZI, R. A. R., 1943. *Leiognathus mediterraneus* nov. sp. *Compte Rendu Société Turquie des Sciences physiques et naturelles*, 10: 49-53.
- <sup>79</sup>ERGEN, Z., 1976. Investigations on the taxonomy and ecology of Polychaeta from Izmir Bay and its adjacent areas. *Scientific Reports of the Faculty of Science Ege University*, 209: 1-73.
- <sup>80</sup>ERGEN, Z., 1992. The latest status of Polychaeta in the soft substrate of Izmir Bay. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 33.
- <sup>81</sup>ERGEN, Z. & ÇINAR, M. E., 1997. Polychaeta of Antalya Bay (Mediterranean coast of Turkey). *Israel Journal of Zoology*, 43: 229-241.
- <sup>82</sup>ERGEN, Z., ÇINAR, M. E. & UNSAL, M., 1998. Polychaetes from the Manavgat River Delta (Turkish Mediterranean coast). *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 35: 536-537.

- <sup>83</sup>ERGEN, Z., ÇINAR, M. E., DAGLI, E. & KURT, G., 2002. Lessepsian polychaete species from the Turkish coasts. p. 50-55. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.
- <sup>84</sup>ERGEN, Z., ÇINAR, M. E., DAGLI, E. & KURT, G., 2006. Seasonal dynamics of soft-bottom polychaetes in Izmir Bay (Aegean Sea, eastern Mediterranean). *Scientia Marina*, in press.
- <sup>85</sup>ERGEV, M. B., ÇINAR, M. E., MUTLU, E. & ERGEN, Z., 2003. Ecological features of the lessepsian migrant *Leonnates persicus* (Polychaeta: Nereididae) from the Levant coast of Turkey. *Journal of the Marine Biological Associations of the United Kingdom*, 83: 1225-1226.
- <sup>86</sup>ERTAN, Ö., TURNA, O. I. & CORMACI, M., 1998. A New Record for the Marine Algal flora of Turkey: *Caulerpa scalpelliformis* (Brown ex Turner) C. Agardh (*Caulerpaceae*, *Chlorophyceae*). *Turkish Journal of Botany*, 22: 285-287.
- <sup>87</sup>ERYILMAZ, L., ÖZULUG, M. & MERİÇ, N., 2003. The Smooth Pufferfish, *Sphoeroides pachygaster* (Müller & Troschel, 1848) (Teleostei: Tetraodontidae), new to the Northern Aegean Sea. *Zoology in the Middle East*, 28: 125-126.
- <sup>88</sup>FALCHI, S., 1974. Molluschi di provenienza Indopacifica lungo le coste Turche. *Conchiglie*, Milano, 10: 89.
- <sup>89</sup>FISCHER, W., BAUCHOT, M.-L. & SCHNEIDER, M., 1987. Fiches FAO d'identification des espèces pour les besoins de la pêche. Méditerranée et mer Noire. Zone de pêche 37. *FAO & EEC*, Rome: 761-1530.
- <sup>90</sup>FISCHER-PIETTE, E., 1960. *Rapana bezoar* I. sur la cote Turquie de la mer Noire. *Hydrobiologie*, Serie B5, 1-2: 51.
- <sup>91</sup>FRITSCH, K., 1899. Beitrag zur flora von Constantinopel. I. Kryptogamen. *Denkschriften der kaiserlichen Akademie der Wissenschaften zu Wien Math*, 68: 219-250.
- GALIL, B., 1986. Red Sea decapods along the Mediterranean coast of Israel, ecology and distribution. p. 179-183. In: *Environmental quality and Ecosystem stability*, edited by Z. Dubinsky & Y. Steinberg, Bar Ilan Pres.
- GALIL, B. S. & LÜTZEN, J., 1995. Biological observations on *Heterosaccus dollfusi* Boschma (Cirripedia: Rhizocephala), a parasite of *Charybdis longicollis* Leene (Decapoda: Brachyura), a Lessepsian migrant to the Mediterranean. *Journal of Crustacean Biology*, 15: 659-670.
- <sup>92</sup>GALIL, B. S., FROGLIA, C. & NOEL, P. Y., 2002. CIESM Atlas of Exotic Crustaceans in the Mediterranean. Volume 2: Crustacean Decapods and Stomatopods, F. Briand (ed). Monaco: CIESM Publishers.
- <sup>93</sup>GELDIAY, R., 1969a. A report on the collection of Natantia (Crustacea, Decapoda) along the coast of Turkey from the Eastern Mediterranean to the vicinity of Izmir. *Scientific Reports of Faculty of Science Ege University*, 74: 1-17.
- <sup>94</sup>GELDIAY, R., 1969b. Izmir Körfezinin baslica balıkları ve muhtemel invasionları. *Ege Üniversitesi Fen Fakültesi Monografileri*, Izmir, 135 pp.
- <sup>95</sup>GELDIAY, R. & KOCATAŞ, A., 1968. The species of Penaeidae family from the Mediterranean Coast of Turkey. *Scientific Reports of Faculty of Science, Ege University*, 58: 1-15.
- <sup>96</sup>GIUNCHI, L., MICALI, P. & TISELLI, M., 2001. Report of two new lessepsian immigrants from Turkey. *La Conchiglia*, 298: 47-48.
- <sup>97</sup>GOKOGLU, M., BODUR, T. & KAYA, Y., 2003. First record of the Red Sea bannerfish (*Heniochus intermedius* Steindachner, 1893) from the Mediterranean Sea. *Israel Journal of Zoology*, 49: 324-325.
- <sup>98</sup>GOKOGLU, M., BODUR, T. & KAYA, Y., 2004. First records of *Hippocampus fuscus* and *Syngnathus rostellatus* (Osteichthyes: Syngnathidae) from the Anatolian coast (Mediterranean Sea). *Journal of*

- the Marine Biological Association of the United Kingdom*, 84: 1093-1094.
- GOLANI, D., 1998. Distribution of Lessepsian migrant fish in the Mediterranean. *Italian Journal of Zoology*, 65: 95-99.
- <sup>99</sup>GRIPPA G., 1982. First record of *Leucosia signata* Paulson, 1875 from the South-Western Turkish coast (Decapoda, Brachyura, Leucosidae). *Quaderni del Laboratorio di Tecnologia della Pesca. Ancona*, 3: 335-338.
- <sup>100</sup>GRUVEL, A., 1928. Répartition géographique de quelques crustacés comestibles sur les côtes d'Égypte et de Syrie. *Comptes rendus de la société de biogéographie*, 5: 45-46.
- <sup>101</sup>GRUVEL, A., 1930. Les richesses marines et fluviales de la Syrie. *Exploitation actuelle Avenir Revue Scientifique*, Paris, 68: 33-41.
- <sup>102</sup>GÜCÜ, A.C., BINGEL, F., AVSAR, D. & UYSAL, N., 1994. Distribution and occurrence of Red Sea fish at the Turkish Mediterranean coast - northern Cilician Basin. *Acta Adriatica*, 34: 103-113.
- <sup>103</sup>GÜNER, H., 1970. Ege Denizi sahil algleri üzerinde taksonomik ve ekolojik araştırma. *Ege Üniversitesi Fen Fakültesi İlimi Raporlar Serisi*, 76: 1-77.
- <sup>104</sup>GÜNER, H. & AYSEL, V., 1977. İzmir Körfezi'nde bulunan bazı *Ulva* türleri (*Chlorophyta*) hakkında taksonomik araştırma. *Ege Üniversitesi Fen Fakültesi Dergisi*, Seri B, 1: 241-251.
- <sup>105</sup>GÜNER, H., AYSEL, V., SUKATAR, A. & ÖZTÜRK, M., 1985. Türkiye Ege Denizi Florası I. Mavi-Yesil, Yesil, Esmer Algler ve Kapalı Tohumlular. *Doga Bilim Dergisi*, A2, 9: 272-282.
- <sup>106</sup>HOLTHUIS, L. B., 1961. Report on a collection of crustacea decapoda and stomatopoda from Turkey and the Balkans. *Zoologische Verhandelingen*, 47: 1-67.
- <sup>107</sup>HOLTHUIS, L. B. & GOTTLIEB, E., 1956. Two interesting cabs (crustacea, decapoda, brachyura) from Mersin Bay, S.E. Turkey. *Zoologische Verhandelingen*, 21: 287-301.
- <sup>108</sup>ISINIBILIR, M., TARKAN A. N. & KIDEYS, A. E., 2004. Decreased levels of the invasive ctenophore *Mnemiopsis* in the Sea of Marmara in 2001. p. 155-165. In: *The Invasion of the Black, Mediterranean and Caspian Seas by the American Ctenophore, Mnemiopsis leidyi Agassiz: A Multidisciplinary Perspective*, edited by H. Dumont, T. Shiganova & U. Niermann, Kluwer Academic Publishers, Dordrecht, The Netherlands.
- <sup>109</sup>KATAGAN, T., 1983. Türkiye'nin Ege Denizi littoral kumase'leri üzerine sistematik ve ekolojik araştırmalar. *Ege University Faculty of Science Journal, Serie B*, 6: 9-18.
- <sup>110</sup>KATAGAN, T., KOCATAŞ, A., ZENGİN, M. & ATES, A. S., 2004. An Indo-Pacific stomatopod from the Sea of Marmara: *Erugosquilla massavensis* (Kossmann, 1880). *Crustaceana*, 77: 381-383.
- <sup>111</sup>KAYA, M., MATER, S. & BENLİ, H. A., 1992. A new Indo-Pacific gobiid fish *Oxyurichthys papuensis* (Val., 1837) for eastern Mediterranean coasts of Turkey. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 33: 298.
- <sup>112</sup>KAYA, M., MATER, S. & KORKUT, A. Y., 1998. A new grey mullet species "*Mugil so-iuy* Basilewsky" (Teleostei: Mugilidae) from the Aegean coast of Turkey. *Turkish Journal of Zoology*, 22: 303-306.
- KIDEYS, A. E., 2002. Fall and rise of the Black Sea ecosystem. *Science*, 297: 1482-1484.
- <sup>113</sup>KIDEYS, A. E. & GÜCÜ, A. C., 1995. *Rhopilema nomadica*: A poisonous indo-pacific scyphomedusa new to the Mediterranean coast of Turkey. *Israel Journal of Zoology*, 41: 615-617.
- <sup>114</sup>KIDEYS, A. E. & NIERMANN, U., 1994. Occurrence of *Mnemiopsis* along the Turkish coasts (from northeastern Mediterranean to Istanbul). *ICES Journal of Marine Science*, 51:423-427.
- <sup>115</sup>KIDEYS, A. E., & ROMANOVA, Z.,



2001. Distribution of gelatinous macrozooplankton in the southern Black Sea during 1996-1999. *Marine Biology*, 139: 535-547.
- <sup>116</sup>KINZELBACH, R., 1985. Lesseps'sche wanderung: neue stationen von muscheln (Bivalvia: Anisomyaria). *Archiv für Molluskenkunde*, 115: 273-278.
- <sup>117</sup>KNIGHT-JONES, P., KNIGHT-JONES, W. & ERGEN, Z., 1991. Sabelliform polychaetes, mostly from Turkey's Aegean coast. *Journal of Natural History*, 25: 837-858.
- KOÇAK, F., ERGEN, Z. & ÇINAR, M. E., 1999. Fouling organisms and their developments in a polluted and an unpolluted marina in the Aegean Sea (Turkey). *Ophelia*, 50:1-20.
- <sup>118</sup>KOCATAŞ, A., 1976. Tanaiidaces et Isopodes recuillis dans le golfe d'Izmir, Turquie. *Crustaceana*, 30: 68-72.
- <sup>119</sup>KOCATAŞ, A., 1978. Contribution a l'étude des peuplements des horizons supérieurs de substrat rocheux du Golfe d'Izmir. *Scientific Monographs of the Faculty of Science, Ege University*, 12: 1-93.
- <sup>120</sup>KOCATAŞ, A., 1981. Liste préliminaire et repartition des Crustacea Decapodes des eaux Turques. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 27: 161-162.
- <sup>121</sup>KOCATAŞ, A. & KATAGAN, T., 1978. Türkiye denizlerinin littoral bentik amphipod'lari ve yayilislari. *TÜBİTAK Temel Bilimler Arastirma Grubu*, Proje No.TBAG-223, 63 p.
- <sup>122</sup>KOCATAŞ, A. & KATAGAN, T., 1994. Impact of Lessepsian Decapod and Stomatopod species on the biodiversity of the Turkish Seas. *XII. Ulusal Biyoloji kongresi*: 290-294.
- <sup>123</sup>KOCATAŞ, A., KORAY, T., KAYA, M. & KARA, O. F., 1993. Review of the fishery resources and their environment in the Sea of Marmara. *General Fisheries Council for The Mediterranean, Studies and Reviews*, 64: 87-143.
- <sup>124</sup>KORAY, T., 1984. The occurrence of red-tides and causative organisms in Izmir Bay. *Ege Universitesi Fen Fakültesi Dergisi*, Seri B, 1: 75-83.
- <sup>125</sup>KORAY, T., 1987. One-celled microplankton species of Izmir Bay (Aegean Sea): A species list and a comparison with the records of adjacent regions. *Doga, Turkish Journal of Biology*, 11: 130-146.
- <sup>126</sup>KOSSWIG, C., 1950. Erythraische fische im Mittelmeer und an der grenze der Agais. *Syllegomena Biologica, Festschrift Kleinschmidt*: 203 -212.
- <sup>127</sup>KOSSWIG, C., 1956. Facts and problems offered by the erythrean invaders into the Mediterranean. *Balık ve Balıkçılık*, 4: 31-37.
- <sup>128</sup>KUMLU, M., EROLDÖGAN, O. T., AKTAS, M. & GÖÇER, M., 2002. New Records: A new shrimp record for the Turkish Seas: *Melicertus hator* (Burkenroad, 1959) (Penaeidae: Crustacea). *Israel Journal of Zoology*, 48: 246-247.
- <sup>129</sup>LINDNER, G., 1987. Interessante Schneckenfunde an der Südtürkischen Mittelmeerküste (Reisemitbringsel der Familie SCHMIDT, Feldkirchen). *Club Conchylia Informationen*, 19 : 32-43.
- MACK, R. N., SIMBERLOFF, D., LONSDALE, W. M., EVANS, H., CLOUT, M. & BAZZAZ, F.A., 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications*, 10: 689-710.
- <sup>130</sup>MATER, S. & BILECENOĞLU, M., 1999. Türkiye deniz balıkları. p. 790-808. In: *Genel Zoocoğrafya ve Türkiye Zoocoğrafyası*, edited by A. Demirsoy, Meteksan Matbaası, Ankara.
- <sup>131</sup>MATER, S. & KAYA, M., 1987. Türkiye'nin Akdeniz sularında yeni kaydedilen üç balık türü, *Sudis hyalina* Rafinesque, *Pelates quadrilineatus* (Bloch), *Apogon nigripinnis* Cuvier (Teleostei) hakkında. *Doga, Türk Zooloji Dergisi*, 11: 45-49.
- <sup>132</sup>MERİÇ, E., AVSAR, N. & BERGIN, F., 2004. Benthic foraminifera of eastern Aegean Sea (Turkey) Systematics and

- Autoecology. *Turkish Marine Research Foundation*, 18, 306 pp.
- <sup>133</sup>MERICİ, E., AVSAR, N., BERGIN, F. & YOKES, B., 2002. The proliferation of *Amphistegina* (Lessepsian migrants) population at three-islands (Uçadalar, Antalya), a new observation from the Turkish Mediterranean coast. p. 27-34. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.
- <sup>134</sup>MICALI, P. & PALAZZI, S., 1992. Contributo Alla Conoscenza Dei Pyramidellidae Della Turchia, Con Segnalazione di Due Nuove Immigrazioni Dal Mar Rosso. *Bolletino Malacologico*, 28: 83-90.
- <sup>135</sup>MIENIS, H. K., 2004. New data concerning the presence of lessepsian and indopacific migrants among the molluscs in the Mediterranean Sea with emphasize on the situation in Israel. p. 117-131. In: *Ist National Malacology Congress*, Izmir, edited by B. Öztürk & A. Salman, Turkish Marine Research Foundation Press.
- <sup>136</sup>MONOD, T., 1930. Über einige indopazifische Decapoden der Meeresfauna Syriens. *Zoologischer Anzeiger*, 92: 135-141.
- <sup>137</sup>MUTLU, E., BINGEL, F., GÜCÜ, A.C., MELNIKOV, V. V., NIERMANN, U., OSTR, N. A. & ZAIKA, V. E., 1994. Distribution of new invader *Mnemiopsis* sp. and the resident *Aurelia aurita* and *Pleurobranchia pileus* populations in the Black Sea in the years 1991-1993. *ICES Journal of Marine Science*, 51: 407-421.
- <sup>138</sup>NICOLAY, K. & ROMAGNA-MANOJA, E., 1983. *Strombus* (*Conomurex*) *decorus raybaudii* n. ssp. *La Conchiglia*, 174-175: 20-21.
- <sup>139</sup>NIEDERHÖFER, H., ENZENROSS, L. & ENZENROSS, R., 1991. Neue Erkenntnisse über die Ausbreitung von "Lesseps'schen Einwanderern" (Mollusca) an der Türkischen Mittelmeerküste. *Club Conchylia Informationen*, 13: 94-108.
- OCCHIPINTI AMBROGI, A., 2001. Transfer of marine organisms: a challenge to the conservation of coastal biocoenoses. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 11: 243-251.
- <sup>140</sup>ØKSNEBJERG, B., ENZENROSS, R. & ENZENROSS, L., 1997. First record of Rhizocephala (Crustacea: Cirripedia) from Turkish waters, with notes on Lessepsian migration. *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)*, 55: 1-7.
- OKUDAN, E. S., YURDABAK, F., TUNCER, S., ERDEMİR, C. C. & AYSEL, V., 2002. The distribution of *Caulerpa racemosa* (Forskål) J. Agard on the coast of Turkey. p. 46-49. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.
- <sup>141</sup>ÖNEN, M., 1983. The qualitative and quantitative investigation of the macrobenthic fauna found in the soft substratum of the Urla Harbour. *Ege University Faculty of Science Journal*, 6: 29-39.
- <sup>142</sup>ÖNEN, M., ERGEN, Z., KATAGAN, T. & ÇINAR, M.E., 1998. Güllük Lagünü bentik organizmalarının mevsimsel dinamiği. *XIV Ulusal Biyoloji Kongresi*, Samsun, 2: 203-226.
- ÖZCAN, T., ÖZCAN, G. & AKYURT, I., 2003. Iskenderun Körfezi kıyısındaki kentsel alanda yengeç tüketim ve satın alma davranışları üzerine bir araştırma. *XII. Ulusal Su Urunleri Sempozyumu*: 546-549.
- <sup>143</sup>ÖZTİG, F., 1957. Erdek sahillerinin deniz vejetasyonu hakkında. *Türk Biyoloji Dergisi*, 7: 12-13.
- <sup>144</sup>ÖZTÜRK, B. & POUTIERS, J. M., 2005. *Fulvia fragilis* (Bivalvia: Cardiidae): a lessepsian mollusc species from Izmir Bay (Aegean Sea). *Journal of the Marine Biological Association of the United Kingdom*, 85: 351-356.
- <sup>145</sup>ÖZTÜRK, M., 1993. Türkiye'nin Ege ve Akdeniz Kıyılarındaki *Chordariales* ve *Sporochyales* (*Phaeophyta*) üyelerinin yayılımı ve taksonomisi. *Turkish Journal*

- of *Botany*, 17: 237-247.
- <sup>146</sup>ÖZTÜRK, M. & GÜNER, H., 1986. Türkiye'nin Ege ve Akdeniz Kıyılarındaki Ectocarpales (Phaeophyta) üyelerinin yayılımı ve taksonomisi. *Turkish Journal of Botany*, 10: 459-472.
- ÖZTÜRK, B. & CAN, A., 2006. Indo-Pacific Gastropod Species in the Levantine and Aegean Seas. *Aquatic Invasions*, 1: in press.
- <sup>147</sup>PALAZZI, S., 1993. Ci sono due *Zafra* in mediterraneo, ma come si chiamano? *Notiziario del CISMA*, 14: 13-18.
- POR, F. D., 1978. Lessepsian migration-the influx of Red Sea biota into the Mediterranean by way of the Suez Canal. Springer, Berlin, 228 pp.
- <sup>148</sup>QUATREFAGES, M. A., 1865. Histoire naturelle des Annélides marines et d'eau douce, Annélides et Gephyriens, 2. Sedentaire (Paris), 562 pp.
- <sup>149</sup>RULLIER, F., 1963. Les Annélides polychètes du Bosphore, de la Mer de Marmara et de la Mer Noire, en relation avec celles de la Méditerranée. *Extrait des Rapports et Procès-verbaux des réunions de la CIESM*, 17: 161-260.
- <sup>150</sup>SALMAN, A., 2002. New report of the loliginid squid *Sepioteuthis lessoniana* Lesson, 1830 in the Mediterranean. *Israel Journal of Zoology*, 48: 249-250.
- <sup>151</sup>SALMAN, A., KATAGAN, T. & BOLETZKY, S. von, 1999. New Cephalopod molluscs in the eastern Mediterranean: previously unnoted species or recent migrants? *Vie et Milieu*, 49: 11-17.
- <sup>152</sup>SCHNIEBS, K., 2000. Erster Nachweis einer Stomatelliden-Art im Mittelmeer (Gastropoda: Prosobranchia). *Malakologische Abhandlungen*, 20: 99-100.
- <sup>153</sup>STOCK, J. H., 1962. Some Turkish pycnogonid records. *Entomologische Berichten*, 22: 218-219.
- <sup>154</sup>SWENNEN, C., 1961. On a collection of Opisthobranchia from Turkey. *Zoologische Mededelingen*, 38: 41-75.
- <sup>155</sup>TARKAN, A. N., ISINIBILIR, M. & ERK, M. H., 2001. Dominant zooplankton species in the waters of the northern Gökçeada. *Ulusal Ege Adaları 2001 Toplantısı Bildiriler Kitabı, Turkish Marine Research Foundation*, Istanbul: 59-70.
- <sup>156</sup>TASKAVAK, E., BILECENOĞLU, M., BASUSTA, N. & MATER, S., 2000. Occurrence of *Pteragogus pelycus* Randall, 1981 (Teleostei: Labridae) and *Petroscirtes ancyloдон* Rüppell, 1838 (Teleostei: Blenniidae) on the eastern Mediterranean coast of Turkey. *Acta Adriatica*, 41: 53-58.
- <sup>157</sup>TOKLU, B. & SARIHAN, E., 2003. The Copepoda and Cladocera (Crustacea) Fauna along the Yumurtalik-Botas coastline in Iskenderun Bay. *Ege University Journal of Fisheries and Aquatic Sciences*, 20: 63-68.
- <sup>158</sup>TRINGALI, L., 1994. Levant Sea: A new lessepsian host of the genus *Sticteulima* F. C. Laceron, 1955. *La Conchiglia*, 273: 57-58.
- <sup>159</sup>TRINGALI, L. & VILLA, R., 1990. Rinvenimenti malacologici dalle coste Turche (Gastropoda, Polyplacophora, Bivalvia). *Notiziario del CISMA*, 11: 33-41.
- <sup>160</sup>TURKOGLU, M. & KORAY, T., 2002. Phytoplankton species' succession and nutrients in the southern Black Sea (Bay of Sinop). *Turkish Journal of Botany*, 26: 235-252.
- <sup>161</sup>TURNA, I., CORMACI, M. & FURNARI, G., 2000. First Record of *Botryocladia madagascariensis* G. Feldmann (*Rhodymeniaceae*, *Rhodophyceae*) from the Gulf of Antalya (Mediterranean coast of Turkey). *Plant Biosystems*, 134: 111-115.
- <sup>162</sup>ÜNAL, A., 1970. Türkiye sahillerinde yetiştirilen deniz alglerinin sistematigi, Ankara, 64 p.
- <sup>163</sup>ÜNAL, E., SHMELEVA, A. A. & KIDEYS, A. E., 2002. Three new species of *Acartia* (Copepoda, Calanoida) from the northeastern Levantine Basin. p. 35-45. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.

- <sup>164</sup>ÜNAL, E., SHMELEVA, A. A., ZAGORODNYAYA, J. & KIDEYS, A. E., 2000. Zooplankton structure and copepod species of the Sea of Marmara in spring 1998. p. 450-460. In: *Proceedings book of symposium on Marmara Sea 2000*, edited by B. Öztürk, M. Kadiroglu & H. Öztürk, TUDAV, Istanbul, 5.
- <sup>165</sup>ÜNSAL, I. & d'HONDT, J. L., 1979. Contribution à la connaissance des bryozoaires marins de Turquie (Eurystomata et Cyclostomata). *Vie Milieu*, 28-29: 613-634.
- <sup>166</sup>ÜNSAL, S., 1992. Türkiye denizleri için yeni bir kefal balığı türü: *Mugil so-ıuy* Basilewsky. *Doga, Turkish Journal of Veterinary and Animal Sciences*, 16: 427-432.
- <sup>167</sup>UYSAL, Z. & MUTLU, E., 1993. Preliminary note on the occurrence and biometry of Ctenophoran *Mnemiopsis leidyi* finally invaded Mersin Bay. *Doga, Turkish Journal of Zoology*, 12: 229-236.
- <sup>168</sup>UYSAL, Z., KIDEYS, A. E., SHMELEVA, A. A., ZAGORODNYAYA, J. A. & GUBANOVA, A. D. 2002. Checklist of copepods (Calonida and Podoplea) from the northern Levantine basin shelf waters. *Hydrobiologia*, 482: 15-21.
- <sup>169</sup>WHITEHEAD, P. J. P., 1984. Clupeidae. p. 268-281. In: *Fishes of the North-eastern Atlantic and the Mediterranean*, edited by P. J. P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, UNESCO, Paris.
- <sup>170</sup>YOKES, B & GALIL, B. S., 2004. New records of alien decapods from the south-western coast of Turkey. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 37: 556.
- <sup>171</sup>YOKES, B. & RUDMAN, W. B., 2004. Lessepsian opisthobranchs from south-western coast of Turkey; five new records for Mediterranean. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 37: 557.
- <sup>172</sup>YOKES, B., DERVISOGLU, R. & KARACIK, B., 2002. Likya kıyılarında denizel biyolojik zenginlik araştırması. *Sualti Bilim ve Teknoloji Toplantısı Bildiriler Kitabı*, Istanbul, p.166-181.
- <sup>173</sup>ZAITSEV, Y. & ÖZTÜRK, B., 2001. Exotic species in the Aegean, Marmara, Black, Azov and Caspian Seas. *Turkish Marine Research Foundation*, Istanbul, 267 pp.
- <sup>174</sup>ZEYBEK, N., 1969. Türkiye'nin Akdeniz algleri 1. Bodrum-Finike Körfezi sahil boyu. 2. Ege Denizi-Edremit Saros Körfezi-Sile. TBAG-124 nolu proje.
- <sup>175</sup>ZEYBEK, N., 1973a. Meeresalgen aus der Türkei. 1. Die Buchten von Edremit und Saros am Aegaeischen meer, 2. Die küste von Igneada bis Sile am Schwarzen meer. *Verhandl der Schweizerischen naturforschenden Gesellschaft*: 95-100.
- <sup>176</sup>ZEYBEK, N., 1973b. Türkiye'nin deniz algleri. *IV Biyoloji Kongresi*, Ankara, 214: 1-3.
- <sup>177</sup>ZEYBEK, N. & GÜNER, H., 1973. Çanakkale Bogazi ve Bozcaada deniz algleri. *Ege Üniversitesi Fen Fakültesi Ilmi Raportlar Serisi*, 145: 1-19.
- <sup>178</sup>ZEYBEK, N., GÜNER, H. & AYSEL, V., 1986. The marine algae of Türkiye *Proceed. 5th OPTIMA Meeting*, Istanbul: 169-197.
- ZIBROWIUS, H., 1992. Ongoing modification of the Mediterranean marine fauna & flora by the establishment of exotic species. *Mésogée*, 51: 83-107.