

## Growth performance of the rabbitfish *Siganus sutor* raised at outdoor rearing ponds in Toliara, SouthWestern Madagascar

Ravelohasina Helga Berjulie<sup>1</sup>, Rasolofonirina Richard<sup>1</sup>, Rougeot Carole<sup>2</sup>, Jaonalison Henitsoa<sup>1</sup>, Mahafina Jamal<sup>1</sup>, Lepoint Gilles<sup>3</sup>, Maminantenaina Nandrianina<sup>1</sup> and Frédéric Bruno<sup>4</sup>

<sup>1</sup> Institut Halieutique et des Sciences Marines (IH.SM), University of Toliara, Madagascar, 171 Rue Fond Des Tawes, 4000 Liège, Belgium

E-mail: helga.ravelohasina@student.uliege.be

<sup>2</sup> Unity of Aquatic Resources Management and Aquaculture (UGeRAA), FOCUS, University of Liège, Belgium, 4000 Liège, Belgium

<sup>3</sup> Laboratory of Trophic and Isotopes Ecology, MARE Center, FOCUS, University of Liège, Belgium, 4000 Liège, Belgium

<sup>4</sup> Laboratory of Evolutionary Ecology, FOCUS, University of Liège, Belgium, 4000 Liège, Belgium

In Madagascar, decrease of fish stock and poverty lead small scale fishermen to deploy destructive fishing gear. At Toliara Reef, fishermen using mosquito seine net catch up to 50% of juvenile fish and up to 42% of these juveniles belong only to the rabbitfish *Siganus sutor* species. This prohibited fishing practice disturbs the structure of adult fish populations and aquaculture may be one of the effective solutions to alleviate the problem of stock depletion. Rabbitfishes possess most of the desirable characteristics for aquaculture (e.g. herbivory and responsive to artificial food, high survival in high rearing densities) and the culture of several *Siganus* species have been tested in various countries. However, no study has been conducted on the rearing of *S. sutor* in Madagascar. Here, we investigated such a possibility regarding the growth performances and survival rate of *S. sutor*. Juveniles (initial mean body weight of  $6.06\text{g} \pm 2.43$  and length of  $6.23\text{cm} \pm 0.8$ ) were successfully retrieved from fishermen catches and we applied fish grow-out experiments in outdoor rearing pond ( $16\text{m}^3$ ) for a duration of five months. The rearing density was 8 fish/ $\text{m}^3$  and the performance of different diets were tested. A locally produced industrial fish food (composed of corn, soya flour, wheat, vitamins, oil...) at a feed ration of 8% of fish biomass were compared with cooked rice at a feed ration of 25% of fish biomass. Dissolved oxygen, temperature, salinity, turbidity and pH were recorded three times a day along the experiment. Fish fed with industrial fish food had a significant higher growth ( $p < 0.05$ ) with a final mean body weight of  $68.11\text{g} \pm 26.71$  and a final mean length of  $13.91\text{cm} \pm 1.69$ . At the end of the experiment, fish fed with cooked rice displayed a mean body weight of  $34.81\text{g} \pm 15$  and a mean length of  $11.15\text{cm} \pm 1.76$ . Feed conversion ratio is 2.99 for industrial food while it reaches up to 20.67 for the cooked rice. Surprisingly, fish fed with industrial food showed a significant lower survival rate ( $p < 0.05$ ) of 47.2%, compared to 95.7% for the cooked rice. Our study highlights the possibility of rearing juvenile rabbitfish in outdoor rearing pond using essentially locally produced industrial food. Controlling the rearing of juvenile rabbitfishes can be suggested as a practical approach to improve the survival rate of these juveniles through sea ranching and for improving livelihood of local communities through alternative activity.

### Keywords

Growth; Survival Performances; *Siganus Sutor*; Siganidae; Juveniles; Aquaculture