

Ecopath model of the Southern Bight of the North Sea (version 2)

Steven Pint¹, Martha Stevens¹, Marleen De Troch², Dick van Oevelen³, Johanna Jacomina Heymans⁴, Gert Everaert¹

¹ Flanders Marine Institute (VLIZ), Wandelaarkaai 7, B-8400, Ostend, Belgium

² Ghent University (Marine Biology), Campus Sterre Krijgslaan 281-S8, B-9000 Ghent, Belgium

³ Royal Netherlands Institute for Sea Research (NIOZ, EDS), Korringaweg 7 4401 NT, Yerseke, Netherlands

⁴ University of the Highlands and Islands (SAMS, UHI), Oban, Argyll PA37 1QA, Scotland

Contents

Contents	1
Preface	4
1. Basic input parameters	4
1.1. Introduction to Southern Bight of the North Sea	4
1.2. Data sources for the Southern Bight of the North Sea model	5
1.3. Functional group input parameters	6
1.3.1. Marine mammals	7
Harbour porpoise	7
Seals	7
1.3.2. Seabirds	8
Discard feeders	8
Non-discard feeders	8
1.3.3. Fish	8
Cod	9
Sharks	9
Rays	9
Whiting	9
Sea Bass	10
Demersal fish	10
Herring	10
Sprat	10
Mackerel	10
Horse mackerel	11
Sandeels	11

Plaice	11
Dab	11
Sole	11
Other flatfish	12
Other gadoids	12
Miscellaneous filter feeding pelagic fish	12
1.3.4. Invertebrates	12
Squid & cuttlefish	12
Large crabs + shrimps	12
Benthos	13
Crangon crangon	13
Blue mussels (aquaculture)	13
1.3.5. Plankton	13
Carnivorous zooplankton	13
Herbivorous & omnivorous plankton (copepods)	13
Gelatinous zooplankton	13
Phytoplankton	14
1.3.6. Detritus	14
Discards	14
1.4. Diets	14
2. Defining fleets and assigning landings and discards	17
2.1. Fleets (commercial) distinguished by STECF gear types	17
2.2. Assigning 1991 landings to fleets	17
2.3. Calculating discards for 1991	17
3. Preliminary results	18
4. Balancing of the Ecopath model	19
4.1. Pre-balance diagnostics	19
4.1.1. Biomass	20
4.1.2. Production/biomass	21
4.1.3. Consumption/biomass	21
4.1.4. Production/consumption	22
4.1.5. Respiration/assimilation	22
4.2. Revision of the input data	23

4.2.1. Herring	23
4.2.2. Sharks	23
4.2.3. Plaice	23
4.2.4. Demersal fish	23
4.2.5. Other gadoids	23
4.2.6. Sprat	23
4.2.7. Benthos	23
4.2.8. Cod	23
4.2.9. Horse mackerel	23
4.2.10. Whiting	23
4.2.11. Large crabs and shrimps	24
4.2.12. Other flatfish	24
4.2.13. Sea bass	24
4.2.14. Squid and cuttlefish	24
4.2.15. Gelatinous zooplankton	24
4.2.16. Herbivorous and omnivorous plankton	24
4.2.17. Dab	24
4.2.18. Sole	24
4.2.19. Sandeels	24
4.2.20. Discards	24
4.3. Balanced model	25
4.4. Post-balance diagnostics	29
4.1.1. Biomass	29
4.1.2. Production/biomass	30
4.1.3. Consumption/biomass	30
4.1.4. Production/consumption	31
Abbreviations	32
Data availability	33
Citation	33
References	33

Preface

In this technical report, we describe the required data, the pre-processing steps taken and the methodologies used to develop a fully calibrated ecosystem model for the southern part of the North Sea. The ecosystem model was developed in both Ecopath with Ecosim (EwE) and Rpath, which is a R implementation of the commonly used EwE software. The model will serve as a tool to explore the ecosystem's natural history and its evolution throughout the years. A first version of the model was developed between July 2020 and February 2021.

Ecopath with Ecosim (EwE) is an ecosystem modelling software (<https://ecopath.org/>) established in 1984 by Polovina (1984) and updated through the years by Christensen and Pauly (1992) and Walters et al. (1997). It has three main components: 1) Ecopath: a mass-balance snapshot of the ecosystem; 2) Ecosim: a time dynamic simulation of the system; and 3) Ecospace: a spatial and time dynamic module.

In this technical report, we describe an Ecopath model for the Southern Bight of the North Sea (SBNS; ICES area IVc) in 1991. The year of the model, 1991, was chosen for three main reasons: 1) it is the year with the most complete diet data, i.e. North Sea Year of the Stomach; 2) it is the base year in the “mother model”, the North Sea model of Mackinson and Daskalov (2007); and 3) the period between 1991 and 2023 is a long time span for time series fitting in Ecosim to understand the influence of parameters, such as vulnerability.

We provide an overview of our basic input parameters, the fisheries included in this model, our data sources, and methods to estimate basic input parameters. Steps undertaken to assess and balance the model are described in this manuscript as well. We are using an ecosystem modelling approach, i.e. an Ecopath with Ecosim model of the SBNS, to address our research question: “What does the food web in the SBNS look like, and how did it evolve over time?”.

1. Basic input parameters

1.1. Introduction to Southern Bight of the North Sea

We developed an ecosystem model for the Southern Bight of the North Sea (SBNS). The SBNS (ICES area IV division c) has a surface area of 63,633.86 km² (ICES Spatial Facility, Fig. 1a). It is a relatively shallow (<50 m) part of the North Sea (Fig. 1b). The southern limit of the SBNS is determined by the English Channel and its northern limit is at the latitude of the island Schiermonniksoog, Netherlands. Apart from extreme observations, surface water temperatures vary seasonally between 5°C and 20°C. The salinity varies between 29 to 35 PSU and is strongly influenced by the inflow of Atlantic water (offshore) and the river plumes of the Scheldt, Rhine, Seine, and Meuse in the coastal zones (Lacroix et al., 2004). An anticlockwise circulation in the North Sea brings water from the North Atlantic along the coast of the UK in the west of the SBNS and water from the English Channel in the east, along the Belgian and Dutch coast (Turrell, 1992). The sediment in SBNS consists mainly of sands (Fig. 1c, ICES, 2008).




Figure 1: The geographical location of our study area (a), the Southern Bight of the North Sea (ICES area IVc, in red) adjusted from SWD/2016/0272 final - 2016/0238 (COD) and the environmental setting, (b) bathymetry of the North Sea (source RIVO – alternative from ICES, 2008) and (c) seabed sediment types of the North Sea. http://www.awi-bremerhaven.de/GEO/Marine_GIS/Margis%20homepage/index.html.

1.2. Data sources for the Southern Bight of the North Sea model

Ecopath models are mass-balance snapshots of the ecosystem, and the input data are expressed in mass-based units. To feed a SBNS specific model, we collected a range of data from multiple sources. Here, we give an overview of how we estimated input parameters, e.g. biomass (wet mass in $\text{t km}^{-2} \text{y}^{-1}$) and Q/B (Fig. 2). To estimate the biomass for our study area, we relied on biomass estimates from stock assessments and catch per unit effort (CPUE) data. For most commercial fish species, biomass data is available from ICES expert group reports such as HAWG, WGwide and WGNSSK. For species that were not included in ICES expert group reports, e.g. sea bass, seabirds and brown shrimp, estimates from research papers were used instead. In the specific case of harbour porpoise, we used an abundance estimate from OSPAR and the average weight per individual (Bjørge and Tolley, 2009). For some groups, such as plankton, we did not find a good estimate of the biomass in the SBNS. Hence, we assumed similar biomass compared to the study area of the other EwE models, the North Sea model (Mackinson and Daskalov, 2007) and the Southern North Sea (SNS; Stäbler et al., 2016). CPUE data was collected from the International Bottom Trawl survey in the North Sea (NS-IBTS, ICES Database of Trawl Surveys (DATRAS), 2020; https://datras.ices.dk/Data_products/Download/Download_Data_public.aspx).




Figure 2: An overview of estimating the biomass and other parameters for the Southern Bight of the North Sea (SBNS) EwE model.

Catch related data was obtained from STECF (mainly for information about fishing gear types, landings, and discards) and ICES (mainly for historical catch data). The diets in the SNS model of Stäbler et al. (2016) were adjusted to our study area. Diets of the species that were not included in the SNS model of Stäbler et al. (2016) were taken from CEFAS fish stomach records (<https://www.cefas.co.uk/data-and-publications/fish-stomach-records/>) and scientific literature. Detailed information concerning data sources for the input data in our model is described in the following sections.

1.3. Functional group input parameters

We determined the functional groups (FGs), i.e. a biomass groups representing one or multiple species, based on the SNS model of Stäbler et al. (2016). The ecosystem model of Stäbler et al. (2016) was focused on ICES areas IVb and IVc. Functional groups that had a low occurrence in our study area (ICES area IVc), based on the ICES ecosystem map (<https://ecosystemdata.ices.dk/Map/index.aspx?Action=AddLayer&DataSet=657&LatN=&LatS=&LonE=&LonW=&Sdate=&Edate>), were left out of our model. Other functional groups, i.e. functional groups that are not within our focus or have less economic value, were clustered together to decrease the complexity in the model. We obtained a total of 38 functional groups, for which an overview of the input data and their sources can be consulted in Table 1.

Table 1: An overview of the input parameters and references per functional group. The grey shaded area indicates a functional group with a multi-stanza.

Functional group	Input parameters			Data sources
Harbour porpoise	0.03	0.02	17.63	Bjørge and Tolley (2009), Hammond et al. (2017; OSPAR-SCAN III), Stäbler et al. (2016)
Seals	0.008	0.09	26.84	Mackinson and Daskalov (2007)
Seabirds (discard feeders)	0.0004	0.1	60.52	Bicknell et al. (2013), Mackinson and Daskalov (2007), Nilsson and Nilsson (1976), Potiek et al. (2019), Reeves and Furness (2002), Sherley et al. (2020), Waggitt et al. (2020)
Seabirds (non-discard feeders)	0.002	1.12	48.2	Bicknell et al. (2013), Mackinson and Daskalov (2007), Nilsson and Nilsson (1976), Potiek et al. (2019), Reeves and Furness (2002), Sherley et al. (2020), Waggitt et al. (2020)
Sharks	0.03	0.54	3.37	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Rays	0.81	0.30	2.89	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Cod				
Juvenile Cod	0.06	1.13	7.05	ICES (2019a)
Cod (adult)	0.16	1.13	3.20	Fishbase – Life history tool*, ICES (2019a; Table 4.11a), ICES – Historical Nominal Catches 1950-2010
Whiting				
Juvenile Whiting		2.80		ICES (2019a)
Whiting (adult)	0.16	1.31	5.41	Fishbase – Life history tool*, ICES (2019a; Table 23.19), ICES Database of Trawl Surveys (DATRAS)
Other gadoids	6.01	0.69	5.23	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Demersal fish	5.07	0.60	4.94	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Herring				
Juvenile Herring		0.87		ICES (2020)
Herring (adult)	0.78	0.87	6.78	Fishbase – Life history tool*, ICES (2020; Table 2.6.3.12), ICES Database of Trawl Surveys (DATRAS)
Sprat	0.21	0.80	9.01	ICES (2020; Table 10.6.4), ICES Database of Trawl Surveys (DATRAS)
Mackerel	0.51	1.34	4.81	ICES (2014; Table 3.6.3.2), ICES Database of Trawl Surveys (DATRAS)
Horse mackerel	0.68	0.34	4.96	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Sandeels	12.81	1.00	9.59	ICES (2020; Table 9.2.10)
Plaice				
Juvenile Plaice		0.49		ICES (2019a)
Plaice (adult)	0.51	0.89	3.63	Fishbase – Life history tool*, ICES (2019a; Table 13.3.4), ICES Database of Trawl Surveys (DATRAS)
Dab	5.14	0.46	5.58	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Other flatfish	0.08	0.84	6.90	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Sole				
Juvenile Sole		0.54		ICES (2019a)
Sole (adult)	0.32	1.15	5.56	Fishbase – Life history tool*, ICES (2019a; Table 17.10.1), ICES Database of Trawl Surveys (DATRAS)
Sea Bass	0.004	0.56	3.81	ICES Database of Trawl Surveys (DATRAS), Sparholt (1990)
Pelagic fish		4.00	10.19	0.98 Mackinson and Daskalov (2007)
Squid & cuttlefish	0.130	11.02	20.00	ICES Database of Trawl Surveys (DATRAS), Mackinson and Daskalov (2007)
Carnivorous zooplankton		4.00		0.32 0.99 Mackinson and Daskalov (2007)
Herbivorous zooplankton	16.00	9.20	30.00	Mackinson and Daskalov (2007)
Gelatinous zooplankton	0.090	2.86	0.45	Stäbler et al. (2016)
Large crabs & shrimp	2.300	1.05	0.2	Mackinson and Daskalov (2007), OBIS**, Stäbler (2016)
Blue mussels (aquaculture)	0.055	0.36	2.67	Horn et al. (2020)
Benthos	164.540	1.06	0.27	BenBioDen database, Stäbler et al. (2016)
C. crangon				
Crangon (below 5cm)		12.50		Hufnagl et al. (2010a), Stäbler et al. (2016)
Crangon (commercial size)	0.360	6.50	10.00	Hufnagl et al. (2010a), Tulp et al. (2016), Stäbler et al. (2016)
Phytoplankton	7.500	286.70		Mackinson and Daskalov (2007)
Detritus	50.000			Mackinson and Daskalov (2007)
Discards	0.00			Stäbler et al. (2016)

* Asila and Ogari (1988); Beddington and Cooke (1983); Beverton and Holt (1957); Beverton and Holt (1964); Beverton (1992); Bluweiss et al. (1978); Froese and Binohlan (2000); Froese et al. (in prep.); Gulland (1971); Musick (1999); Palomares (1991); Palomares and Pauly (1999); Pauly (1979); Pauly (1980); Pauly (1984); Pauly (1986); Pauly (1989); Pauly and Christensen (1998); Pauly et al. (1998); Ricker (1975); Taylor (1958)
 ** Brodie et al. (2013); Buhl-Mortensen (2014); DFO (2016); Hassel (2014); ICES (2010); IFREMER (2016); Libby (2014); MBA (2016); Meurisse and Semal (2020); Miller et al. (2014); National Museum of Natural History (2001); NOAA's National Marine Fisheries Service (2005); Northeast Fisheries Science Center (2010); OBIS Canada (2011); Parr (n.d.); Rees et al. (n.d.); Santos (2016); Swedish county administration boards et al. (2017); The Norwegian Oil Industry Association (2001); Turkey (n.d.); Van Guelpen (2016); Van Guelpen and Pohle (2014); VLIZ (2004)

1.3.1. Marine mammals

Three marine mammals regularly occur in the SBNS: harbour porpoise (*Phocoena phocoena*), common seal (*Phoca vitulina*) and grey seal (*Halichoerus grypus*). Other species such as the bottlenose dolphin and minke whale are uncommon in this area and sighted rarely. As they are irregular inhabitants of the SBNS and more likely to just travel through or arrive by accident, they are not included in the model.

Harbour porpoise

A density of 0.607 individuals km⁻² was obtained from OSPAR – SCANS III (Hammond et al., 2017) and multiplied with the average weight per individual of about 0.055 t (Bjørge and Tolley, 2009). We calculated a biomass of 0.03 t km⁻² y⁻¹ for our model. Production-biomass ratio (P/B) of 0.02 y⁻¹ and consumption-biomass ratio (Q/B) of 17.63 y⁻¹ for harbour porpoise were assumed to be similar to the values of toothed whales from the SNS model of Stäbler et al. (2016).

Seals

We assumed that the parameters, such as biomass, P/B, and Q/B for seals, i.e. grey and common seal were the same as in the North Sea model of Mackinson and Daskalov (2007). Thus, the biomass for seals was 0.008 t km⁻² y⁻¹, P/B 0.09 y⁻¹ and Q/B 26.84 y⁻¹.

1.3.2. Seabirds

We divided the seabirds into two functional groups, i.e. discard feeders and non- discards feeders. We did this because discards can be a large portion of some species' diets. Large flock of seabirds are often observed feeding on discards thrown overboard by fishermen. Other seabirds, such as divers and northern gannets do not feed on discards (or do so to a limited extent).

Discard feeders

Discard feeders consist of seabirds whose diet relies for a large part on discards. The following species were included in this group: European herring gull (*Larus argentatus*), great black-backed gull (*Larus marinus*), lesser black-backed gull (*Larus fuscus*), common gull (*Larus canus*), black-legged kittiwake (*Rissa tridactyla*), and great skua (*Stercorarius skua*). The biomass of discard feeders was estimated to be 0.0004 t km⁻² y⁻¹ based on scientific literature from Bicknell et al. (2013), Potiek et al. (2019), Reeves and Furness (2002), Sherley et al. (2020) and Waggitt et al. (2020). P/B was taken from the North Sea model of Mackinson and Daskalov (2007) and estimated to be 0.10 y⁻¹, whereas Q/B was estimated using the empirical formula of Nilsson and Nilsson (1976) resulting in 60.52 y⁻¹.

Non-discard feeders

Non-discard feeders consist of seabirds where discards do not (or to a limited extend) contribute to their diet. The following species were included in this group: Great crested grebe (*Podiceps cristatus*), Northern gannet (*Morus bassanus*), little gull (*Hydrocoloeus minutus*), common tern (*Sterna hirundo*), common murre (*Uria aalge*), razorbill (*Alca torda*), scoter sp. (*Melanitta sp.*), loon (Gaviidae) and tern (Sternidae). The biomass of non-discard feeders was estimated at 0.002 t km⁻² y⁻¹ based on several research papers: Bicknell et al. (2013), Potiek et al. (2019), Reeves and Furness (2002), Sherley et al. (2020) and Waggitt et al. (2020). P/B (1.12 y⁻¹) was taken from the North Sea model (Mackinson and Daskalov, 2007), and Q/B was estimated with the empirical formula of Nilsson and Nilsson (1976) resulting in 48.20 y⁻¹.

1.3.3. Fish

The functional groups containing fish were chosen based on their importance for fisheries, both commercial and recreational. The P/B of fish was calculated using equation 1. The total mortality (Z) equals fishing mortality (F) plus the natural mortality (M, Eq. 1). Fishing mortality is equal to the catch (C) over biomass (B, Eq. 2). The natural mortality is calculated according to the empirical formula of Pauly (1980) using asymptotic length (L_{inf} ; Eq. 3) or asymptotic weight (W_{inf} ; Eq. 4) depending on which parameter was known.

$$\frac{P}{B} = Z = F + M \quad (1)$$

$$F = \frac{C}{B} \quad (2)$$

$$\log M = -0.0066 - 0.279 \log L_{inf} + 0.6543 \log K + 0.4634 \log T \quad (3)$$

$$\log M = -0.2107 - 0.0824 \log W_{inf} + 0.6757 \log K + 0.4627 \log T \quad (4)$$

In equation 3 and 4, K is the growth coefficient and T is the temperature (°C), i.e. 10°C. Consumption-biomass ratio (Q/B; Eq. 5) was estimated using the empirical formula of Pauly et al. (1990). Pf (apex and/or pelagic predators and/or zooplankton feeders) is 1 for top predators and zooplanktivores and 0 for detritivores and herbivores. Hd (herbivores and detritivores) is 1 for herbivores and 0 for carnivores. The values for the parameters; K, L_{inf} and W_{inf} were obtained from Fishbase (Froese and Pauly 2023) through the Life history tool (Asila and Ogari, 1988); Beddington and Cooke, 1983; Beverton and Holt, 1957; Beverton and Holt, 1964; Beverton, 1992; Blueweiss et al., 1978; Froese and Binohlan, 2000; Froese et al., in prep.; Gulland, 1971; Musick, 1999; Palomares, 1991; Palomares and Pauly, 1999; Pauly, 1979; Pauly,

1980; Pauly, 1984; Pauly, 1986; Pauly, 1989; Pauly and Christensen, 1998; Pauly et al., 1998; Ricker, 1975; Taylor, 1958) for each species. For FGs composed of multiple species, we took a weighted mean of the species' parameters with their biomass as weight.

$$\log \frac{Q}{B} = 6.37 - 1.5045 \left(\frac{1000}{(T+273.1)} \right) - 0.168 \log W_{inf} + 0.1399 \log Pf + 0.2765 \log Hd \quad (5)$$

Table 1 provides an overview of all model parameter estimates.

Cod

We divided cod (*Gadus morhua*) into two stanza groups as they are a target species for fisheries, which focus on adult individuals. For adult cod (2+) we estimated the biomass to be 0.16 t km⁻² y⁻¹. We obtained this biomass estimate by multiplying the spawning stock biomass of cod in the greater North Sea (North Sea, English Channel and Skagerrak; Table 4.11a in ICES, 2019a) with the ratio of cod landings in the SBNS compared to those in the greater North Sea (ICES – Historical Nominal Catches 1950-2010). P/B and Q/B were calculated as 1.14 y⁻¹ and 3.20 y⁻¹ for adult cod. Total mortality of juvenile cod was calculated using F and M from ICES (2019a) and was estimated at 1.13 y⁻¹. For multi-stanza functional groups such as Cod, weight at maturity over asymptotic weight ($W_{maturity} / W_{infinity}$) which were retrieved from Fishbase – Life history tool (Asila and Ogari, 1988); Beddington and Cooke, 1983; Beverton and Holt, 1957; Beverton and Holt, 1964; Beverton, 1992; Blueweiss et al., 1978; Froese and Binohlan, 2000; Froese et al., in prep.; Gulland, 1971; Musick, 1999; Palomares, 1991; Palomares and Pauly, 1999; Pauly, 1979; Pauly, 1980; Pauly, 1984; Pauly, 1986; Pauly, 1989; Pauly and Christensen, 1998; Pauly et al., 1998; Ricker, 1975; Taylor, 1958), is required as basic input in Ecopath. For this, a value of 0.137 was obtained.

Sharks

The most common sharks in the SBNS are the spurdog (*Squalus acanthias*), school shark (*Galeorhinus galeus*), small-spotted catshark (*Scyliorhinus canicula*) and smooth-hound (*Mustelus spp.*). We estimated their biomass following the approach of Sparholt (1990). We calculated the shark-cod ratio of the SBNS using data from the International Bottom Trawl survey in the North Sea (NS-IBTS, ICES Database of Trawl Surveys (DATRAS)). We then multiplied the shark-cod ratio with the calculated biomass of cod. We obtained a biomass of 0.03 t km⁻² y⁻¹. The P/B was estimated to be 0.54 y⁻¹ and Q/B 3.37 y⁻¹.

Rays

Ray species included in this functional group are thornback ray (*Raja clavata*), spotted ray (*Raja montagui*), starry ray (*Amblyraja radiata*, low abundance), rajids unid., blue skate (*Dipturus batis*, low abundance) and cuckoo ray (*Leucoraja naevus*, low abundance). We estimated their biomass according to the approach of Sparholt (1990). We calculated the ray-plaice ratio of the SBNS using data of the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). Then we multiplied the ray-plaice ratio with the calculated total stock biomass of plaice. We obtained a biomass of 0.81 t km⁻² y⁻¹. The P/B was estimated to be 0.30 y⁻¹ and Q/B 2.90 y⁻¹.

Whiting

We divided whiting (*Merlangius merlangus*) in two stanza groups as they are a target species for fisheries. For adult whiting (2+), we estimated the biomass to be 0.16 t km⁻² y⁻¹. This estimate was obtained based on the ratio of whiting landings in the SBNS compared to the greater North Sea (NS-IBTS; ICES Database of Trawl Surveys (DATRAS)), which was then multiplied with the total spawning whiting stock biomass in the greater North Sea (North Sea, English Channel and Skagerrak; Table 23.19 in ICES, 2019a). P/B and Q/B were calculated as 1.31 y⁻¹ and 5.41 y⁻¹ for adult whiting. Total mortality of juvenile whiting was calculated using F and M from ICES (2019a) and was estimated at 2.80 y⁻¹. Maturity over asymptotic weight ($W_{maturity} / W_{infinity}$) for whiting was estimated at 0.182 (Fishbase – Life history tool; Asila and Ogari, 1988);

Beddington and Cooke, 1983; Beverton and Holt, 1957; Beverton and Holt, 1964; Beverton, 1992; Blueweiss et al., 1978; Froese and Binohlan, 2000; Froese et al., in prep.; Gulland, 1971; Musick, 1999; Palomares, 1991; Palomares and Pauly, 1999; Pauly, 1979; Pauly, 1980; Pauly, 1984; Pauly, 1986; Pauly, 1989; Pauly and Christensen, 1998; Pauly et al., 1998; Ricker, 1975; Taylor, 1958).

Sea Bass

We estimated sea bass' (*Dicentrarchus labrax*) biomass following the approach of Sparholt (1990). We calculated the 'sea bass-cod' ratio of the SBNS using data of the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). We then multiplied the 'sea bass-cod' ratio with the calculated biomass of cod. We obtained a biomass of 0.004 t km⁻² y⁻¹. The P/B was estimated to be 0.56 y⁻¹ and Q/B 3.81 y⁻¹.

Demersal fish

The demersal fish FG includes demersal fish, gurnards, and dragonets. The most common species in this FG are john dory (*Zeus faber*), eelpout (*Zoarces viviparus*), shorthorn sculpin (*Myoxocephalus Scorpius*), longspined bullhead (*Taurulus bubalis*), hooknose (*Agonus cataphractus*), striped sea snail (*Liparis liparis*), greater weever (*Trachinus draco*), lesser weever (*Echiichthys vipera*) and surmullet (*Mullus surmuletus*). We estimated their biomass following the approach of Sparholt (1990). We calculated the demersal-cod ratio of the SBNS using data from the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). Then, we multiplied the demersal-cod ratio with the calculated biomass of cod. We obtained a biomass of 5.07 t km⁻² y⁻¹. The P/B was estimated to be 0.60 y⁻¹ and Q/B 4.94 y⁻¹.

Herring

We divided herring (*Clupea harengus*) into two stanza groups as they are a target species for fishery. For adult herring (1+), we estimated the biomass to be 0.78 t km⁻² y⁻¹. This estimate was obtained based on the ratio of herring landings in the SBNS compared to the greater North Sea (NS-IBTS; ICES Database of Trawl Surveys (DATRAS)), which was then multiplied with the total spawning stock biomass of herring in the greater North Sea (North Sea, English Channel and Skagerrak; Table 2.6.3.12 in ICES, 2020). P/B and Q/B were calculated as 0.87 y⁻¹ and 6.78 y⁻¹ for adult herring. Total mortality of juvenile herring was calculated using F and M from ICES (2020) and was 0.87 y⁻¹. Maturity over asymptotic weight ($W_{maturity} / W_{infinity}$) for herring was 0.207 obtained from FishBase – Life history tool (Asila and Ogari, 1988); Beddington and Cooke, 1983; Beverton and Holt, 1957; Beverton and Holt, 1964; Beverton, 1992; Blueweiss et al., 1978; Froese and Binohlan, 2000; Froese et al., in prep.; Gulland, 1971; Musick, 1999; Palomares, 1991; Palomares and Pauly, 1999; Pauly, 1979; Pauly, 1980; Pauly, 1984; Pauly, 1986; Pauly, 1989; Pauly and Christensen, 1998; Pauly et al., 1998; Ricker, 1975; Taylor, 1958).

Sprat

Biomass was estimated based on the ratio of sprat (*Sprattus sprattus*) landings in the SBNS compared to the greater North Sea (NS-IBTS; ICES Database of Trawl Surveys (DATRAS)), which was then multiplied with the total spawning stock biomass of sprat in the greater North Sea (North Sea, English Channel and Skagerrak; Table 10.6.4 in ICES, 2020). We calculated a biomass of 0.20 t km⁻². P/B and Q/B were calculated as 0.80 y⁻¹ and 9.01 y⁻¹.

Mackerel

Mackerel (*Scrombus scrombus*) biomass was estimated based on the ratio of sprat landings in the SBNS compared to the greater North Sea (NS-IBTS; ICES Database of Trawl Surveys (DATRAS)), which was then multiplied with the total spawning stock biomass of sprat in the greater North Sea (North Sea, English Channel and Skagerrak). The greater North Sea spawning stock biomass required for this calculation was estimated for the North Sea by taking the North Sea portion (stock) of the spawning stock biomass in the

Atlantic (Table 3.6.3.2 in ICES, 2014). This resulted in a biomass estimate of $0.51 \text{ t km}^{-2} \text{ y}^{-1}$. P/B and Q/B were calculated as 1.34 y^{-1} and 4.81 y^{-1} .

Horse mackerel

We estimated the biomass of horse mackerel (*Trachurus trachurus*) following the approach of Sparholt (1990). We calculated the horse mackerel-whiting ratio of the SBNS using data from the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). Then we multiplied the horse mackerel-whiting ratio with the calculated biomass of whiting. We obtained a biomass of $0.68 \text{ t km}^{-2} \text{ y}^{-1}$. The P/B was estimated to be 0.34 y^{-1} and Q/B 4.96 y^{-1} .

Sandeels

Sandeel (*Ammodytidae*) biomass was estimated based on the ratio of sandeel landings in the SBNS compared to the North Sea (sandeel area 1r; ICES, 2020), which was then multiplied with the total stock biomass of sandeels in the North Sea (sandeel area 1r; Table 9.2.10 in ICES, 2020). This resulted in a biomass estimate of $12.81 \text{ t km}^{-2} \text{ y}^{-1}$. P/B and Q/B were calculated as 1.00 y^{-1} and 9.59 y^{-1} .

Plaice

We divided plaice (*Pleuronectes platessa*) into two stanza groups, as they are a target species for fisheries. For adult plaice (2+), we estimated the biomass to be $0.51 \text{ t km}^{-2} \text{ y}^{-1}$. This estimate was based on the ratio of plaice landings in the SBNS compared to the North Sea (NS-IBTS; ICES Database of Trawl Surveys (DATRAS)), which was then multiplied with the total stock biomass of plaice in the North Sea (North Sea and Skagerrak; Table 13.3.4 in ICES, 2019a). P/B and Q/B were calculated as 0.89 y^{-1} and 3.64 y^{-1} for adult plaice. Total mortality of juvenile plaice was calculated using F and M from ICES (2019a) and was 0.49 y^{-1} . A $W_{\text{maturity}} / W_{\text{infinity}}$ value of 0.147 was estimated for plaice using FishBase – Life history tool (Asila and Ogari, 1988); Beddington and Cooke, 1983; Beverton and Holt, 1957; Beverton and Holt, 1964; Beverton, 1992; Blueweiss et al., 1978; Froese and Binohlan, 2000; Froese et al., in prep.; Gulland, 1971; Musick, 1999; Palomares, 1991; Palomares and Pauly, 1999; Pauly, 1979; Pauly, 1980; Pauly, 1984; Pauly, 1986; Pauly, 1989; Pauly and Christensen, 1998; Pauly et al., 1998; Ricker, 1975; Taylor, 1958).

Dab

We estimated dab's (*Limanda limanda*) biomass following the approach of Sparholt (1990). We calculated the dab-plaice ratio of the SBNS using data from the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). Then, we multiplied the dab-plaice ratio with the calculated total stock biomass of plaice. We obtained a biomass of $5.14 \text{ t km}^{-2} \text{ y}^{-1}$. The P/B was estimated to be 0.46 y^{-1} and Q/B 5.58 y^{-1} .

Sole

We divided sole (*Solea solea*) in two stanza groups as they are a target species for fisheries. For adult sole (2+), we estimated the biomass to be $0.32 \text{ t km}^{-2} \text{ y}^{-1}$. This estimate was based on the ratio of sole landings in the SBNS compared to the North Sea (NS-IBTS; ICES Database of Trawl Surveys (DATRAS)), which was then multiplied with the total stock biomass of sole in the North Sea (North Sea; Table 17.10.1 in ICES, 2019a). P/B and Q/B were calculated at 1.15 y^{-1} and 5.56 y^{-1} for adult sole. Total mortality of juvenile sole was calculated using F and M from ICES (2019a) and was 0.54 y^{-1} . A $W_{\text{maturity}} / W_{\text{infinity}}$ value of 0.185 was estimated for sole using FishBase – Life history tool (Asila and Ogari, 1988); Beddington and Cooke, 1983; Beverton and Holt, 1957; Beverton and Holt, 1964; Beverton, 1992; Blueweiss et al., 1978; Froese and Binohlan, 2000; Froese et al., in prep.; Gulland, 1971; Musick, 1999; Palomares, 1991; Palomares and Pauly, 1999; Pauly, 1979; Pauly, 1980; Pauly, 1984; Pauly, 1986; Pauly, 1989; Pauly and Christensen, 1998; Pauly et al., 1998; Ricker, 1975; Taylor, 1958).

Other flatfish

The other common flatfish in the SBNS (besides plaice, dab and sole) are European flounder (*Platichthys flesus*), lemon sole (*Microstomus kitt*), turbot (*Scophthalmus maxima*) and brill (*Scophthalmus rhombus*). We estimated their biomass according to the approach of Sparholt (1990). We calculated the 'other flatfish-plaice' ratio of the SBNS using data from the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). Then, we multiplied the 'other flatfish-plaice' ratio with the calculated total stock biomass of plaice. We obtained a biomass of 0.08 t km⁻² y⁻¹. The P/B was estimated to be 0.83 y⁻¹ and Q/B 6.90 y⁻¹.

Other gadoids

The FG of the other gadoids consists of fivebeard rockling (*Ciliata mustela*), poor cod (*Trisopterus minutus*), pouting (*Trisopterus luscus*), three-bearded rockling (*Gaidropsarus vulgaris*, low abundance), fourbeard rockling (*Enchelyopus cimbrius*, low abundance) and silvery pout (*Gadiculus argenteus*, low abundance). We estimated their biomass following the approach of Sparholt (1990). We calculated the 'other gadoid-cod' ratio of the SBNS using data from the NS-IBTS (ICES Database of Trawl Surveys (DATRAS)). We then multiplied the 'other gadoid-cod' ratio with the calculated biomass of cod. We obtained a biomass of 6.01 t km⁻² y⁻¹. The P/B was estimated to be 0.69 y⁻¹ and Q/B 5.23 y⁻¹.

Miscellaneous filter feeding pelagic fish

The miscellaneous filter feeding group consist of shad (*Alosa* sp.), anchovy (*Engraulis encrasicolus*) and European pilchard (*Sardina pilchardus*). We could not estimate the biomass of miscellaneous filter feeding pelagic fish with the data available to us. Instead, we took the values of the North Sea model of Mackinson and Daskalov (2007) for which P/B is 4.00 y⁻¹ and the Q/B is 10.19 y⁻¹. Trophic efficiency was set to 0.98.

1.3.4. Invertebrates

Squid & cuttlefish

We included three species of squid and cuttlefish in this FG, i.e. veined squid (*Loligo forbesii*), European squid (*Loligo vulgaris*) and common cuttlefish (*Sepia officinalis*). We estimated the biomass for squid and cuttlefish by calculating the mean ratio from 1999 until 2018 between the abundance in the SBNS and the greater North Sea based on the ICES NS-IBTS data and multiplying this ratio with the total biomass of the North Sea model (Mackinson and Daskalov, 2007). The resulting biomass was 0.13 t km⁻² y⁻¹. We based P/B (11.02 y⁻¹) and Q/B (20.00 y⁻¹) on the North Sea model (Mackinson and Daskalov, 2007).

Large crabs + shrimps

Common large crabs with commercial value in the SBNS are edible crab (*Cancer pagurus*), common spider crab (*Maja brachydactyla*) and velvet swimming crab (*Necora puber*). Note that European lobster (*Homarus gammarus*) is also included in this FG. Shrimp species included are *Crangon allmanni*, *Eualus pusiolus*, *Pandalus montagui*, *Spirontocaris lilljeborgi*, *Processa nouveli*, and *Pandalina* spp. We estimated the biomass of shrimp species by calculating the abundance ratio of the SBNS and North Sea (Brodie et al., 2013; National Museum of Natural History, 2001; Northeast Fisheries Science Center, 2010; Türkay, n.d.; NOAA's National Marine Fisheries Service, 2005; DFO, 2016; Hassel, 2014; Swedish county administration boards et al., 2017; Buhl-Mortensen, 2014; Parr, n.d.; Rees et al., n.d.; Miller et al., 2014; ICES, 2010); Van Guelpen, 2016; OBIS Canada, 2011; Santos, 2016; Meurisse and Semal, 2020; MBA, 2016; IFREMER, 2016; VLIZ, 2004; Libby, 2014; Van Guelpen and Pohle, 2014; The Norwegian Oil Industry Association, 2001), which we then multiplied with the best estimate biomass of Mackinson and Daskalov (2007, Table 11.8). *Pandalus borealis* and nephrops (*Nephrops norvegicus*) are not included in our model as they have a low occurrence in our study area. The total biomass of all shrimp species was added to the total large crab biomass, which we assumed to be equal to the entire North Sea (Mackinson and Daskalov, 2007). This resulted in an estimated biomass of 2.30 t km⁻² y⁻¹. For P/B, a weighted mean was taken from all relevant functional groups from Stäbler et al. (2016) (with biomass as weight) which resulted in an

estimate of 1.05 y^{-1} and production-consumption (P/Q) ratio, 0.20. For these estimates, we assumed that the ratio between the functional groups in the SBNS is similar to the ratio in the North Sea.

Benthos

Total dry weight of all benthos, (i.e. epifaunal macrobenthos, infaunal macrobenthos, small mobile epifauna, small infauna, sessile epifauna and meiofauna) was obtained from the BenBioDen database for the SBNS (Stratmann et al., 2020) and multiplied with a weight-to-weight conversion factor of 11.41 based on Ricciardi and Bourget (1998). This resulted in an estimated biomass for benthos of $164.54 \text{ t km}^{-2} \text{ y}^{-1}$. A weighted mean was taken from all relevant functional groups from Stäbler et al. (2016) (with biomass as weight) for P/B, which gave an estimate of 1.06 y^{-1} and a P/Q of 0.27. We assumed that the ratio between the functional groups in the SBNS is similar to the ratio in the North Sea.

Crangon crangon

We divided brown shrimp (*Crangon crangon*) into two stanza groups as they are a target species for fisheries. *Crangon crangon* larger than 50 mm were considered adults, because this is the length at recruitment for fishery. For adult *Crangon crangon* (> 50 mm), we based the biomass on a research study of Tulp et al. (2016) to be $0.36 \text{ t km}^{-2} \text{ y}^{-1}$. The total mortality (Z) of the adults, 6.50 y^{-1} , was based on Stäbler et al. (2016), after Hufnagl et al. (2010a). Z for juveniles, 12.50 y^{-1} , was estimated in the same manner (Hufnagl et al., 2010a. Stäbler et al., 2016). The Q/B was set to 10.00 y^{-1} (Stäbler et al., 2016). The von Bertalanffy growth function K for *Crangon crangon* was found to be 1.17 (Hufnagl et al., 2010a). The asymptotic length of *Crangon crangon* is 79.32 mm, length at 'maturity' is 50 mm length, i.e. length at recruitment to fisheries. From both, we calculated the corresponding weight according to Hufnagl et al. (2010b; section 3.1.4). The resulting weight at maturity over asymptotic weight ($W_{\text{maturity}} / W_{\text{infinity}}$) required as basic input to Ecopath is 0.24.

Blue mussels (aquaculture)

This functional group was introduced in the model for a case study in a later stage. We directed our attention to the cultivation of mussels in Belgium, commencing with a minimal initial biomass, i.e. 0.055 t km^{-2} . This was because there was no existing mussel aquaculture in Belgium in 1991. The P/B and Q/B were taken from an Ecopath model of the northern Wadden Sea (Horn et al., 2020), and are 0.36 y^{-1} and 2.67 y^{-1} respectively.

1.3.5. Plankton

Carnivorous zooplankton

In this FG we included krill species of the order Euphausiacea, e.g. *Thysanoessa inermis*, *Meganyctiphanes norvegica*. We assumed that the carnivorous zooplankton P/B, production-consumption ratio and ecotrophic efficiency in the SBNS were the same as in the North Sea (Mackinson and Daskalov, 2007). The P/B is 4.00 y^{-1} , the production-consumption ratio is 0.32, and the ecotrophic efficiency is 0.99.

Herbivorous & omnivorous plankton (copepods)

The herbivorous & omnivorous plankton consist of several copepod species, e.g. *Pseudocalanus elongatus*, *Paracalanus parvus*, *Microcalanus pusillus*, *Acartia spp.* and *Temora longicornis*. We assumed that the herbivorous & omnivorous zooplankton biomass, P/B and Q/B in the SBNS were similar to those of the North Sea model (Mackinson and Daskalov, 2007). The biomass is $16.00 \text{ t km}^{-2} \text{ y}^{-1}$, the P/B is 9.20 y^{-1} and the Q/B is 30.00 y^{-1} .

Gelatinous zooplankton

The gelatinous zooplankton are jellyfish, i.e. moon jellyfish (*Aurelia aurita*), blue jellyfish (*Cyanea lamarckii*) and lion's mane jellyfish (*Cyanea capillata*). We assumed that the gelatinous zooplankton

biomass, P/B and P/Q ratio in the SBNS were the same as in the SNS model (Stäbler et al., 2016). The biomass is $0.09 \text{ t km}^{-2} \text{ y}^{-1}$, the P/B is 2.86 y^{-1} and the production-consumption ratio is 0.45.

Phytoplankton

We assumed that the phytoplankton biomass and P/B in the SBNS were the same as for the entire North Sea (Mackinson and Daskalov, 2007). The biomass is $7.50 \text{ t km}^{-2} \text{ y}^{-1}$ and the P/B is 286.70 y^{-1} .

1.3.6. Detritus

We combined the two detritus groups (POM and DOM) from the North Sea model of Mackinson and Daskalov (2007). The sum of the biomass of these two groups resulted in the detritus biomass in our model: $50.00 \text{ t km}^{-2} \text{ y}^{-1}$.

Discards

The biomass of the discards in the SBNS were estimated at $0.0001 \text{ t km}^{-2} \text{ y}^{-1}$. We assumed that the discards biomass in our study area was equal to that of the SNS (Stäbler et al., 2016).

1.4. Diets

We based the diets of our functional groups on Stäbler et al. (2016), which were adjusted to fit our functional groups. To achieve this, the diets for some functional groups from Stäbler et al. (2016) were merged by taking the weighted mean of the functional groups' diets assuming that the biomass ratio between functional groups remained equal. The weight used for calculating the weighted mean was the biomass of the corresponding functional groups. For example, we grouped gurnards, dragonets, and small demersal fish into a single functional group, i.e. demersal fish, and thus a single diet. We used their corresponding biomass' as the weight for taking the weighted mean for each prey item. Several functional groups of the SNS model of Stäbler et al. (2016) did not occur in our model. We eliminated these functional groups from the diets of our functional groups and recalculated the diet proportions back to a total of one while keeping the diet items ratio equal. Note that there is one exception to our methodology to estimate diets: the diet of blue mussels. Their diet was based on Horn et al. (2020) and as filter feeders, they mainly feed on phytoplankton, detritus, and herbivorous zooplankton. An overview of all diets is provided in Table 2.

Table 2: An overview of the diets in the Southern Bight of the North Sea.

Prey \ predator	1	2	3	4	5	6	7	8	9	10	11	12
1 Harbour porpoise												
2 Seals												
3 Seabirds (discard)												
4 Seabirds (non-discard)												
5 Sharks					3.53E-02							
6 Rays					1.08E-02							8.58E-06
7 Juvenile Cod	5.68E-02	3.82E-02		2.44E-03		7.62E-03	6.22E-03	1.11E-03	9.83E-03	2.26E-03	1.11E-03	5.72E-03
8 Cod (adult)		7.69E-02		2.44E-03				9.52E-03				
9 Juvenile Whiting	4.72E-02	5.20E-03			2.17E-03	4.46E-03	1.75E-02	5.40E-02	9.61E-04	1.93E-02	3.12E-04	1.67E-02
10 Whiting (adult)	1.45E-01	8.34E-02			1.04E-01	4.04E-03		8.71E-02		1.34E-04		
11 Other gadoids	1.08E-01	3.36E-02	1.36E-01	2.91E-01	1.17E-01	2.98E-02	2.92E-02	2.67E-02	1.09E-03	7.57E-03	1.76E-02	1.68E-02
12 Demersal fish	2.12E-01	1.38E-01	1.45E-03	1.47E-01	5.45E-02	4.10E-02	5.33E-02	3.59E-02	6.08E-02	2.14E-02	1.96E-02	4.73E-02
13 Juvenile Herring	1.11E-03	1.04E-05	2.34E-02	2.44E-02	1.72E-03	3.27E-03	4.67E-03	1.69E-02	2.44E-03	2.90E-02		3.95E-04
14 Herring (adult)	3.34E-03	7.38E-03	2.34E-02	2.44E-02	1.84E-01	1.17E-02	3.22E-03	1.58E-01				
15 Sprat	1.06E-01	3.39E-03	4.69E-02	2.44E-03	3.74E-02	4.00E-03	1.87E-02	1.10E-02	1.43E-02	2.92E-01	4.05E-03	1.05E-02
16 Mackerel			4.69E-02	2.44E-03	9.92E-02			4.98E-05				3.37E-04
17 Horse mackerel		1.50E-02			1.68E-02		8.40E-04	2.99E-03	1.53E-03	9.88E-03		4.41E-05
18 Sandeels	1.83E-01	3.65E-01	7.83E-02	4.25E-01	3.49E-02	3.01E-01	1.07E-01	4.78E-02	1.62E-01	3.26E-01	1.04E-03	3.23E-01
19 Juvenile Plaice	1.90E-04			2.44E-03			1.73E-05	3.76E-02				
20 Plaice (adult)	9.45E-05	7.44E-02			5.22E-02	1.85E-03		4.63E-03				
21 Dab	2.41E-03	5.98E-02		2.44E-03	1.76E-02	1.16E-02	3.58E-03	2.17E-01	6.99E-04	1.19E-04	5.20E-03	2.67E-02
22 Other flatfish	3.62E-05	5.57E-02		2.44E-03				3.18E-03	3.27E-06			
23 Juvenile Sole	2.81E-04					1.07E-03	1.88E-03	3.49E-03	1.23E-03	5.33E-04		
24 Sole (adult)	1.37E-05	4.44E-02			1.34E-04	2.79E-03						3.26E-04
25 Sea Bass	1.63E-04	1.06E-04	1.11E-06	1.13E-04	4.19E-05	3.15E-05	4.10E-05	2.76E-05	4.67E-05	1.64E-05	1.50E-05	3.64E-05
26 Pelagic fish	1.09E-01		1.79E-02	4.93E-02	4.56E-04		1.16E-04	5.55E-05	4.93E-04	1.77E-05		
27 Squid & cuttlefish	2.32E-02			4.88E-03	5.88E-02	2.64E-03	6.14E-03	1.07E-02	8.37E-02	9.60E-03	8.32E-04	2.19E-02
28 Carnivorous zooplankton			1.23E-02	2.44E-03	5.89E-03	4.41E-03	2.26E-02	7.91E-04	1.32E-01	4.86E-02	1.05E-01	1.83E-02
29 Herbivorous plankton (copepods)			1.23E-02	2.44E-03		6.50E-07	6.50E-06		6.55E-03	3.09E-03	3.12E-01	5.98E-02
30 Gelatinous zooplankton					3.39E-02	0.00E+00	1.79E-03	1.00E-03	4.57E-03	4.05E-03		
31 Large crabs + shrimps					3.69E-02	1.97E-01	1.57E-01	1.08E-01	1.33E-01	4.39E-02	1.78E-02	1.04E-01
32 Blue mussels (aquaculture)		1.15E-04	1.00E-04	1.00E-04		1.06E-04		1.13E-04		1.10E-04		1.01E-04
33 Benthos	3.03E-03		6.53E-02	9.76E-03	8.86E-02	3.48E-01	4.00E-01	1.39E-01	3.01E-01	1.65E-01	5.17E-01	3.41E-01
34 Crangon (below 5cm)					5.91E-03	1.71E-02	1.26E-01	9.44E-03	7.48E-02	1.51E-02	2.57E-04	5.93E-03
35 Crangon (commercial size)					2.26E-03	6.54E-03	3.96E-02	3.44E-03	1.74E-02	3.21E-03	9.79E-05	1.83E-03
36 Phytoplankton										2.54E-05		
37 Detritus												
38 Discards			4.69E-01									
Import			6.64E-02	2.44E-03								
Sum	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
(1 - Sum)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 2: An overview of the diets in the Southern Bight of the North Sea. (continued)

Prey \ predator	13	14	15	16	17	18	19	20	21	22	23	24
1 Harbour porpoise												
2 Seals												
3 Seabirds (discard)												
4 Seabirds (non-discard)												
5 Sharks												
6 Rays												
7 Juvenile Cod				1.98E-03						5.16E-03		
8 Cod (adult)												
9 Juvenile Whiting				5.73E-05	3.33E-02					1.01E-03		
10 Whiting (adult)												
11 Other gadoids				3.96E-04	4.28E-03					3.05E-03		
12 Demersal fish				1.45E-03	7.15E-02					2.08E-02	1.28E-02	
13 Juvenile Herring				1.27E-02	1.31E-01		2.03E-04			1.26E-04		
14 Herring (adult)												
15 Sprat				2.92E-02	3.26E-02					1.53E-02		
16 Mackerel				8.04E-06								
17 Horse mackerel				1.02E-01	8.44E-02							
18 Sandeels				1.20E-01	2.00E-02		7.63E-03			1.63E-02		
19 Juvenile Plaice												
20 Plaice (adult)												
21 Dab					6.07E-03					5.33E-02		
22 Other flatfish										3.42E-02		
23 Juvenile Sole												
24 Sole (adult)												
25 Sea Bass				1.12E-06	5.49E-05					1.60E-05	9.82E-06	
26 Pelagic fish												
27 Squid & cuttlefish				8.10E-03	1.14E-01							
28 Carnivorous zooplankton	2.73E-01	3.09E-01		6.60E-02	1.70E-01	6.62E-02				4.01E-02		
29 Herbivorous plankton (copepods)	6.75E-01	6.70E-01	8.89E-01	4.85E-01	5.89E-02	6.41E-01			1.25E-02			
30 Gelatinous zooplankton				8.49E-02	8.48E-05			3.46E-03		5.55E-02		
31 Large crabs + shrimps				2.28E-02	7.10E-02			1.69E-02	1.09E-01	1.07E-02	4.44E-02	5.44E-02
32 Blue mussels (aquaculture)						1.05E-04	1.00E-04			1.00E-04	1.08E-04	
33 Benthos	5.19E-02	2.06E-02		4.36E-02	2.01E-01	2.70E-01	9.72E-01	8.81E-01	9.76E-01	7.07E-01	9.87E-01	9.46E-01
34 Crangon (below 5cm)				3.95E-04	2.24E-03		5.90E-05	8.60E-03	2.47E-04	2.38E-03		
35 Crangon (commercial size)					9.58E-04		1.00E-05	1.52E-03	9.50E-05	9.10E-04		
36 Phytoplankton			1.11E-01	2.21E-02		2.28E-02						
37 Detritus												
38 Discards												
Import												
Sum	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
(1 - Sum)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Prey \ predator	25	26	27	28	29	30	31	32	33	34	35	
1 Harbour porpoise												
2 Seals												
3 Seabirds (discard)												
4 Seabirds (non-discard)												
5 Sharks												
6 Rays												
7 Juvenile Cod				1.13E-02								
8 Cod (adult)												
9 Juvenile Whiting	2.92E-03		9.51E-04									
10 Whiting (adult)	2.05E-03											
11 Other gadoids	1.95E-02											
12 Demersal fish	2.07E-02		5.70E-03							5.07E-03		
13 Juvenile Herring			9.51E-04									
14 Herring (adult)												
15 Sprat			1.13E-02									
16 Mackerel	1.33E-01											
17 Horse mackerel	6.65E-02		1.13E-02									
18 Sandeels	2.98E-03		4.51E-02									
19 Juvenile Plaice										4.52E-04		
20 Plaice (adult)				5.70E-03								
21 Dab				5.70E-03								
22 Other flatfish	1.43E-02		1.14E-02							1.62E-03		
23 Juvenile Sole	1.07E-03											
24 Sole (adult)			5.70E-03									
25 Sea Bass	1.59E-05		4.38E-06							3.89E-06		
26 Pelagic fish	6.87E-02	5.31E-02	2.24E-02									
27 Squid & cuttlefish	8.77E-03	5.60E-03	5.63E-02			7.36E-02						
28 Carnivorous zooplankton		1.49E-01	2.31E-01	5.83E-02		2.94E-01	5.15E-03					
29 Herbivorous plankton (copepods)	2.09E-01	6.95E-01	3.05E-01	8.27E-01		2.94E-01	1.03E-02	1.00E-02		2.88E-03		
30 Gelatinous zooplankton		3.32E-02										
31 Large crabs + shrimps	3.71E-01	9.48E-03	1.44E-02			4.85E-02	2.56E-03			1.21E-02		
32 Blue mussels (aquaculture)												
33 Benthos	1.05E-03	4.99E-02	1.92E-01	1.15E-01		1.18E-01	8.98E-01		3.81E-01	9.37E-01	9.96E-01	
34 Crangon (below 5cm)	6.11E-02	3.56E-03	5.40E-03			1.82E-02	5.34E-04			4.14E-02	4.31E-03	
35 Crangon (commercial size)	9.03E-03	1.36E-03	2.07E-03			6.96E-03	2.04E-04					
36 Phytoplankton			5.63E-02		9.47E-01	1.47E-01		8.50E-01	5.19E-02			
37 Detritus					5.26E-02			2.78E-02	1.40E-01	5.68E-01		
38 Discards								5.52E-02				
Import	8.80E-03											
Sum	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
(1 - Sum)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

2. Defining fleets and assigning landings and discards

2.1. Fleets (commercial) distinguished by STECF gear types

Nine commercial EwE fleets were distinguished using STECF gear types. In total, 17 gear types were included in the STECF data for the SBNS (STECF, 2017; <https://stecf.jrc.ec.europa.eu/dd/effort/graphs-annex>). These gear types were allocated to their corresponding EwE fleet (Table 3).

Table 3: Structure of the EwE fleets in the Southern Bight of the North Sea model and the allocation of gear types

Gear description	Gear types	EwE fleet
Beam trawls ≥ 120 mm	BT1	
Beam trawls ≥ 80 mm and < 120 mm	BT2	beam
Beam trawls > 31 mm and < 80 mm or missing mesh size	BEAM	
Bottom trawls and seines ≥ 100 mm	TR1	
Bottom trawls and seines ≥ 16 mm and < 32 mm	TR3	
Bottom trawls and seines ≥ 70 mm and < 100 mm	TR2	demersal
Danish seine ≥ 90 mm	DEM_SEINE	
OTTER ≥ 32 mm and < 70 mm or missing mesh size	OTTER	
Dredges	DREDGE	dredge
Longlines	LL1	hooks
Gill nets, entangling nets	GN1	nets
Trammel nets	GT1	
Unspecified gear type	NONE	other
Pelagic seine (all mesh size)	PEL_SEINE	pelagic
Pelagic trawls (all mesh size)	PEL_TRAWL	
Pots	POTS	pots
Beam trawls < 31 mm and < 90 mm	BEAM	shrimp

2.2. Assigning 1991 landings to fleets

Catch data, i.e. landings and discards at age, from 2003 – 2017 for the SBNS (ICES are IVc) was obtained from STECF FDI (STECF, 2017; <https://stecf.jrc.ec.europa.eu/dd/effort/graphs-annex>). Historical catch data from ICES (<https://www.ices.dk/data/dataset-collections/Pages/Fish-catch-and-stock-assessment.aspx>) was obtained for the catches of 1991 in the SBNS. We applied methods used by Bentley et al. (2018) and Stähler et al. (2016) to estimate the landings and discards for the SBNS in 1991. Catches of both 1991 and 2003 were first allocated to the EwE FGs of the model. Next, the contribution of each fleet to the catches of 2003 was calculated. Because this fleet data was not available for 1991, the landings of the 1991 EwE FGs were multiplied with the contribution of fleets in 2003. This resulted in total landings biomass for each functional group and fleet. Note that for multi-stanza FGs, we assumed that all landings belonged to the adult group of that multi-stanza.

2.3. Calculating discards for 1991

We estimated the discards for 1991 by calculating the ratio of landings and discards in 2003 per fleet and per FG. We multiplied the landing-discard ratio of 2003 with the landings of 1991 per FG and per fleet to estimate the discards in 1991 assuming the ratio remained the same from 1991 to 2003. We also assumed that all discards in multi-stanza FGs belonged to the juvenile group of that FG.

2.4. Recreational fishery

We included Belgian recreational fisheries based on Verleye et al. (2019). We incorporated the recreational fleets (i.e. anglers (boat), trawlers, anglers (land), nets and others) grouped according to Verleye et al. (2019). As only data from 2018 onwards is available for Belgian recreational fisheries, the recreational catch for target species, such as cod, sea bass, mackerel, plaice, sole and brown shrimp was limited in the model.

3. Preliminary results

In this section, we present an overview of the preliminary food web obtained using the input data described in previous sections before the model was mass-balanced (Fig. 3).

At the bottom of the food web, i.e. trophic level 1, we have primary producers (phytoplankton), detritus and discards. At trophic level 2, we only have copepods that graze on the phytoplankton. A little higher in the food web, we have benthic invertebrates and zooplankton, i.e. carnivorous and gelatinous. As supported by scientific literature and expert knowledge, the benthos FG has a high biomass. In the middle of the food web, we observe fish species, such as plaice, withing and sea bass, and fish related FGs, e.g. flatfish, other gadoids and miscellaneous filter feeding fish. Adult cod has a higher trophic level than most fish related to his predatory diet. Top predators, at the top left of the food web (Fig. 3) occupy higher trophic levels, with exception of the discard-feeding seabirds. Discard-feeding seabirds have a lower trophic level because they feed on discards, i.e. a FG with a low trophic level. Fisheries act as predators in the model by preying on fish and other FGs. Fisheries will have a trophic level respective to their target species.




Figure 3: An illustration of the unbalanced Southern Bight of the North Sea food web model. Trophic level is indicated on the y-axis, ranging from 1 to 5. The size of circles represents the biomass included in that functional group. The ellipses show related functional groups. A. The functional groups at the bottom of the food web. B. The invertebrates in the food web with a trophic level higher than 2. C. Fish related functional groups in the food web. D. The top predators in the model have a trophic level between 4 and 5. E. The fishing fleets included in the food web.

4. Balancing of the Ecopath model

The previous chapter describes an unbalanced EwE model of the SBNS. Initially thirteen FGs were unbalanced when using the basic input parameters from chapter 1. Energy ‘imbalance’ is determined by examining each group’s ecotrophic efficiency (EE). A group’s EE exceeding 1 indicates that the energy demand placed upon that group exceeds its P/B and therefore needs to be reduced.

4.1. Pre-balance diagnostics

We used pre-balance (PREBAL) diagnostics as described in Link (2010) to identify ecological issues in the model and to balance it obeying the rules of thermodynamics and ecology. Group biomass should span 5 to 7 orders of magnitude and decline by 5 to 10% with each increasing trophic level. P/B and Q/B ratios should also show a general decline with increasing trophic level, whilst P/Q ratios should fall between 0.1 and 0.3 (Darwall et al., 2010, Heymans et al., 2016). Respiration/assimilation should be below 1 as the proportion of biomass lost through respiration cannot exceed the biomass of food assimilated (Heymans et al., 2016).

Table 4: Basic estimates from the initial unbalanced model of the Southern Bight of the North Sea. Numbers highlighted in red indicate unbalances, i.e. the ecotrophic efficiency is too high (>1).

Functional group	Input parameters						
	TL	B (t km ⁻²)	Z (y ⁻¹)	P/B (y ⁻¹)	Q/B (y ⁻¹)	EE	P/Q (y ⁻¹)
Harbour porpoise	4.6	0.03		0.02	24.64	0.00	0.0008
Seals	4.7	0.01		0.09	26.84	0.00	0.0034
Seabirds (discard)	3.1	0.0004		0.10	60.52	0.17	0.0017
Seabirds (non-discard)	4.5	0.0016		1.12	48.20	0.02	0.02
Sharks	4.6	0.03		0.54	3.37	0.87	0.16
Rays	4.1	0.81		0.30	2.89	0.13	0.10
Cod							
Juvenile Cod	4.2	0.06	1.13		7.05	2.04	0.16
Cod (adult)	4.5	0.16	1.13		3.20	0.91	0.35
Whiting							
Juvenile Whiting	4.2	0.43	2.80		13.41	0.99	0.21
Whiting (adult)	4.1	0.16	1.31		5.41	2.13	0.24
Other gadoids	3.5	0.09		0.85	5.23	3.98	0.16
Demersal fish	4.1	0.07		2.49	4.94	7.03	0.50
Herring							
Juvenile Herring	3.3	0.04	0.87		18.00	16.53	0.05
Herring (adult)	3.4	0.78	0.87		6.78	0.48	0.13
Sprat	2.9	0.21		0.80	9.01	4.42	0.09
Mackerel	3.7	0.51		1.34	4.81	0.72	0.28
Horse mackerel	4.3	0.68		0.34	4.96	2.77	0.07
Sandeels	3.2	12.81		1.00	9.59	0.23	0.10
Plaice							
Juvenile Plaice	3.6	0.06	0.49		9.39	8.62	0.05
Plaice (adult)	3.7	0.51	0.89		3.63	1.24	0.25
Dab	3.6	5.14		0.46	5.58	0.13	0.08
Other flatfish	3.9	0.08		0.84	6.90	2.09	0.12
Sole							
Juvenile Sole	3.6	0.14	0.54		9.74	0.39	0.06
Sole (adult)	3.7	0.32	1.15		5.56	0.57	0.21
Sea Bass	4.3	0.004		0.56	3.81	0.84	0.15
Pelagic fish	3.4	0.06		4.00	10.19	0.98	0.39
Squid & cuttlefish	3.7	0.13		11.03	20.00	0.78	0.55
Carnivorous zooplankton	3.1	3.88		4.00	12.50	0.99	0.32
Herbivorous plankton (copepods)	2.0	16.00		9.20	30.00	0.87	0.31
Gelatinous zooplankton	3.5	0.09		2.86	6.35	1.13	0.45
Large crabs + shrimps	3.5	2.30		1.05	5.27	1.11	0.20
Blue mussels (aquaculture)	2.0	0.06		0.36	2.67	0.83	0.13
Benthos	2.6	164.54		1.06	3.98	2.04	0.27
C. crangon							
Crangon (below 5cm)	3.7	0.58	12.50		25.14	0.17	0.50
Crangon (commercial size)	3.6	0.36	6.51		10.00	0.12	0.65
Phytoplankton	1.0	7.50		286.70		0.23	
Detritus	1.0	50.00				0.22	
Discards	1.0	0.00				0.32	

4.1.1. Biomass

The biomass (log scale) in the unbalanced SBNS model spans seven orders of magnitude, falling within the ecological range (5-7 orders) and declines by 3.8 %, falling below the ecological range (5-10%) suggested by Link (2010; Fig. 4). Several groups show variance from the trend line. The PREBAL ecological rules of thumb from Link et al. (2010) are applicable to poikilotherms and not so much to homeotherms. Marine mammals and seabirds consequently show large divergence. Benthos has a high biomass which can be explained by our choice of grouping all benthos together. Sandeels and dab also seem to have a high biomass and may be overestimated.




Figure 4: PREBAL diagnostic biomass. Functional groups are ordered by increasing trophic level.

4.1.2. Production/biomass

P/B (log scale) in the unbalanced SBNS model spans five orders of magnitude, falling within the ecological range (5-7 orders) and declines by 2.7 %, falling below the ecological range (5-10%) suggested by Link (2010; Fig. 5). Overall, there is little variance from the trend line apart from seabirds, marine mammals, phytoplankton, juvenile brown shrimp (*Crangon crangon*), and squid and cuttlefish. The homeotherms, i.e. seabirds and marine mammals have higher consumptive demands per unit of body mass than poikilotherms.




Figure 5: PREBAL diagnostic production/biomass. Functional groups are ordered by increasing trophic level.

4.1.3. Consumption/biomass

Q/B is expected to decline with increasing trophic level following basic energetic assumptions. Q/B (log scale) in the unbalanced SBNS model spans two orders of magnitude and declines by 0.3 % (Fig. 6). Blue mussels (aquaculture) and benthos have low Q/B ratios. The opposite is true for seabirds and marine mammals being homeotherms, which demands a high amount of energy. These deviances may affect the trend line.




Figure 6: PREBAL diagnostic consumption/biomass. Functional groups are ordered by increasing trophic level.

4.1.4. Production/consumption

There are several FGs falling outside the suggested P/Q range of 0.1 - 0.3 (Fig. 7). However, most fish groups fall within or close to the range, with few exceptions such as pelagic fish and demersal fish. There are invertebrate species such brown shrimp (*Crangon crangon*), squid and cuttlefish that have high P/Q. These ratios are similar to values found by Stähler et al. (2016). Nevertheless, it is advised to revise the ratios of the FGs that fall outside the advised range. Marine mammals and seabirds are slow producing and the energy from consumption is prioritised for other physiological processes such as thermoregulation.




Figure 7: PREBAL diagnostic production/consumption. Functional groups are ordered by increasing trophic level.

4.1.5. Respiration/assimilation

All respiration/assimilation values fall below 1.

4.2. Revision of the input data

By altering the basic input parameters and group diet composition based on the PREBAL diagnostics, we balanced the basic estimates. Groups with the highest EE estimates were adjusted first as suggested by (Heymans et al., 2016). By doing so, other imbalances may improve as well. The diets of functional groups were adjusted in case high cannibalism was present in the FG. The vast majority of imbalances in the model were corrected by adjusting functional group input parameters, either to fix ecological issues or to reduce EE estimates. The adjustments to our input parameters were performed keeping the mother models, i.e. North Sea (Mackinson and Daskalov, 2007), and SNS (Stähler et al., 2016), in mind. The changes made to the input parameters are listed below. The balanced food web is illustrated in Figure 8. Below in Table 5, we provide the input parameters of the balanced model of the SBNS. In Table 6, the revised diet composition is provided. In addition, the basic estimates are shown in Table 7.

4.2.1. Herring

Juvenile herring's total mortality was increased from 0.871 y^{-1} to 2.95 y^{-1} . Biomass adult herring increased from 0.778 t km^{-2} to 1.501 t km^{-2} and its total mortality from 0.869 y^{-1} to 1.55 y^{-1} .

4.2.2. Sharks

Biomass was increased from $0.00009 \text{ t km}^{-2}$ to 0.012 t km^{-2} and P/B from 0.542 y^{-1} to 1.012 y^{-1} .

4.2.3. Plaice

The total mortality of juvenile plaice increased from 0.462 y^{-1} to 1.75 y^{-1} and adult plaice from 0.893 y^{-1} to 1.12 y^{-1} .

4.2.4. Demersal fish

Increased biomass from 0.14 t km^{-2} to 1.24 t km^{-2} . P/B was decreased from 2.49 y^{-1} to 1.552 y^{-1} . Its diet composition was adjusted to decrease the cannibalism.

4.2.5. Other gadoids

P/B was increased from 0.772 y^{-1} to 1.6 y^{-1} and biomass from 0.16 t km^{-2} to 0.29 t km^{-2} .

4.2.6. Sprat

P/B was adjusted from 0.798 y^{-1} to 2.659 y^{-1} and its biomass was increased from 0.205 t km^{-2} to 0.5 t km^{-2} .

4.2.7. Benthos

The biomass of benthos was increased from 164.5 t km^{-2} to 300 t km^{-2} . Its diet composition was altered decreasing the cannibalism from 0.381 to 0.15.

4.2.8. Cod

The juvenile cod's Z was increased from 1.128 y^{-1} to 1.96 y^{-1} . Its Q/B was also increased from 7.05 y^{-1} to 8.079 y^{-1} . The total mortality of adult cod was increased from 1.13 y^{-1} to 1.135 y^{-1} .

4.2.9. Horse mackerel

The P/B of horse mackerel was increased from 0.2944 y^{-1} to 0.67 y^{-1} . In addition, its biomass was increased to 1.551 t km^{-2} .

4.2.10. Whiting

The biomass of adult whiting was increased from 0.162 t km^{-2} to 0.33 t km^{-2} and the P/B was increased from 1.309 y^{-1} to 1.352 y^{-1} .

4.2.11. Large crabs and shrimps

The biomass of this FG was increased from 2.3 t km⁻² to 5.1 t km⁻².

4.2.12. Other flatfish

The biomass and P/B of flatfish were increase from 0.0773 t km⁻² and 0.843 y⁻¹ to 0.1 t km⁻² and 1.4 y⁻¹ respectively.

4.2.13. Sea bass

Sea bass P/Q was set to 0.2 y⁻¹ to estimate its P/B by Ecopath.

4.2.14. Squid and cuttlefish

The diet composition of this FG was adjusted to decrease the high cannibalism from 0.0563 to 0.025. In addition, the diets of its predators were adjusted as well to decrease predation pressure. P/B was decreased to 4.5 and Q/B to 15.

4.2.15. Gelatinous zooplankton

We increased the biomass from 0.0912 t km⁻² to 0.125 t km⁻².

4.2.16. Herbivorous and omnivorous plankton

The biomass was increased from 16 t km⁻² to 17.2 t km⁻².

4.2.17. Dab

We increased the P/B from 0.463 y⁻¹ to 0.6 y⁻¹ as its P/Q was below 0.1.

4.2.18. Sole

We increased the P/B of juvenile sole from 0.5 y⁻¹ to 1.1 y⁻¹ as its P/Q was below 0.1.

4.2.19. Sandeels

We increased the P/B from 1.0 y⁻¹ to 2.1 y⁻¹ as its P/Q was low for a forage fish species. The biomass of sandeels was decreased from 12.81 t km⁻² to 5.565 t km⁻² as the initial estimation was high.

4.2.20. Discards

Biomass from 0.0001 t km⁻² to 50 t km⁻², using the biomass from Mackinson and Daskalov (2007).

4.3. Balanced model




Figure 8: Visual representation of the balanced food web model for the Southern Bight of the North Sea. Each functional group and fisheries fleet is represented with a circle indicating its estimated biomass, with trophic level represented on the y-axis. Boxes highlight related functional groups and fisheries fleets: (A) apex predators, (B) fish species, (C) zooplankton, (D) crabs and shrimp, (E) recreational fisheries and (F) commercial fisheries including mussel aquaculture.

Table 5: The input parameters per functional group of the balanced Southern Bight of the North Sea Ecopath model. The grey shaded area indicates a functional group with a multi-stanza.

Functional group	Input parameters					
	B (t km ⁻²)	Z (y-1)	P/B (y-1)	Q/B (y-1)	EE	P/Q (y-1)
Harbour porpoise	0.03		0.02	24.64		
Seals	0.01		0.09	26.84		
Seabirds (discard)	0.0004		0.10	60.52		
Seabirds (non-discard)	0.0016		1.12	48.20		
Sharks	0.01		1.01	3.37		
Rays	0.81		0.30	2.89		
Cod						
Juvenile Cod	0.11	1.96		8.08		
Cod (adult)	0.16	1.13		3.20		
Whiting						
Juvenile Whiting	0.93	2.80		13.33		
Whiting (adult)	0.33	1.35		5.41		
Other gadoids	0.29		1.60	5.23		
Demersal fish	1.24		1.55	4.94		
Herring						
Juvenile Herring	0.40	2.95		17.21		
Herring (adult)	1.50	1.55		6.78		
Sprat	0.50		2.66	9.01		
Mackerel	0.51		1.34	4.81		
Horse mackerel	1.55		0.67	4.96		
Sandeels	5.57		2.10	9.59		
Plaice						
Juvenile Plaice	0.21	1.75		10.12		
Plaice (adult)	0.51	1.12		3.63		
Dab	5.14		0.60	5.58		
Other flatfish	0.10		1.40	6.90		
Sole						
Juvenile Sole	0.19	1.10		10.46		
Sole (adult)	0.32	1.15		5.56		
Sea Bass	0.004			3.81		0.20
Pelagic fish			4.00	10.19	0.98	
Squid & cuttlefish	0.13		11.03	20.00		
Carnivorous zooplankton			4.00		0.99	0.32
Herbivorous plankton (copepods)	17.20		9.20	30.00		
Gelatinous zooplankton	0.13		2.86			0.45
Large crabs + shrimps	5.10		1.05			0.20
Blue mussels (aquaculture)	0.06		0.36	2.67		
Benthos	300.00		1.06			0.27
C. crangon						
Crangon (below 5cm)	0.58	12.50		25.14		
Crangon (commercial size)	0.36	6.51		10.00		
Phytoplankton	7.50		286.70			
Detritus	50.00					
Discards	50.00					

Table 6: The diet composition of the balanced Southern Bight of the North Sea Ecopath model.

Prey \ predator	1	2	3	4	5	6	7	8	9	10	11	12
1 Harbour porpoise												
2 Seals												
3 Seabirds (discard)												
4 Seabirds (non-discard)												
5 Sharks												
6 Rays								1.11E-03				3.70E-05
7 Juvenile Cod	5.68E-02	3.82E-02		2.44E-03		7.69E-03	6.22E-03	9.83E-03	2.26E-03	1.11E-03		6.97E-03
8 Cod (adult)		7.69E-02		2.44E-03				9.52E-03				
9 Juvenile Whiting	4.72E-02	5.20E-03			1.40E-03	4.51E-03	1.75E-02	5.40E-02	9.61E-04	1.93E-02	3.12E-04	2.03E-02
10 Whiting (adult)	1.45E-01	8.34E-02			6.73E-02	4.07E-03		8.71E-02		1.34E-04		
11 Other gadoids	1.08E-01	3.36E-02	1.36E-01	2.91E-01	7.53E-02	3.01E-02	2.92E-02	2.67E-02	1.09E-03	7.57E-03	1.76E-02	2.05E-02
12 Demersal fish	2.12E-01	1.38E-01	1.45E-03	1.47E-01	8.05E-03	9.59E-04	5.33E-02	3.59E-02	6.08E-02	2.14E-02	1.96E-02	1.00E-02
13 Juvenile Herring	1.11E-03	1.04E-05	2.34E-02	2.44E-02	1.11E-03	3.30E-03	4.67E-03	1.69E-02	2.44E-03	2.90E-02		4.81E-04
14 Herring (adult)	3.34E-03	7.38E-03	2.34E-02	2.44E-02	1.19E-01	1.18E-02	3.22E-03	1.58E-01				
15 Sprat	1.06E-01	3.39E-03	4.69E-02	2.44E-03	2.42E-02	4.04E-03	1.87E-02	1.10E-02	1.43E-02	2.92E-01	4.05E-03	1.27E-02
16 Mackerel			4.69E-02	2.44E-03	6.40E-02			4.98E-05				4.11E-04
17 Horse mackerel		1.50E-02			1.08E-02		8.40E-04	2.99E-03	1.53E-03	9.88E-03		5.37E-05
18 Sandeels	1.83E-01	3.65E-01	7.83E-02	4.25E-01	2.25E-02	3.04E-01	1.07E-01	4.78E-02	1.82E-01	3.26E-01	1.04E-03	4.55E-01
19 Juvenile Plaice				2.44E-03			1.73E-05	3.76E-02				
20 Plaice (adult)	9.45E-05	7.44E-02			3.37E-02	1.87E-03		4.63E-03				
21 Dab	2.41E-03	5.98E-02			2.44E-03	1.13E-02	1.18E-02	3.58E-03	2.17E-01	6.99E-04	1.19E-04	5.20E-03
22 Other flatfish	3.62E-05	5.57E-02			2.44E-03				3.18E-03	3.27E-06		
23 Juvenile Sole	2.81E-04					1.08E-03	1.88E-03	3.49E-03	1.23E-03	5.33E-04		
24 Sole (adult)	1.37E-05	4.44E-02			8.65E-05	2.82E-03						3.97E-04
25 Sea Bass	1.63E-04	1.06E-04	1.11E-06	1.13E-04	6.19E-06	7.37E-07	4.10E-05	2.76E-05	4.67E-05	1.64E-05	1.50E-05	5.50E-05
26 Pelagic fish	1.09E-01		1.79E-02	4.93E-02	2.94E-04		1.16E-04	5.55E-05	4.93E-04	1.77E-05		
27 Squid & cuttlefish	2.32E-02			4.88E-03	3.79E-02	2.67E-03	6.14E-03	1.07E-02	3.00E-02	9.60E-03	8.32E-04	2.67E-02
28 Carnivorous zooplankton			1.23E-02	2.44E-03	3.80E-03	4.45E-03	2.26E-02	7.91E-04	1.66E-01	4.86E-02	1.05E-01	2.23E-02
29 Herbivorous plankton (copepods)			1.23E-02	2.44E-03		6.56E-07	6.50E-06		6.55E-03	3.09E-03	3.12E-01	7.28E-02
30 Gelatinous zooplankton					2.19E-02		1.79E-03	1.00E-03	4.57E-03	4.05E-03		
31 Large crabs + shrimps					5.83E-02	1.39E-01	1.57E-01	1.08E-01	1.33E-01	4.39E-02	1.78E-02	2.53E-01
32 Blue mussels (aquaculture)		1.15E-04	1.00E-04	1.00E-04		1.07E-04		1.13E-04		1.10E-04		1.23E-04
33 Benthos	3.03E-03		6.53E-02	9.76E-03	4.34E-01	4.42E-01	4.00E-01	1.39E-01	3.01E-01	1.65E-01	5.17E-01	5.70E-02
34 Crangon (below 5cm)					3.81E-03	1.73E-02	1.26E-01	9.44E-03	7.48E-02	1.51E-02	2.57E-04	7.23E-03
35 Crangon (commercial size)					1.46E-03	6.61E-03	3.96E-02	3.44E-03	1.74E-02	3.21E-03	9.79E-05	2.22E-03
36 Phytoplankton										2.54E-05		
37 Detritus												
38 Discards			4.69E-01									
Import			6.64E-02	2.44E-03								
Sum	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
(1 - Sum)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Prey \ predator	13	14	15	16	17	18	19	20	21	22	23	24
1 Harbour porpoise												
2 Seals												
3 Seabirds (discard)												
4 Seabirds (non-discard)												
5 Sharks												
6 Rays												
7 Juvenile Cod				1.98E-03						5.52E-03		
8 Cod (adult)												
9 Juvenile Whiting				5.73E-05	3.33E-02					1.08E-03		
10 Whiting (adult)												
11 Other gadoids				3.96E-04	4.28E-03					3.26E-03		
12 Demersal fish				1.45E-03	7.15E-02					1.33E-02		1.28E-02
13 Juvenile Herring				1.27E-02	1.31E-01			2.03E-04		1.34E-04		
14 Herring (adult)												
15 Sprat				2.92E-02	3.26E-02					1.64E-02		
16 Mackerel					8.04E-06							
17 Horse mackerel					1.02E-01	8.44E-02						
18 Sandeels				1.20E-01	2.00E-02			7.63E-03		1.74E-02		
19 Juvenile Plaice												
20 Plaice (adult)												
21 Dab					6.07E-03					5.70E-02		
22 Other flatfish										5.78E-02		
23 Juvenile Sole												
24 Sole (adult)												
25 Sea Bass				1.12E-06	5.49E-05					1.02E-05		9.82E-06
26 Pelagic fish												
27 Squid & cuttlefish				8.10E-03	9.00E-02							
28 Carnivorous zooplankton	2.73E-01	3.09E-01		6.60E-02	1.70E-01	6.62E-02				4.29E-02		
29 Herbivorous plankton (copepods)	6.75E-01	6.70E-01	8.89E-01	4.85E-01	5.89E-02	6.41E-01			1.25E-02			
30 Gelatinous zooplankton				8.49E-02	8.48E-05			3.46E-03		5.93E-02		
31 Large crabs + shrimps				2.28E-02	7.10E-02		1.09E-01	1.69E-02	1.07E-02	6.99E-02	5.44E-02	
32 Blue mussels (aquaculture)						1.05E-04		1.00E-04	1.00E-04	1.15E-04		
33 Benthos	5.19E-02	2.06E-02		4.36E-02	2.24E-01	2.70E-01	8.81E-01	9.72E-01	9.76E-01	6.52E-01	9.46E-01	9.87E-01
34 Crangon (below 5cm)				3.95E-04	2.24E-03		8.60E-03	5.90E-05	2.47E-04	2.54E-03		
35 Crangon (commercial size)					9.58E-04		1.52E-03	1.00E-05	9.50E-05	9.73E-04		
36 Phytoplankton			1.11E-01	2.21E-02		2.28E-02						
37 Detritus												
38 Discards												
Import												
Sum	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
(1 - Sum)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 6 (continued): The diet composition of the balanced Southern Bight of the North Sea Ecopath model.

Prey \ predator	25	26	27	28	29	30	31	32	33	34	35
1 Harbour porpoise											
2 Seals											
3 Seabirds (discard)											
4 Seabirds (non-discard)											
5 Sharks											
6 Rays											
7 Juvenile Cod			1.13E-02								
8 Cod (adult)											
9 Juvenile Whiting	2.92E-03		9.51E-04								
10 Whiting (adult)	2.05E-03										
11 Other gadoids	1.95E-02										
12 Demersal fish	2.07E-02		5.70E-03							5.07E-03	
13 Juvenile Herring			9.51E-04								
14 Herring (adult)											
15 Sprat			1.13E-02								
16 Mackerel	1.33E-01										
17 Horse mackerel	6.65E-02		1.13E-02								
18 Sandeels	2.98E-03		4.51E-02								
19 Juvenile Plaice										4.52E-04	
20 Plaice (adult)			5.70E-03								
21 Dab			5.70E-03								
22 Other flatfish	1.43E-02		1.14E-02							1.62E-03	
23 Juvenile Sole	1.07E-03										
24 Sole (adult)			5.70E-03								
25 Sea Bass	1.59E-05		4.38E-06							3.89E-06	
26 Pelagic fish	6.87E-02	5.31E-02	2.24E-02								
27 Squid & cuttlefish	8.77E-03	5.60E-03	2.50E-02			7.36E-02					
28 Carnivorous zooplankton		1.49E-01	2.31E-01	5.83E-02		2.94E-01	5.00E-03				
29 Herbivorous plankton (copepods)	2.09E-01	6.95E-01	3.37E-01	8.27E-01		2.94E-01	1.00E-02	1.00E-02		2.88E-03	
30 Gelatinous zooplankton		3.32E-02									
31 Large crabs + shrimps	3.71E-01	9.48E-03	1.44E-02			4.85E-02	2.00E-03			1.21E-02	
32 Blue mussels (aquaculture)											
33 Benthos	1.05E-03	4.99E-02	1.92E-01	1.15E-01		1.18E-01	9.29E-01		1.50E-01	9.96E-01	9.37E-01
34 Crangon (below 5cm)	6.11E-02	3.56E-03	5.40E-03			1.82E-02	5.18E-04			4.31E-03	4.14E-02
35 Crangon (commercial size)	9.03E-03	1.36E-03	2.07E-03			6.96E-03	1.98E-04				
36 Phytoplankton			5.63E-02		9.47E-01	1.47E-01		8.50E-01	2.87E-01		
37 Detritus					5.26E-02			1.40E-01	5.63E-01		
38 Discards							5.36E-02				
Import	8.80E-03										
Sum	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
(1 - Sum)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 7: Basic estimates from the balanced Ecopath model of the Southern Bight of the North Sea.

Functional group	Input parameters									
	TL	B (t km-2)	Z (y-1)	P/B (y-1)	Q/B (y-1)	EE	P/Q (y-1)	BA (t km-2)	BA rate (y-1)	
Harbour porpoise	4.52	0.03			0.02	24.64	0.00	0.00		
Seals	4.48	0.01			0.09	26.84	0.00	0.00		
Seabirds (discard)	3.03	0.0004			0.10	60.52	0.17	0.00		
Seabirds (non-discard)	4.31	0.0016			1.12	48.20	0.02	0.02		
Sharks	3.87	0.01			1.01	3.37	0.94	0.30		
Rays	3.72	0.81			0.30	2.89	0.12	0.10		
Cod										
Juvenile Cod	3.82	0.11	1.96			8.08	0.92	0.24		
Cod (adult)	4.22	0.16	1.13			3.20	0.91	0.35		
Whiting										
Juvenile Whiting	3.88	0.93	2.80			13.33	0.58	0.21		
Whiting (adult)	3.94	0.33	1.35			5.41	0.99	0.25		
Other gadoids	3.30	0.29		1.60		5.23	1.00	0.31		
Demersal fish	4.01	1.24		1.55		4.94	1.00	0.31		
Herring										
Juvenile Herring	3.31	0.40	2.95			17.21	0.99	0.17		
Herring (adult)	3.34	1.50	1.55			6.78	0.13	0.23		
Sprat	2.89	0.50		2.66		9.01	0.98	0.29		
Mackerel	3.59	0.51		1.34		4.81	0.71	0.28		
Horse mackerel	3.97	1.55		0.67		4.96	0.98	0.14		
Sandeels	3.10	5.57		2.10		9.59	0.63	0.22		
Plaice										
Juvenile Plaice	3.29	0.21	1.75			10.12	0.71	0.17		
Plaice (adult)	3.20	0.51	1.12			3.63	0.98	0.31		
Dab	3.18	5.14		0.60		5.58	0.17	0.11		
Other flatfish	3.56	0.10		1.40		6.90	0.94	0.20		
Sole										
Juvenile Sole	3.23	0.19	1.10			10.46	0.18	0.11		
Sole (adult)	3.20	0.32	1.15			5.56	0.57	0.21		
Sea Bass	4.06	0.004		0.76		3.81	0.99	0.20		
Pelagic fish	3.31	0.06		4.00		10.19	0.98	0.39		
Squid & cuttlefish	3.47	0.13		4.50		15.00	0.92	0.30		
Carnivorous zooplankton	3.08	4.24		4.00		12.50	0.99	0.32		
Herbivorous plankton (copepods)	2.00	17.20		9.20		30.00	0.62	0.31		
Gelatinous zooplankton	3.36	0.13		2.86		6.35	0.96	0.45		
Large crabs + shrimps	3.12	5.10		1.05		5.27	0.99	0.20		
Blue mussels (aquaculture)	2.01	0.06		0.36		2.67	0.05	0.13		
Benthos	2.18	300.00		1.06		3.98	0.91	0.27		
C. crangon										
Crangon (below 5cm)	3.18	0.58	12.50			25.14	0.20	0.50		
Crangon (commercial size)	3.24	0.36	6.51			10.00	0.19	0.65		
Phytoplankton	1.00	7.50		286.70		0.39				
Detritus	1.00	50.00				0.38		1117.91	22.36	
Discards	1.00	50.00				0.68				

4.4. Post-balance diagnostics

4.1.1. Biomass

The biomass (log scale) in the balanced SBNS model spans seven orders of magnitude and declines by 6.2 %, falling within the ecological range suggested by Link (2010; Fig. 8). Still, some groups show variance from the trend line, i.e. blue mussels (aquaculture) and benthos. A possibility to improve the benthos group is to split this FG in multiple FGs.




Figure 9: POSTBAL diagnostic biomass. Functional groups are ordered by increasing trophic level.

4.1.2. Production/biomass

P/B (log scale) in the balanced SBNS model spans five orders of magnitude and declines by 2.8 % (Fig. 9). This decrease falls still below the suggested decline (Link, 2010). Overall, there is little variance from the trend line except for phytoplankton, juvenile brown shrimp (*Crangon crangon*), and squid and cuttlefish. However, these P/B values are similar to the P/B values of Stäbler et al. (2016) for the central and SNS and Mackinson and Daskalov (2007) for the entire North Sea.




Figure 10: POSTBAL diagnostic production/biomass. Functional groups are ordered by increasing trophic level.

4.1.3. Consumption/biomass

Q/B is expected to decline with increasing trophic level following basic energetic assumptions. Q/B (log scale) in the balanced SBNS model spans two orders of magnitude and declines by 0.1 % (Fig. 10). It did not improve, but it follows the decline with increasing trophic levels.




Figure 11: POSTBAL diagnostic consumption/biomass. Functional groups are ordered by increasing trophic level.

4.1.4. Production/consumption

Most FGs fall inside the suggested P/Q range of 0.1 - 0.3 (Fig. 11). The groups that fall outside the ideal range, were compared to the mother models' values and are in line with the P/Q values of the mother models (Mackinson and Daskalov, 2007; Stäbler et al., 2016). Some invertebrate species such brown shrimp (*Crangon crangon*), squid and cuttlefish that have high P/Q.




Figure 12: POSTBAL diagnostic production/ consumption. Functional groups are ordered by increasing trophic level.

Abbreviations

B:	Biomass
C:	Catch
CEFAS:	Centre for Environment, Fisheries, and Aquaculture Science
CPUE:	Catch Per Unit Effort
EwE:	Ecopath with Ecosim
F:	Fishing mortality
FG:	Functional group; plural FGs: functional groups
HAWG:	Herring Assessment Working Group for the area south of 62° N
Hd:	Herbivores and detritivores
ICES:	International Council for the Exploitation of the Sea
K:	von Bertalanffy growth function
L_{inf} :	asymptotic length, i.e. the length fish of a population would reach if they were to grow indefinitely
M:	Natural mortality
NS-IBTS:	International Bottom Trawl Survey North Sea
OBIS:	Ocean Biodiversity Information System
OSPAR:	the mechanism by which 15 Governments & the EU cooperate to protect the marine environment of the North-East Atlantic. (OSlo and PARis convention)
OSPAR - SCANS III:	Small Cetacean Abundance in the European Atlantic and North Sea
P/B:	Production over biomass
P/Q:	Production-consumption ratio
Pf:	Apex and/or pelagic predators and/or zooplankton feeders
Q/B:	Consumption
SBNS:	Southern Bight of the North Sea
SNS:	Southern North Sea
SSB:	Spawning Stock Biomass
STECF:	Scientific, Technical and Economic Committee for Fisheries
STECF FDI:	Fisheries-Dependent Information
T:	Temperature
TSB:	Total Stock Biomass
WGNSSK:	Working Group on the assessment of Demersal Stocks in the North Sea and Skagarrak
WGWIDE:	Working Group on Widely Distributed Stocks
W_{inf} :	Asymptotic weight, i.e. the weight fish of a population would reach if they were to grow indefinitely
$W_{maturity}$:	The mean weight at which fish of a given population mature for the first time
Z:	Total mortality

Data availability

The input data, Rscripts and ecopath model can be accessed by <https://doi.org/10.14284/638> and should be cited as follows:

Pint, S.; Stevens, M.; De Troch, M.; van Oevelen, D.; Heymans, J.J.; Everaert, G.; Flanders Marine Institute; Ghent University; Royal Netherlands Institute for Sea Research; University of the Highlands and Islands; (2023): Data - Ecopath model of the Southern Bight of the North Sea. Marine Data Archive. <https://doi.org/10.14284/638>

Citation

Pint S., Stevens M., De Troch M., van Oevelen D., Heymans J. J., Everaert G.: Ecopath model of the Southern Bight of the North Sea (version 2), <https://dx.doi.org/10.48470/23>, 2023.

References

- Asila, A.A. and J. Ogari.: Growth parameters and mortality rates of Nile perch (*Lates niloticus*) estimated from length-frequency data in the Nyanza Gulf (Lake Victoria), p. 272-287. In S. Venema, J. Möller-Christensen and D. Pauly (eds.) Contributions to tropical fisheries biology: papers by the participants of the FAO/DANIDA follow-up training courses. FAO Fish. Rep., (389): 519 p., 1988.
- Beddington, J.R. and Cooke, J.G.: The potential yield of fish stocks. FAO Fish. Tech. Pap. (242), 47 p., 1983.
- Bentley, J. W., Serpetti, N., Fox, C., Reid, D. and Heymans, J. J.: Modelling the food web in the Irish Sea in the context of a depleted commercial fish community. Part 1: Ecopath Technical Report., Scottish Assoc. Mar. Sci. Oban, U.K., (294), 147, 2018.
- Beverton, R.J.H. and Holt, S.J.: A review of methods for estimating mortality rates in fish populations, with special references to sources of bias in catch sampling. Rapp. P.-V. Réun. CIEM 140:67-83, 1956.
- Beverton, R.J.H. and Holt, S.J.: On the dynamics of exploited fish populations. Fish. Invest. Ser. II. Vol. 19, 533 p., 1957.
- Beverton, R.J.H. and Holt, S.J.: Table of yield functions for fisheries management. FAO Fish. Tech. Pap. 38, 49 p., 1964.
- Beverton, R.J.H.: Patterns of reproductive strategy parameters in some marine teleost fishes. J. Fish Biol. 41(B):137-160, 1992.
- Bicknell, A. W. J., Oro, D., Camphuysen, K. C. J. and Votier, S. C.: Potential consequences of discard reform for seabird communities, J. Appl. Ecol., 50(3), 649–658, doi:10.1111/1365-2664.12072, 2013.
- Bjørge, A. and Tolley, K. A.: Harbor Porpoise: *Phocoena phocoena*, Encycl. Mar. Mamm., 530–533, doi:10.1016/B978-0-12-373553-9.00125-5, 2009.
- Blueweiss, L., Fox, H., Kudzman, V., Nakashima, D., Peters, R., and Sams, S.: Relationships between body size and some life history parameters. Oecologia 37:257-272, 1978
- Brodie, B., Mowbray, F. and Power, D.: DFO Newfoundland and Labrador Region Ecosystem Trawl Surveys. Version 1 In OBIS Canada Digital Collections. Bedford Institute of Oceanography, Dartmouth, NS, Canada. Published by OBIS, Digital <http://www.iobis.org/>, 2013, Accessed on 29/06/2020.
- Buhl-Mortensen, L.: MAREANO - Base-line mapping of hyperbenthic crustacea fauna obtained with RP-sledge. Institute of Marine Research, Norway <https://doi.org/10.15468/gecvl4>, 2014
- Christensen, V. and Pauly, D.: ECOPATH II - a software for balancing steady-state ecosystem models and

- calculating network characteristics, *Ecol. Modell.*, 61(3–4), 169–185, doi:10.1016/0304-3800(92)90016-8, 1992.
- Christensen V., Walters C.J., Pauly D. and Forrest R.: *Ecopath with Ecosim version 6 User Guide*, 2008.
- Corrales X., Coll M., Ofir E., Piroddi C., and others: Hindcasting the dynamics of an Eastern Mediterranean marine ecosystem under the impacts of multiple stressors. *Mar Ecol Prog Ser* 580:17-36. <https://doi.org/10.3354/meps12271>, 2017.
- DFO: Central and Arctic Multi-Species Stock Assessment Surveys Version 6 In OBIS Canada Digital Collections. Bedford Institute of Oceanography, Dartmouth, NS, Canada. Published by OBIS, Digital <http://www.iobis.org/>, 2016, Accessed on 29/06/2020.
- Flanders Marine Institute (VLIZ). Taxonomic Information System for the Belgian coastal area. 10 Aug 2004, Oostende, Belgium, Accessed on 29/06/2020.
- Froese, R., and Binohlan, C.: Empirical relationships to estimate asymptotic length, length at first maturity and length at maximum yield per recruit in fishes, with a simple method to evaluate length frequency data. *J. Fish Biol.* 56:758-773, 2000.
- Froese, R., Binohlan, C., and Pauly, D.: Empirical equations to estimate natural mortality in fishes. (In prep.).
- Froese, R. and Pauly, D. (Editors): *FishBase*. World Wide Web electronic publication. www.fishbase.org, version (10/2023), 2023
- Gulland, J.A.: *The fish resources of the oceans*. FAO/Fishing News Books, Surrey, UK, 1971.
- Hammond, P. S., Lacey, C., Gilles, A., Viquerat, S., Börjesson, P., Herr, H., Macleod, K., Ridoux, V., Santos, M. B., Scheidat, M., Teilmann, J., Vingada, J., Øien, N., Gillespie, D., Leaper, R., Sveegaard, S., & Antonio Vázquez, J.: *Estimates of cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys*. 2017.
- Hassel, A.: MAREANO - Base-line mapping of epifauna obtained with Beamtrawl. Institute of Marine Research, Norway <https://doi.org/10.15468/iomgfj>, 2014.
- Heymans, J. J., Coll, M., Link, J. S., Mackinson, S., Steenbeek, J., Walters, C. and Christensen, V.: Best practice in Ecopath with Ecosim food-web models for ecosystem-based management, *Ecol. Modell.*, 331, 173–184, doi:10.1016/j.ecolmodel.2015.12.007, 2016.
- Horn, S., Coll, M., Asmus, H. and Dolch, T., Food web models reveal potential ecosystem effects of seagrass recovery in the northern Wadden Sea. *Restor Ecol*, 29: e13328. <https://doi.org/10.1111/rec.13328>, 2021.
- Hufnagl, M., Temming, A., Siegel, V., Tulp, I. and Bolle, L.: Estimating total mortality and asymptotic length of *Crangon crangon* between 1955 and 2006, *ICES J. Mar. Sci.*, 67(5), 875–884, doi:10.1093/icesjms/fsq003, 2010a.
- Hufnagl, M., Temming, A., Dänhardt, A. and Perger, R.: Is *Crangon crangon* (L. 1758, Decapoda, Caridea) food limited in the Wadden Sea?, *J. Sea Res.*, 64(3), 386–400, doi:10.1016/j.seares.2010.06.001, 2010b.
- ICES: Greater North Sea Ecoregion – fisheries overview. ICS Advice: Fisheries Overviews. Report. <https://doi.org/10.17895/ices.pub.4647>, 2018
- ICES Historical Plankton Dataset (1901-1912). The International Council for the Exploration of the Sea, Copenhagen. Online source: <http://ecosystemdata.ices.dk/>, 2010.
- ICES: North Sea Ecosystem overview Ecosystem components, *North Sea Ecosyst. Overv.*, 1–24, 2008.
- ICES: Report of the Benchmark Workshop on Pelagic Stocks (WKPELA), (43), 341, 2014.
- ICES: Report of the Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak (WGNSSK)., 2019a.
- ICES: Report of the Herring Assessment Working Group for the Area South of 62° N (HAWG)., 2020.
- ICES: Working Group on Widely Distributed Stocks (WGWIDE)., 2019b.
- Institut Français de Recherche pour l'Exploitation de la Mer – IFREMER: COMARGIS: Information System

- on Continental Margin Ecosystems <https://doi.org/10.15468/0djslr>, 2016.
- Lacroix, G., Ruddick, K., Ozer, J. and Lancelot, C.: Modelling the impact of the Scheldt and Rhine/Meuse plumes on the salinity distribution in Belgian waters (southern North Sea), *J. Sea Res.*, 52(3), 149–163, doi:10.1016/J.SEARES.2004.01.003, 2004.
- Libby, D. A.: Maine Department of Marine Resources Inshore Trawl Survey, 2000 – 2009. Maine Department of Marine Resources, PO Box 8, West Boothbay Harbor, Maine 04575, 2010.
- Link, J. S.: Adding rigor to ecological network models by evaluating a set of pre-balance diagnostics: A plea for PREBAL, *Ecol. Modell.*, 221(12), 1580–1591, doi:10.1016/j.ecolmodel.2010.03.012, 2010.
- Mackinson, S. and Daskalov, G.: An ecosystem model of the North Sea to support an ecosystem approach to fisheries management: description and parameterisation, *Sci. Ser. Tech. Rep.*, (142), 196, 2007.
- Marine Biological Association of the UK (MBA): DASSH: The UK Archive for Marine Species and Habitats Data, 2016.
- Meurisse, L., Semal, P.: Royal Belgian Institute of Natural Sciences Crustacea collection, 2020.
- Miller, R, Nozères, C, Kennedy, M.: DFO Quebec Region MLI museum collection. Version 2. OBIS Canada Digital Collections. Published by OBIS <http://www.iobis.org/>, 2014. Accessed on 29/06/2020.
- Musick, J.A.: Criteria to define extinction risk in marine fishes. *Fisheries* 24(12):6-14, 1999.
- National Museum of Natural History, Smithsonian Institution NMNH Invertebrate Zoology Collection Database. National Museum of Natural History, Smithsonian Institution, 10th and Constitution Ave. N.W., Washington, DC 20560-0193, Version 3.2.04 (0802221), 2001.
- Nilsson, S. G. and Nilsson, I. N.: Numbers, Food Consumption, and Fish Predation by Birds in Lake Möckeln, Southern Sweden, *Scand. J. Ornithol.*, 7(1), 61–70, 1976.
- NOAA's National Marine Fisheries Service (NMFS) Northeast Fisheries Science Center.: Northeast Fisheries Science Center Bottom Trawl Survey Data. NOAA's National Marine Fisheries Service (NMFS) Northeast Fisheries Science Center. Woods Hole, Massachusetts, United States of America, 2005.
- Northeast Fisheries Science Center, National Marine Fisheries Service, NOAA, U.S. Department of Commerce. NEFSC Benthic Database. Northeast Fisheries Science Center, 166 Water Street, Woods Hole Laboratories, Woods Hole, MA 02543. Retrieved from <http://www.usgs.gov/iobis-usa/>, 2010.
- OBIS Canada: FRB Eastern Arctic Investigations: The Calanus Series. OBIS Canada Digital Collections. OBIS Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada, Version 2, Digital, retrieved from <http://iobis.org/>, 2011.
- Parr, J. Marine Life Information Network (MarLIN) marine survey data (Professional). Marlin, Collated Marine Life Survey Datasets, Marine Biological Association of the UK, Plymouth, UK
- Palomares, M.L.D.: La consommation de nourriture chez les poissons: étude comparative, mise au point d'un modèle prédictif et application à l'étude des réseaux trophiques. Ph.D. Thesis, Institut National Polytechnique de Toulouse, France, 1991.
- Palomares, M.L. and Pauly, D.: A multiple regression model for predicting the food consumption of marine fish populations. *Aust. J. Mar. Freshwat. Res.* 40:259-273, 1989.
- Palomares, M.L.D. and Pauly, D.: Predicting the food consumption of fish populations as functions of mortality, food type, morphometrics, temperature and salinity. *Mar. Freshwat. Res.* 49:447-453, 1999.
- Pauly, D.: Gill size and temperature as governing factors in fish growth: a generalization of von Bertalanffy's growth formula. *Ber. Inst. f. Meereskunde Univ. Kiel.* No 63, xv + 156 p., 1979.
- Pauly, D.: On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks, *ICES J. Mar. Sci.*, 39(2), 175–192,

- doi:10.1093/icesjms/39.2.175, 1980.
- Pauly, D.: A mechanism for the juvenile-to-adults transition in fishes. *J. Cons. CIEM* 41:280-284, 1984.
- Pauly, D.: A simple method for estimating the food consumption of fish populations from growth data and food conversion experiments. *Fish. Bull. (US)* 84:827-840, 1986.
- Pauly, D.; Food consumption by tropical and temperate fish populations: some generalizations. *J. Fish Biol.* 35 (Suppl. A):11-20, 1989.
- Pauly, D. and Christensen, V.: Trophic levels of fishes, p. 155. *In* R. Froese and D. Pauly (eds.) *FishBase 1998: concepts, design and data sources*. ICLARM, Manila, Philippines. 293 p., 1998.
- Pauly, D., Moreau, J. and Gayanilo, F.C. Jr.: Auximetric analyses, p. 130-134. *In* R. Froese and D. Pauly (eds.) *FishBase 1998: concepts, design and data sources*. ICLARM, Manila, Philippines. 293 p., 1998.
- Pauly, D. Christensen, V. and Sambilay, V.: Some features of fish food consumption estimatives used by ecosystem modellers, *ICES Counc. Meet.*, (657), 1–8, 1990.
- Polovina, J. J.: Model of a coral reef ecosystem, *Coral Reefs*, 3(1), 23–27, doi:10.1007/bf00306137, 1984.
- Potiek, A., Vanermen, N., Middelveld, R. P., de Jong, J., Stienen, E. W. M. and Fijn, R. C.: Spatial and temporal distribution of different age classes of seabirds in the North Sea Analysis of ESAS database., 2019.
- Rees, H.L. et al.: A comparison of benthic biodiversity in the North Sea, English Channel and Celtic Seas - Epifauna. Centre for Environment, Fisheries and Aquaculture Science; Burnham Laboratory, 12 Apr 2005, Essex, UK.
- Reeves, S. A. and Furness, R. W.: Net Loss – Seabirds Gain? Implications of fisheries management for seabirds scavenging discards in the northern North Sea, (June 2014), 74 pp., 2002.
- Ricciardi, A. and Bourget, E.: Weight-to-weight conversion factors for marine benthic macroinvertebrates, *Mar. Ecol. Prog. Ser.*, 163, 245–251, doi:10.3354/meps163245, 1998.
- Ricker, W.E.: Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Can.* (191), 382 p., 1975.
- Santos, R.: Zariquiey Collection. Biological Reference Collections ICM CSIC. Institute of Marine Sciences (ICM-CSIC). <https://dx.doi.org/10.15470/cqtmj1>, 2016.
- Scientific, Technical and Economic Committee for Fisheries (STECF) – Fisheries Dependent - Information – Classic (STECF-17-09). Publications Office of the European Union, Luxembourg, ISBN 978-92-79-67481-5, doi:10.2760/561459, JRC107598, 2017.
- Scott E., Serpetti N., Steenbeek J., Heymans J.J.: A Stepwise Fitting Procedure for automated fitting of Ecopath with Ecosim models, *SoftwareX*, 5, 25-30, <https://doi.org/10.1016/j.softx.2016.02.002>, 2016.
- Sherley, R. B., Ladd-Jones, H., Garthe, S., Stevenson, O. and Votier, S. C.: Scavenger communities and fisheries waste: North Sea discards support 3 million seabirds, 2 million fewer than in 1990, *Fish Fish.*, 21(1), 132–145, doi:10.1111/faf.12422, 2020.
- Sparholt, H.: An estimate of total biomass of fish in the North Sea, *J. Cons. int. Explor. Mer.*, 46, 200–210, doi:10.1093/icesjms/40.2.161, 1990.
- Stäbler, M., Kempf, A., Mackinson, S., Poos, J. J., Garcia, C. and Temming, A.: Combining efforts to make maximum sustainable yields and good environmental status match in a food-web model of the southern North Sea, *Ecol. Modell.*, 331, 17–30, doi:10.1016/j.ecolmodel.2016.01.020, 2016.
- Stratmann, T., van Oevelen, D., Martínez Arbizu, P., Wei, C. L., Liao, J. X., Cusson, M., Scrosati, R. A., Archambault, P., Snelgrove, P. V. R., Ramey-Balci, P. A., Burd, B. J., Kenchington, E., Gilkinson, K., Belley, R. and Soetaert, K.: The BenBioDen database, a global database for meio-, macro- and megabenthic biomass and densities, *Sci. Data*, 7(1), 1–12, doi:10.1038/s41597-020-0551-2, 2020.

- Swedish county administration boards, Swedish municipalities, Swedish coalitions of water conservation, Swedish Meteorological and Hydrological Institute, et al.: SHARK - Regional monitoring and monitoring projects of Epibenthos in Sweden since 1994, 2017.
- Taylor, C.C.: Cod growth and temperature. *J. Cons. CIEM* 23:366-370, 1958.
- The Norwegian Oil Industry Association: Offshore reference stations, North/Norwegian sea. The Norwegian Oil Industry Association (OLF), Akvaplan-niva and Det Norske Veritas, Norway, 2001.
- Tulp, I., Chen, C., Haslob, H., Schulte, K., Siegel, V., Steenbergen, J., Temming, A. and Hufnagl, M.: Annual brown shrimp (*Crangon*) biomass production in Northwestern Europe contrasted to annual landings, *ICES J. Mar. Sci. J. du Cons.*, 73(10), 2539–2551, doi:10.1093/icesjms/fsw141, 2016.
- Türkay, M.: Senckenbergisches Sammlungsverwaltungssystem, SeSam. Senckenbergische Naturforschende Gesellschaft, Frankfurt, Germany. <http://sesam.senckenberg.de/>
- Turrell, W. R.: New hypotheses concerning the circulation of the northern North Sea and its relation to North Sea fish stock recruitment, *ICES J. Mar. Sci.*, 49(1), 107–123, doi:10.1093/icesjms/49.1.107, 1992.
- Van Guelpen L and Pohle, G.: Bay of Fundy Species List. v1.2. Canadian node of the Ocean Biogeographic Information System (OBIS Canada). Dataset/Occurrence. <http://ipt.iobis.org/obiscanada/resource?r=bofetf&v=1.2>, 2014.
- Van Guelpen, L.: Atlantic Reference Centre Museum of Canadian Atlantic Organisms - Invertebrates and Fishes Data. Version 4 In OBIS Canada Digital Collections. Bedford Institute of Oceanography, Dartmouth, NS, Canada. Published by OBIS, Digital <http://www.iobis.org/>, 2016. Accessed on 29/06/2020.
- Verleye, T. J., Dauwe, S., van Winsen, F. and Torreele, E.: Recreatieve zeevisserij in België anno 2018 - Feiten cijfers., 2019.
- Waggitt, J. J., Evans, P. G. H., Andrade, J., Banks, A. N., Boisseau, O., Bolton, M., Bradbury, G., Brereton, T., Camphuysen, C. J., Durinck, J., Felce, T., Fijn, R. C., Garcia-Baron, I., Garthe, S., Geelhoed, S. C. V., Gilles, A., Goodall, M., Haelters, J., Hamilton, S., Hartny-Mills, L., Hodgins, N., James, K., Jessopp, M., Kavanagh, A. S., Leopold, M., Lohrengel, K., Louzao, M., Markones, N., Martínez-Cedeira, J., Ó Cadhla, O., Perry, S. L., Pierce, G. J., Ridoux, V., Robinson, K. P., Santos, M. B., Saavedra, C., Skov, H., Stienen, E. W. M., Sveegaard, S., Thompson, P., Vanermen, N., Wall, D., Webb, A., Wilson, J., Wanless, S. and Hiddink, J. G.: Distribution maps of cetacean and seabird populations in the North-East Atlantic., 2020.
- Walters, C., Christensen, V. and Pauly, D.: Structuring dynamic models of exploited ecosystems from trophic mass-balance assessments, *Rev. Fish Biol. Fish.*, 7(2), 139–172, doi:10.1023/A:1018479526149, 1997.