

## Morphology and Phylogenetic Position of Sargassum Nudibranch *Scyllaea fulva* Quoy & Gaimard, 1824 (Nudibranchia: Scyllaediae): First Record in Hong Kong

Sam King Fung Yiu<sup>1,2,\*</sup>, Thomas Ka Tung Leung<sup>2</sup>, Gabriel Yeung Lee<sup>2</sup>, and Meng Yan<sup>2,\*</sup>

<sup>1</sup>*School of Biological Science, The University of Hong Kong, Hong Kong, China. \*Correspondence: E-mail: cyclesamyiu@hotmail.com (Yiu)*

<sup>2</sup>*State Key Laboratory of Marine Pollution, City University of Hong Kong, Hong Kong, China. \*Correspondence: E-mail: mengyan@cityu.edu.hk (Yan)*

*E-mail: thomas1126199716@gmail.com (Leung); auytfff1357924680@gmail.com (Lee)*

ORCID:

Sam King Fung Yiu: <https://orcid.org/0000-0001-6212-7339>

Thomas Ka Tung Leung: <https://orcid.org/0000-0009-0871-9048>

Gabriel Yeung Lee: <https://orcid.org/0000-0002-2814-4191>

Meng Yan: <https://orcid.org/0000-0002-5992-3171>

*(Received 13 July 2023 / Accepted 27 December 2023 / Published -- 2024)*

*Communicated by James D. Reimer*

Scyllaediae is a small group of nudibranchs comprising three genera (*Scyllaea*, *Notobryon*, and *Crosslandia*) with striking morphological similarities, making their identification challenging based on external features alone. Previous studies have highlighted the significance of central radular teeth in distinguishing *Notobryon* from *Scyllaea* and *Crosslandia*. The genus *Scyllaea*, commonly known as the sargassum nudibranch, currently consists of only two valid species, *Scyllaea pelagica* and *Scyllaea fulva*. These species inhabit *Sargassum* spp., feeding on hydroids. During a biodiversity survey conducted in April 2023, seven individuals of sargassum nudibranch were collected from the seaweed *Sargassum* spp. at a depth of 2 meters in Tai She Wan through SCUBA diving. Initially, the specimens were misidentified based on their resemblance to *Notobryon wardi* and previous local records. However, thorough morphological and molecular examinations confirmed them to be *Scyllaea fulva*, representing the first record of this species in Hong Kong. Notably, our specimens lacked the blue spots observed in specimens from Thailand and the West Pacific Ocean, as reported in previous studies. Internally, a pineapple-like structure formed a honeycomb pattern on the surface of the masticatory edge of the jaw flap, with the presence of central radular teeth. A Maximum Likelihood tree analysis revealed *Crosslandia* to be the sister group of *Scyllaea*. Comparative analysis of intra-specific distances between individuals from the Philippines, French Polynesia, and

Hong Kong indicated a close relationship between the Hong Kong specimens and those from the Philippines. Furthermore, we provide a detailed description of the external and internal morphology of *Scyllaea fulva* in this paper, integrating valuable morphological information for future species identification purposes.

**Key words:** Scyllaeidae, Sea slug, Mollusc, Subtropical reef, South China Sea

Citation: Yiu SKF, Leung TKT, Lee GY, Yan M. 2024. Morphology and phylogenetic position of sargassum Nudibranch *Scyllaea fulva* Quoy & Gaimard, 1824 (Nudibranchia: Scyllaeidae): first record in Hong Kong. Zool Stud 63:04.

## BACKGROUND

Scyllaeidae Alder and Hancock in 1855 is a small group of nudibranchs that consists of *Crosslandia* Eliot, 1902, *Notobryon* Odhner, 1936, and *Scyllaea* Linnaeus, 1758. They bear a striking resemblance to algae based on the body colour and wing-like lobes and primarily feed on epiphytic hydroids (Gosliner et al. 2008). Notably, they possess the ability to swim through the flexing of their rhinophoral sheaths and lateral appendages (Gosliner et al. 2008). Morphologically, Scyllaeidae can be distinguished by their dorsal margin, which extends into large lobes. These lobes, whether continuous or separated, often exhibit traces on the rhinophoral sheaths and are adorned with branchial tufts. The presence of an indistinct velum and the absence of frontal papillae are additional distinguishing features. Their rhinophore club is perfoliate, and the anus is positioned laterally or laterodorsally (Odhner 1936). However, due to the striking resemblance among Scyllaeidae members, the high similarity in their external morphology has led to significant confusion and misidentification (Pola et al. 2012).

*Scyllaea*, commonly known as sargassum nudibranch, can be distinguished from other scyllaeids by a radula with a central tooth and two pairs of well-separated dorsal lobes (Odhner 1936). *Scyllaea* comprises two recognised species including *Scyllaea pelagica* Linnaeus, 1758 and *Scyllaea fulva* Quoy & Gaimard, 1824 (MolluscaBase 2023) which both inhabit the seaweed *Sargassum* spp. *Scyllaea fulva*, a poorly studied species, was previously synonymized as *Nerea punctata* Lesson, 1830, *Scyllaea dracaena* Kelaart, 1858 and *Scyllaea quoyi* Gray, 1850. All of these species had been later synonymized as *Scyllaea pelagica* after Odhner (1936) considered *Scyllaea fulva* to be a spurious species. Pola et al. (2012) demonstrated that specimens of *Scyllaea pelagica* collected in the Indo-Pacific region were genetically distinct from those found in the Atlantic. The genetic difference between the Atlantic and Indo-Pacific specimens was determined to

be 7% in the cytochrome *c* oxidase subunit I (*COI*) gene. This significant genetic distinction provided strong evidence for considering them as separate species. Therefore, the name *Scyllaea fulva* was re-erected. Their study underscores the crucial role of molecular analysis in discerning between morphologically similar organisms.

In April 2023, we conducted a biodiversity survey in the *Sargassum* habitat. Seven individuals of *Sargassum*-like nudibranchs were found firmly adhering to *Sargassum* thalli. Upon initial examination, we identified the specimens as *Notobryon wardi* Odhner, 1936, based on a thorough review of external morphological features only (Rudman 2002b; Picton 2002; Chow et al. 2022). The external morphology (body colour, the shape of dorsal lobes, etc.) exhibited striking similarities between our specimens and those described in the literature. Furthermore, previous records had documented the presence of this species in Hong Kong waters (Rudman 2002b; Picton 2002; Chow et al. 2022). However, to ensure accurate identification, we employed molecular analysis to confirm the identities of our specimens. Surprisingly, the results revealed that our specimens were not *Notobryon wardi*, but instead belonged to the species *Scyllaea fulva*. It highlights the potential challenges of relying solely on external morphology for the identification of scyllaeids. Interestingly, the absence of previous records of the genus *Scyllaea* in Hong Kong literature (Jensen 1998; Chow et al. 2022; Astudillo et al. 2023) further emphasizes the significance of our findings. Additionally, we observed a significant dearth of well-documented morphological information (particularly internal morphology) pertaining to *Scyllaea fulva* in existing literature (Quoy and Gaimard 1824; Lesson 1830; Gray 1850; Kelaart 1858; Odhner 1936). Hence, we initiated this study to present the first recorded occurrence of *Scyllaea fulva* in Hong Kong. This study aims to investigate both the external and internal morphology of this species. Furthermore, we seek to elucidate its phylogenetic position by analyzing three genetic markers: the cytochrome *c* oxidase subunit I (*COI*), the 16S ribosomal RNA (*16S rRNA*), and the Histone 3 (*H3*) genes. The data generated through this study will provide valuable insights into the morphology of scyllaeids, facilitating more accurate identification by researchers in the future.

## MATERIALS AND METHODS

### Sampling

Seven individuals of *Scyllaea fulva* were collected from *Sargassum* spp. at Tai She Wan (22°21'32.7"N, 114°20'15.1"E) by SCUBA diving on 28th March 2023 and 11st April 2023. The specimens were preserved in 95% ethanol for morphological and molecular analysis.

## Morphological analysis

External morphological characteristics were examined under a Motic SMZ-171 stereomicroscope (Motic, China). Seven specimens were dissected to extract the buccal mass and reproductive system. The buccal masses were dissolved in 20% diluted bleach for 30 min at room temperature to remove connective tissues and muscles. The jaws and the radula were isolated. The jaws and the radula were dried, coated with gold, mounted on a stub, examined and photographed under a LEO 1530 field emission scanning electron microscope (SEM) (Zeiss, Germany), and the reproductive system was observed and was draw under the stereomicroscope.

## Molecular analysis

Genomic DNA of foot tissues from two specimens (NMMB-M011734 to 735) were extracted using the Chelex 100 solution (Walsh et al. 1991). Concentration and purity of the DNA samples were determined using a NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA), and DNA integrity was checked using electrophoresis on 1.0% agarose gel. Polymerase chain reaction (PCR) reactions were then conducted using the extracted DNA as templates to amplify the *COI*, *16S rRNA* and *H3* genes. Primer sequences for partial sequences of *COI* were sourced from Folmer et al. (1994) using pairs LCO1490 (5'-GGTCAACAAATCATAAAGATATTGG-3') and HC02198 (5'-TAAACTTCAGGGTGACCAAAAATCA-3'). Partial sequences of the *16S rRNA* region were amplified using the pairs 16Sar-L (5'-CGCCTGTTTATCAAAAACAT-3') (Palumbi et al. 1991) and 16s-xH (5'-CCGGTYTGAAMYYAGATCACGTAGG-3') (Mehrotra et al. 2020). Primers for the *H3* region were taken from Colgan et al. (2000) using the primers H3F (5'-ATGGCTCGTACCAAGCAGACVGC-3') and H3R (5'-ATATCCTTRGGCATRATRGTGAC-3'). The following polymerase chain reaction (PCR) program was used for the *COI* gene: 1 min at 95°C (initial denaturation); 35 cycles of 30 s at 95°C (denaturation), 30 s at 52°C (annealing), and 45 s at 72°C (elongation); and 7 min at 72°C (final extension). The PCR reactions for the *16S rRNA* and *H3* genes were conducted using a similar programme, except that the annealing temperature was 50°C. PCR products were sent to BGI Hong Kong for sequencing on an ABI 310 Genetic Analyzer. All new sequences were deposited in GenBank (Table S1).

Alignments of the three genes were conducted separately and trimmed manually to 661 bp for *COI*, 466 bp for *16S rRNA* and 328 bp for *H3* using MEGA 7 (Kumar et al. 2016). The sequence of *COI*, *16S rRNA* and *H3* genes from other scyllaeids and a distantly related outgroup (*Melibe viridis*) were downloaded from GenBank (Table S1). Sequences were concatenated using

SequenceMatrix v.1.7.8 (Vaidya et al. 2011) and then imported to the website version of IQ-Tree (<http://iqtree.cibiv.univie.ac.at/>; Nguyen et al. 2015) for Maximum Likelihood tree reconstruction with 1,000 ultrafast bootstrap pseudoreplicates (Hoang et al. 2017). The best fit model (TPM3u+F+I+G4) was determined by IQ-Tree server. The phylogenetic trees were visualized and edited using FigTree v1.4.4. Pairwise genetic distances for the respective *COI*, *16S* and *H3* genes were estimated using MEGA 7 separately based on the *p*-distance method using the bootstrap method with 10,000 pseudoreplicates for variance estimation. Rates among sites were gamma distributed with invariant sites (G+I) and the gamma parameter was set to four.

## RESULTS

### TAXONOMIC ACCOUNT

#### Class Gastropoda Cuvier, 1795

#### Order Nudibranchia Cuvier, 1817

#### Family Scyllaeidae Alder & Hancock, 1855

#### Genus *Scyllaea* Linnaeus, 1758

#### Species *Scyllaea fulva* Quoy & Gaimard, 1824

#### Synonyms


*Nerea punctata* (Lesson 1830), *Scyllaea dracaena* (Kelaart 1858), *Scyllaea quoyi* (Gray 1850), *Scyllaea pelagica* (Pola et al. 2012).

*Materials examined*: NMMB-M011734: 0.9 cm in length; NMMB-M011735: 0.85cm in length; NMMB-M011736: 1.4 cm in length; NMMB-M011737: 0.95 cm in length; NMMB-M011738: 0.5 cm in length; NMMB-M011739: 1.3 cm in length; NMMB-M011740: 2.9 cm in length.


*Locality*: Seven specimens were collected from *Sargassum* spp. at Tai She Wan (22°21'32.7"N, 114°20'15.1"E) at 2 m on 28th March 2023 and 11th April 2023.

*Type locality*: New Guinea. Geographic distribution (Fig. 1): Hong Kong (this study), Indo-Pacific region including Japan (Baba 1949), Papua New Guinea, Philippines (Pola et al. 2012), Réunion (Cadet 2012), Gump Station, Cook's Bay, Moorea, French Polynesia (Goodheart et al. 2017 2018); Mozambique (Tibiriçá et al. 2017) and Thailand (Mehrotra et al. 2021).

*Habitat*: Inhabit on the seaweed *Sargassum* spp. (Fig. 2A).



**Fig. 1.** Map showing the currently known distribution of *Scyllaea fulva* and *Scyllaea pelagica* in the world.




**Fig. 2.** Selected photographs showing the sargassum habitat and *Scyllaea fulva*. A, Sargassum bed at Tai She Wan; B, *In situ* swimming *Scyllaea fulva*; C, Dorsal view of *Scyllaea fulva* in a beaker; D, Lateral view of *Scyllaea fulva* in a beaker. Scale bar = 0.5 cm.


*External morphology* (Fig. 2B–D and Fig. 3A–B): The body colour varies from transparent to semi-transparent green or light yellow. The body is slender, soft, flaccid, and elevated, but laterally compressed. The body surface is smooth. There are some black or brown spots on the body. The foot is narrow. The posterior crest is moderately large with an entire margin. The edge of posterior crest is brown. The front of the head is expanded in a semicircular veil. The rhinophores are perfoliate with 3–4 lamellae. There are two pairs of equal sized dorsolateral lobes. The lobes are wing-like denticulated. Each dorsolateral lobe bears four to five large and transparent dendritic ‘gills’ on their upper surface and two on the tail as well as four on each side of tail.

*Internal morphology* (Fig. 3C–F and Fig. 4): The buccal mass contained a pair of thin and triangular jaws. The masticatory edge of the jaws is expanded into a wing-like flap. Over the edge of this flap is a series of pineapple-like structure. These structures form a honey-comb pattern over the entire surface of the masticatory edge, the number of rodlets on pineapple-like structure ranged from two to five. Central radular teeth were present. Each primary denticle bear two to five secondary denticles. The radula formula was  $11 \times 15.1.15$ . The reproductive system contained a small bursa copulatrix, a big size of female gland mass, a duct connected between female gland mass and a prostate with the ampulla.

*Remarks:* *Scyllaea fulva* can be distinguished from *Scyllaea pelagica* based on the radular formula. The radular formula of *Scyllaea pelagica* is  $16-24 \times 24-54.1.24-54$  (Odhner 1936), which exhibits a higher number of denticles compared to *Scyllaea fulva*.




**Fig. 3.** Drawing showing the external and internal morphology of *Scyllaea fulva*. A, Dorsal view; B, Lateral view; C, Jaw; D, Radula; E, Teeth; F, Reproductive system. bc, bursa copulatrix; fmg, female gland mass; pr, prostate; amp, ampulla; hd, hermaphroditic duct. Scale bars: A and B =0.5 cm; C = 0.5 mm; D = 250 μm; E = 20 μm; F = 0.2 mm.



**Fig. 4.** SEM photos showing the jaws and radula. A, Overview of a jaw; B, Jaw element; C, Central teeth presented on the radula; D, Close-up image of the denticles. Scale bars: A = 500 μm; B = 20 μm; C = 500 μm; D = 20 μm.

### Molecular analysis

Three gene sequences were obtained from each of the two specimens (NMMB-M011734 to 735). Alignment and concatenation resulted in a dataset of 1455 bp (661 bp for *COI*, 466 bp for *16S* rRNA and 328 bp for *H3*). Our phylogenetic analyses showed that the two specimens we collected from the field were *Scyllaea fulva* (Fig. 5). Pair-wise sequence comparisons were conducted to determine the inter- and intra-specific *p*-distances (Table 1; Table S1). Interspecific *p*-distances amongst *Scyllaea fulva* and *Scyllaea pelagica* were 6.90%–7.20% for *COI*, 2.20%–2.50% for *16S* and 0.00% for *H3*. Intraspecific *p*-distances in the *Scyllaea fulva* were in general very small, with 0.00%–0.60% for *COI*, 0.00%–0.20% for *16S*, and 0.00% for *H3*.



**Fig. 5.** Phylogenetic tree of the concatenated *COI/16S/H3* sequence dataset constructed using the maximum likelihood method. Bootstrap values >50 are shown in the nodes.

**Table 1.** *Scyllaea* inter- and intra-specific uncorrected *p*-distance for *COI/16S/H3*. (P) indicates the specimens from Philippine (Pola et al. 2012); (FP) indicates the specimen from French Polynesia (Goodheart et al. 2017; 2018); (HK) indicates the specimens used in this study. Detail can be referred to table S2–4

Species	<i>COI</i>	16S	H3
<i>Scyllaea pelagica</i> vs <i>Scyllaea fulva</i>	6.90%–7.20%	2.20%–2.50%	0.00%
<i>Scyllaea fulva</i> (P) vs <i>Scyllaea fulva</i> (HK)	0.30%–0.60%	0.00%–0.20%	0.00%
<i>Scyllaea fulva</i> (FP) vs <i>Scyllaea fulva</i> (HK)	0.60%	0.00%	-
<i>Scyllaea fulva</i> (HK1) vs <i>Scyllaea fulva</i> (HK2)	0.30%	0.00%	0.00%

## DISCUSSION

We have successfully confirmed the specimens to be *Scyllaea fulva*, commonly called sargassum nudibranch. After reviewing Astudillo et al. (2023), Jensen (1998), and Chow et al. (2022), we have documented the occurrence of *Scyllaea fulva* as the first record in Hong Kong waters, which is consistent with the known distribution of the species. Morphologically, our specimens did not bear the blue spots on the body, which is different from the individuals found by Gosliner et al. (2008), Goodheart et al. 2017 2018) and Mehrotra et al. (2021). This indicates that external morphological variation occurs within the species. Moreover, our specimens closely correspond to the original descriptions of *Scyllaea fulva* and its synonyms, exhibiting an elongated body, tawny color, grooved underside, and four slender wings, with the tentacles widened at the tip (Quoy and Gaimard 1824–1826; Lesson 1831; Gray 1850; Kelaart 1858). However, the literature

did not provide detailed information on the morphology of *Scyllaea fulva*, particularly its internal morphology. Therefore, our findings on internal morphology of *Scyllaea fulva* is important for integrating the morphological information of the species.

Before the work of Pola et al. (2012), *Scyllaea fulva* was considered a spurious species (Odhner 1936). Pola et al. (2012) demonstrated a 7% genetic distance for *COI* among specimens from the Atlantic and Indo-Pacific regions confidently supported its recognition as a distinct species. Thus, *Scyllaea fulva* was re-established. Additionally, Odhner (1936) noted that scyllaeids can be differentiated at the genus level by the presence or absence of a central tooth on the radula. Rudman's (2002abc) SEM photos of scyllaeids revealed that *Scyllaea pelagica* and *Crosslandia viridis* also had a central radular tooth, but *Notobryon wardi* did not. Several *Notobryon* species described by Pola et al. (2012) and Caballer and Ortea (2014) also lacked the central radular tooth, which is consistent with Odhner (1936) and Rudman's (2002abc) conclusions.

To precisely identify scyllaeids, Pola et al. (2012) demonstrated that molecular analysis is a powerful tool for delineating species boundaries. In the present study, we also used molecular analysis to avoid misidentification and accurately identify our specimens. Our findings on inter- and intra-specific *p*-distances revealed that *H3* lacked a phylogenetic signal, but *COI* is a useful gene for delimiting the species. Compared with the genetic distance (0.30%-0.60%) for *COI* among *Scyllaea fulva* specimens from the Philippines, French Polynesia, and Hong Kong, the Hong Kong *Scyllaea fulva* is closely related to the species in the Philippines.

This occurrence of *Scyllaea fulva* expands the diversity of nudibranchs in Hong Kong, making it the second member of Scyllaeidae to occur in Hong Kong since Rudman (2002b) confirmed the Hong Kong specimen (AMC139151) stored in the Australian Museum to be *Notobryon wardi*. Additionally, the present study also reports the external and internal morphology to integrate the morphological data for *Scyllaea fulva*. These findings provide insights for the morphological identification of scyllaeids in the future.

## CONCLUSIONS

Our study reports the first documented occurrence of *Scyllaea fulva* in Hong Kong waters. The confirmed presence of this species expands the diversity of nudibranchs in the region. The observed external morphological variation, coupled with detailed morphological data, provides valuable insights for future identification of scyllaeids and contributes to our understanding of *Scyllaea fulva*'s biology. This study highlights the importance of ongoing research to further explore the phenotypic and genetic diversity of marine organisms.

**Acknowledgments:** This study was financially support by the project (AFCD SQ 314 21C) Provision of Services for Conducting Night Fisheries Resources Surveys. We would like to thank Kwok Leung for field assistance. We would also like to thank two anonymous reviewers' useful comments for manuscript improvement.

**Authors' contributions:** SKFY and MY initiated the study, SKFY, TKTL and GYL conducted sampling, TKTL conducted the drawing, SKFY performed molecular and morphological analysis as well as drafted the manuscript.

**Competing interests:** SKFY, TKTL, GYL and MY declare they have no conflict of interest.

**Availability of data and materials:** The accession numbers of specimens had been deposited on the GenBank. The examined specimens are deposited at National Museum of Marine Biology and Aquarium in Taiwan.

**Consent for publication:** All of the authors agreed to publish the paper.

**Ethics approval consent to participate:** Not applicable.

## REFERENCES

- Astudillo JC, Williams GA, Leung KMY, Cannicci S, Yasuhara M, Yau C, Qiu JW, Ang PO, To AWL, Shea SKH. 2023. Hong Kong Register of Marine Species.  
<https://www.marinespecies.org/hkrms>. Accessed on 30 June 2023.
- Baba K. 1949. Opisthobranchia of Sagami Bay: collected by His Majesty the Emperor of Japan. Iwanami Shoten, Tokyo, Japan.
- Caballer M, Ortea J. 2014. A new sibling species of Notobryon (Gastropoda, Nudibranchia) from the Caribbean Sea. J Mar Biolog Assoc UK **94(7)**:1465–1470.  
doi:10.1017/S0025315414000605.
- Cadet C. 2012. *Scyllaea fulva* Quoy & Gaimard, 1824 or *pelagica*. South-west Indian Ocean Sea slug site. [http://seaslugs.free.fr/nudibranche/a\\_scyllaea\\_pelagica.htm](http://seaslugs.free.fr/nudibranche/a_scyllaea_pelagica.htm). Accessed on 29 June 2023.

- Chow LH, Yu VPF, Kho ZY, See GCL, Wang A, Baker DM, Tsang LM. 2022 An Updated Checklist of Sea Slugs (Gastropoda, Heterobranchia) from Hong Kong Supported by Citizen Science. *Zool Stud* **61**:52. doi:10.6620/ZS.2022.61-52.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotech* **3**:294–299.
- Gray JE. 1850. Figures of molluscous animals, selected from various authors. Longman, Brown, Green and Longmans, London, UK.
- Goodheart JA, Bazinet AL, Valdes A, Collins AG, Cummings MP. 2017. Prey preference follows phylogeny: evolutionary dietary patterns within the marine gastropod group Cladobranchia (Gastropoda:Heterobranchia: Nudibranchia). *BMC Evol. Biol* **17**(1):221. doi:10.1186/s12862-017-1066-0.
- Goodheart JA, Bleidissel S, Schillo D, Strong EE, Ayres DL, Preisfeld A, Collins AG, Cummings MP, Wagele H. 2018. Comparative morphology and evolution of the cnidosac in Cladobranchia (Gastropoda: Heterobranchia: Nudibranchia). *Front Zool* **15**:43. doi:10.1186/s12983-018-0289-2.
- Gosliner TM, Behrens DW, Valdés Á. 2008. Indo-Pacific Nudibranchs and Sea Slugs. Sea Challengers/California Academy of Sciences, San Francisco, CA, USA.
- Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2017. UFBoot2: Improving the ultrafast bootstrap approximation. *Mol Biol Evol* **35**:518–522. doi:10.1093/molbev/msx281.
- Jensen KR. 1998. Zoogeographic affinities of Hong Kong Opisthobranchia (Mollusca, Gastropoda). *In*: Morton, B. (Ed.), *The Marine Biology of the South China Sea III*. Hong Kong University Press, Hong Kong.
- Kelaart EF. 1858. Description of new and little known species of Ceylon nudibranchiate molluscs, and zoophytes. *Journal of the Ceylon Branch of the Royal Asiatic Society* **3**(9):84–139.
- Lesson RP. 1831. Chapitre XI. Mollusques, Annélides et Vers; par R.-P. Lesson. pp. 239–456 [15 Nov 1831]. *In*: Lesson R.P. 1830–1831. *Voyage autour du monde, Exécuté par Ordre du Roi, sur La Corvette de Sa Majesté, La Coquille, pendant les années 1822, 1823, 1824 et 1825, sous le ministere et conformément aux Instructions de S.E.M. le Marquis de Clermont-Tonnerre, Ministre de la Marine; Et publié sous les auspices de son Excellence Mgr le Cte de Chabrol, Ministre de la Marine et des Colonies, par M.L.I. Duperrey, Capitaine de Frégate, Chevalier de Saint-Louis et Membre de la Légion d'Honneur, Commandant de l'Expedition. Zoologie, par M. Lesson. Tome Second. 1re Partie*. Paris, Arthus Bertrand, Libraire-Editeur, Imprimerie de Firmin Didot.

- Mehrotra R, Arnold S, Wang A, Chavanich S, Hoeksema BW, Caballer M. 2020. A new species of coral-feeding nudibranch (Mollusca: Gastropoda) from the Gulf of Thailand. *Mar Biodivers* **50**:36. doi:10.1007/s12526-020-01050-2.
- Mehrotra R, Caballer Gutiérrez MA, Scott CM, Arnold S, Monchanin C, Viyakarn V, Chavanich S. 2021. An updated inventory of sea slugs from Koh Tao, Thailand, with notes on their ecology and a dramatic biodiversity increase for Thai waters. *ZooKeys* **1042**:73–188. doi:10.3897/zookeys.1042.64474.
- MolluscaBase. 2023. MolluscaBase. *Scyllaea* Linnaeus, 1758. World Register of Marine Species. <https://www.marinespecies.org/aphia.php?p=taxdetails&id=138468>. Accessed on 10 July 2023.
- Nguyen LT, Schmidt HA, Von haeseler A, Minh BQ. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**:268–274. doi:10.1093/molbev/msu300.
- Odhner NH. 1936. Nudibranchia Dendronotacea. A revision of the System. in: Melanges Paul Pelseneer. Mémoires du Musée Royal d'Histoire Naturelle de Belgique. (2)**3**:1057–1128.
- Palumbi SR, Kessing B, Martin A. 1991. The simple fool's guide to PCR, 2nd ed. Department of Zoology, University of Hawaii, Honolulu.
- Picton B. 2002. Notobryon from Hong Kong. Sea Slug Forum. Australian Museum, Sydney. <http://www.seaslugforum.net/find/5948>. Accessed on 29 June. 2023.
- Pola M, Camacho-Garcia YE, Gosliner TM. 2012. Molecular data illuminate cryptic nudibranch species: the evolution of the Scyllaeidae (Nudibranchia: Dendronotina) with a revision of *Notobryon*. *Zool J Linn Soc* **165**:311–336. doi:10.1111/j.1096-3642.2012.00816.x.
- Quoy JRC, Gaimard JP. 1824–1826. Zoologie. In: de Freycinet L (ed.) Voyage au tour du monde fait par ordre du roi, sur les corvettes de S. M: l'Uranie et la Physicienne pendant les années 1817 à 1820.
- Rudman WB. 2002a. Radula of *Crosslandia viridis*. Sea Slug Forum. Australian Museum, Sydney. <http://www.seaslugforum.net/find/7937>. Accessed on 29 June. 2023.
- Rudman WB. 2002b. Radula of *Notobryon wardi*. Sea Slug Forum. Australian Museum, Sydney. <http://www.seaslugforum.net/find/7938>. Accessed on 29 June. 2023.
- Rudman WB. 2002c. Radula of *Scyllaea pelagica*. Sea Slug Forum. Australian Museum, Sydney. <http://www.seaslugforum.net/find/7939> Accessed on 29 June. 2023.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**(7):1870–1874. doi:10.1093/molbev/msw054.

- Tibiricá Y, Pola M, Cervera JL. 2017. Astonishing diversity revealed: an annotated and illustrated inventory of Nudipleura (Gastropoda: Heterobranchia) from Mozambique. *Zootaxa* **4359**:1–133. doi:10.11646/zootaxa.4359.1.1.
- Vaidya G, Lohman DJ, Meier R. 2011. SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* **27(2)**:171–180. doi:10.1111/j.1096-0031.2010.00329.x.
- Walsh PS, Metzger DA, Higuchi R. 1991. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques* **10(4)**:506–513.

## Supplementary materials

**Table S1.** Accession numbers of scyllaeids for three genes used in the phylogenetic analysis. (download)

**Table S2.** Uncorrected *COI* *p*-distances (%) calculated based on individual gene of the five available *Scyllaea* species in the NCBI and the sequences of specimens from this study. (download)

**Table S3.** Uncorrected *16S* *p*-distances (%) calculated based on individual gene of the five available *Scyllaea* species in the NCBI and the sequences of specimens from this study. (download)

**Table S4.** Uncorrected *H3* *p*-distances (%) calculated based on individual gene of the four available *Scyllaea* species in the NCBI and the sequences of specimens from this study. (download)