

EGU24-9993, updated on 15 Mar 2024

<https://doi.org/10.5194/egusphere-egu24-9993>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Understanding the paleogeographic evolution of the North Axial Channel, Southern North Sea

Morgan Vervoort¹, Despina Kyriakoudi², Ruth Plets², Thomas Mestdagh², Tine Missiaen², and Marc De Batist¹

¹Ghent University, Ghent, Belgium

²Flanders Marine Institute (VLIZ), Ostend, Belgium

During the last 500 000 years, ice sheets occupied parts of the North Sea during three major glaciations. The existence of these ice sheets was accompanied by a large fall in sea level, causing the Southern North Sea to emerge and become isolated from the Atlantic. In this area a complex drainage system was created by river water of the West-European rivers (e.g., Thames, Rhine, Meuse and Scheldt) and glacial meltwater. Furthermore, most offshore studies support the idea of the formation of large proglacial lakes in front of these ice sheets, which may have caused high-magnitude outburst floods at the end of each glacial period. The existence of such a proglacial lake is used in the argument that glacial outburst floods during the Elsterian (500-450 ka) created erosional features still preserved nowadays in the Dover Strait.

A remnant of this large, complex fluvial and glacial drainage system is the (North) Axial Channel, a prominent geomorphological feature seen on the present-day sea floor of the Southern North Sea. Its formation and evolution, however, are still uncertain. Previous studies state that the Axial Channel forms the northern extension of the Lobourg Channel, located in the Strait of Dover, which was formed during Middle Miocene times. Further erosion is assumed to have occurred during the Pliocene and Pleistocene, as sediments within the Murray Pit (located in the Axial Channel, about 100 km northeast of the Lobourg Channel) are assumed to be Early Pliocene, and no Quaternary infilled sediments have been identified. A series of NE-SW oriented scarps are identifiable from bathymetric and seismic reflection data and have been attributed to different Pleistocene incisional events. However, currently only a relative chronology of potential events has been established, with large uncertainties. Understanding the paleogeographic changes that affected the region also increases the knowledge on how early humans may have settled in and/or migrated through the region.

In the framework of the WALDO project (“**W**here are **A**ll the (proglacial) **L**ake **s**ediments in the **N**orth Sea Basin?”), a survey has been conducted in October 2023 during which high-resolution geophysical data (multibeam bathymetry and backscatter, acoustic and seismic data) combined with ground-truth data (vibrocores) have been acquired. One of the reflection-seismic grids was conducted ~40 km east of the East of England coast, over the western edge of the North Axial Channel, where also four sediment cores were taken. Here, we present the first interpretation of

these new data, which allow us to evaluate, update and improve the relative chronology of the formation of the (North) Axial Channel.□