

# Contributions on a small collection of the former Subulinidae Fischer & Crosse, 1877 (Eupulmonata, Achatinoidea) with catalogue of the *Glessula* and *Rishetia* species recorded from Myanmar

Nem Sian Man<sup>1,2</sup>, Jonathan D. Ablett<sup>3</sup>, Ngwe Lwin<sup>4</sup>, Chirasak Sutcharit<sup>1</sup>, Somsak Panha<sup>1,5</sup>

1 Animal Systematics Research Unit, Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok, 10330, Thailand

2 Department of Zoology, University of Yangon, University Avenue Road, Kamayut Township 11041, Yangon, Myanmar

3 Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

4 Fauna and Flora International, No. 35, 3<sup>rd</sup> Floor, Shan Gone Condo, Myay Ni Gone Market Street, Sanchaung Township, Yangon, Myanmar

5 Academy of Science, The Royal Society of Thailand, Bangkok 10300, Thailand

Corresponding authors: Chirasak Sutcharit ([chirasak.s@chula.ac.th](mailto:chirasak.s@chula.ac.th), [jirasak4@yahoo.com](mailto:jirasak4@yahoo.com)); Somsak Panha ([somsak.pan@chula.ac.th](mailto:somsak.pan@chula.ac.th), [somsak\\_panha@yahoo.com](mailto:somsak_panha@yahoo.com))



Academic editor: Menno Schilthuizen

Received: 22 November 2023

Accepted: 12 June 2024

Published: 30 July 2024

ZooBank: <https://zoobank.org/0215503D-E8B4-4179-89F6-FB2CCC59F37E>

**Citation:** Man NS, Ablett JD, Lwin N, Sutcharit C, Panha S (2024) Contributions on a small collection of the former Subulinidae Fischer & Crosse, 1877 (Eupulmonata, Achatinoidea) with catalogue of the *Glessula* and *Rishetia* species recorded from Myanmar. ZooKeys 1208: 173–239. <https://doi.org/10.3897/zookeys.1208.116083>

**Copyright:** © Nem Sian Man et al.  
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Abstract

The taxonomy of subulinid snails in Myanmar has been evaluated, resulting in the recognition of 40 species and subspecies across nine genera: *Allopeas*, *Bacillum*, *Curvella*, *Glessula*, *Opeas*, *Paropeas*, *Rishetia*, *Tortaxis*, and *Zootecus*. Nine species are re-described based on recently collected specimens, and two new species, *Glessula mandalayensis* Man & Panha, **sp. nov.** from Mandalay Region and *Tortaxis cylindropsis* Man & Panha, **sp. nov.** from Shan State are introduced. The genitalia and radula of *Zootecus pullus* was studied for the first time. This study also presents a comprehensive list of all subulinid species recorded to date from Myanmar. The type specimens and authenticated museum specimens have been illustrated with accompanying taxonomic remarks and nine species formerly assigned in *Glessula* are now placed in *Rishetia*: *R. akouktoungensis*, *R. baculina*, *R. basseinensis*, *R. burrailensis maxwelli*, *R. kentungensis*, *R. limborgi*, *R. nathiana*, *R. pertenuis*, and *R. pertenuis major*.

**Key words:** Conservation, molluscs, Southeast Asia, systematics, taxonomy, type specimen

## Table of contents

Introduction .....	175
Materials and methods.....	176
Sampling and morphological studies.....	176
Institutional abbreviations.....	177
Systematic account .....	178
Genus <i>Allopeas</i> Baker, 1935.....	178
1 <i>Allopeas gracile</i> (Hutton, 1834) .....	180
Genus <i>Bacillum</i> Theobald, 1870 .....	181
2 <i>Bacillum obtusum</i> (Blanford, 1869).....	181
3 <i>Bacillum theobaldi</i> (Hanley, 1870) .....	182
Genus <i>Curvella</i> Chaper, 1885 .....	182
4 <i>Curvella plicifera</i> (Blanford, 1865) .....	183
5 <i>Curvella pusilla</i> (Blanford, 1865).....	184
6 <i>Curvella puta</i> (Benson, 1857).....	184
7 <i>Curvella scrobiculata</i> (Blanford, 1865) .....	185
Genus <i>Glessula</i> von Martens, 1860 .....	188
8 <i>Glessula blanfordiana</i> Nevill, 1877 .....	190
9 <i>Glessula crassilabris</i> (Benson, 1836) .....	191
10 <i>Glessula feddeni</i> Godwin-Austen, 1920 .....	192
11 <i>Glessula gemma</i> (Reeve, 1850) .....	193
12 <i>Glessula inedita</i> Godwin-Austen, 1920.....	193
13 <i>Glessula latestriata</i> von Möllendorff, 1899 .....	194
14 <i>Glessula mandalayensis</i> Man & Panha, sp. nov. ....	195
15 <i>Glessula orophila</i> (Reeve, 1849) .....	196
16 <i>Glessula peguensis</i> (Blanford, 1865) .....	197
17 <i>Glessula perlevis</i> Godwin-Austen, 1920.....	197
18 <i>Glessula ponsiensis</i> Godwin-Austen, 1920.....	198
19 <i>Glessula woodthorpei</i> Godwin-Austen, 1920.....	200
20 <i>Glessula yuangensis</i> Godwin-Austen, 1920.....	203
Genus <i>Opeas</i> Albers, 1850 .....	206
21 <i>Opeas filiforme</i> von Möllendorff, 1894.....	206
22 <i>Opeas innocens</i> Preston, 1910 .....	207
Genus <i>Paropeas</i> Pilsbry, 1906 .....	207
23 <i>Paropeas swettenhami</i> (de Morgan, 1885).....	208
24 <i>Paropeas terebralis</i> (Theobald, 1870) .....	209
25 <i>Paropeas turricula</i> (von Martens, 1860).....	209
26 <i>Paropeas walkeri</i> (Benson, 1863).....	210
Genus <i>Rishetia</i> Godwin-Austen, 1920 .....	212
27 <i>Rishetia akouktoungensis</i> (Godwin-Austen, 1920), comb. nov. ....	213
28 <i>Rishetia baculina</i> (Blanford, 1871), comb. nov. ....	213
29 <i>Rishetia basseinensis</i> (Godwin-Austen, 1920), comb. nov. ....	214
30 <i>Rishetia burrailensis maxwelli</i> (Godwin-Austen, 1920), comb. nov. ...	214
31 <i>Rishetia hastula</i> (Benson, 1860).....	216
32 <i>Rishetia kentungensis</i> (Godwin-Austen, 1920), comb. nov. ....	217
33 <i>Rishetia limborgi</i> (Godwin-Austen, 1920), comb. nov. ....	217
34 <i>Rishetia nathiana</i> (Godwin-Austen, 1920), comb. nov. ....	218
35 <i>Rishetia pertenuis pertenuis</i> (Blanford, 1865), comb. nov. ....	218

36 <i>Rishetia pertenuis major</i> (Blanford, 1865), comb. nov. ....	219
37 <i>Rishetia tenuispira</i> (Benson, 1836).....	222
Genus <i>Tortaxis</i> Pilsbry, 1906.....	223
38 <i>Tortaxis cylindropsis</i> Man & Panha, sp. nov. ....	225
Genus <i>Zootecus</i> Westerlund, 1887.....	226
39 <i>Zootecus insularis</i> (Ehrenberg, 1831) .....	226
40 <i>Zootecus pullus</i> (Gray, 1834).....	229
Conclusions.....	230
Acknowledgements .....	231
Additional information.....	231
References.....	232

## Introduction

The Subulinidae Fischer & Crosse, 1877 is a highly diverse land snail family comprised of more than 800 species and approximately 80 genera which are native to tropical and subtropical regions in South and Central America, Africa, and South and Southeast Asia (Gude 1914; Naggs 1994; Schileyko 1999; D'ávila et al. 2020; Horsák et al. 2020). In general, subulinid snails are characterised by having a small to medium-sized (6–66 mm in height), ovate to slender, elongated conical shell, a mostly turreted spire, smooth to strong sculptures, and a concave or straight and truncated or continuous columella (Pilsbry 1906; Godwin-Austen 1920; Schileyko 1999; Do and Do 2014). The genitalia consist of a simple and slender to muscular penis, with (sometimes variously developed flagella) or without epiphallus (Gude 1914; Godwin-Austen 1920; Schileyko 1999; Budha 2017; Budha et al. 2017; D'ávila et al. 2018).

The earliest overview of molecular phylogenetic studies of the stylommatophorans revealed the 'achatinoideid' clade, which principally comprises the Achatinoidea Swainson, 1840 [including the subulinid snails] and the Streptaxoidea Gray, 1860 (Wade et al. 2001, 2006). It confirms the traditional and widely accepted hypothesis that the Subulinidae belong within the Achatinoidea. However, the systematic classification of the achatinoideans has varied considerably and has been subject to change throughout time. The name 'Subulininae' was initially nominated, and then was subsequently raised to the family level under the 'Achatinacea' [= Achatinoidea] (Fischer and Crosse 1877; Thiele 1931: 549). This classification was later followed by Zilch (1959) and Vaught (1989). At the same time, some studies considered 'subulinids' as a subfamily of the Achatinidae (Inkhavilay et al. 2019; Neubert and Bochud 2020; MolluscaBase 2023). Whereas Schileyko (1999) reclassified the 'subulinids' as its own superfamily consisting of the Subulinidae Glessulidae Godwin-Austen, 1920, Micractaeonidae Schileyko, 1999 and Ferrussaciidae Bourguignat, 1883, and recognised nine subfamilies within the Subulinidae. Recently, the multi-gene phylogeny of the Achatinoidea revealed the (sub)family as classified in Zilch (1959), Vaught (1989) and Schileyko (1999) are polyphyletic (Fontanilla et al. 2017). More recently, Bouchet et al. (2017) treated 'Subulinidae sensu by earlier authors' as a subfamily of the Achatinidae. Additionally, taxonomic placements within the Achatinoidea have also varied substantially since the revisions based on exhaustive anatomical characters had not been provided. Therefore, systematic classification of the Achatinoidea

remains a source of discussion but surpasses the scope of this work, which aims to highlight the Subulinidae from Myanmar, as classified in Zilch (1959) and Vaught (1989), and to attract and encourage further systematic research.

The subulinid snails are highly diverse and primarily found in Africa and also in Southeast Asia. In Myanmar, six genera are known: *Bacillum* Theobald, 1870, *Curvella* Chaper, 1885, *Glessula* von Martens, 1860, *Opeas* Albers, 1850, *Prosopeas* Mörch, 1876, and *Zootecus* Westerlund, 1887 with a total of 37 nominal species being documented (Gude 1914; Godwin-Austen 1920). Most of these nominal species have been described based solely on shell morphology, with uncertain taxonomic status and distribution due to the scarcity of detailed illustrations of their respective type specimens and a lack of newly collected samples for over a century since the work of Gude (1914) and Godwin-Austen (1920). Therefore, this work focuses on examining the historical museum specimens, both type and non-type, and also newly collected materials, with the aim to update the taxonomic knowledge of achatinoid snails from Myanmar. Recent specimen collections were conducted in collaboration with the Forest Department of Myanmar and Fauna and Flora International (FFI), as described in our previously published taxa (Man et al. 2022, 2023). This paper will enhance the knowledge of the Subulinidae diversity in Myanmar, contribute to our understanding of land snail biogeography and will serve as a valuable resource for the improvement of the existing taxonomy of the subulinid taxa.


## Materials and methods

### Sampling and morphological studies

Between 2015 and 2016, malacofauna surveys of the limestone habitats in Myanmar were resumed through collaborative efforts involving the Animal Systematics Research Unit (ASRU), the Forest Department of Natural Resources and Environmental Conservation and Forestry, Myanmar, and Fauna and Flora International (FFI), under the framework of an MOU (Letter No. 0092). During field trips, small samples of subulinid snails were primarily collected in Shan State and the Mandalay Region within the northeastern part of the country, as well as in Kayin and Mon states and the Tanintharyi Region in the southeast, as shown in Fig. 1.

These newly collected samples were deposited in the Chulalongkorn University, Museum of Zoology (**CUMZ**), Thailand. The specimens were examined for shell morphology (Fig. 2) and reproductive system structure (when available), and the terminology used in this study follows Gude (1914), Godwin-Austen (1920), Schileyko (1999), and Budha et al. (2017). Specimens were compared to the type specimens and available authenticated collections. Adult and intact shells were measured with a digital calliper for shell height (**SH**; largest length from the apex to the base of aperture), shell width (**SW**; widest diameter from one side of the last whorl to the outermost side of aperture), and whorl counts (Kerney and Cameron 1979; Table 1).

The following species list was produced using both the literature and collections, the primary type specimens (i.e., holotype, lectotype, syntype(s), and neotype) and secondary type specimens [paratype(s) and paralectotype(s)]. The taxa are arranged in alphabetical order according to their current taxonomic status. The references for the usage of each taxon name are provided herein. The name



**Figure 1.** Approximate collecting localities of the subulinids from Myanmar examined in this study (right), inset figure (upper right) indicates the collecting site from Shan State, and living snails (lower right) of *Glessula mandalayensis* sp. nov., paratype CUMZ 13073 from Pyinyaung Village, Mandalay Region, Myanmar (SH ~ 14 mm). The numbers correspond to localities listed in Table 1.

in the original combination is given with bibliographic information on the original description. The type locality is given, and if possible, the modern name and/or regional names of the type locality are provided in square brackets. Where necessary, remarks are given on the status of type specimens, authorships, availability of name, notes on the type locality, and further valuable comments.

### Institutional abbreviations

- ANSP** Academy of Natural Science of Philadelphia, Drexel University, Philadelphia
- B.M.** used in old labels for the British Museum (now The Natural History Museum (NHM), London)
- CUMZ** Chulalongkorn University Museum of Zoology, Bangkok
- MNHN** Muséum National d'Histoire Naturelle, Paris
- NHMK** The Natural History Museum, London
- SM** Forschungsinstitut und Naturmuseum Senckenberg, Frankfurt am Main
- UMZC** University Museum of Zoology Cambridge, Cambridge
- ZMB** Museum für Naturkunde, Humboldt University, Berlin

**Table 1.** Shell measurements of subulinid species examined in this study. SH: shell height; SW: shell width. The localities numbers are indicated in Fig. 1.

Species, locality no., and CUMZ no.	No. of specimens	Ranges, mean $\pm$ S.D. in mm		SH/SW ratio	No. of whorls
		Shell height	Shell width		
<i>Allopeas gracile</i>					
1. Pyinyaung Village, Meiktila, Mandalay (13066)	8	9.8–12.1 10.76 $\pm$ 0.92	2.9–3.3 3.08 $\pm$ 0.169	3.37–3.78 3.48 $\pm$ 0.17	7½–8
<i>Glessula blanfordiana</i>					
2. Aik Kham Cave, Taunggyi, Shan (13067)	3	7.7–8.6 8.16 $\pm$ 0.47	3.7–3.9 3.81 $\pm$ 0.08	2.05–2.26 2.141 $\pm$ 0.10	6–6½
<i>Glessula feddeni</i>					
1. Pyinyaung Village, Meiktila, Mandalay (13068)	8	7.8–10.8 9.72 $\pm$ 1.14	4.0–4.8 4.47 $\pm$ 0.33	1.95–2.25 2.16 $\pm$ 0.11	6½–7
<i>Glessula gemma</i>					
3. Montawa Cave, Taunggyi, Shan (13069)	5	8.7–9.2 8.99 $\pm$ 0.28	3.4–3.9 3.61 $\pm$ 0.22	2.37–2.58 2.49 $\pm$ 0.09	6½–7
<i>Glessula latestriata</i>					
4. Thale Cumon Temple, Kalaw City, Shan (13070)	4	8.4–10.8	4.0–4.7	1.86–2.31	6½–7
5. Ywangan Village, near Lin Way Monastery, Kalaw, Shan (13071)		9.22 $\pm$ 1.14	4.51 $\pm$ 0.18	2.04 $\pm$ 0.23	
<i>Glessula mandalayensis</i> sp. nov.					
1. Pyinyaung Village, Meiktila, Mandalay (13072, 13073)	30	13.3–16.0 14.32 $\pm$ 0.72	6.7–7.8 7.19 $\pm$ 0.30	1.87–2.12 1.99 $\pm$ 0.07	6–7
<i>Opeas filiforme</i>					
6. Phra (Buddha) Cave, Tanintharyi (13074)	15	8.1–11.2 9.51 $\pm$ 1.00	2.4–3.4 2.82 $\pm$ 0.30	3.32–3.49 3.37 $\pm$ 0.05	7½–8
<i>Paropeas turricula</i>					
6. Phra (Buddha) Cave, Tanintharyi (13075)	3	11.7–15.0	3.00–3.90	3.78–3.9	8, 8½
7. Lampane Village, Tanintharyi (13076)		13.00 $\pm$ 1.72	3.4 $\pm$ 0.45	3.84 $\pm$ 0.58	
<i>Paropeas walkeri</i>					
2. Aik Kham Cave, Taunggyi, Shan (13077)	5	8.4–15.3	3.3–3.8	2.54–4.02	8–8½
5. Ywangan Village, near Lin Way Monastery, Kalaw, Shan (13078)		12.65 $\pm$ 1.76	3.54 $\pm$ 0.22	3.54 $\pm$ 0.56	
<i>Tortaxis cylindropsis</i> sp. nov.					
8. Parpant area, Taunggyi, Shan (13079, 13080)	35	9.3–11.7 10.29 $\pm$ 1.06	2.1–2.5 2.27 $\pm$ 0.16	4.11–5.00 4.53 $\pm$ 0.39	8–9½
<i>Zootecus pullus</i>					
9. Dhammayazaka Pagoda, Bagan, Mandalay (13081, 13082)	90	11.2–13.3 11.99 $\pm$ 0.75	4.3–4.8 4.55 $\pm$ 0.16	2.48–2.80 2.63 $\pm$ 0.14	8–9½

## Systematic account

### Family Subulinidae Fischer & Crosse, 1877

#### Genus *Allopeas* Baker, 1935

*Lamellaxis* (*Allopeas*) Baker, 1935: 84. Zilch 1959: 349.

*Allopeas*—Schileyko 1999: 509.

**Type species.** *Bulimus gracilis* Hutton, 1834, by original designation.

**Diagnosis.** Shell slender and conical; spire high and gradually attenuated; embryonic whorls pointed and smooth; subsequent whorls with striations.

Aperture vertical, broad, and oblong; columella straight, and columellar margin expanded near umbilicus. Penis long, fusiform shape at base then narrowing towards epiphallus, and flagellum absent; vagina cylindrical and narrow tube, and ~ 1/2 of penis length.

**Remarks.** The genus is sometimes confused with *Paropeas* Pilsbry, 1906 and *Opeas* Albers, 1850. *Allopeas* can be distinguished from *Paropeas* by its less turreted shell, finer striations, smooth embryonic whorls, straight columella, and columellar margin expanded (Table 2). *Paropeas* possesses a mostly turreted shell, stronger irregular striations throughout and entire whorls, concave columella, and columellar margin less expanded. Additionally, *Allopeas* can be differentiated from *Opeas* by having larger and broader shell, attenuated spire, and stronger striations (Schileyko 1999). By contrast, *Opeas* has a smaller and narrower shell, cylindrical or less attenuated spire, and finer striations (Pilsbry 1906).

The genus *Allopeas* consists of ~ 25 species distributed across tropical regions of Asia, Europe, Africa, and America (Schileyko 1999; MolluscaBase 2023). In Southeast Asia, Vietnam has recorded four species (Schileyko 2011), while Cambodia, Laos, Myanmar, and Thailand have reported only one species, namely *Allopeas gracile* (Gude 1914; Inkhavilay et al. 2019; Sutcharit et al. 2020b).

**Table 2.** Comparison of shell morphology among subulinid genera recorded in Myanmar. The superscript numbers are references; <sup>1</sup> = Pilsbry (1906), <sup>2</sup> = Gude (1914), <sup>3</sup> = Godwin-Austen (1920), <sup>4</sup> = Naggs (1994), <sup>5</sup> = Schileyko (1999), and <sup>6</sup> = Budha et al. (2017).

Genus (type species)	Shell shape	Spire	Sculpture		Aperture shape	Columella
			Embryonic whorl	Spire whorl		
<i>Allopeas (Bulimus gracilis</i> Hutton, 1834) <sup>5</sup>	slender conical	high and gradually attenuated	smooth	striations	vertically broad; oblong	straight
<i>Bacillum (Achatina cassiaca</i> Reeve, 1849) <sup>1</sup>	slender conical	high, turreted, and gradually attenuated	striations	obliques striations or ribs	obliquely narrow; ovate	concave and truncated
<i>Curvella (Curvella sulcata</i> Chaper, 1885) <sup>1,5</sup>	oblong-conical	low or high conical and rapidly attenuated	smooth	equally or irregularly spaced striations or ribs	vertically broad; ovate; oblong	straight
<i>Glessula (Achatina gemma</i> Reeve, 1850) <sup>3,6</sup>	ovate-conical	low conical, and gradually attenuated	smooth or striations	equally spaced of striations; grooves; radial ribs	obliquely narrow or broad; ovate	concave and truncated
<i>Rishetia (Achatina tenuispira</i> Benson, 1836) <sup>3,6</sup>	slender conical	high, turreted, and gradually attenuated	smooth or striations	equally or irregularly spaced striations or ribs	obliquely, narrow; ovate	concave and truncated
<i>Opeas (Helix goodallii</i> Miller, 1822) <sup>1,5</sup>	slender conical	low or high, turreted, and gradually attenuated	smooth	smooth; fine striations and growth lines	vertically narrow; oblong	concave or straight
<i>Paropeas (Bulimus acutissimus</i> Mousson, 1857) <sup>1,2,4</sup>	slender conical	high, turreted, and gradually attenuated	striations	irregular, dense, fine or coarse striations	obliquely narrow or broad; ovate	concave or straight
<i>Tortaxis (Achatina erecta</i> Benson, 1842) <sup>1</sup>	cylindrical to slender conical	high, mostly turreted, narrowly or cylindrically attenuated	smooth	striations and growth lines	vertically narrow; oblong	concave or straight and with spiral fold below
<i>Zootecus (Pupa insularis</i> Ehrenberg, 1831) <sup>5</sup>	pupiform	high, broad, and cylindrical	smooth	irregular, dense, fine or coarse striations	obliquely broad; oblong; rounded	straight

### 1 *Allopeas gracile* (Hutton, 1834)

Fig. 3A–D, Table 1

*Bulimus* (?) *gracilis* (?) Hutton, 1834: 84, 85, 93. Type locality: Mirzapoor, Futtehpoor Sikra, between Agra and Neemuch [Uttar Pradesh and Madhya Pradesh states, India].

*Bulimus* (*Opeas*) *gracilis*—Pfeiffer 1856: 156.

*Stenogyra* (*Opeas*) *gracilis*—von Martens 1860: 265. Pfeiffer and Clessin 1881: 321.

*Spiraxis gracilis*—Blanford 1861: 362.

*Stenogyra gracilis*—de Morgan 1885: 389, 390.

*Opeas gracilis*—Theobald 1878: 146. Godwin-Austen 1895: 443.

*Opeas gracile*—von Möllendorff 1894: 151. Pilsbry 1906: 125–132, pl. 18, figs 3–6. Gude 1914: 355–357. van Benthem Jutting 1952: 378–380.

*Lamellaxis gracilis*—van Benthem Jutting 1959: 131, 132.

*Lamellaxis* (*Allopeas*) *gracile*—Solem 1966: 94.

*Allopeas gracilis*—Maassen 2001: 81, 82.

*Allopeas gracile*—Schileyko 2011: 9. Do and Do 2014: 452, fig. 1a. Raheem et al. 2014: 117, 118, fig. 73d–f. Inkhavilay et al. 2019: 50, fig. 21a–c. Sutcharit et al. 2020b: 19.

**Type specimens.** **Lectotype** NHMUK 1856.9.15.68/1 (Fig. 3A) ex. Hutton collection from Mirzapore, designated in Raheem et al. (2014: 118). **Paralectotypes** NHMUK 1856.9.15.68/2–11 (10 shells; Fig. 3B).

**Other material.** Limestone hills (Apache Cement Factory), Pyinyaung Village, Meiktila District, Mandalay Region, Myanmar (20°49'39.1"N, 96°23'35.1"E): CUMZ 13066 (8 shells; Fig. 3C, D).

**Description.** Shell conically elongated, narrow, translucent, glossy, pale yellowish colour, and with 7½–8 whorls. Apex blunt; protoconch ~ 2 whorls and smooth; subsequent whorls with obliquely more or less fine and crowded riblets. Spire gradually tapering; whorls flatly convex; suture wide and shallow; last whorl largest. Aperture high and oblong shape; peristome thin and columellar margin near umbilicus is little expanded; columella straight. Umbilicus narrowly opened.

**Distribution.** *Allopeas gracile* is distributed worldwide in America, Africa, Europe, and Asia (Gude 1914; Horsák et al. 2020). In South Asia, it has been reported from India, Pakistan, Sri Lanka, Nepal, and all countries in Southeast Asia (van Benthem Jutting 1952; Panha 1998; Schileyko 2011; Do and Do 2014; Raheem et al. 2014; Budha et al. 2015; Foon et al. 2017; Inkhavilay et al. 2019; Nurinsiyah and Hausdorf 2019; Sutcharit et al. 2020b).

This species was previously reported in Myanmar from several localities, including Kayin State, Mon State, Kachin State, Rakhine State, Yangon Region, and Bago Region (see Gude 1914) and presently in the Mandalay Region.

**Remarks.** Our material from the Mandalay Region matches well with the type specimen of this species. *Allopeas gracile* is a globally distributed alien species introduced to numerous countries and exhibits significant variability in both size and shape, influenced by its expansive distribution range. This species is commonly found in public parks, irrigation areas, greenhouse environments, and residential campuses (Pilsbry 1906; Neubert and Bochud 2020; Kalita 2022).

## Genus *Bacillum* Theobald, 1870

*Achatina* (*Bacillum*) Theobald in Hanley and Theobald 1870: 17.

*Bacillum*—Pilsbry 1906: 1. Gude 1914: 343. Godwin-Austen 1920: 7. Zilch 1959: 346. Schileyko 1999: 534.

**Type species.** *Achatina cassiaca* Reeve, 1849a, subsequent designation by Pilsbry (1906: 1).

**Diagnosis.** Shell slender and conical in shape; spire high, turreted, and gradually attenuated; embryonic whorls cylindrically rounded, and with or without radial striations; subsequent whorls with equally spaced thick or fine radial striations. Aperture oblique and narrowly ovate; columella concave and truncated, and columellar margin simple or slightly expanded.

**Remarks.** *Bacillum* can be differentiated from *Allopeas* in being cylindrically rounded and with striations on the embryonic whorls, whilst the columella is concave and truncated (Pilsbry 1906; Schileyko 1999). While *Allopeas* has a narrowly attenuated and smooth embryonic whorls, columella straight, and columellar margin near umbilicus expanded (Table 2).

At present, this genus contains seven species mainly distributed in India, and among these, two species are known from Myanmar (Blanford 1869; Pilsbry 1906; Gude 1914; Ramakrishna et al. 2010; MolluscaBase 2023). All *Bacillum* species are known only from shell morphology, and none of the reproductive anatomy has been published so far. Godwin-Austen (1920) stated, ‘...extended knowledge of the animals of *Bacillum* and *Glessula* shows that the two genera come next each other...’, which suggests Godwin-Austen had dissected *Bacillum*, but his findings were never published.

### 2 *Bacillum obtusum* (Blanford, 1869)

Fig. 3E–H

*Achatina* (*Glessula*) *obtusum* Blanford, 1869: 449. Type locality: Bhamo in regno Avaë [Bhamo District, Kachin State, Myanmar].

*Achatina* (*Bacillum*) *obtusum*—Hanley and Theobald 1870: 17, pl. 36, fig. 6.

*Achatina obtusum*—Pfeiffer 1876: 290.

*Glessula obtusum*—Nevill 1877: 25.

*Stenogyra* (*Subulina*) *obtusum*—Pfeiffer and Clessin 1881: 327.

*Bacillum obtusum*—Pilsbry 1906: 1, 2, pl. 1, fig. 1. Gude 1914: 347.

**Type specimens.** **Syntypes** NHMUK 1906.2.2.349 [re-registered in error as 19850143] (4 shells; Fig. 3E, F) ex. Blanford collection from Bhamo, Upper Burmah.

**Other material.** NHMUK 1903.7.1.1725 (2 shells + 1 juvenile; Fig. 3G, H) ex. Godwin-Austen collection from Bhamo, Upper Burma. NHMUK 18/88.12.4.1012–1014 (3 shells) from Bhamo. NHMUK (1 shell) ex. TV Oldham collection from Bhamo, Burma. SMF 145947/2 (2 shells) ex. von Möllendorff collection from Bhamo.

**Diagnosis.** Shell slender, elongate turreted and apical whorls rapidly attenuated; apex rounded, blunt and very large embryonic shell; subsequent whorls with fine radial striae. Suture shallow and whorls flattened. Aperture obliquely subovate; columella curved and truncated.

**Distribution.** This species is known only from the type locality.

**Remarks.** No new material of this species was collected in this study, but the syntypes are illustrated for the first time.

### 3 *Bacillum theobaldi* (Hanley, 1870)

Fig. 3I

*Achatina (Electra) theobaldi* Hanley in Hanley and Theobald 1870: 9, pl. 17, fig. 5. Type locality: Near the Salwen [near Salween River, Myanmar].

*Achatina (Glessula) theobaldiana* [sic]—Theobald 1870: 395.

*Achatina theobaldi*—Pfeiffer 1876: 290.

*Stenogyra (Glessula) theobaldiana*—Nevill 1878: 172.

*Stenogyra (Subulina) theobaldi*—Pfeiffer and Clessin 1881: 327.

*Bacillum theobaldi*—Pilsbry 1906: 4, pl. 1, fig. 8. Gude 1914: 344, 345. Ramakrishna et al. 2010: 150.

**Other material.** NHMUK 1912.4.16.121 (1 shell; Fig. 3G) ex. Beddome collection from Burmah.

**Diagnosis.** Shell elongate turreted and gradually attenuated; apex rounded, blunt and very large embryonic shell; subsequent whorls with strong and equally spaced radial ribs throughout. Aperture ovate; columella curved and truncated.

**Distribution.** This species was recorded from near the Salween River and in Shan State, Myanmar (Hanley and Theobald 1870; Pilsbry 1906). The additional report from 'India' by Ramakrishna et al. (2010) is possibly erroneous.

**Remarks.** No new material of this species was examined. *Bacillum theobaldi* was initially proposed without a proper description other than comparing it with *B. cassiacum* (Reeve, 1849a) and with an imprecise type locality noted as 'Near the Salwen' (Hanley and Theobald 1870). Pilsbry (1906) confined the type locality as near to the Salween River in Shan State.

This species is superficially similar to *B. obtusum* in shell shape. However, *B. theobaldi* has equally spaced radial ridges (Fig. 3I), whereas *B. obtusum* processes finer radial striations (Fig. 3H).

### Genus *Curvella* Chaper, 1885

*Bulimus (Hapalus)* Albers, 1850: 140. [non Illiger 1801 (Coleoptera)]. Type species *Bulimus grateloupi* Pfeiffer, 1846. Albers 1860: 238.

*Curvella*—Chaper, 1885: 49. Pilsbry 1906: 46. Gude 1914: 348. Zilch 1959: 351. Schileyko 1999: 515. Budha et al. 2017: 120.

**Type species.** *Curvella sulcata* Chaper, 1885, by original designation in Chaper (1885: 48).

**Diagnosis.** Shell oblong-conical; spire low or high and rapidly attenuated; embryonic whorls smooth; subsequent whorls with equally or irregularly spaced thick or fine radial striations or ribs. Aperture vertical, broad-ovate, or oblong,

and somewhat pointed above; columella straight; columellar margin expanded near umbilicus. Penis simple and short papillate or short tube; epiphallus and flagellum absent; vagina large, muscular, and nearly equal to penis length.

**Remarks.** *Curvella* is clearly distinct from other subulinid genera such as *Allopeas*, *Bacillum*, and *Opeas* by its oblong-conical shell, short spire, high and wide aperture, and its much broader last whorls (Table 2). Regarding the genitalia, this genus displays a more rounded, muscular, short penis and a thicker vagina than *Allopeas* and *Opeas* (Schileyko 1999). In comparison, the two latter genera have a slender, narrow, more elongated penis and vagina compared to *Curvella* (Schileyko 1999; Budha 2017). However, only genitalia of *Curvella sikkimensis* (Reeve, 1850) is known for this genus at present (Budha 2017).

This genus is widely distributed from Africa to South Asia and China, and ~ 95 species have been reported (Pilsbry 1906; Gude 1914; Schileyko 1999; Ramakrishna et al. 2010; Budha et al. 2017; MolluscaBase 2023). There are scattered reports of four *Curvella* species in Myanmar, while one species, *C. tonkiniana* Jaeckel, 1950, has been reported in Vietnam and one species, *C. jousseaumei* (de Morgan, 1885), from Peninsular Malaysia (Gude 1914; Maassen 2001; Schileyko 2011; Vermeulen et al. 2015).

#### 4 *Curvella plicifera* (Blanford, 1865)

Fig. 4A

*Bulimus plicifer* Blanford, 1865: 77. Type locality: Thayet Myo, Pegu. Pfeiffer 1868: 151. Hanley and Theobald 1874: 34, pl. 80, fig. 8.

*Bulimina (Hapalus) plicifera*—Pfeiffer and Clessin 1881: 300.

*Buliminus* (?) *plicifer*—Kobelt 1901: 688, 689, pl. 103, fig. 22.

*Curvella plicifera*—Pilsbry 1906: 63, pl. 9, fig. 45. Gude 1914: 352.

**Type specimens. Syntypes** NHMUK 1906.2.2.235 (2 shells; Fig. 4A) ex. Blanford collection from Thyetmyo, Pegu? Akouktoung.

**Diagnosis.** Shell ovate conic; spire low; apex bluntly obtuse; subsequent whorls with prominent growth lines throughout. Suture shallow and whorls flattened. Aperture broadly ovate and somewhat pointed posteriorly; columella straight and expanded; parietal callus thin and with small parietal lamella. Umbilicus narrow.

**Distribution.** This species is restricted to Myanmar and is known only from the type locality.

**Remarks.** No new material of this species was found in this study, and only the available syntypes are examined here. The modern name of the type locality is Thayet District, Magway Region, central Myanmar [not in the Bago Region].

Among the four *Curvella* species from Myanmar, this species is similar to *C. pusilla* (Blanford, 1865) in shell shape. However, *C. plicifera* has a small parietal tooth, straight columella, and narrowly opened umbilicus, whereas *C. pusilla* possesses a smooth parietal wall (without tooth), slightly twisted columella plait, and closed umbilicus. Further confirmation from additional specimens is necessary to verify their distinctions.

### 5 *Curvella pusilla* (Blanford, 1865)

Fig. 4B, C

*Spiraxis pusilla* Blanford, 1865: 78. Type locality: Prome district, Pegu [Pyay District, Bago Region, Myanmar]. Pfeiffer 1868: 192. Hanley and Theobald 1874: 34, pl. 79, fig. 8.

*Hapalus pusillus*—Nevill 1878: 174.

*Stenogyra* [*Spraxis* (*Euspiraxis*)] *pusilla*—Pfeiffer and Clessin 1881: 324.

*Curvella pusilla*—Pilsbry 1906: 64, pl. 9, fig. 48. Gude 1914: 351.

**Type specimens. Possible syntypes** NHMUK 1906.2.2.234 (2 shells; Fig. 4B) ex. Blanford collection from Akouktong, Pegu.

**Other material.** NHMUK 1888.12.4.1003–1004 (1 shell identified as *C. pusilla*; Fig. 4C [another shell identified as *C. puta*]) ex. W. Theobald collection from Pegu.

**Diagnosis.** Shell ovate conic; spire low; apex bluntly obtuse; subsequent whorls with prominent growth lines. Suture shallow and whorls slightly flattened. Aperture broadly ovate and pointed posteriorly; columella twisted and expanded; parietal callus thin. Umbilicus closed.

**Distribution.** This species seems restricted to Myanmar and is known only from the type locality. The modern administrations of the type locality are 'Prome district' [Pyay District] and 'Pegu' [Bago Region].

**Remarks.** No new material of this species was found in this study, and only the probable syntypes were examined. These are considered possible syntype material since the locality data (Akouktong, Pegu) does not precisely match that which was given in the original description (Prome district, Pegu).

Blanford (1865) noted that *C. pusilla* resembles the young specimen of *C. puta* (Benson, 1857), but this species has a closed umbilicus, while *C. puta* possesses an opened umbilicus. We recognise *C. pusilla* as a valid species following Blanford's description, and no new materials of this species were examined in this study.

### 6 *Curvella puta* (Benson, 1857)

Fig. 4D, E

*Bulimus putus* Benson, 1857: 330. Type locality: Tavoy [Dawei, Tanintharyi Region, Myanmar]. Pfeiffer 1859: 502. Blanford 1865: 94. Hanley and Theobald 1874: 34, pl. 80, fig. 9.

*Hapalus putus*—Nevill 1878: 178.

*Bulimina* (*Hapalus*) *puta*—Pfeiffer and Clessin 1881: 299.

*Buliminus* (?) *putus*—Kobelt 1901: 689, pl. 103, fig. 23.

*Curvella puta*—Pilsbry 1906: 63, 64, pl. 9, fig. 46. Gude 1914: 351, 352. Preece et al. 2022: 130, fig. 55f.

**Type specimen. Holotype** UMZC I.102795 (Fig. 4D; after Preece et al. 2022: fig. 55f) ex. R. McAndrew ex. Benson collection from Tavoy, Birmah.

**Other material.** NHMUK 1888.12.4.1003–1004 (1 shell identified as *C. puta*; Fig. 4E [another shell identified as *C. pusilla*]) ex. W. Theobald collection from Pegu. NHMUK 1906.1.1.1033 (1 shell) ex. Godwin-Austen collection from Bassein, Pegu.

**Diagnosis.** Shell conical; spire high; apex obtuse; subsequent whorls with strong and prominent growth lines. Suture impressed and whorls slightly convex. Aperture semi-ovate and somewhat pointed above; columella straight and dilate; parietal callus thin. Umbilicus narrow.

**Distribution.** This species was reported from the Tanintharyi and Bago regions of Myanmar and Thailand (Blanford 1865; Gude 1914; Panha 1998).

**Remarks.** *Curvella puta* was described based on a single bleached specimen collected from 'Tavoy' in Myanmar. No new material of this species was found during this survey, but the type specimens and authenticated museum specimens are illustrated herein.

There is a mixed-species lot, NHMUK 1888.12.4.1003–1004, ex. the W. Theobald collection, labelled '*Hapalus putus* Benson' from 'Pegu', consisting of two shells. The first shell (Fig. 4E) has a high, turreted, and pointed spire, narrowly opened umbilicus, and strong radial striations, which agree with the holotype of *C. puta* from 'Tavoy' (Fig. 4D). The other shell (Fig. 4C) has a lower and convex spire, fine radial striations, narrowly opened umbilicus, and without a parietal tooth, which more resembles *C. pusilla*. In comparison, *C. pusilla* possesses a closed umbilicus without a parietal tooth (Fig. 4B), while *C. plicifera* has an open umbilicus and a small parietal tooth (Fig. 4A).

This mixed-species lot suggests several interpretations: i) an extended distribution of *C. puta* beyond its type locality to the Bago Region in central Myanmar, and ii) the presence or absence of a parietal tooth and an open or closed umbilicus are possibly intraspecific variation rather than diagnostic characters distinguishing between *C. plicifera* and *C. pusilla*. Additional evidence and further specimens from a wider geographic range will clarify this issue.

### 7 *Curvella scrobiculata* (Blanford, 1865)

Fig. 4F

*Bulimus scrobiculatus* Blanford, 1865: 77. Type locality: Pegu, west of Irawady [Bago Region, west of Ayeyarwady River, Myanmar]. Pfeiffer 1868: 151. Hanley and Theobald 1874: 34, pl. 79, fig. 9.

*Bulimina (Hapalus) scrobiculata*—Pfeiffer and Clessin 1881: 300.

*Hapalus scrobiculatus*—Nevill 1878: 175.

*Curvella scrobiculata*—Pilsbry 1906: 64, 65, pl. 9, fig. 49.


*Curvella scrobiculatus*—Gude 1914: 350, 351.

**Type specimens. Possible syntypes** NHMUK 1906.2.2.226 (2 shells; Fig. 4F) ex. Blanford collection from Akouktong, Pegu.

**Diagnosis.** Shell conical; spire high; apex obtuse; subsequent whorls with strong and equally spaced radial ridges throughout. Suture impressed and whorls flattened convex. Aperture truncate and somewhat pointed posteriorly; columella straight and thickened; parietal callus thin. Umbilicus narrowly open.


**Distribution.** This species is known only from the type locality and is probably an endemic species in Myanmar.

**Remarks.** No new material of this species was found in this study, and the probably syntype is illustrated here for the first time. Blanford (1865) described three *Curvella* species, *C. plicifera*, *C. pusilla*, and *C. scrobiculata*, from Myanmar




**Figure 2.** Schematic shell morphology of the subulinid snails **A** protoconch sculptures as recognised in this study: smooth and striate **B** crenulate suture **C** columella shape as recognised in this study: straight, truncate and fold **D** shell surface sculpture of the *Glessula* as recognised in this study: smooth, groove, and ribbed.

consecutively in the same publication with their type localities as 'Pegu' [now in the Bago Region]. Compared to the authenticated museum specimens, *C. scrobiculata* is distinct from *C. puta* by having strong radial ribs, flatter whorls, and a more turreted, pointed, and higher spire. Furthermore, *C. scrobiculata* is easily distinguishable from *C. plicifera* and *C. pusilla* by its higher, pointed, turreted spire and prominent radial ribs. In contrast, the two latter species exhibit



**Figure 3.** **A–D** *Allopeas gracile* **A** lectotype NHMUK 1856.09.15.68/1 from Mirzapore **B** paralectotypes NHMUK 1856.09.15.68/2–11 and **C, D** specimen CUMZ 13066 from Mandalay Region, Myanmar with embryonic whorls and aperture **E–H** *Bacillum obtusum* **E, F** syntype NHMUK 1906.2.2.349 from Bhamo, Upper Burmah **G, H** specimen NHMUK 1903.7.1.1725 from Bhamo, Upper Burma and **H** juvenile shell with last whorl **I** *Bacillum theobaldi*, syntype NHMUK 1912.4.16.121 from Burmah with last whorl.

a lower and more convex spire with finer radial striations to nearly smooth shell surfaces. These are considered possible syntype material since the locality data (Akouktong, Pegu) does not exactly match that which was given in the original description (Pegu, west of Irawady).



**Figure 4.** **A** *Curvella plicifera*, possible syntype NHMUK 1906.2.2.235 from Thyetmyo, Pegu; Akouktoung **B, C** *Curvella pusilla* **B** possible syntype NHMUK 1906.2.2.234 from Akouktong, Pegu and **C** specimen NHMUK 1888.12.4.1003–1004 from Pegu **D, E** *Curvella puta* **D** holotype UMZC I.102795 from Tavoy, Birmah (after Preece et al. 2022: fig. 55f) and **E** specimen NHMUK 1888.12.4.1003–1004 from Pegu **F** *Curvella scrobiculata*, possible syntype NHMUK 1906.2.2.226 from Akouktong, Pegu.

#### Genus *Glessula* von Martens, 1860

*Electra* Albers, 1850: 194. [non Lamouroux 1816 (Bryozoa)]. Type species: *Achatina ceylanica* Pfeiffer, 1845. Adams and Adams 1855: 105. Pfeiffer 1856: 168. *Cionella* (*Glessula*)—von Martens in Albers 1860: 254. Pfeiffer and Clessin 1881: 329.

*Stenogyra* (*Glessula*)—Nevill 1878: 166.

*Glessula*—Tenison-Woods 1888: 1053. Beddome 1906: 160. Pilsbry 1908: 50. Gude 1914: 377. Zilch 1959: 343. van Benthem Jutting 1952: 373, 374. Schileyko 2011: 11.

**Type species.** *Achatina gemma* Reeve, 1850 by original designation.

**Diagnosis.** Shell ovate-conical in shape; spire low or high conical, and regularly attenuated; embryonic whorls smooth or with striations; subsequent whorls with equally or irregularly spaced radial striations or ribs. Aperture oblique, narrow to wide, and ovate; columella concave and truncated, and columellar margin not expanded. Penis large, thick, and moderately long; epiphallus short, stout, terminally pointed, and curved; flagellum present (rarely absent) with comb-like or hand-like structures; epiphallic caecum absent; vagina muscularly and nearly equal to  $\sim 1/5$ – $2/5$  of penis length.

**Remarks.** Superficially, *Glessula* and *Rishetia* are similar in having a concave and truncated columella, but they differ in their shell shape and genital structure (Godwin-Austen 1920; Raheem et al. 2014; Budha et al. 2017). In terms of shell morphology, *Glessula* can be differentiated by its ovate-conical shape, short spire, large and blunt embryonic whorls, while *Rishetia* possesses a slender, conical-shaped elongate spire which narrow and rather acute embryonic whorls (Table 2). Regarding the genital characters, *Rishetia* exhibits a simple, tubular-shaped flagellum with the epiphallic caecum present (Table 3), while *Glessula* displays a well-developed comb-like flagellum (appendage) and the epiphallic caecum absent (Budha et al. 2017); except *G. mandalayensis* sp. nov. which has no flagellum. Likewise, *Glessula* is also distinct from *Curvella* by its concave and truncated columella, not expanded columellar margin, and obliquely narrow and ovate aperture, while *Curvella* possesses a straight columella, expanded columellar margin, and more vertical, higher, and broader oblong aperture.

**Table 3.** Comparison of terminal part of male genitalia between *Glessula* and *Rishetia*. References: 1 = Semper (1874); 2 = Godwin-Austen (1918); 3 = Godwin-Austen (1920); 4 = Schileyko and Kuznetsov (1996); 5 = Schileyko (1999); 6 = Budha et al. (2017). n/a = data not available (not mentioned in the description or could not differentiated from the original figure).

Species	Epiphallic caecum	Flagellum	References
<b><i>Glessula</i> von Martens, 1860</b>			
<i>G. hebetata</i> Godwin-Austen, 1920	absent	comb-like with numerous notches	6
<i>G. inornata</i> (Pfeiffer, 1853)	absent	comb-like with numerous notches	3
<i>G. oakesi</i> Godwin-Austen, 1918	absent	comb-like with three notches	2, 3
<i>G. ochracea</i> Godwin-Austen, 1918	absent	comb-like with numerous notches	2, 3
<i>G. orobia</i> (Benson, 1860)	absent	comb-like with five notches	3, 6
<i>G. orophila</i> (Reeve, 1849)	absent	comb-like with numerous notches	1
<i>G. serena</i> (Benson, 1860)	absent	comb-like with numerous notches	5
<i>G. tamakoshi</i> Budha & Backeljau, 2017	absent	comb-like with numerous notches	6
<i>G. mandalayensis</i> sp. nov.	absent	absent	This study
<b><i>Rishetia</i> Godwin-Austen, 1920</b>			
<i>R. garoense</i> (Godwin-Austen, 1920)	n/a	long cylindrical	3
<i>R. hastula</i> (Benson, 1860)	short knob-like	short tubular	6
<i>R. kathmandica</i> Budha & Backeljau, 2017	short cylindrical	long cylindrical	6
<i>R. longispira</i> (Godwin-Austen, 1920)	n/a	long cylindrical	3, 6
<i>R. mastersi</i> (Godwin-Austen, 1920)	short cylindrical	long cylindrical	6
<i>R. nagarjunensis</i> Budha & Naggs, 2017	short knob	short knob	6
<i>R. rishikeshi</i> Budha & Naggs, 2017	short cylindrical	long cylindrical	6
<i>R. subulata</i> Budha & Naggs, 2017	short cylindrical	long cylindrical	6
<i>R. tribhuvana</i> Budha, 2017	short cylindrical	long cylindrical	6
<i>R. tenuispira</i> (Benson, 1936)	short cylindrical	long cylindrical	4, 5

*Glessula* is widely distributed throughout the Indian subcontinent, and nearly 100 species have been described (Godwin-Austen 1920; Schileyko 1999; Ramakrishna et al. 2010; MolluscaBase 2023). Among the sporadic species recorded in Southeast Asia, Myanmar has 12 species (Gude 1914; Godwin-Austen 1920), while two species, *G. latestriata* von Möllendorff, 1899 and *G. paviei* Morlet 1893, have been recorded from Vietnam, Laos, and Thailand, and another two species, *G. sumatrana* (von Martens, 1864) and *G. wallacei* (Pfeiffer, 1856) from Indonesia (Tenison-Woods 1888; van Benthem Jutting 1952; Solem 1966; Schileyko 2011; Inkhavilay et al. 2019).

In this study, only five species including a new species, *G. blanfordiana* Nevill, 1877, *G. feddeni* Godwin-Austen, 1920, *G. gemma* (Reeve, 1850), *G. latestriata*, and *G. mandalayensis* sp. nov. have been collected and re-described.

### 8 *Glessula blanfordiana* Nevill, 1877

Fig. 5A–D, Table 1

*Glessula blanfordiana* Nevill, 1877: 26. Type locality: Pensee, Yunnan [Pensee, Kahkyen Hills, Yunnan Province, China]. Beddome 1906: 171. Pilsbry 1909: 98, pl. 13, fig. 11. Gude 1914: 437, 438. Godwin-Austen 1920: 56, pl. 164, fig. 21. *Stenogyra (Glessula) blanfordiana*—Nevill 1881: 138, pl. 5, fig. 12.

**Type specimens. Syntypes** NHMUK 1988150 (2 shells; Fig. 5A, B) ex. Godwin-Austen collection from Pensee, Yunnan, W. China.

**Other material.** Aik Kham Cave, Taunggyi Township, Taunggyi District, Shan State, Myanmar (20°49'07.0"N, 97°13'42.0"E): CUMZ 13067 (3 shells; Fig. 5C, D).

**Description.** Shell ovately conical, solid, brownish colour, and with 6–6½ whorls. Apex rounded; protoconch ~ 2 whorls with weak radial striations; subsequent whorls with equally and widely spaced ribs, more raised, and coarser near suture of last whorl. Spire regularly attenuated, slightly turreted; whorls convex; suture deep; last whorl largest. Aperture obliquely and narrowly ovate; peristome white and thickened; columella short, concave, slightly twisted and abruptly truncated. Umbilicus closed.

**Distribution.** This species was originally described from Yunnan, China, and subsequently reported from Bhamo, Kachin State, Myanmar (Nevill 1881). In this study, new material has been collected from Shan State, which borders Yunnan to the east and Bhamo to the north.

**Remarks.** When nominating this species, Nevill (1877: 26) compared it with *G. peguensis* (Blanford, 1865) from Bago and stated that *G. blanfordiana* has slender (less convex) whorls, wavy or curving, and stronger radial ribs. Nevertheless, the type specimens of these two species, *G. blanfordiana* (height 8–9 mm, Fig. 5A, B) and *G. peguensis* (height ~ 7 mm, Fig. 10B, C), exhibit considerable similarity in terms of shell shape and shell size. Therefore, more specimens from the Bago Region are necessary to ascertain whether the difference in shell sculptures represents variations or are taxonomically informative characters separating these two species.

*Glessula blanfordiana* examined herein are noticeably distinct from their congeners recorded in Shan State and Mandalay Region, namely *G. feddeni*

Godwin-Austen, 1920 and *G. latestriata* von Möllendorff, 1899. Specifically, *G. blanfordiana* has a shorter and bluntly attenuated spire, more bulging whorls, deeper suture, and stronger and densely radial ribbed sculptures. In contrast, the other species exhibit higher and pointed spires, more flattened whorls, and smooth to radial grooves sculptures (see Fig. 2D).

*Glessula blanfordiana* was described based on two specimens collected by J. Anderson during the Expedition to Yunnan and Upper Burma. The original description was very brief, without shell dimensions and illustrations. However, under the note of *G. ponsiensis* Godwin-Austen, 1920, states, 'The type from Pensee, Yunnan (Plate CLXIV. fig. 20, apex) [= *G. blanfordiana*], has been sent me from Calcutta by the Director of the Zoological Survey of India, Dr. N. Annandale...' (Godwin-Austen 1920: 56). The NHMUK type collections contain a single lot with an original label stating 'Type' from 'Dr. J. Anderson', and collection locality from 'Pensee'. These two specimens match well with the species description; both are figured herein for the first time. It is possible that N. Annandale may have gifted this type specimen lot to Godwin-Austen for comparison with other species, as was the case with other specimens from the Zoological Survey of India.

### 9 *Glessula crassilabris* (Benson, 1836)

Fig. 5E, F

*Achatina crassilabris* Benson, 1836: 353. Type locality: N.E. Frontier of Bengal.  
 Pfeiffer 1848: 261. Reeve 1850: *Achatina*, pl. 21, fig. 81. Pfeiffer 1853: 493.  
 Pfeiffer 1860: 313, pl. 25, figs 12, 13. Benson 1860: 464. Blanford 1865: 95.  
*Oleacina (Electra) crassilabris*—Adams and Adams 1855: 105.  
*Achatina (Electra) crassilabris*—Pfeiffer 1856: 168. Hanley and Theobald 1870:  
 17, pl. 36, fig. 1.  
*Cionella (Glessula) crassilabris*—von Martens 1860: 254.  
*Stenogyra (Glessula) crassilabris*—Nevill 1878: 170.  
*Glessula crassilabris*—Godwin-Austen 1876: 315. Pfeiffer and Clessin 1881:  
 330. Beddome 1906: 169. Pilsbry 1909: 96, pl. 10, figs 14, 16. Gude 1914:  
 426. Godwin-Austen 1920: 39, 40, pl. 160, figs 14, 17–20, pl. 164, figs 16, 17.  
 Ramakrishna et al. 2010: 157. Preece et al. 2022: 122, fig. 51e.

**Type specimen. Syntype** UMZC I.102430 (1 shell; Fig. 5E after Preece et al. 2022: fig. 51e) from Bengal.

**Other material.** NHMUK (1 shell; Fig. 5F) from Cherrapoonji (M10604/2). NHMUK 20200349 (2 shells) from N.E. Bengal. NHMUK 20200350 (2 shells) M.10695/2 from Khasi.

**Diagnosis.** Shell oblong ovate and regularly attenuated; spire high conical; apex rounded; subsequent whorls with weak and irregularly spaced radial ridges, slightly stronger near suture, and fine radial grooves present. Suture impressed and whorls slightly convex. Aperture ovate; columella strong, concave, and truncated.

**Distribution.** This species is known from India and Rakhine State, Myanmar (Godwin-Austen 1920).

**Remarks.** The taxonomic history and clarification status of the type specimens have recently been published in Preece et al. (2022). Blanford (1865: 95) reported a small variety of *Glessula crassilabris* based on specimens from 'Arakan' [Rakhine State] and 'Shan hills near Ava' [Shan Hills near Innwa], and also suggested that the latter specimen lot were possibly distinct from *G. crassilabris* s. s., although they were closely related. The following year, Godwin-Austen (1920) introduced the Shan hills specimens as *G. feddeni* Godwin-Austen, 1920, but the small variety from 'Arakan' has never been made available.

### 10 *Glessula feddeni* Godwin-Austen, 1920

Fig. 6, Table 1

*Glessula feddeni* Godwin-Austen, 1920: 58, pl. 162, fig. 15. Type locality: Shan Hills [Shan State, Myanmar].

*Glessula feddeni* var.—Godwin-Austen 1920: 58, pl. 163, fig. 14.

**Type specimens. Syntypes** NHMUK 1906.2.2.261 [re-registered in error as 1985142] (5 shells; Fig. 6A, B) ex. Blanford ex. Godwin-Austen collection from the Shan Hills, E. of Ava.

**Other material.** NHMUK 1906.3.3.41 (1 shell; Fig. 6C) ex. Blanford ex. Fedden collection labelled as 'var.' from the Shan Hills. Limestone hills (Apache Cement Factory), Pyinyaung Village, Meiktila District, Mandalay Region, Myanmar (20°49'39.1"N, 96°23'35.1"E): CUMZ 13068 (8 shells; Fig. 6D–F).

**Description.** Shell globosely conical, solid, glossy, brownish colour, and with 6½–7 whorls. Apex slightly pointed; protoconch ~ 2 whorls with smooth surface; subsequent whorls with strong, oblique, and equally spaced radial grooves that more distinct near suture. Spire attenuated, turreted and pointed; whorls flatly convex; suture deep; last two whorls almost equal and largest. Aperture narrowly and obliquely ovate; parietal wall straight; peristome white and thickened; columella short, concave, slightly twisted, and truncated. Umbilicus closed.

**Distribution.** This species was originally described from 'Shan Hills', but the precise type locality was not provided. In this study, *G. feddeni* was located in the Mandalay Region adjacent to the Shan Hills.

**Remarks.** Godwin-Austen (1920) clearly states the catalogue number 'Type No. 261.06.2.2' (NHMUK 1906.2.2.261) from the Shan Hills. The original description included an illustration and two sets of measurements. The Godwin-Austen's type lot in the NHMUK consists of five shells with an original handwritten label stating 'Type'. The two specimens with the largest shell height closest to the given measurements are here illustrated together with another shell from the syntype lot.

Godwin-Austen (1920: 58, pl. 163, fig. 14) also recognised another variety as 'var.' and illustrated specimen lot 'No. 41.06.3.3' (NHMUK 1906.3.3.41). Nevertheless, this entity has never been properly made available. The specimen lot NHMUK 1906.3.3.41 (Fig. 6C) labelled as 'var.' is excluded from the type series of this nominal species (ICZN 1999: Art. 72.4.1). However, this specimen has a more slender shell than the type series (Fig. 6A, B) and is possibly more closely related to *G. latestriata*, which is also distributed in the Shan State.

### 11 *Glessula gemma* (Reeve, 1850)

Fig. 7A–D, Table 1

*Achatina gemma* Reeve, 1850: pl. 22, fig. 123 (Benson MSS). Type locality: Barrackpore, Bengal. Pfeiffer 1853: 496. Benson 1860: 464.

*Oleacina (Electra) gemma*—Adams and Adams 1855: 105.

*Achatina (Electra) gemma*—Pfeiffer 1856: 168. Hanley and Theobald 1870: 17, pl. 36, fig. 7.

*Cionella (Glessula) gemma*—von Martens 1860: 254.

*Stenogyra (Glessula) gemma*—Nevill 1878: 170.

*Glessula gemma*—Pfeiffer and Clessin 1881: 331. Beddome 1906: 169. Pilsbry 1909: 97, pl. 13, figs 1, 3. Gude 1914: 428. Godwin-Austen 1920: 22. Raheem et al. 2014: 124, figs 78b, 79c, d. Preece et al. 2022: 235, 236.

**Type specimen.** The type specimen could not be located in the UMZC nor in the NHMUK collections (Raheem et al. 2014; Preece et al. 2022).

**Other material.** NHMUK 1946.10.16.45–53 (9 shells; Fig. 7A, B) ex. Benson collection from Bengal. Montawa Cave, Taunggyi City, Taunggyi District, Shan State, Myanmar. (20°45'15.9"N, 97°1'3.4"E): CUMZ 13069 (5 shells; Fig. 7C, D).

**Description.** Shell elongate ovate, solid, purple-black or yellowish colour, and with 6½–7 whorls. Apex blunt; protoconch ~ 2 whorls with smooth surface; subsequent whorls with smooth and glossy to weakly radial ribs. Spire grows evenly, is high and obtuse; whorls flatly convex; suture wide and shallow; last two whorls almost equal and largest. Aperture obliquely ovate; peristome moderately thickened and white; columella short, concave, and truncated. Umbilicus closed.

**Distribution.** *Glessula gemma* was originally described from India and then its range extended to Bangladesh (Gude 1914; Godwin-Austen 1920; Raheem et al. 2014). In Myanmar, this species has been recorded from the Rakhine State and now in the Shan State.

**Remarks.** The historical specimens NHMUK 1946.10.16.45–53 ex. Benson collection from Bengal is illustrated herein as an example.

Our specimens match well with the authenticated Reeve specimen of this species, although it is far from the type locality and previously reported region in the Rakhine State, Myanmar. Among all the *Glessula* species from Myanmar, *G. gemma* is superficially similar to *G. latestriata* from Shan State. In comparison, *G. gemma* has smooth shell surfaces, less turreted spire, and a shallow suture, while *G. latestriata* possesses well-developed grooves, a turreted and more attenuated spire, and a deeper suture.

### 12 *Glessula inedita* Godwin-Austen, 1920

Fig. 7E, F

*Glessula ineditus* Godwin-Austen, 1920: 58, 59. Type locality: Shan Hills [Shan State, Myanmar].

**Type specimens. Syntypes** NHMUK 1906.5.5.88 [re-registered in error as 1985228] (3 shells; Fig. 7E, F) ex. W.T. Blanford collection from Shan Hills.

**Diagnosis.** Shell oblong turreted and regularly attenuated; spire high conical; apex rounded; subsequent whorls with very weak irregularly spaced radial ribs. Suture deep and whorls convex. Aperture widely ovate; columella concave and truncated.

**Distribution.** This species is known to occur only from the type locality.

**Remarks.** The original description did not include an illustration, and one set of shell measurements was given. Godwin-Austen (1920: 58) clearly states, 'Type No. 88.06.5.5' (NHMUK 1906.5.5.88) is from the Blanford collection. This syntype lot consists of three shells, and the original label states 'Type'. The largest specimen matches well with the given measurements, and another shell from the same syntype lot is illustrated herein for the first time. No new materials have been reported to date.

### 13 *Glessula latestriata* von Möllendorff, 1899

Fig. 8A–D, Table 1

*Glessula latestriata* von Möllendorff, 1899: 166. Type locality: Kalow, Shan State [Kalaw Township, Taunggyi District, Shan State, Myanmar]. Beddome 1906: 172. Pilsbry 1909: 100. Gude 1914: 443. Godwin-Austen 1920: 59. Solem 1966: 93, 94, pl. 2, fig. e. Zilch 1973: 110, pl. 5, fig. 26. Inkhavilay et al. 2019: 49, fig. 20c, d.

**Type specimens.** **Lectotype** SMF 145919/1 (Fig. 8A) designated in Zilch (1973: pl. 5, fig. 26) from Kalow, Shan States. **Paralectotypes** SMF 227513/2 (2 shells) and NHMUK 1926.2.3.19–20 (2 shells; Fig. 8B) from Kalow, Shan States.

**Other material.** Thale Cumon Temple, Kalaw City, Shan State, Myanmar (20°43'24.1"N, 96°35'38.9"E): CUMZ 13070 (3 shells; Fig. 8C). Ywangan Village, near Lin Way Monastery, Kalaw Township, Taunggyi District, Shan State, Myanmar (21°13'43.3"N, 96°33'19.2"E): CUMZ 13071 (1 shell; Fig. 8D).

**Description.** Shell ovate conic, solid, glossy, brownish colour, and with 6½–7 whorls. Apex blunt; protoconch ~ 2 whorls with smooth to shallow radial grooves; subsequent whorls with distinctly thickened and equally spaced grooves. Spire gradually attenuated, slightly turreted, and bluntly pointed; whorls convex; suture deep; last two whorls almost equal and largest. Aperture narrow and obliquely ovate; peristome white and thickened; columella short, concave, slightly twisted, and truncated. Umbilicus closed.

**Distribution.** This species is known from Shan State in Myanmar, northern Thailand, and Laos (Gude 1914; Solem 1966; Inkhavilay et al. 2019).

**Remarks.** von Möllendorff (1899) described this species based on specimens collected by B Strubell from Shan State, Myanmar. The original description did not include any illustrations, and only one set of measurements was given. The species description was not explicitly based on one specimen. Nevertheless, Zilch (1973: 110) later used the terms 'Holotypus SMF 145919' and 'Paratypen SMF 227513/2' for the von Möllendorff type series in the SMF collection. This is not a valid holotype designation, but we consider it an inadvertent lectotype designation (ICZN 1999: Arts. 73.1, 74.5, and Recommendation 73F). The other specimens labelled as 'Paratypen SMF 227513/2' and 'Co-types NHMUK 1926.2.3.19–20' are considered paralectotypes.

Although both *G. latestriata* and *G. feddeni* possess radial grooves striations, the former has a slender shell, gradually attenuated spire, and blunter apex; in contrast, *G. feddeni* has a larger and broader shell, turreted, intermediately attenuated spire and pointed apex with abruptly truncate columella.

**14 *Glessula mandalayensis* Man & Panha, sp. nov.**

<https://zoobank.org/01347449-D63E-469B-826C-BEE5D0D56B8E>

Figs 8E, F, 9, Tables 1, 3

**Type specimens.** *Holotype* CUMZ 13072 (height 14.6 mm, width 7.1 mm; Fig. 8E), *paratypes* CUMZ 13073 (29 shells; Fig. 8F), NHMUK 20230922 (2 shells) and SMF (2 shells).

**Type locality.** Limestone Hills (Apache Cement Factory), Pyinyaung Village, Meiktila District, Mandalay Region, Myanmar (20°49'39.1"N, 96°23'35.1"E).

**Etymology.** The specific name *mandalayensis* is the name of region, where the type specimens of this species were collected.

**Diagnosis.** Shell large, globosely ovate; spire broad, almost cylindrically attenuated, turreted; last whorl largest; shell surface glossy, smooth to shallow grooves; aperture ovately rounded and broad; peristome thin and white.

**Description.** Shell globosely ovate, solid, glossy, brown to bright ochraceous colour, and with 6–7 whorls. Apex blunt and large; protoconch ~ 2 whorls with smooth surface; subsequent whorls with smooth to fine radial striations or sometimes with shallow grooves. Spire broad, cylindrically attenuated, turreted; whorls flatly convex; suture deep; last whorl largest. Aperture ovately rounded and broad; peristome white and thin; columella short, concave, and truncated. Umbilicus closed.

**Genitalia** ( $n = 1$ ). Atrium very short. Penis evenly broad, stout, and thick muscularly. Epiphallus short, stout, slightly curved triangular shape, thick muscularly, smooth surface, and ~ 1/4 of penis length; flagellum absent. Penial retractor muscle long, thickened and attaches laterally at junction of penis and epiphallus. Vas deferens long distinct tube connected between tip of epiphallus and free oviduct and held in position to penis and vagina with weak connective tissue (Fig. 9A).

Vagina muscularly and nearly as long as penis. Gametolytic duct short and enlarged at base; gametolytic sac small and elliptical in shape. Free oviduct short and stout, and spermoviduct enlarged.

**Radula.** Each row contains ~ 70+ teeth with half-row formula: central-lateral-marginal teeth (1–(19–20)–(15–16+)). Central tooth relatively small, symmetrical tricuspid with pointed central cusp and very small lateral cusps. Lateral teeth bicuspid: endocone long, slender and with pointed tip; ectocone small, pointed tip and located at base of teeth. Latero-marginal teeth gradually reduced in size and asymmetrically bicuspid: endocone short with pointed tip; ectocone small triangle on shape and situated around middle of tooth height. Outermost teeth small and becoming tricuspid; mesocone short and blunt tip; endocone and ectocone very small and pointed tips (Fig. 9B).

**Distribution.** This new species is known only from the type locality.

**Differential diagnosis.** *Glessula mandalayensis* sp. nov. can be distinguished from other congeners in Myanmar by having the largest and most globosely ovate shell, obtuse and broad spire, wide aperture, and thinner peristome. In particular,

this new species differs from a sympatric species, *G. feddeni*, by having a larger size (height 13.3–16.0 mm; Table 1), more rounded shell, obtuse and broader turreted spire, wider aperture, thinner peristome, and smoother shell surface. In comparison, *G. feddeni* is smaller in size (height 7.8–10.8 mm; Table 1), with a more solid, slender conical shell, pointed and narrow spire, smaller aperture, thicker peristome, and with distinct grooves on the shell surface. *Glessula mandalayensis* sp. nov. can also be separated from *G. latestriata* by its broader shell, wider spire, and blunt apex with smoother shell surface. Whereas *G. latestriata* has a slenderer shell, narrowly attenuated spire and pointed apex with distinct radial grooves striations. This new species also differs from *G. gemma* by having a larger and globose ovate shell (height 13.3–16.0; Table 1), more turreted spire, convex whorls, vertically elongated aperture, and deeper suture. Whereas *G. gemma* possesses a smaller and slender shell (height 8.7–9.2; Table 1), spire high conical, flatly convex whorls, more rounded aperture, and shallow suture.

Additionally, this new species differs from *G. crassilabris* s. s. from India and its variety from Rakhine State, Myanmar, in having a smoother shell, obtuse, and lower and broader spire. While *G. crassilabris* s. s. (Fig. 5E, F) displays stronger striations and a narrowly pointed spire (Blanford 1865).

**Remark.** So far, the genitalia of eight *Glessula* species has. The terminal part of the male genitalia of *G. mandalayensis* sp. nov. (Fig. 9A) clearly differs from these eight species in having no flagellum, while a flagellum is well-developed with a comb-like shape in the others (Table 3). However, we examined only a single specimen; therefore, the extent of variability within the species remains unknown.

### 15 *Glessula orophila* (Reeve, 1849)

Fig. 10A

*Achatina orophila* Reeve, 1849a: *Achatina*, pl. 19, species 105. Type locality: Neilgherry Hills, India [Nilgiri Hills, India]; Colombo, Ceylon. Benson 1860: 465.

*Oleacina (Electra) orophila*—Adams and Adams 1855: 106.

*Cionella (Glessula) orophila*—von Martens 1860: 254.

*Stenogyra (Glessula) orophila*—Nevill 1881: 137, pl. 5, fig. 19.

*Glessula orophila*—Beddome 1906: 168. Pilsbry 1909: 79, pl. 10, fig. 10. Gude 1914: 423. Raheem et al. 2014: 128, fig. 83a.

**Type specimen.** The type specimen could not be located from the UMZC and the NHMUK collections (Raheem et al. 2014; Preece et al. 2022).

**Other material.** NHMUK (3 shells + 2 juveniles; Fig. 10A) ex. Blanford collection from Mahableshwar.

**Diagnosis.** Shell ovate conical and regularly attenuated; spire low conical; apex rounded; subsequent whorls with very weak irregularly spaced radial ridges, and fine radial grooves present. Suture shallow and whorls flattened. Last whorl large and ovate; aperture ovate; columella concave and truncated.

**Distribution.** This species was first introduced from India and later reported from Sri Lanka and Pegu [Bago Region, Myanmar] (Gude 1914: 423).

**Remarks.** *Glessula orophila* is listed in the doubtful name by Raheem et al. (2014). No Myanmar material has been examined, and only the literature record is known.

### 16 *Glessula peguensis* (Blanford, 1865)

Fig. 10B, C

*Achatina peguensis* Blanford, 1865: 78. Type locality: Irawady valley, Pegu [Ayeyarwady valley, Bago Region, Myanmar]. Pfeiffer 1868: 228. Hanley and Theobald 1874: 41, pl. 102, fig. 6.

*Glessula peguensis*—Theobald and Stoliczka 1872: 334. Pfeiffer and Clessin 1881: 331. Beddome 1906: 171. Pilsbry 1909: 99, pl. 13, fig. 12. Gude 1914: 438. Godwin-Austen 1920: 55, 56, pl. 162, figs 20, 21.

*Stenogyra* (*Glessula*) *peguensis*—Nevill 1878: 171.

**Type specimen.** The type specimen could not be located in the NHMUK collection.

**Other material.** NHMUK 1909.3.15.8 (3 shells; Fig. 10B, C) ex. Blanford collection from Thyet-Myo. NHMUK 20230920 ex. H.B.F. collection (4 shells) from Thyet Myo. SMF 235657/2 (2 shells) from Pegu.

**Diagnosis.** Shell oblong ovate and regularly attenuated; spire high conical; apex rounded and blunt; subsequent whorls with prominent regularly spaced radial ribs. Suture deeply impressed and whorls convex. Aperture widely ovate; columella strong, concave, and truncated.

**Distribution.** *Glessula peguensis* is distributed in Bago Region, Magway Region, and Rakhine State in Myanmar, as well as in Chittagong, Bangladesh (Gude 1914; Godwin-Austen 1920).

**Remarks.** The specimen from 'Thyet-myo', which is present in the Blanford collection with the original label as 'authentic', is illustrated herein as an example of this species. According to Godwin-Austen (1920: 56), this species is commonly found in Pegu.

The original type locality of this species was from a broad geographical range, 'Pegu', and the museum label stated Thyet-Myo as the collection locality. The reference to 'Pegu' and 'Thyet-Myo' on the original label needs to be clarified as to whether they refer to the same area or if there is a specific location in 'Thyet-Myo, Pegu'. The current administration area now includes 'Thyet-Myo' [Thayet Myo] as a district of the Magway Region located in central Myanmar, formerly part of 'Pegu, Lower Burma' during the British colonial period. Furthermore, Godwin-Austen (1920: 56) stated that additional specimens from 'Arakan near Tongoop' (No. 262.06.2.2) tended to have a larger shell size than the type specimen.

### 17 *Glessula perlevis* Godwin-Austen, 1920

Fig. 10D, E

*Glessula perlevis* Godwin-Austen, 1920: 59. Type locality: Shan Hills [Shan State, Myanmar].

**Type specimens. Syntypes** NHMUK 1906.5.5.89 [re-registered in error as 1986023] (5 shells; Fig. 10D, E) ex. Godwin-Austen collection from Shan Hills, Burma.

**Diagnosis.** Shell oblong ovate and regularly attenuated; spire high conical; apex rounded and blunt; subsequent whorls nearly smooth with very weak growth lines, and somewhat strong irregularly spaced ridges near suture.

Suture impressed and whorls slightly convex. Aperture ovate; columella concave and truncated.

**Distribution.** This species is known only from the type locality in the Shan Hills near Mandalay. The type locality noted by Godwin-Austen seems ambiguous because it was described as ‘probably comes from the Shan State near Mandalay collected by Fedden’ (Godwin-Austen, 1920: 59).

**Remarks.** Godwin-Austen (1920) described this species based on five specimens in the Blanford collection. The species description included one set of shell measurements, the number of specimens examined and the W.T. Blanford catalogue number ‘Type No. 89.06.5.5’ (NHMUK 1906.5.5.89), and there was no illustration of the shell. The NHMUK collections contain a type specimen lot consisting of five shells (2 matures + 3 immatures) with an original label stating ‘Types’. The mature shells correspond well with the shell dimensions given in the original description and are illustrated herein for the first time.

### 18 *Glessula ponsiensis* Godwin-Austen, 1920

Figs 10F, 11A, B

*Stenogyra* (*Glessula*) *pyramis* var. *major* Nevill, 1878: 169 [nomen nudum]. Type locality: Pensee.

*Glessula ponsiensis*—Godwin-Austen, 1920: 56, pl. 164, fig. 19. Type locality: Pensee, Yunnan.


**Type specimens. Syntypes** NHMUK 1989042 (1 shell; Fig. 10F) and NHMUK 1989043 (3 shells; Fig. 11A), both specimen lots collected by J. Anderson from Pensee.

**Other material.** NHMUK 1912.4.16.747 (2 shells; Fig. 11B) ex. Beddome collection from Pensee, Yunnan.

**Diagnosis.** Shell oblong turreted and regularly attenuated; spire high conical; apex rounded and blunt; subsequent whorls nearly smooth with fine growth lines, and strong, equally spaced radial ridges near suture. Suture impressed and whorls slightly convex. Aperture ovate; columella strong, concave, and truncated.

**Distribution.** *Glessula ponsiensis* was first described from Pensee, Yunnan, and later discovered in Bhamao, Kachin State, situated on the upper part of Ayeyarwady River.

**Remarks.** The original description included a set of shell measurements and an illustration of the apex. Godwin-Austen (1920: 56) clearly states that this species was described based on specimens from the Indian Museum collection (now Zoological Survey of India (ZSI), Kolkata) collected by J. Anderson from Pensee, which was misidentified as ‘*Stenogyra* (*Glessula*) *pyramis* Bs., var. *major* Nevill, 1878’. The name ‘var. *major*’ by Nevill (1878: 169) is unavailable (ICZN 1999: Art. 12). Under the note of the species, Godwin-Austen states, ‘I am fortunate in getting the type-shells for examination’. It seemed that he received on loan the type specimen (or specimens that were examined by Nevill (1878: 169)). The NHMUK type collections contain a specimen lot with two registration numbers, M10905/2 and 10906/2, which are ex-Indian Museum collection numbers where the original label states ‘*Stenogyra* (*Glessula*) *pyramis* Benson,



**Figure 5.** A–D *Glessula blanfordiana* A, B syntypes NHMUK 1988150 from Pensee, Yunnan, W. China and C, D specimen CUMZ 13067 from Taunggyi, Shan State, Myanmar E, F *Glessula crassilabris* E syntype UMZC I.102430 from Bengal (after Preece et al. 2022: fig. 51e) and F specimen NHMUK from Cherrapoonji (M10604/2).

var. *major*', collection of 'Dr. J. Anderson' and with collection locality from 'Ponsee'. Interestingly, a species-group name was struck through and rewritten with red ink by Godwin-Austen, handwritten as '*ponsiensis*, G-A.' and with 'Type and three other one takes for B.M.'. It is probable that these specimens were presented to Godwin-Austen for examination. The specimen NHMUK 1989042 (1 shell has blue wool inside the shell; Fig. 10F) has a subsequent label stating 'holotype', and NHMUK 1989043 (3 shells; Fig. 11A) originates from the same Nevill type lot and has a subsequent label stating 'paratypes'. This is not a valid holotype designation (ICZN 1999: Arts 73.1, 73.2 and Recommendation 73F) and these specimens are considered as the syntypes, illustrated herein for the first time.




Figure 6. A–F *Glessula feddeni* A, B syntypes NHMUK 1906.2.2.261 from Shan Hills, E. of Ava C specimen NHMUK 1906.3.3.41 from Shan Hills and D–F specimen CUMZ 13068 from Pyinyaung Village, Mandalay Region, Myanmar.


### 19 *Glessula woodthorpei* Godwin-Austen, 1920

Fig. 11C–E

*Glessula woodthorpei* Godwin-Austen, 1920: 58, pl. 162, fig. 19. Type locality: Shan States [Shan State, Myanmar].

**Type specimens.** *Syntypes* NHMUK 1903.7.1.1628 [re-registered in error as 1986051] (9 shells; Fig. 11C–E) collected by Woodthorpe ex. Godwin-Austen collection from Shan States.

**Other material.** NHMUK 1903.7.1.3655 (2 shells) collected by Woodthorpe ex. Godwin-Austen collection from Siam N.W. Boundary.




**Figure 7.** A–D *Glessula gemma* A, B specimen NHMUK 1946.10.16.45–53 from Bengal and C, D specimen CUMZ 13069 from Taunggyi, Shan State, Myanmar E, F *Glessula inedita*, syntypes NHMUK 1906.5.5.88 from Shan Hills.

**Diagnosis.** Shell oblong conical and regularly attenuated; spire high conical; embryonic whorls rounded; subsequent whorls nearly smooth and equally spaced radial ridges appeared near suture. Suture impressed and whorls slightly convex. Aperture ovate; columella strong, concave, and truncated.

**Distribution.** This species is known from Shan State, Myanmar with an additional record from Thailand.


**Remarks.** Godwin-Austen (1920) indicated that nine specimens with the catalogue number 'Type No. 1682 B.M.' were examined. The species description included one set of measurements and an illustration of one shell. The NHMUK type collections contain a lot of nine specimens from the



**Figure 8.** **A–D** *Glessula latestriata* **A** lectotype SMF 145919 from Kalow, Shan States **B** paralectotypes NHMUK 1926.2.3.19–20 from Kalow, Shan States **C** specimen CUMZ 13070 from Thale Cumon Temple, Kalow, Shan State, Myanmar and **D** specimen CUMZ 13071 from Ywangan Village, Kalow, Shan State, Myanmar **E, F** *Glessula mandalayensis* sp. nov. **E** holotype CUMZ 13072 from Pyinyaung Village, Mandalay Region, Myanmar and **F** paratypes CUMZ 13073 from the type locality. Photograph: Gojšina & Páll-Gergely (**A**).

Woodthorpe collection with a label in Godwin-Austen's handwriting stating 'Type'. The measurements of the specimen with red wool inside the shell are close to the measurements given in the original description. It likely corresponds to the illustration of the species provided in the original description and it herein figured (Fig. 11C) along with other shells (Fig. 11D, E) from the same syntype lot.

Godwin-Austen separately mentioned (1920: 58) another bleached specimen lot NHMUK 1903.7.1.3655 (2 shells) from Thailand. However, this specimen lot does not form part of the type series of this species.




**Figure 9.** Genitalia and radula of *Glessula mandalayensis* sp. nov., paratype CUMZ 13073 from Pyinyaung Village, Mandalay Region, Myanmar **A** reproductive anatomy and **B** radula morphology: yellow colour and 'C' indicates central teeth row and numbers indicate tooth order from lateral to marginal end. Abbreviations: **at**, atrium; **ep**, epiphallus; **fo**, free oviduct; **gd**, gametolytic duct; **gs**, gametolytic sac; **p**, penis; **pr**, penial retractor muscle; **so**, spermoviduct; **v**, vagina; **vd**, vas deferens.

### 20 *Glessula yuangensis* Godwin-Austen, 1920

Fig. 11F

*Glessula yuangensis* Godwin-Austen, 1920: 59, pl. 162, fig. 18. Type locality: Yuang Ha, Siam Boundary [Mong Yawng Township, Tachileik District, Shan State, Myanmar].

**Type specimen.** *Holotype* NHMUK 1903.7.1.1312 [re-registered in error as 1986052] (Fig. 11F) collected by Woodthorpe ex. Godwin-Austen collection from Yuang Ha, Shan States.




**Figure 10.** **A** *Glessula orophila*, specimen NHMUK ex. Blanford collection from Mahableshtar **B, C** *Glessula peguensis*, specimen NHMUK 1909.3.15.8 from Thyet-Myo **D, E** *Glessula perlevis*, syntypes NHMUK 1906.5.5.89 from Shan Hills, Burma **F** *Glessula ponsiensis*, syntypes NHMUK 1989042 from Pensee.

**Diagnosis.** Shell oblong turreted and regularly attenuated; spire high conical; apex rounded and blunt; subsequent whorls with prominent wide spaced radial ribs, and radial groves present. Suture impressed and whorls slightly convex. Aperture ovate; columella strong, concave, and truncated.

**Distribution.** The species is currently known only from its type locality.

**Remarks.** Godwin-Austen clearly stated that this species was described based on only one specimen collected by R. Woodthorpe. The description included a set of shell dimensions and a shell illustration. There is one Godwin-Austen specimen lot, NHMUK 1903.7.1.1312, that has an original label stating 'Type' and is from



**Figure 11.** **A, B** *Glessula ponsiensis* **A** syntypes NHMUK 1989043 from Pensee and **B** specimen NHMUK 1912.4.16.747 from Pensee, Yunnan **C–E** *Glessula woodthorpei*, syntypes NHMUK 1903.7.1.1628 from Shan States **F** *Glessula yuagensis*, holotype NHMUK 1903.7.1.1312 from Yuang Ha, Shan States.

the Woodthorpe collection. We recognise this specimen as the holotype fixed by monotypy. Additionally, Godwin-Austen seemed to consider another specimen lot, 'No. 1156 B.M.' (NHMUK 1903.7.1.1156) from Kentung State [Kengtung Township, Shan State], as a distinct entity and it is therefore excluded from the type series.

Originally, the type locality was said to be from 'Yuang Ha, Siam Boundary'; however, this town currently is found in the Shan State, Myanmar. The original type locality is now Mong Yawng Township, Shan State ( $\sim 21^{\circ}10'42.7''\text{N}$ ,  $100^{\circ}21'24.3''\text{E}$ ), Myanmar, with a further record from Kengtung, Shan State (Godwin-Austen 1920: 59).

### Genus *Opeas* Albers, 1850

*Bulimus* (*Opeas*) Albers, 1850: 175, 176.

*Stenogyra* (*Opeas*)—von Martens 1860: 265, 266.

*Opeas*—Fischer and Crosse 1877: 592. Pilsbry 1906: 122. Gude 1914: 354. Zilch 1959: 351, 352. van Benthem Jutting 1952: 378. Schileyko 1999: 492.

**Type species.** *Helix goodallii* Miller, 1822 [junior homonym of Férussac (1821)] accepted as *Opeas hannense* (Rang, 1831), subsequent designation by von Martens in Albers (1860: 265).

**Diagnosis.** Shell small, slender, and conical; spire high, turreted, cylindrically, and gradually attenuated; embryonic whorls smooth; subsequent whorls with fine radial striations or growth lines. Aperture vertical, narrow, oblong, columellar margin expanded, and columella straight or concave. Umbilicus narrowly opened or closed. Penis cylindrical tube with short epiphallus, and flagellum and penial sheath absent; vagina long ~ 1/2 of penis length.

**Remarks.** *Opeas* and *Tortaxis* Pilsbry, 1906 generally resemble one another in having slender conical or cylindrical shells, nearly smooth to fine striations, and flatly convex whorls (Table 2). However, *Opeas* has a smaller size, straight columella, and mostly oblong aperture, whereas *Tortaxis* displays mostly larger size, nearly straight or slightly concave columella with a spiral fold, and more or less obliqued aperture. Likewise, *Opeas* is distinctly differentiated from *Bacillum* in that it is smaller in size, has a straight columella, fine shell sculptures and embryonic whorls which are narrowly rounded. In contrast, *Bacillum* exhibits a larger shell size, truncated and concave columella, stronger shell sculptures, and the embryonic whorls cylindrically rounded.

This genus is distributed in tropical and subtropical regions in Europe, Asia, Africa, and North America, and comprises nearly 200 species (Schileyko 1999; MolluscaBase 2023). In Myanmar, two species are recorded (Gude 1914).

### 21 *Opeas filiforme* von Möllendorff, 1894

Fig. 12A–C, Table 1

*Opeas filiforme* von Möllendorff, 1894: 151, pl. 16, fig. 11. Type locality: Gulf von Siam: Samui-Inseln [Samui Island, Suratthani Province, Thailand]. Pilsbry 1906: 161, pl. 19, fig. 22. Zilch 1973: 120.

*Prosopeas filiforme*—Panha 1998: 32.

**Type specimens.** **Lectotype** SMF 227532 (Fig. 12A) von Möllendorff ex. C. Roebelen from Gulf von Siam: Samui-Inseln, designated in Zilch (1973: 120).

**Paralectotypes** SMF 145657/7 (7 shells), SMF 227533/7 (7 shells).

**Other material.** Phra (Buddha) Cave, Tanintharyi Region, Myanmar (11°14'01.5"N, 99°10'42.8"E): CUMZ 13074 (15 shells; Fig. 12B, C).

**Description.** Shell slender and conical, translucent, whitish colour, and with 7½–8 whorls. Apex rounded; protoconch ~ 2 whorls, rounded and smooth surface; subsequent whorls generally nearly smooth with fine growth lines. Spire gradually and cylindrically tapering and largely turreted; whorls flatly convex

and last whorl largest; suture deep and weakly crenulated. Aperture oblong; peristome thin; columella straight; columellar margin slightly expanded. Umbilicus narrowly opened to closed.

**Distribution.** This species occurs in peninsular Thailand and is newly recorded from the Tanintharyi Region, southern Myanmar.

**Remarks.** New discovered of this species were found in Tanintharyi, which borders the type locality of Thailand. The type specimen (Fig. 12A) possesses less turreted and more convex whorls than our specimens; however, this differentiation is herein considered an intraspecific variation. Compared to *O. innocens* Preston, 1910 from Mawlamyine, *O. filiforme* from Tanintharyi can generally be differentiated by its slender and more attenuated shell, slightly convex whorls, and turreted spire. However, a comprehensive examination of *O. innocens* specimens from the type locality is necessary to confirm this distinction.

## 22 *Opeas innocens* Preston, 1910

*Opeas innocens* Preston, 1910: 33, 34, fig. 2. Type locality: Khayon Cave, near Moulmein, Lower Burma [Khayon or Farm Cave, Mawlamyine Township, Mon State, Myanmar]. Gude 1914: 358.

**Distribution.** This species is known only from the type locality.

**Remarks.** No new specimens were found during our sampling conducted in Mawlamyine, Mon State. Furthermore, the authenticated and type specimens could not be located in the NHMUK collection, and currently, this species is known solely from the original description.

## Genus *Paropeas* Pilsbry, 1906

*Prosopeas* (*Paropeas*) Pilsbry, 1906: 14. Zilch 1959: 353.  
*Paropeas*—Naggs 1994: 175–191. Schileyko 1999: 508.

**Type species.** *Bulimus acutissimus* Mousson, 1857 by original designation.

**Diagnosis.** Shell slender and conical; spire high, turreted, and gradually attenuated; embryonic and subsequent whorls with irregularly dense, fine, or coarse radial striations. Aperture oblique, narrow or broadly ovate; columella concave; columellar margin expanded or not expanded. Penial simple, relatively long, slender, epiphallus short, flagellum absent, and penial sheath present; vagina ~ 1/2 of penis length and wider than penis.

**Remarks.** *Paropeas* can be distinguished from *Bacillum* and *Rishetia* by its irregularly coarse radial striations throughout the shell, less concave or straighter columella, and narrowly tapering and pointed embryonic whorls (Table 2). While *Bacillum* has stronger and more evenly spaced radial striations, deeper concave and truncate columella, and cylindrically tapering and obtuse embryonic whorls. Likewise, *Rishetia* has a glossy and smoother shell, broader whorls, more concave columella, and rounded and wider aperture (Pilsbry

1906; Gude 1914; Naggs 1994). In terms of genitalia, *Rishetia* is obviously distinct from *Paropeas* by having a tubular-shaped flagellum, while *Paropeas* has no flagellum (Naggs 1994; Schileyko 1999).

*Paropeas* and *Prosopeas* are very similar in terms of shell form and sculpture, but *Paropeas* possess stronger and more compact striations on embryonic whorls (Gude 1914; Naggs 1994). However, distinguishing between these two genera is still challenging, and precise identification requires further evidence, such as data on genitalia and molecular phylogeny.

This genus is mainly distributed in Southeast Asia. Seven recognised species are present, four are reported in Myanmar (Gude 1914; Naggs 1994; Schileyko 1999; MolluscaBase 2023).

### 23 *Paropeas swettenhami* (de Morgan, 1885)

Fig. 12D

*Stenogyra swettenhami* de Morgan, 1885: 389, pl. 6, fig. 6. Type locality: G.

Tchöra, près d'Ipoh (Kinta) [Gunung Cheroh, Ipoh, Perak State, Malaysia].

*Prosopeas swettenhami*—Pilsbry 1906: 32, pl. 4, figs 11, 12. Laidlaw 1933: 216.

*Paropeas swettenhami*—Naggs 1994: 188.

**Type specimen.** **Syntype** MNHN-IM-2000-38930 (1 shell; Fig. 12D) from Malacca, G. Tchéhèl.

**Diagnosis.** Shell slender, elongate turreted and rapidly attenuated; apex pointed; subsequent whorls coarse with dense and fine radial striae. Suture wide and shallow, and whorls flattened. Aperture elongate ovate; columella concave.

**Distribution.** This species has been reported from Malaysia and Shan State, Myanmar (Naggs 1994).

**Remarks.** de Morgan (1885) describes *P. swettenhami* and *P. tchehelense* consecutively in the same publication and from a very close geographical area within Perak State, Peninsular Malaysia. Then von Möllendorff (1891: 337) suggests '*Stenogyra swettenhami* de Morgan, 1885' as a slight variation of *P. tchehelense*; therefore, some subsequent authors recognise the former as a synonym of the latter (i.e., Maassen 2001; MolluscaBase 2023). In contrast, Pilsbry (1906: 32) argues that they differ in shell shape and structure. We have examined the syntypes of both species and agree with Pilsbry (1906) that *P. swettenhami* (Fig. 12D) differs from *P. tchehelense* by having a slimmer shell, narrower aperture, cylindrical and rounded embryonic whorls. *Prosopeas tchehelense* (Fig. 12E) possesses a large, broad shell, wide aperture, convex, and rounded embryonic whorls.

No new materials of this species were found in this survey, and the type specimen is illustrated herein. Naggs (1994) provisionally reported the occurrence of *P. swettenhami* from Shan State, Myanmar. Considering the vast geographical distance between Perak and Shan State, it may be that the specimens Naggs (1994) regarded as *P. swettenhami* are, in fact, *P. terebralis* or another species. However, collecting topotypical material from Shan State, Myanmar, is necessary to confirm the existence of this species in Myanmar and the systematic relationship with *P. tchehelense*.

## 24 *Paropeas terebralis* (Theobald, 1870)

Fig. 12F

*Stenogyra* (*Opeas*) *terebralis* Theobald, 1870: 401. Type locality: Shan States [Shan State, Myanmar]. Nevill 1878: 166. Pfeiffer and Clessin 1881: 321.

*Bulimus* (*Stenogyra*) *terebralis*—Pfeiffer 1876: 133.

*Prosopeas terebrale*—Pilsbry 1906: 31. Gude 1914: 363.

*Paropeas terebralis*—Naggs 1994: 188.

**Type specimen. Syntype** NHMUK 1888.12.4.1081 (1 shell; Fig. 12F) ex. Theobald collection from Shan States, Burma.

**Diagnosis.** Shell slender, elongate turreted and rapidly attenuated; apex obtuse; subsequent whorls coarse with dense growth lines throughout. Suture wide and shallow, and whorls slightly convex. Aperture elongate ovate and narrow; columella curved; peristome little acute.

**Distribution.** This species is only known from the type locality.

**Remarks.** No recent material of this species was collected in this survey, and the syntype is illustrated here for the first time. Gude (1914) recognised this species as belonging to the *Prosopeas*, but Naggs (1994) provisionally transferred this species to the *Paropeas* after comparing it with the type specimen.

*Paropeas terebralis* can be distinguished from *P. swettenhami* by its broader and more rounded embryonic whorls, deeper suture and slightly convex whorls, and coarse shell surface with dense growth lines. Whereas *P. swettenhami* possesses narrower and pointed embryonic whorls, shallower suture and flattened whorls, and shell surface with dense and fine radial striae.

## 25 *Paropeas turricula* (von Martens, 1860)

Fig. 12G, H, Table 1

*Stenogyra turricula* von Martens, 1860: 9. Type locality: Siam [Thailand]. von Martens 1867: 82, 83, pl. 22, fig. 7.

*Paropeas turricula*—Naggs 1994: 188. Maassen 2001: 81.

*Prosopeas turricula*—Pilsbry 1906: 30, 31, pl. 3, figs 95, 96. Inkhavilay et al. 2019: 52.

**Type specimens. Syntypes** NHMUK 1859.8.1.10 (3 shells; Fig. 12G) from Siam.

**Other material.** Phra (Buddha) Cave, Tanintharyi Region, Myanmar (11°14'01.5"N, 99°10'42.8"E): CUMZ 13075 (1 shell). The limestone karsts located close to Lampane Village, Tanintharyi Region, Myanmar (11°40'18.1"N, 99°13'30.1"E): CUMZ 13076 (2 shells; Fig. 12H).

**Description.** Shell slender, conical, translucent, whitish colour, and with 8–8½ whorls. Apex rounded; protoconch ~ 2 whorls, turreted and with nearly smooth to fine radial striations; subsequent whorls with dense and fine radial striations. Spire high, turreted, gradually, and cylindrically tapering; whorls flatly convex, last three whorls nearly equal; suture deep. Aperture narrowly ovate and elongate; peristome thin; columella truncated and concave; columellar margin slightly expanded. Umbilicus narrowly opened to closed.

**Distribution.** This species was first described in Thailand, then subsequently recorded in Laos and Malaysia (Naggs 1994; Maassen 2001; Inkhavilay et al. 2019), and is here reported from the Tanintharyi Region, Myanmar.

**Remarks.** *Paropeas turricula* can be differentiated from *P. tchehelense* by its slender shell, cylindrically turreted spire, and finer striations, while *P. tchehelense* has a broader and larger shell, a more rounded and broader spire, and stronger radial striations. This species can also be separated from *P. terebralis* from Shan State by having finer sculpture, flatly convex whorls, shallower suture, and embryonic whorls that are more turreted with nearly smooth to weaker striations, whereas *P. terebralis* displays a more elongated shell, thicker sculptures, more convex whorls, deeper suture, and embryonic whorls more rounded with stronger striations. *Paropeas swettenhami* has a larger shell, stronger striations, and wider and more deeply concave columella.

## 26 *Paropeas walkeri* (Benson, 1863)

Fig. 12I–L, Table 1

*Spiraxis walkeri* Benson, 1863: 90. Type locality: ad Portum Blair [Port Blair, Andaman Islands, India]. Pfeiffer 1868: 189. Hanley and Theobald 1873: 34, pl. 79, fig. 4.

*Opeas walkeri*—Theobald 1870: 395. Godwin-Austen 1895: 443. Blanford 1903: 280.

*Stenogyra* (*Opeas*) *walkeri*—Nevill 1878: 165.

*Stenogyra* [*Spiraxis* (*Euspiraxis*)] *walkeri*—Pfeiffer and Clessin 1881: 323.

*Prosopeas walkeri*—Pilsbry 1906: 29, 30, pl. 6, fig. 70. Gude 1914: 363, 364. Ramakrishna et al. 2010: 183. Preece et al. 2022: 130, 131, fig. 56e.


**Type specimens.** **Syntypes** UMZC I.103115.A (5 shells; Fig. 12I, after Preece et al. 2022: fig. 56e) ex. R. McAndrew collection from Andaman Island.

**Other material.** NHMUK 1885.2.18.13–18 ex E.S. Berkeley collection Viper Id. Andamans: (6 shells; Fig. 12J). Aik Kham Cave, Taunggyi Township, Taunggyi District, Shan State, Myanmar (20°49'07.0"N, 97°13'42.0"E): CUMZ 13077 (3 shells; Fig. 12K). Ywangan Village, near Lin Way Monastery, Kalaw Township, Taunggyi District, Shan State, Myanmar (21°13'43.3"N, 96°33'19.2"E): CUMZ 13078 (2 shells; Fig. 12L).

**Description.** Shell slender, conical, translucent, whitish to pale yellowish colour, and with 8–8½ whorls. Apex rounded; protoconch ~ 2 whorls, rounded and with fine radial striations on entire whorls. Spire high, gradually tapering, and turreted; whorls flatly convex and last three whorls nearly equal; suture narrow and deep. Aperture narrowly ovate and elongate; peristome thin; columella straight or slightly concave; columellar margin simple to slightly expanded. Umbilicus narrowly opened to closed.

**Distribution.** This species was originally described from the Andaman Islands, India, and was later recorded in Shan State, Myanmar and Thailand (Gude 1914; Panha 1998).

**Remarks.** In Myanmar, *Paropeas terebralis* and *P. walkeri* are both recorded from Shan State. However, *P. walkeri* possesses broader, fewer, and more convex whorls, a straight columella with a slightly reflected columellar margin,



**Figure 12.** **A–C** *Opeas filiforme* **A** lectotype SMF 227532 from Gulf von Siam: Samui-Inseln **B, C** specimen CUMZ 13074 from Tanintharyi Region, Myanmar **D** *Paropeas swettenhami*, syntype MNHN-IM-2000-38930 from Malacca **E** *Paropeas tchehelense*, syntype MNHN-IM-2000-38932 from Pérak, Mont Tchéhèl **F** *Paropeas terebralis*, syntype NHMUK 1888.12.4.1081 from Shan States, Burma **G, H** *Paropeas turricula* **G** syntypes NHMUK 1859.8.1.10 from Siam and **H** specimen CUMZ 13076 from Tanintharyi Region, Myanmar **I–L** *Paropeas walkeri* **I** syntypes UMZC I.103115.A from Andaman Island (after Preece et al. 2022: fig. 56e) **J** specimen NHMUK 1885.2.18.13–18 from Viper Id. Andamans **K** specimen CUMZ 13077 from Taunggyi Township, Shan State, Myanmar with embryonic whorls and aperture and **L** specimen CUMZ 13078 from Kalaw Township, Shan State, Myanmar with embryonic whorls and aperture. Photographs: V Gojšina & B Páll-Gergely (**A**), P Bourguignon (**D, E**).

weaker radial striations, shallow suture, and embryonic whorls rounded. By contrast, *P. terebralis* has slimmer, higher, and flatter whorls, more concave columella, strong radial striations, deeper suture, and more pointed embryonic whorls. Additionally, *P. walkeri* also differs from *P. turricula* by its more convex whorls, straight columella, slightly reflected columellar margin, finer

and crowded radial striations, deeper suture, and embryonic whorls rounded with stronger radial striations. Furthermore, *P. walkeri* can be differentiated from *P. swettenhami* by its slenderer shell, convex and closely grow whorls, deeper suture, more rounded protoconch whorls, distinctly turreted spire, and straight columella. *Prosopeas swettenhami* displays a broader shell, flatter whorls, shallow suture, convex protoconch whorls, less turreted spire, and more concave columella.

Among the specimens examined from Shan State, we observed shell variations, such as the columellar margin being either expanded or not expanded and the columella being straight or slightly concave. In addition, the specimen identified as *P. walkeri* from the Andaman Islands (Fig. 12J) exhibits a broader shell and a more expanded columellar margin compared to the syntype specimen (Fig. 12I).

### Genus *Rishetia* Godwin-Austen, 1920

*Glessula* (*Rishetia*) Godwin-Austen, 1920: 7.

*Rishetia*—Schileyko 1999: 532. Raheem et al. 2014: 138, 139. Budha et al. 2017: 137. Preece et al. 2022: 127.

**Type species.** *Achatina tenuispira* Benson, 1836 by original designation.

**Diagnosis.** Shell slender and conical; spire high, turreted, and regularly attenuated; embryonic whorls smooth or with striations and subsequent whorls have thick or fine and equally or irregularly spaced radial ribs. Aperture oblique, narrow to broad, and ovate shape; columella concave and truncated; columellar margin not expanded. Penis large, thick, and moderately long; epiphallus relatively long; flagellum present with tubular shape; epiphallic caecum present; vagina enlarged, short ~ 1/2 of penis length.

**Remarks.** Originally, Godwin-Austen (1920) proposed *Rishetia* as a subgenus of *Glessula*, which Schileyko (1999) raised to the generic level, followed by Raheem et al. (2014), Budha et al. (2017), and Preece et al. (2022). *Rishetia* can be distinguished from *Glessula* by having a slender elongate-conical shell with a tubular-shaped flagellum and epiphallic caecum present, while *Glessula* processes an ovate-conical shell with a comb-like flagellum and epiphallic caecum absent (Tables 2, 3).

*Rishetia* can generally be differentiated from *Bacillum* by having a narrowly attenuated shell, convex and smaller apex, straight columella, and weak striations; *Bacillum* has a cylindrical shell, large and rounded apex, more concave columella, and stronger radial striations (Table 2). More information on the genitalia as well as a molecular phylogeny based on multiple species will clarify the systematic relationship of these two genera.

This genus has been documented in India, Sri Lanka, Bangladesh, and Nepal, with its presence in Southeast Asia limited to Myanmar (Godwin-Austen 1920; Schileyko 1999; Raheem et al. 2014; Budha et al. 2015, 2017). Currently, 23 species are recognised; among these, eleven species and subspecies have been documented from Myanmar (Godwin-Austen 1920; MolluscaBase 2023).

**27 *Rishetia akouktoungensis* (Godwin-Austen, 1920), comb. nov.**

Fig. 13A–C

*Glessula akouktoungensis* Godwin-Austen, 1920: 55. Type locality: Akouktoung on Irawady, Pegu [Akauk Taung (hill), Pyay District, Bago Region, Myanmar].

**Type specimens. Syntypes** NHMUK 1906.1.1.2207 [re-registered in error as 1985138] (6 shells; Fig. 13A–C). ex. W.T. Blanford ex. Godwin-Austen collection from Akouktoung, Pegu.

**Diagnosis.** Shell elongate, turreted, and regularly attenuated; apex rounded; subsequent whorls nearly smooth and with coarse radial ridges near suture. Suture impressed and whorls flattened convex. Aperture broadly ovate; columella strong, concave, and truncated.

**Distribution.** The species is currently known from its type locality, in which the modern name is Akauk Taung [hill, ~ 18°30'33.7"N, 95°06'48.5"E], south of Htone Bo (or Tonbo) Village, Padaung Township, Pyay District, Bago Region].

**Remarks.** The original description did not include an illustration, and only one set of measurements was given. Godwin-Austen's description was based on one lot of specimens which clearly stated 'Type No. 2207.06.1.1 B.M.' (NHMUK 1906.1.1.2207). There are six specimens from the NHMUK type collection with the original label stating 'Type'. The specimen that has a similar shell measurements as given in the original description and with pink wool inside the shell is figured herein (Fig. 13A). Other shells from the same syntype lot are also illustrated (Fig. 13B, C).

**28 *Rishetia baculina* (Blanford, 1871), comb. nov.**

Fig. 13D

*Glessula baculina* Blanford, 1871: 43, 44, pl. 2, fig. 6. Type locality: Khersiong Himalayae Sikkimensis. Beddome 1906: 160. Gude 1914: 379, 380. Godwin-Austen 1920: 14–16, pl. 159, fig. 7. Ramakrishna et al. 2010: 153.

*Achatina (Electra) baculina*—Hanley and Theobald 1873: 33, pl. 78, fig. 6.

*Achatina baculina*—Pfeiffer 1876: 291.

*Stenogyra (Glessula) baculina*—Nevill 1878: 170.

*Stenogyra (Subulina) baculina*—Pfeiffer and Clessin 1881: 327.

*Glessula tenuispira* var. *baculina*—Pilsbry 1909: 88, 89, pl. 9, fig. 2.

**Type specimens. Syntypes** NHMUK 1909.3.15.9 [re-registered in error as 19850144] (3 shells; Fig. 13D) ex. Godwin-Austen ex. Blanford collection from Kursiong, Darjiling.

**Diagnosis.** Shell slender, elongate turreted and rapidly attenuated; apex rounded; subsequent whorls with fine and equally spaced radial ribs throughout. Suture impressed and whorls slightly convex. Aperture obliquely ovate; columella strong, concave, and truncated.

**Distribution.** This species was originally discovered in India and later reported from the Magway Region, Myanmar (Godwin-Austen 1920).

**Remarks.** Godwin-Austen (1920: 15) stated that specimens identified to this species in the Beddome collection were collected from 'Thyet Myo' [Thayet

District, Magway Region]. However, the Myanmar specimens show noticeable differences from the type specimens in having a more attenuated apex, and larger and broader aperture.

**29 *Rishetia basseinensis* (Godwin-Austen, 1920), comb. nov.**

Fig. 13E, F

*Achatina pertenuis* var. *major* Blanford, 1865: 79 [in part].

*Glessula basseinensis*—Godwin-Austen 1920: 54, pl. 161, fig. 3, pl. 164, fig. 12.

Type locality: Bassein, Pegu and Pyema Khyoung, Bassein.

**Type specimens. Syntypes** NHMUK 1909.3.15.19 [re-registered in error as 1985139] (3 shells; Fig. 13E, F) ex. Blanford ex. Godwin-Austen collection from Bassein, Pegu.

**Diagnosis.** Shell slender, elongate turreted and regularly attenuated; apex rounded and blunt; subsequent whorls with fine and equally spaced radial ribs, which more prominent near suture. Suture impressed and whorls flattened convex. Aperture ovate; columella strong, concave, and truncated.

**Distribution.** *Rishetia basseinensis* appears to be restricted within Myanmar and is known only from the type locality. Based on the current administrative boundary, 'Bassien (or Pathein), Pegus' is in the Pathein Township, Pathein District, Ayeyarwady Region, rather than Pegu [Bago Region].

**Remarks.** Godwin-Austen (1920: 53) stated that when examining the type series of '*Glessula pertenuis* var. *major*' ex. Blanford collection, he found it very different from the nominotypical taxon, and described this specimen lot under a different name, '*Glessula basseinensis*'. In the original description, Godwin-Austen (1920: 54) provided two collection localities, namely 'Bassein, Pegu' and 'Pyema Khyoung, Bassein' and comparing three and six specimens, respectively. The description clearly states a catalogue number 'Type No. 19.9.3.15 B.M.' and includes an illustration of specimens from 'Bassein, Pegu'. The NHMUK collection contains this specimen lot NHMUK 1909.3.15.19 consisting of three shells with an original label in Godwin-Austen's handwriting stating 'Type' from the H.F. Blanford collection. The specimen with a similar shell measurement given in the original description and with pink wool inside the shell is figured herein (Fig. 13E). Additionally, one further shell from the same syntype lot is also illustrated (Fig. 13F).

The specimen lot NHMUK 1909.1.1.2208 is excluded from the type series of this species (see also under the *Rishetia pertenuis major*).

**30 *Rishetia burrailensis maxwelli* (Godwin-Austen, 1920), comb. nov.**

Figs 13G–I, 14A–C

*Glessula* (*Rishetia*) *burrailensis*, var. *maxwelli* Godwin-Austen, 1920: 45, 46, pl. 160, figs 5, 6. Type locality: Naga Hills, east of Kohima and Somra, Khulen Post. West of Kyendwin or Chindwin River, Upper Burma.

*Glessula burrailensis*, var. *maxwelli*—Godwin-Austen 1920: 61. Type locality: Somra Tracts, Somra Khulen Post, Upper Burma.

**Type specimens.** *Holotype* NHMUK 1903.7.1.1717/1 (Fig. 13G) ex. Godwin-Austen collection from Naga Hills, East of Kohima. *Paratypes* NHMUK 1903.7.1.1717/2–4 (3 shells; Fig. 13I) ex. Godwin-Austen collection from the location same as the holotype.

**Other material.** NHMUK 1903.7.1.3742 (7 shells + 1 broken shell; Fig. 14A, B) collected by F. Ede ex. Godwin-Austen collection from Somra Tracts, Somra Khulen Post, Burma (labelled as 'var. *somraensis*'). NHMUK 1903.7.1.3744 (2 shells; Fig. 14C) collected by F. Ede ex. Godwin-Austen collection from Somra Tracts, Somra Khulen Post, Burma (labelled as 'var. *somraensis*'). NHMUK 1988147 (4 shells; Fig. 13H) collected by L.R. Mawson from Somra Tracts, Somra Khulen Post, west of Kyengdwen River, Burma.

**Diagnosis.** Shell elongate, turreted, and regularly attenuated; apex rounded and blunt; subsequent whorls nearly smooth with fine irregularly spaced radial ridges throughout, and coarser ridges appearing near suture. Suture impressed and whorls flattened. Aperture ovate; columella concave and truncated.

**Distribution.** This subspecies was originally described from the Naga Hills. The locality reported in Myanmar is Somra Town (~25°21'43.6"N, 94°41'14.4"E), a mountainous area in Leshi or Layshi Township, Sagaing Region in northwestern Myanmar. It is neighbouring Nagaland to the west and Kachin State to the east and along the Chindwin River.

**Remarks.** The name *maxwelli* was stated twice in the same publication, on pages 45 and 61. The first mention is '*Glessula (Rishetia) burrailensis*, var. *maxwelli*'; when proposing this name, Godwin-Austen (1920: 45) recognised two specimen lots from Naga Hills... (collected by Col. H. St. P. Maxwell) and from Somra... Upper Burma (collected by F. Ede). Although the species description is brief, Godwin-Austen provides the catalogue number 'No. 1717 B.M.', two sets of shell dimensions, and illustrations of two shells from Naga Hill. We have examined this specimen lot, NHMUK 1903.7.1.1717, from Naga Hills, which consists of four shells. In the original description, Godwin-Austen (1920: 46) further states, 'Size: Fig. 5. Type' and in the figure caption as 'Fig. 5. (1<sup>st</sup> Type) from Naga Hills' and 'Fig. 6. (2<sup>nd</sup> Type) from Naga Hills'. It seemed that Godwin-Austen intended to designate the specimen figured in plate 160, fig. 5, as the unique name-bearing type, which we consider as the holotype. The other three specimens from the same lot are therefore paratypes.

The second mention is '*Glessula burrailensis*, var. *maxwelli*', for which Godwin-Austen (1920: 61) provided a very detailed description together with specimen lot number 'No. 3742 B.M.', two sets of shell measurements, from 'Somra Tracts, Somra Khulen Post, Upper Burma, but without illustration. The NHMUK collections contain specimen lot NHMUK 1903.7.1.3742, consisting of eight specimens with the original label stating 'var. *somraensis*, GA.'. This manuscript name has never properly been published and made available (ICZN 1999: Art. 10), the specimen lot NHMUK 1903.7.1.3742, with its original label stating 'var. *somraensis*' is considered as the syntype. Therefore, this second combination becomes an available name (ICZN 1999: Arts 11, 12), and it is considered a junior (primary) homonym (ICZN 1999: Arts 24.2.1, 24.2.2, 53.3, 57.2). Additionally, syntypes also include, NHMUK 1988147 (4 shells), from Somra, Khulen Post, West of Kyendwin River, Upper Burmah, as Godwin-Austen (1920: 61) recognised in the original description.

After examining the type specimens carefully, we found no differences in the diagnostic characters between the senior homonym (NHMUK 1903.7.1.1717/1 and NHMUK 1903.7.1.1717/2–4) and junior homonym (NHMUK 1903.7.1.3742, NHMUK 1903.7.1.3744 and NHMUK 1988147) specimens, especially shell shape, shell sculpture, and protoconch sculpture. The only detected difference is the brownish or yellowish brown periostracum, which we consider as intra-specific variation. Therefore, it seems unnecessary to propose a new replacement name for this junior homonym.

### 31 *Rishetia hastula* (Benson, 1860)

Fig. 14D

*Achatina hastula* Benson, 1860: 461. Type locality: ad Pankabari, prope Darjiling [Pankhabari, near Darjiling, West of Bengal, India]. Pfeiffer 1868: 235.

*Achatina (Electra) hastula*—Hanley and Theobald 1870: 9, pl. 18, fig. 4.

*Stenogyra (Glessula) hastula*—Nevill 1878: 169.

*Stenogyra (Subulina) hastula*—Pfeiffer and Clessin 1881: 327.

*Glessula hastula*—Theobald and Stoliczka 1872: 334. Beddome 1906: 167. Pilsbry 1909: 93, pl. 12, fig. 12. Gude 1914: 414. Ramakrishna et al. 2010: 161

*Glessula (Rishetia) hastula*—Godwin-Austen 1920: 16, 17, pl. 161. figs 16, 17, pl. 163. figs 9, 9a, 10.

*Rishetia hastula*—Budha et al. 2017: 139, figs 2d, 6. Preece et al. 2022: 127, 128, fig. 55a.

**Type specimen. Neotype** NHMUK 1906.1.1.880 (designated in Preece et al. 2022: 127, fig. 55a) (Fig. 14D) ex. Blanford collection from Darjiling.

**Diagnosis.** Shell elongate, turreted, and regularly attenuated; apex rounded and blunt; subsequent whorls coarse with fine equally spaced radial ridges throughout. Suture somewhat impressed and whorls convex. Aperture broadly ovate; columella strong, concave, and truncated.

**Distribution.** The species was originally described from India and later recorded from Nepal (Ramakrishna et al. 2010; Budha et al. 2015; Kalita 2022). In Myanmar, it was reported from Kumah Hill and Maii [Ma-ei Town ~ 19°20'36.5"N, 94°08'21.9"E], which are in the Thandwe District, Rakhine State in the westernmost part of Myanmar (Theobald and Stoliczka 1872; Gude 1914; Godwin-Austen 1920).

**Remarks.** To clarify the taxonomic status of this species, Preece et al. (2022) have designated a neotype based on the specimen figured in Hanley and Theobald (1870, pl. 18, fig. 4). However, the image used to represent the neotype in Preece et al. (2022: fig. 55a) is an erroneous repetition of '*Achatina leptospira* Benson, 1865' (see Preece et al. 2022: fig. 51f). Therefore, the correct image of the neotype NHMUK 1906.1.1.880 is illustrated herein.

Theobald and Stoliczka (1872: 334) noted that the specimens from Arakan tended to differ from the type specimen in having a larger shell size. Nevertheless, Arakan and Sikkim are non-adjacent regions and far apart, which raises doubts about whether they are the same or distinct species.

**32 *Rishetia kentungensis* (Godwin-Austen, 1920), comb. nov.**

Fig. 14E, F

*Glessula kentungensis* Godwin-Austen, 1920: 57, 58. Type locality: Mong Sing, Siam Boundary [Muang Sing District, Luang Namtha Province, Laos]. Inkhavilay et al. 2019: 49, fig. 20b.

**Type specimens. Syntypes** NHMUK 1903.7.1.3650 [re-registered in error as 1986002] (3 shells + 1 juvenile; Fig. 14E, F) collected by Woodthorpe ex. Godwin-Austen collection from Mong Sing, Siam Boundary.

**Other material.** NHMUK 1903.7.1.3748 (4 shells) collected by Woodthorpe ex. Godwin-Austen collection from the Mekong River.

**Diagnosis.** Shell elongate, turreted, and regularly attenuated; apex rounded, blunt and with very large embryonic shell; subsequent whorls nearly smooth throughout, and prominent irregularly spaced radial ridges near suture. Suture impressed and whorls slightly flattened. Aperture broadly ovate; columella strong, concave, and truncated.

**Distribution.** This species was initially found in Laos; however, as noted by Godwin-Austen (1920), its presence in the Mekong River region is assumed to likely extend to the East Shan State.

**Remarks.** The original description gave the measurements for only one shell and did not include an illustration. Godwin-Austen (1920) clearly states the collection locality and the catalogue number 'Type No. 3650 B.M.'. The NHMUK collections contain a lot of four specimens from the Godwin-Austen ex. Woodthorpe collection with an original label stating 'Type' and giving the collection locality as 'Mong Sing, Siam Boundary'. The two specimens match well with the original description, and shell dimensions are illustrated herein. Another specimen, lot 'No. 3748 B.M.' from the 'Mekong River', is excluded from the type series of this nominal species.

Originally, the type locality was said to be from 'Mong Sing, Siam Boundary'; currently, this locality refers to Muang Sing District, Luang Namtha Province, Laos. The specimen NHMUK 1903.7.1.3748, collected from the 'Mekong River' by Colonel R. G. Woodthorpe, has a vague locality that could possibly encompass the East Shan State (formerly Keng Tung State) and the Bokeo and Luang Namtha provinces in Laos.

**33 *Rishetia limborgi* (Godwin-Austen, 1920), comb. nov.**

Fig. 14G

*Glessula limborgi* Godwin-Austen, 1920: 56, 57. Type locality: Tenasserim [Tanintharyi Region, Myanmar].

**Type specimen. Holotype** NHMUK 1903.7.1.3200 [re-registered in error as 1985219] (Fig. 14G) collected by Limborg ex. Godwin-Austen collection from Tenasserim.

**Diagnosis.** Shell elongate, turreted and regularly attenuated; apex rounded; subsequent whorls with fine and equally spaced radial ridges throughout. Suture impressed and whorls flattened. Aperture broadly ovate; columella strong, concave, and truncated.

**Distribution.** This species is known only from the type locality and is likely endemic to that region.

**Remarks.** Godwin-Austen (1920) clearly states that this taxon was described based on only a single specimen collected by O. Limborg. The original description included only a single set of shell measurements but was without illustration. The NHMUK collections contain the type specimen lot, NHMUK 1903.7.1.3200 ex. Godwin-Austen ex. O. Limborg consisting of a single shell with an original label stating 'Type'. Therefore, we considered this shell as the holotype fixed by monotypy.

### **34 *Rishetia nathiana* (Godwin-Austen, 1920), comb. nov.**

Fig. 14H, I

*Glessula* (*Rishetia*) *nathiana* Godwin-Austen, 1920: 54, 55. Type locality: Bassein District [Patheingyi District, Ayeyarwady Region, Myanmar].

**Type specimens. Syntypes** NHMUK 1906.1.1.2206 [re-registered in error as 1986018] (5 shells; Fig. 14H, I) ex. Blanford collection from Bassein District.

**Diagnosis.** Shell elongate turreted and regularly attenuated; apex rounded and blunt; subsequent whorls with very fine equally spaced radial ridges, which coarser near suture. Suture impressed and whorls flattened convex. Aperture broadly ovate; columella concave and truncated.

**Distribution.** *Rishetia nathiana* is known only from the type locality and is likely endemic to that region.

**Remarks.** Godwin-Austen clearly stated that the original description was based on a specimen lot, ex. W.T. Blanford collection with the catalogue number 'Type No. 2206.06.1.1 B.M.'. The NHMUK collections contain a lot of five specimens from the Godwin-Austen ex. Blanford collection with original label 'Type' and 'ex. duplicate collection'. The specimen matches well with the given shell measurements, and the original description is illustrated herein for the first time.

### **35 *Rishetia pertenuis pertenuis* (Blanford, 1865), comb. nov.**

Fig. 15A–E

*Achatina pertenuis* Blanford, 1865: 79. Type locality: Tongoop, Arakan [Taungup Township, Rakhine State, Myanmar]. Pfeiffer 1868: 237.

*Achatina* (*Electra*) *pertenuis*—Hanley and Theobald 1870: 9, pl. 18, fig. 5.

*Stenogyra* (*Glessula*) *pertenuis*—Nevill 1878: 169.

*Stenogyra* (*Subulina*) *pertenuis*—Pfeiffer and Clessin 1881: 327.

*Glessula pertenuis*—Beddome 1906: 160. Gude 1914: 380. Godwin-Austen 1920: 52–54, pls 164, fig. 11, pl. 159, figs 1, 2. Ramakrishna et al. 2010: 170.

*Glessula tenuispira* var. *pertenuis*—Pilsbry 1909: 89, 90, pl. 9, fig. 3.

**Type specimen. Possible syntypes** NHMUK 1906.2.2.239 (7 shells; Fig. 15A, B) ex. Blanford collection from Henzada, Pegu and Tongoop, Arakan.

**Other material.** NHMUK 1888.12.4.1229–31 (3 shells; Fig. 15D, E) from Pegu. NHMUK 20230921 (5 shells) from Tongoop, Arakan (Fig. 15C).

**Diagnosis.** Shell slender, elongate turreted, and rapidly attenuated near apical whorls; apex rounded; subsequent whorls nearly smooth, fine growth lines, and with equally spaced ridges near suture. Suture impressed and whorls slightly convex. Aperture ovate; columella concave and truncated.

**Distribution.** This species was reported from several places in Myanmar: Henzada [Hinthada City, Ayeyarwady Region], Thayet Myo [Thayet City, Thayet District, Magway Region], Pegu [Bago Region], and Tongoop, Arakan [Taungup Township, Thandwe District, Rakhine State] (Nevill 1878; Gude 1914). It was also reported from Meghalaya State, India (Ramakrishna et al. 2010).

**Remarks.** The original description does not clearly state how many specimens were available to the author, although only one set of measurements was given. The NHMUK collection contains a lot of seven specimens from the Blanford collection with this species name. However, this is a mixed specimen lot of the same species as it consists of two collection localities, 'Henzada, Pegu' and 'Tongoop, Arakan', where only the latter was mentioned in the original description as the type locality. Therefore, this specimen lot NHMUK 1906.2.2.239 is considered as possible syntypes, from which two specimens are illustrated herein.

### 36 *Rishetia pertenuis major* (Blanford, 1865), comb. nov.

Fig. 15F, G

*Achatina pertenuis* var. *major* Blanford, 1865: 79. Type locality: Pyema Khyoung, Bassein District, Pegu [PyayMa Khaung or Pyinma Khuang, Pathein Township, Pathein District, Ayeyarwady Region, Myanmar].

*Glessula pertenuis* var. *major*—Gude 1914: 381. Godwin-Austen 1920: 53.


**Type specimen. Possible syntypes** NHMUK 1906.1.1.2208 (6 shells; Fig. 15F, G) ex. Blanford collection from Pyema Khyoung, Bassein; label state '*Glessula basseinensis* G-A.'

**Diagnosis.** Shell elongate, turreted and regularly attenuated; apex rounded and blunt; subsequent whorls with fine equally spaced radial ridges throughout, which coarser near suture. Suture impressed and whorls slightly flattened. Aperture broadly ovate; columella strong concave and truncated.

**Distribution.** This subspecies was known solely from the type locality.


**Remarks.** When describing '*Achatina pertenuis*', Blanford (1865) recognised the larger shells as a distinct variety and gave them the name 'var. *major*'. This taxon name was made available, although without description, because the two sets of shell dimensions and materials from 'Pyema Khyoung, Bassein district, Pegu' confer the indication (ICZN 1999: Arts 11.9 and 12).

The type series of the taxa could not be traced. Only one specimen lot from the Godwin-Austen collection belongs to W.T. Blanford's original type series., i.e., NHMUK 1906.1.1.2208, with the original label in Godwin-Austen's handwriting marked 'ex. duplicate collection Blf., intermediate between *A. tenuispira* and *A. pertenuis*', with a species name '*Glessula basseinensis* G-A.', and collection



**Figure 13.** **A–C** *Rishetia akouktoungensis*, syntypes NHMUK 1906.1.1.2207 from Akouktoung, Pegu **D** *Rishetia baculina*, syntypes NHMUK 1909.3.15.9 from Kursiong, Darjiling **E, F** *Rishetia basseinensis*, syntypes NHMUK 1909.3.15.19 from Bassein **G–I** *Rishetia burrailensis maxwelli* **G** holotype NHMUK 1903.7.1.1717/1 from Naga Hills, East of Kohima **H** specimen NHMUK 1988147 from Somra Tracts, Somra Khulen Post, west of Kyengdwen River, Burma and **I** paratypes NHMUK 1903.7.1.1717/2–4 from Naga Hills, East of Kohima.

locality 'Pyema Khyoung, Bassein'. Therefore, this specimen lot is here considered as possible syntype material, and the two shells closest to the shell measurements given in the original description are illustrated herein.



**Figure 14.** **A–C** *Rishetia burrailensis maxwelli* **A, B** specimen NHMUK 1903.7.1.3742 from Somra Tracts, Somra Khulen Post, Burma and **C** specimen NHMUK 1903.7.1.3744 from Somra Tracts, Somra Khulen Post, Burma **D** *Rishetia hastula*, neotype NHMUK 1906.1.1.880 from Darjiling **E, F** *Rishetia kentungensis*, syntypes NHMUK 1903.7.1.3650 from Mong Sing, Siam Boundary **G** *Rishetia limborgi*, holotype NHMUK 1903.7.1.3200 from Tenasserim **H, I** *Rishetia nathiana*, syntypes NHMUK 1906.1.1.2206 from Bassein District.

The modern name of 'Pyema Khyoung, Bassein District, Pegu' is the PyayMa Khaung or Pyinma Khuang (~ 18°38'51.7"N, 95°45'46.9"E), Pathein Township, Pathein District, Ayeyarwady Region, Myanmar.




Figure 15. A–E *Rishetia pertenuis* A, B possible syntypes NHMUK 1906.2.2.239 from Henzada, Pegu and Tongoop, Arakan C specimen NHMUK from Tongoop, Arakan and D, E specimen NHMUK 1888.12.4.1229–31 from Pegu F, G *Rishetia pertenuis major*, possible syntypes NHMUK 1906.1.1.2208 from Pyema Khyoung, Bassein H *Rishetia tenuispira*, specimen NHMUK 1903.7.1.3332 from Garo Hills.

### 37 *Rishetia tenuispira* (Benson, 1836)

Fig. 15H

*Achatina tenuispira* Benson, 1836: 353. Type locality: N.E. Frontier of Bengal.  
 Pfeiffer 1848: 262. Reeve 1849a: *Achatina*, pl. 16. Benson 1860: 464. Pfeiffer  
 1860: 310, 311 pl. 25, figs 6, 7. Blanford 1865: 95.

*Subulina tenuispira*—Adams and Adams 1855: 110.  
*Achatina (Subulina) tenuispira*—Pfeiffer 1856: 169.  
*Achatina (Electra) tenuispira*—Hanley and Theobald 1870: 17, pl. 36, fig. 8.  
*Stenogyra (Glessula) tenuispira*—Nevill 1878: 169.  
*Stenogyra (Subulina) tenuispira*—Pfeiffer and Clessin 1881: 327.  
*Glessula tenuispira*—Beddome 1906: 160. Pilsbry 1909: 88, pl. 9, figs 1, 4. Gude 1914: 378. Godwin-Austen 1920: 31, 32.  
*Rishetia tenuispira*—Raheem et al. 2014: 138, fig. 90c. Budha et al. 2017: fig. 14c. Preece et al. 2022: 128, fig. 55b.

**Type specimen.** The type specimen could not be located in the UMZC collection (Raheem et al. 2014; Preece et al. 2022). However, Budha et al. (2017: fig. 14a) figured a possible syntype of this species based on specimens ex. Benson collection from ‘Teria Ghat’, this specimen lot was not considered to form part of the type series by Preece et al. (2022).

**Other material.** NHMUK 1946.10.16.1–7 (7 shells) ex. R. McAndrew collection from Pegu. NHMUK 1903.7.1.3332 (1 shell; Fig. 15H) labelled as ‘var.’ ex. Godwin-Austen collection from Garo Hills. UMZC I.102045 from Teria Ghat (Budha et al. (2017: fig. 14a) recognised this specimen as a possible syntype).

**Diagnosis.** Shell elongate, turreted, and regularly attenuated; apex rounded; subsequent whorls with fine equally spaced radial ridges throughout and more prominent near suture. Suture impressed and whorls slightly flattened. Aperture elongate ovate; columella concave and truncated.

**Distribution.** This species is broadly distributed in Bangladesh, India, and Nepal (Budha et al. 2015; Preece et al. 2022). It was also reported from several localities in the southwest and eastern Myanmar: Ayeyarwady, Bago, and Tanintharyi regions (Benson 1860; Blanford 1865; Beddome 1906).

**Remarks.** Benson (1860: 464) reported that Theobald found a variety of ‘*Achatina tenuispira*’ on the banks of Irrawaddy [Ayeyarwady River] and from Phie Than [probably in Tanintharyi Region]. Blanford (1865) recorded the presence of a small-sized species in Akoutoung [Akauk Taung, Ayeyarwady Region] and further south, while Beddome (1906) noted its occurrence in Pegu [Bago Region].


### Genus *Tortaxis* Pilsbry, 1906

*Tortaxis* Pilsbry, 1906: 5, 6. Zilch. 1959: 347. Schileyko 1999: 534. Schileyko 2011: 10. Do and Do 2014: 455.

**Type species.** *Achatina erecta* Benson, 1842, by original designation.

**Diagnosis.** Shell slender, cylindrically shaped; spire high, mostly turreted, and gradually attenuated; embryonic whorls smooth, and subsequent whorls with nearly smooth to strong radial ribs. Aperture vertical, narrow, and oblong; columella concave or straight and with spiral fold below; columellar margin with or without expansion near umbilicus. Umbilicus narrowly opened or closed.

**Remarks.** *Tortaxis* can be distinguished from other subulinid taxa from Myanmar, namely *Allopeas*, *Opeas*, and *Bacillum*, by having a slender, cylindrical, and turreted spire, flatter whorls, smooth embryonic whorls, a large and rounded apex, and a distinct spiral fold on the columella (Table 2).



**Figure 16.** **A** *Tortaxis erectus*, neotype NHMUK 1991104A from Nanking, China **B** *Tortaxis elongatissimus*, syntype MNHN-IM-2000-4705 from Pac-Kha **C** *Tortaxis lubricus*, holotype ANSP 66056 from near Haiphong **D** *Tortaxis papillosa*, MNHN-IM-2000-33971 from Nam-Nhang **E** *Tortaxis permira*, syntype MNHN-IM-2000-4675 from That-Khé, Tonkin **F** *Tortaxis pilsbryi*, syntype MNHN-IM-2000-4678 from Bac-Khan et That-Khé **G–I** *Tortaxis cylindrotis* sp. nov. **G** holotype CUMZ 13078 from Taunggyi, Shan State, Myanmar with embryonic whorls and aperture **H, I** paratypes CUMZ 13079 from the type locality **H** probably juvenile shell with 7 whorls and **I** adult shell. Photographs: P Bourguignon & D Brabant (**B, D–F**).

The genus is mainly distributed in Southeast Asia and southern China and comprises 14 extant species (Pilsbry 1906; Schileyko 1999; MolluscaBase 2023) and one Miocene-amber fossil from China (Yu et al. 2023). In Indochina, nine species have been reported from Vietnam, and only one species from

Thailand (Panha 1998; Schileyko 2011; Do and Do 2014). In this study, we propose one new species which marks the first record of the genus in Myanmar.

**38 *Tortaxis cylindropsis* Man & Panha, sp. nov.**

<https://zoobank.org/084E4BD5-300A-4391-AF52-A6A8F6ADD951>

Fig. 16G–I, Table 1

**Type specimens.** *Holotype* CUMZ 13079 (height 9.6 mm, width 2.3 mm; Fig. 16G), *paratypes* CUMZ 13080 (35 shells; Fig. 16H, I), NHMUK 20230919 (2 shells) and SMF (2 shells).

**Type locality.** Parpant area, Taunggyi City, Taunggyi District, Shan State, Myanmar (20°15'3.7"N, 97°14'23.9"E).

**Etymology.** The specific name *cylindropsis* is from the Latin word for cylinder, and the Greek suffix *-opsis* means 'having the appearance of or like'. It refers to the cylindrical shell shape of this species.

**Diagnosis.** Shell slender cylindrical, suture deeply impressed, spire distinctly turreted, protoconch rounded and smooth, subsequent whorls with fine growth lines and a distinct spiral fold on columellar margin.

**Description.** Shell (height 9.3–11.7 mm) slender cylindrical in shape, translucent, whitish to pale yellowish colour, and with 8–9½ whorls. Apex rounded; protoconch ~ 2 whorls, rounded and with smooth surface; subsequent whorls with fine growth lines more distinct on last whorl. Spire high, grows evenly and is largely turreted; whorls flatly convex; last whorls slightly larger than preceding whorls; suture narrow, deep, and weakly crenulated. Aperture vertical, narrowly ovate, and elongate; peristome thin; columella straight; columellar margin slightly expanded with distinct spiral fold. Umbilicus closed.

**Distribution.** This species is known only from its type locality.

**Differential diagnosis.** Comparing this new species with the Vietnamese species, *T. comaensis* Do, 2014 has a much larger and taller shell (height 56.1–66.4 mm), an attenuated spire, coarser sculptures, a broad aperture, and a thickened peristome, whereas this new species display a much smaller shell (height 9.3–11.7 mm), grows evenly and has a cylindrical spire, smooth shell surface, narrow aperture, and thin peristome (Do and Do 2014). *Tortaxis erectus* (Benson, 1842) has a less turreted shell, wide and flatly convex whorls, and a shallow suture (Fig. 16A), while *T. cylindropsis* sp. nov. has a distinctly turreted, slender shell with narrow and convex whorls, and a deep suture. *Tortaxis elongatissimus* Bavay & Dautzenberg, 1909 possesses a more elongated shell with stronger radial ribs and a more attenuated spire (Fig. 16B) than this new species. *Tortaxis lubricus* Pilsbry, 1906 has a broad last whorl, attenuated spire, wider aperture, and shallow suture (Fig. 16C); in contrast, the shell of *T. cylindropsis* sp. nov. presents narrowly and even whorls, cylindrical, slender, and turreted spire, narrow aperture, and deep suture. *Tortaxis papillosa* Dautzenberg & Fischer, 1908 shows a broad shell with an attenuated spire, rounded embryonic whorls, flatter whorls, and a shallow suture (Fig. 16D), whereas *T. cylindropsis* sp. nov. has a slender shell, with cylindrical spire, convex whorls, and deeper suture. *Tortaxis permira* (Ancey in Bavay & Dautzenberg, 1904) and *T. pilsbryi* (Ancey in Bavay & Dautzenberg, 1904) have a wider shell, with a shallower suture, gradually attenuated spire, and an aperture wider than that of *T. cylindropsis* sp. nov. (Fig. 16E, F).

**Remarks.** This genus is reported for the first time in Myanmar, and the present finding expands the distribution range of the genus from China, Vietnam, Laos, and Thailand to include Myanmar.

### Genus *Zootecus* Westerlund, 1887

*Zootecus* Westerlund, 1887: 75. Kobelt 1902: 1022, 1032. von Martens 1895: 103. Pilsbry 1906: 104. Gude 1914: 366. Zilch 1959: 355. Schileyko 1999: 519, fig. 678. Neubert 2003: 154.

*Obeliscella*—Jousseau, 1889: 359.

*Chilogymnus*—Jousseau, 1894: 289.

**Type species.** *Pupa insularis* Ehrenberg, 1831, subsequent designation by Kobelt (1902: 1022).

**Diagnosis.** Shell pupiform; spire high, broad, cylindrical, and apex pointed; embryonic whorls smooth surface; subsequent whorls with irregularly dense, fine, coarse, or weak radial striations. Aperture oblique, broad, and oblong or rounded; columella straight. Penis long slender tube, and with slightly thickened wall and conical at base; vagina very much larger and muscularly thicker than male organ.

**Remarks.** *Zootecus* can be distinctly distinguished from all subulinid taxa in Myanmar by its pupiform shell, broad spire, straight columella, thickened whitish peristome, and irregular coarse striations (Table 2).

The genus is distributed from the Cabo Verde Islands and the Sahara, extending eastwards to Arabia, Socotra Island, India, and Southeast Asia (Gude 1914; Solem 1966; Schileyko 1999; Neubert 2003). Currently, the genus consists of nine species, and two are known from Myanmar (Gude 1914; MolluscaBase 2023).

### 39 *Zootecus insularis* (Ehrenberg, 1831)

Fig. 17A–C

*Pupa insularis* Ehrenberg, 1831: 13. Type locality: In insula Cameran, quae prope Maris rubri ostium australe inter Loheiam et Moccham iuxta Arabiae felicis littus sita est [Cameran Island, Red Sea between Loheia and Mocha, near coast of Arabia]. Pfeiffer 1848: 307.

*Bulimus insularis*—Pfeiffer 1853: 403. Hanley and Theobald 1870: 11, pl. 22, fig. 10.

*Pupa (Cylindrus) insularis*—Nevill 1877: 22.

*Stenogyra insularis*—von Martens 1895: 106, pl. 8, figs 5, 6.

*Zootecus insularis*—Pilsbry 1906: 106–108, pl. 26, figs 21–25, 29–33. Gude 1914: 367, 368. Schileyko 1999: 519, 520, fig. 678. Neubert 2003: 154, 155, figs 1, 2. Raheem et al. 2014: 118, fig. 74c.

**Type specimen.** **Lectotype** ZMB 109990 (Fig. 17A, after Raheem et al. 2014: fig. 74c).

**Other material.** NHMUK 1875.12.4.16 (5 shells; Fig. 17B, C) ex. Beddome collection from Burma. SMF 296651/2 (2 shells) ex. Ehrmann collection from Burma.

**Diagnosis.** Shell subcylindrical; spire high and dome-shaped; apex pointed; subsequent whorls with fine and wavy radial striae, which stronger near suture. Aperture semi-ovate; columella short and straight. Umbilicus narrow.





Figure 17. **A–C** *Zootecus insularis* **A** lectotype ZMB 109990 (after Raheem et al. 2014: fig. 74c) and **B, C** specimen NHMUK 1875.12.4.16 from Burma **D–H** *Zootecus pullus* **D** lectotype NHMUK 1986252/1 and **E** paralectotype NHMUK 1986252/2 from Banks of the Ganges, South Asia and **F–H** specimen CUMZ 13081 from Bagan, Mandalay Region, Myanmar.

**Distribution.** *Zootecus insularis* has a wide geographic distribution, spanning across the Cape Verde Islands, northeastern Africa, and the Arabian Peninsula. Its range also extends to South Asia, including India, Pakistan, Sri Lanka, and Afghanistan (Pilsbry 1906; Gude 1914; Neubert 2003; Raheem et al. 2014). In Myanmar, it was recorded from ‘Pagan’ [Bagan] in the Mandalay Region (Blanford 1865; Gude 1914).



**Figure 18.** Genitalia and radula of *Zootecus pullus*, specimen CUMZ 13082 from Bagan, Mandalay Region, Myanmar **A, B** reproductive anatomy with inset of the male organ and **C** radula morphology with inset of central teeth: yellow colour and 'C' indicates central teeth row, green colour indicates lateral teeth in the transition to marginal teeth, and numbers indicate tooth order from lateral to marginal end. Abbreviations: **fo**, free oviduct; **gd**, gametolytic duct; **gs**, gametolytic sac; **p**, penis; **pr**, penial retractor muscle; **so**, spermoviduct; **v**, vagina; **vd**, vas deferens.

**Remarks.** No new specimens of this species were collected during this survey. However, the specimens collected from Bagan in the Mandalay Region are here identified as *Z. pullus* rather than *Z. insularis* (see under *Z. pullus* for further comparison). The historical museum specimens with brief locality records such as 'Burma' (Fig. 17B, C) match well with the respective type specimens of this species (Fig. 17A). As a result, the existence of *Z. insularis* in Myanmar requires further evidence from newly collected specimens with precise collection locality data, otherwise it must be excluded from the faunal list.

#### 40 *Zootecus pullus* (Gray, 1834)

Figs 17D–H, 18, Table 1

*Bulimus pullus* Gray, 1834: 66. Type locality: India Orientali ad ripas Gangis [Eastern India, banks of the Ganges River]. Pfeiffer 1848: 162. Reeve 1849b:

*Bulimus* pl. 67, species 476. Blanford 1865: 94.

*Bulimus (Opeas) pullus*—Albers 1850: 175.

*Pupa (Cylindrus) pulla*—von Martens 1860: 297.

*Bulimina (Mastus) pulla*—Pfeiffer and Clessin 1881: 293.

*Rumina pulla*—Ancey 1886: 61.

*Stenogyra pulla*—von Martens 1895: 106, 107, pl. 8, figs 7, 8.

*Zootecus insularis* var. *pullus*—Pilsbry 1906: 110, pl. 26, figs 26–28.

*Zootecus pullus*—Gude 1914: 371, 372. Raheem et al. 2014: 118, 119, fig. 74e, f.

**Type specimens.** **Lectotype** NHMUK 1986252/1 (Fig. 17D; designated in Raheem et al. 2014). **Paralectotype** NHMUK 1986252 (1 shell; Fig. 17E) from Banks of the Ganges, South Asia.

**Other material.** Dhammayazaka Pagoda, Pwasaw Village, Bagan City, Mandalay Region, Myanmar (21°08'40.3"N, 94°52'58.0"E): CUMZ 13081 (70 shells; Fig. 17F–H), CUMZ 13082 (20 specimens in ethanol).

**Description.** Shell subcylindrical, solid, glossy, pale grey colour, slightly thick, and with 8–9½ whorls. Apex slightly elevated; protoconch ~ 2 whorls, dome-shaped and nearly smooth with fine radial striations; subsequent whorls with dense but fine, wavy, radial striations, stronger near suture. Spire grows evenly; whorls flatly convex; suture wide and shallow. Aperture nearly rounded and wide; columella straight; peristome relatively thickened, expanded, and white. Umbilicus narrowly opened.

**Genitalia** ( $n = 5$ ). Atrium undifferentiated. Penis very narrow, slender, almost same length with vagina, and slightly bulging at base. Penial retractor slender, long and attached at junction of penis and vas deferens. Epiphallus very short or indistinct. Vas deferens long, slender tube, and connected between penis/epiphallus to free oviduct (Fig. 18A, B).

Vagina much larger than penis and cylindrical shape. Gametolytic duct short and slender tube; gametolytic sac distinct and bulbous shape. Free oviduct almost the same diameter as vagina; spermooviduct enlarged.

**Radula.** Each row contains ~ 43+ teeth with half-row formula: central-lateral-marginal teeth (1–(11–13)–(6–7+)). Central tooth relatively small, tricuspid with pointed central cusp, and small, rounded, lateral cusps. Lateral teeth bicuspid: endocone large rhomboid in shape and with pointed to dull tip; ectocone small, pointed tip and located at middle of tooth height. Marginal teeth asymmetrically tricuspid starting approximately at tooth number 11–13: mesocone large, triangular, and curved to blunt tip; endocone small and located near tip of mesocone; ectocone triangular, pointed tip and located near tooth base. Outermost teeth small and polycuspid (Fig. 18C).

**Distribution.** Apart from Myanmar, this species is likely to occur in India and Bangladesh (Raheem et al. 2014).

**Remarks.** In Myanmar, the previous records of *Z. pullus* were from 'Ava' [Innwa in Mandalay Region]. Our newly collected specimens from Bagan (~ 150 km southwest of Innwa) are identified as this species, but they present some differences such as having a long and narrower shell and a blunt apex. *Zootecus*

*pullus* has a penis nearly equal in length to the vagina, and the penis is slightly enlarged with a conical shape at base near the atrium (Fig. 18A, B), while *Z. insularis* possesses a penis shorter than the vagina, and the evenly slender penis (see Schileyko 1999: fig. 678 from a near type locality in the Red Sea). Further investigation is required to determine whether these two Myanmar species are distinct or simply demonstrate variation in shell morphology.

## Conclusions

This study elucidates the historically known species of the Subulinidae in Myanmar, identifying nine genera and 40 taxa, including 2 newly described species. Of these, 17 species are restricted only to Myanmar, while the remaining species are also found in India, China, and other Southeast Asian countries. The genera *Glessula* and *Rishetia* represent the highest number of species, 12 and 10 species, respectively, and they are predominantly found in Myanmar rather than other countries in Indochina. The globally distributed *Allopeas gracile* and the newly described species *Tortaxis cylindropsis* sp. nov. are presently documented as the only species of these genera in Myanmar.

Except for *Opeas innocens*, all other unique name-bearing types of the subulinid species recorded from Myanmar have been examined, revised, and are illustrated herein. Our revised taxonomy is grounded in literature records; however, the classification of one particular taxa that has never been illustrated or for which there are no available types remains somewhat arbitrary. Additionally, in regards to the distribution range of certain species, for instance e.g., *G. crassilabris*, *G. orophila*, and *R. hastula*, which extend well beyond their original type locality in India, further data is needed to confirm their occurrence in Myanmar.

The distribution of these subulinid snails sheds light on their historical prevalence. According to the literature, they are primarily found in regions such as Bago, Ayeyarwady, Magway, Rakhine, and Shan rather than in Mon and Kayin states, which are known for their rich limestone outcrops (Gude 1914; Godwin-Austen 1920). Species belonging to the genera *Glessula*, *Opeas*, *Paropeas*, *Tortaxis*, and *Zootecus* were collected in Shan State, Mandalay, and Tanintharyi regions. However, our recent survey did not encounter representative taxa for *Bacillum*, *Curvella*, and *Rishetia*. Instead, species like *Paropeas turricula* and *Opeas filiforme* have expanded their distribution from Thailand into Myanmar, and the genus *Tortaxis* is recorded for the first time. In contrast, other families like Ariophantidae (Pholyotha et al. 2020; Sutcharit and Pholyotha 2023), Clausiliidae (Man et al. 2023), Helicarionidae (Sutcharit et al. 2020a; Pholyotha et al. 2022a, b), Hypselostomatidae (Tongkerd et al. 2024), and Streptaxidae (Man et al. 2022) have yielded several species in Mon and Kayin states during these surveys. Hence, subsequent surveys should focus on regions such as Bago, Ayeyarwady, Magway, and Rakhine, aiming to confirm the identity of uncertain species and to undertake a thorough revision using freshly collected material. Moreover, the occurrence of new records suggests that additional taxa are likely to be discovered.

Budha et al. (2017) highlighted that shell sculptures may serve as valuable characters in distinguishing *Glessula* and *Rishetia*. Likewise, our limited new sample of *Glessula* demonstrates a range of shell surface variations from smooth to ribbed sculpture. Species like *G. mandalayensis* sp. nov. and

*G. gemma* exhibit smooth to fine striations, while grooves are observed in *G. feddeni* and *G. latestriata*, and ribbed sculpture is characteristic of *G. blanfordiana*. Similarly, *Paropeas* is characterised by rough and compact radial striations, while *Allopeas* has finer striations compared to *Paropeas*. On the other hand, *Opeas* and *Tortaxis* display the very fine striations or smooth surfaces among these two genera. However, future research in subulinid taxonomy still requires the integration of genetic data, genital anatomy, and shell morphological traits for a more comprehensive understanding of their systematics.

## Acknowledgements

The senior author is grateful to all members of the Animal Systematics Research Unit (ASRU), Chulalongkorn University, for their kind help during field trips in Myanmar. The authors are indebted to V Héros, P Bourguignon, and D Brabant (project ERECOLNAT: ANR-11-INBS-0004, MNHN, Paris); K Webb and H Taylor (NHMUK, London), R Janssen, J. Sigwart, K-O Nagel, and S Hof (SMF, Frankfurt), and B Páll-Gergely (Centre for Agricultural Research, Budapest) for kindly providing the authors with the shell photographs of material housed in their type collections. We thank the personnel of the Apache Cement Company for sponsoring our surveys in the Mandalay Region. We also thank the Ministry of Natural Resources and the Environmental Conservation Forest Department, Myanmar, for the collection and export permits. We also thank DJ Anderson for his initial linguistic help in the original manuscript. We gratefully acknowledge the reviewers, F Naggs, B Páll-Gergely, JK Foon, S D'ávila and an anonymous reviewer for their helpful and constructive comments, which significantly improved the manuscript.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

The work in Thailand was funded by the Thailand Research Fund (TRF-DPG628001), the NSRF via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation (B42G670038), National Research Council of Thailand (NRCT-N35E660138) and Centre of Excellence on Biodiversity (MHESI), and additional support by CU-ASEAN Scholarships and the 90<sup>th</sup> Anniversary of Chulalongkorn University Fund to NSM. The field trips in Myanmar were partially funded through grants received from Fauna and Flora International (FFI) and Thailand Research Fund (TRF-RTA 5880002).

### Author contributions

NSM, NL, CS, and SP collected and prepared specimens in the field. NL, CS, and SP provided financial and intellectual support. NSM and CS prepared specimens and wrote the manuscript. NSM and CS conceived, designed, supervised the study, prepared figures, and approved and edited the final manuscript. JDA reviewed, advised, and approved the final manuscript.

## Author ORCIDs

Nem Sian Man  <https://orcid.org/0000-0002-4453-734X>  
Jonathan D. Ablett  <https://orcid.org/0000-0002-7277-1934>  
Ngwe Lwin  <https://orcid.org/0000-0002-4394-3741>  
Chirasak Sutcharit  <https://orcid.org/0000-0001-7670-9540>  
Somsak Panha  <https://orcid.org/0000-0002-4431-2458>

## Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Adams H, Adams A (1855) [1854–1858] The Genera of Recent Mollusca; Arranged According to their Organization. Volume 2. van Voorst, London, 661 pp. [pp 93–284 (1855)]
- Albers JC (1850) Die Heliceen nach natürlicher Verwandtschaft systematisch geordnet. Enslin, Berlin, 262 pp. <https://doi.org/10.5962/bhl.title.11507>
- Albers JC (1860) Die Heliceen nach Natürlicher Verwandtschaft Systematisch Geordnet, 2<sup>nd</sup> edn. (manuscript edited by E. von Martens). Wilhelm Englemann, Leipzig, 359 pp. <https://doi.org/10.5962/bhl.title.11218>
- Ancey CF (1886) Essai monographique sur le *Buliminus* de l'Asie Centrale Russe et de l'Afghanistan. Bulletin de la Société Malacologique de France 3: 15–64.
- Baker HB (1935) Jamaican land snails, 3. The Nautilus 48(3): 83–88. <https://doi.org/10.5962/bhl.part.11532>
- Bavay A, Dautzenberg P (1904) Description de coquilles nouvelles de l'Indo-Chine. (3e suite). Journal de Conchyliologie 51[1903]: 201–236.
- Bavay A, Dautzenberg P (1909) Molluscorum terrestrum tonkinorum diagnoses. Journal de Conchyliologie 56[1908]: 229–251. <https://www.biodiversitylibrary.org/page/16298225>
- Beddome CR (1906) Notes on Indian and Ceylonese species of *Glessula*. The Journal of Molluscan Studies 6(3): 160–172. <https://doi.org/10.1093/oxfordjournals.mollus.a066160>
- Benson WH (1836) Descriptive catalogue of terrestrial and fluviatile Testacea, chiefly from the North-East frontier of Bengal. Journal of the Asiatic Society of Bengal 5(54): 350–358. <https://biodiversitylibrary.org/page/37188459>
- Benson WH (1842) Mollusca. In: T. Cantor, General features of Chusan, with remarks on the flora and fauna of that island. Annals and Magazine of Natural History, Series 1 9(60): 486–490. <https://www.biodiversitylibrary.org/page/18684698>
- Benson WH (1857) New species of *Bulimus* from India, Burma, and Mauritius. Annals & Magazine of Natural History Series 2 19(112): 327–330. <https://doi.org/10.1080/00222935708693934>
- Benson WH (1860) On new land-shells from Darjiling, with a series of new Indian species of *Achatina*. Annals & Magazine of Natural History Series 3 5(30): 460–466. <https://doi.org/10.1080/00222936008697255>
- Benson WH (1863) Characters of new land-shells from the Andaman Islands, Burmah, and Ceylon, and of the animal of *Sophina*. Annals & Magazine of Natural History Series 3 11(65): 318–323. <https://doi.org/10.1080/00222936308681435>
- Benson WH (1865) New land-shells from Travancore, western and northern India. Annals & Magazine of Natural History Series 3 15(85): 11–15. <https://doi.org/10.1080/00222936508681753>
- Blanford WT (1861) Contributions to Indian malacology, No. II. Journal of the Asiatic Society of Bengal 30(4): 347–366. <https://www.biodiversitylibrary.org/item/207709>

- Blanford WT (1865) Contributions to Indian Malacology, No. 5. Description of new land shells from Arakan, Pegu, and Ava; with notes on the distributions of described species. *Journal of the Asiatic Society of Bengal* 34(2): 66–105. <https://www.biodiversitylibrary.org/item/114408>
- Blanford WT (1869) Descriptions of new land and freshwater molluscan species collected by Dr. John Anderson in Upper Burma and Yunnan. *Proceedings of the Zoological Society of London* 37(1): 444–450. <https://doi.org/10.1111/j.1469-7998.1869.tb07354.x>
- Blanford WT (1871) On some undescribed species of *Camptoceras* and other land shells. *Journal of the Asiatic Society of Bengal* 40: 39–45. <https://www.biodiversitylibrary.org/page/35630849>
- Blanford WT (1903) Notes on Mr. WM Daly's collections of land and fresh-water Mollusca from Siam. *Proceedings of the Malacological Society of London* 5(4): 274–284. <https://doi.org/10.1093/oxfordjournals.mollus.a065982>
- Bouchet P, Rocroi JP, Hausdorf B, Kaim A, Kano Y, Nützel A, Strong EE (2017) Revised classification, nomenclator and typification of gastropod and monoplacophoran families. *Malacologia* 61(1–2): 1–526. <https://doi.org/10.4002/040.061.0201>
- Bourguignat J-R (1883) Histoire malacologique de l'Abyssinie. *Annales des Sciences Naturelles (Zoologie et Paléontologie)* (6)15(2): 1–162. <https://www.biodiversitylibrary.org/bibliography/46223#/summary>
- Budha PB (2017) First record of *Curvella sikkimensis* (Reeve, 1850) (Pulmonata: Subulinidae: Subulininae) from Nepal and its reproductive organs. *Nepalese Journal of Biosciences* 7(1): 120–124. <https://doi.org/10.3126/njbs.v7i1.41801>
- Budha PB, Naggs F, Backeljau T (2015) Annotated checklist of the terrestrial gastropods of Nepal. *ZooKeys* 492: 1–48. <https://doi.org/10.3897/zookeys.492.9175>
- Budha PB, Naggs F, Backeljau T (2017) Conchological differentiation and genital anatomy of Nepalese Glessulinae (Gastropoda, Stylommatophora, Subulinidae), with descriptions of six new species. *ZooKeys* 675: 129–156. <https://doi.org/10.3897/zookeys.675.13252>
- Chaper M (1885) Description de quelques espèces et genres nouveaux de coquilles vivantes de diverses provenances. *Bulletin de la Société Zoologique de France* 10: 42–54. <https://doi.org/10.5962/bhl.part.14302>
- D'ávila S, Medeiros C, Vargas T, Mendonça CLF (2018) Life history of *Subulina octona* (Brugüière) (Gastropoda: Pulmonata: Subulinidae) based on four-year laboratory observations and a comparative histological analysis of egg-retaining and ovoviviparous subulinids. *Journal of Natural History* 52(23–24): 1551–1569. <https://doi.org/10.1080/00222933.2018.1478996>
- D'ávila S, Simone LRL, De Oliveira LFC, Charles L, Maestrati P (2020) Rediscovery of *Obeliscus agassizi* Pilsbry, 1906 (Gastropoda, Subulinidae, Obeliscinae), annotated checklist of species of *Obeliscus* Beck, 1837 and first description of the anatomy for the genus. *Zoosystema* 42(12): 159–172. <https://doi.org/10.5252/zoosystema20v42a12>
- Dautzenberg P, Fischer H (1908) Liste des mollusques récoltés par M. Mansuy en Indo-Chine et description d'espèces nouvelles. *Journal de Conchyliologie* 56: 169–217.
- de Morgan J (1885) Mollusques terrestres et fluviatiles de Royaume de Perak et des pays voisins (presque l'île Malaise). *Bulletin de la Société Zoologique de France* 10: 353–429. <https://doi.org/10.5962/bhl.part.14301>
- Do DS, Do VN (2014) The land snail family Subulinidae (Gastropoda: Pulmonata) from Sonla, Vietnam, with description of two new species. *Tapchi Sinh Hoc* 36(11): 451–459. <https://doi.org/10.15625/0866-7160/v36n4.6174>
- Ehrenberg CG (1831) *Animalia evertebrata. Symbolae physicae seu icones et descriptiones naturalium novorum aut minus cognitorum quae ex itineribus Lybiam Aegyptum*

- Nubiam Dongalam Syriam Arabiam et Habessinian. Pars Zoologica. Abhandlungen der deutschen Akademie der Wissenschaften. 4: (unpaginated). <https://www.biodiversitylibrary.org/page/48517192>
- Férussac AE d'Audebard de baron de (1821) [1821–1822] Tableaux systématiques des animaux mollusques suivis d'un Prodre général pour tous les mollusques ou fluviatiles, vivantes ou fossiles, 114 pp. [pp 1–114 (1821); pp i–xlvii (1822)] <https://www.biodiversitylibrary.org/bibliography/10558>
- Fischer P, Crosse H (1877) Études sur les mollusques terrestres et fluviatiles du Mexique et du Guatemala. Mission scientifique au Mexique et dans l'Amérique Centrale. Recherches Zoologiques, Partie 7, Volume 1, 702 pp. <https://www.biodiversitylibrary.org/item/216870>
- Fontanilla IK, Naggs F, Wade CM (2017) Molecular phylogeny of the Achatinoidea (Mollusca: Gastropoda). *Molecular Phylogenetics and Evolution* 114: 382–385. <https://doi.org/10.1016/j.ympev.2017.06.014>
- Foon JK, Clements GR, Liew TS (2017) Diversity and biogeography of land snails (Mollusca, Gastropoda) in the limestone hills of Perak, Peninsular Malaysia. *ZooKeys* 682: 1–94. <https://doi.org/10.3897/zookeys.682.12999>
- Godwin-Austen (1876) On the Helicidae collected during the expedition into the Dafla Hills, Assam. *Journal of the Asiatic Society of Bengal* 45(4): 311–184. <https://www.biodiversitylibrary.org/item/114186>
- Godwin-Austen HH (1895) List and distribution of the Land-Mollusca of the Andaman and Nicobar Islands, with descriptions of some supposed new species. *Proceedings of the Zoological Society of London* 63(30): 438–457. <https://www.biodiversitylibrary.org/page/30983429>
- Godwin-Austen HH (1918) Zoological results of the Abor Expedition, 1911–1912. XLIX. Mollusca, IX. Records of the Indian Museum 8(12): 601–621. <https://doi.org/10.26515/rzsi/v8/i12/1918/163139>
- Godwin-Austen HH (1920) Land and freshwater Mollusca of India, including South Arabia, Baluchistan, Afghanistan, Kashmir, Nepal, Burmah, Pegu, Tenasserim, Malay Peninsula, Ceylon, and other islands of the Indian Ocean. Supplementary to Messrs. Theobald and Hanley's *Conchologia Indica*. London, Taylor & Francis, Volume 3, 1–65. [pls 159–165] <https://www.biodiversitylibrary.org/item/57585>
- Gray JE (1834) Exhibition of an extensive series of land and freshwater shells regarded as hitherto undescribed. *Proceedings of the Zoological Society of London* 2: 63–68. <https://biodiversitylibrary.org/page/12859659>
- Gray JE (1860) On the arrangement of the land pulmoniferous Mollusca into familie. *Annals & Magazine of Natural History Series* 3 6(34): 267–269. <https://doi.org/10.1080/00222936008697321>
- Gude GK (1914) Mollusca-II. Trochomorphidae-Janellidae. In: Shipley AE, Marshall GAK (Eds) *The Fauna of British India, including Ceylon and Burma*. Taylor & Francis, London, 520 pp. <https://doi.org/10.5962/bhl.title.12891>
- Hanley SCT, Theobald W (1870, 1873, 1874) [1870–1876] *Conchologia Indica: Illustrations of the Land and Freshwater Shells of British India*. London, 1–65. [pls 1–160] [pp 1–18, pls 1–40 (1870); pp 19–28, pls 41–60 (1872); pp 29–34, pls 61–80 (1873). pp 35–48, pls 81–120 (1874); pp 49–56, pls 121–140 (1875); pp 57–65, pls 141–160 (1876)] <https://www.biodiversitylibrary.org/item/50391>
- Horsák M, Naggs F, Bäckeljau T (2020) *Paropeas achatinaceum* (Pfeiffer, 1846) and other alien Subulinine and Opeatine land snails in European greenhouses (Gastropoda, Achatinidae). *Malacologia* 63(1): 123–130. <https://doi.org/10.4002/040.063.0112>

- Hutton T (1834) On the land shells of India. *Journal of the Asiatic Society of Bengal* 3(26): 81–93.
- Illiger JCW (1801) *Namen der Insekten-Gattungen, ihr Genitiv, ihr grammatisches Geschlecht, ihr Silbenmass, ihr Herleitung; zugleich mit den Deutschen Benennungen.* *Magazin für Insektenkunde* 1: 125–155.
- Inkhavilay K, Sutcharit C, Bantaowong U, Chanabun R, Siriwit W, Srisonchai R, Pholyotha A, Jirapatrasilp P, Panha S (2019) Annotated checklist of the terrestrial molluscs from Laos (Mollusca, Gastropoda). *ZooKeys* 834: 1–166. <https://doi.org/10.3897/zookeys.834.28800>
- International Commission on Zoological Nomenclature (ICZN) (1999) *International Code of Zoological Nomenclature (4<sup>th</sup> edn.)* International Trust for Zoological Nomenclature, London, 306 pp.
- Jaeckel S (1950) Die Mollusken eines tropischen Flußgenistes aus Tonkin. *Archiv für Molluskenkunde* 79: 15–20.
- Jousseaume F (1889) Espèces nouvelles des environs d'Aden suivies d'un aperçu sur la faune malagologique de la Péninsule Arabique. *Bulletins de la Société Malacologique de France* 6: 345–362. <https://www.biodiversitylibrary.org/page/16141631>
- Jousseaume F (1894) Mollusques recueillis à Ceylan par M.E. Simon, et révision générale des espèces terrestres et fluvio-lacustres de cette île. *Mémoires de la Société Zoologique de France* 7: 264–330. <https://www.biodiversitylibrary.org/page/36340927>
- Kalita G (2022) Land snails of Guwahati, Assam, India. *Journal of Threatened Taxa* 14(9): 21845–21852. <https://doi.org/10.11609/jott.8019.14.9.21845-21852>
- Kerney MP, Cameron RAD (1979) *A Field Guide to the Land Snails of Britain and North-west Europe.* Collins, London.
- Kobelt W (1901, 1902) [1899–1902] Die Familie Buliminidae. In: *Systematisches Conchylien-Cabinet von Martini und Chemnitz* 1, 13(2): 397–1051. [pls 71–133] [pp 397–620, pls 71–94 (1899); pp 621–684, pls 95–103 (1900); pp 685–836, pls 104–123 (1901); pp 837–1051, pls 124–133 (1902)] <https://www.biodiversitylibrary.org/item/106746>
- Laidlaw FF (1933) A list of the land and fresh-water Mollusca of the Malay Peninsula. *Journal of the Malaysian Branch of the Royal Asiatic Society* 11: 211–234.
- Lamouroux JVF (1816) *Histoire des Polypiers Coralligènes Flexibles, Vulgairement Nommés Zoophytes.* F. Poisson, Caen, 558 pp. <https://www.biodiversitylibrary.org/page/11606092>
- Maassen WJM (2001) A preliminary checklist of the non-marine molluscs of West Malaysia. *De Kreukel* 2001: 1–155.
- Man NS, Siriboon T, Lin A, Sutcharit C, Panha S (2022) Revision of the carnivorous land snail family Streptaxidae (Stylommatophora, Achatinina) in Myanmar, with description of four new species. *ZooKeys* 1110: 39–102. <https://doi.org/10.3897/zookeys.1110.85399>
- Man NS, Lwin N, Sutcharit C, Panha S (2023) Further notes on the taxonomy of the land snail family Clausiliidae Gray, 1855 (Stylommatophora, Helicina) from Myanmar with description of two new species. *ZooKeys* 1160: 1–59. <https://doi.org/10.3897/zookeys.1160.98022>
- Miller JS (1822) A list of the freshwater and landshells occurring in the environment of Bristol, with observations. *Annals of Philosophy, New Series* 3(5): 376–381. <https://www.biodiversitylibrary.org/page/2498134>
- MolluscaBase eds (2023) *MolluscaBase.* <https://www.molluscabase.org> [in 2023–07–17] <https://doi.org/10.14284/448>
- Mörch OAL (1876) Revision des mollusques terrestres des îles Nicobar. *Journal de Conchyliologie* 24: 353–367. <https://www.biodiversitylibrary.org/page/16071668#page/375>

- Morlet L (1893) Descriptions d'espèces nouvelles provenant de l'Indochine. *Journal de Conchyliologie* 40[1892]: 315–329.
- Mousson A (1857) Novitates Zollingerianae (1). *Journal de Conchyliologie* 6(2): 154–164.
- Naggs F (1994) The reproductive anatomy of *Paropeas achatinaceum* and a new concept of *Paropeas* (Pulmonata: Achatinoidea: Subulinidae). *The Journal of Molluscan Studies* 60(2): 175–191. <https://doi.org/10.1093/mollus/60.2.175>
- Neubert E (2003) The continental malacofauna of Arabia and adjacent areas. II. The genus *Zootecus* Westerlund, 1887. *Archiv für Molluskenkunde der Senckenbergischen Naturforschenden Gesellschaft* 132(1/2): 153–160. <https://doi.org/10.1127/arch.moll/132/2003/153>
- Neubert E, Bochud E (2020) The continental malacofauna of Arabia and adjacent areas, VII. The family Achatinidae (Coeliacinae, Subulininae) in the Socotran Archipelago (Gastropoda, Pulmonata). *Rendiconti Lincei. Scienze Fisiche e Naturali* 31(3): 655–676. <https://doi.org/10.1007/s12210-020-00920-y>
- Nevill G (1877) List of the Mollusca brought back by Dr. J. Anderson from Yunnan and Upper Burma, with descriptions of new species. *Journal of the Asiatic Society of Bengal* 46(2): 14–41. <https://www.biodiversitylibrary.org/item/110105>
- Nevill G (1878) Hand list of Mollusca in the Indian Museum, Calcutta. Part I. Gastropoda. Indian Museum, Calcutta, 338 pp. <https://doi.org/10.5962/bhl.title.11957>
- Nevill G (1881) New or little-known Mollusca of the Indo-Malayan Fauna. *Journal of the Asiatic Society of Bengal* 50(2): 125–167. <https://www.biodiversitylibrary.org/item/110180>
- Nurinsiyah AS, Hausdorf B (2019) Listing, impact assessment and prioritization of introduced land snail and slug species in Indonesia. *The Journal of Molluscan Studies* 85(1): 92–102. <https://doi.org/10.1093/mollus/eyy062>
- Panha S (1998) A checklist and classification of the terrestrial pulmonate snails of Thailand. *Walkerana* 8[1995–1996](19): 31–40.
- Pfeiffer L (1845) Diagnosen einiger neuer Heliceen. *Zeitschrift für Malakozoologie* 2: 152–158. <https://biodiversitylibrary.org/page/16291740>
- Pfeiffer L (1846) Descriptions of nine new species of Helicea, collected by Hugh Cum-  
ing, Esq. *Proceedings of the Zoological Society of London* 14: 41–43.
- Pfeiffer L (1848) *Monographia Heliceorum Viventium: sistens descriptiones system-  
aticas et criticas omnium huius familiae generum et specierum hodie cognitarum.*  
Volumen Secundum. Lipsiae: F.A. Brockhaus, 594 pp. <https://doi.org/10.5962/bhl.title.124010>
- Pfeiffer L (1853) *Monographia Heliceorum Viventium: sistens descriptiones systemati-  
cas et criticas omnium huius familiae generum et specierum hodie cognitarum.* Vol-  
umen Tertium. Lipsiae: F.A. Brockhaus, 711 pp. <https://www.biodiversitylibrary.org/item/47349>
- Pfeiffer L (1856) Versuch einer Anordnung der Heliceen nach natürlichen Gruppen. *Mal-  
akozoologische Blätter* 2: 112–185. <https://www.biodiversitylibrary.org/item/53876>
- Pfeiffer L (1859) *Monographia Heliceorum Viventium: sistens descriptiones system-  
aticas et criticas omnium huius familiae generum et specierum hodie cognitarum.*  
Volumen Quartum. Lipsiae: F.A. Brockhaus, 920 pp. <https://www.biodiversitylibrary.org/page/11648889>
- Pfeiffer L (1860) [1840–1865] [in HC Küster, L Pfeiffer] Die Bulimiden und Achatinen,  
In *Abbildungen nach der Natur. Die gattungen Bulimus, Partula, Achatinella, Acha-  
tina und Azeca.* In: Küster HC, Ed. *Systematisches Conchylien-Cabinet von Martini  
und Chemnitz.* Neu herausgegeben und vervollständigt. Erster Bandes, dreizehnte

- Abtheilung. 1 (13) [(1, 124)]: iii–xix, 1–359. [pls 1–70] [pp 305–336 (1860), pls 25–29 (1863)] <https://www.biodiversitylibrary.org/item/106739>
- Pfeiffer L (1868) Monographia Heliceorum Viventium: sistens descriptiones systematicas et criticas omnium huius familiae generum et specierum hodie cognitarum. Volumen Sextum. Lipsiae: F.A. Brockhaus, 598 pp. <https://www.biodiversitylibrary.org/item/46666>
- Pfeiffer L (1876) [1876–1877] Monographia Heliceorum Viventium: sistens descriptiones systematicas et criticas omnium huius familiae generum et specierum hodie cognitarum. Volumen Octavum. Lipsiae: F.A. Brockhaus, 729 pp. [pp 1–480 (1876); pp 481–729 (1877)] <https://www.biodiversitylibrary.org/item/47348>
- Pfeiffer L, Clessin S (1881) Nomenclator Heliceorum Viventium: quo continentur nomina omnium hujus familiae generum et specierum hodie cognitarum, disposita ex affinitate naturali. Kassel, Fischer, 617 pp. <https://www.biodiversitylibrary.org/item/47266>
- Pholyotha A, Sutcharit C, Tongkerd P, Lin A, Panha S (2020) Taxonomic revision of the land snail genera *Macrochlamys* Gray, 1847 and *Sarika* Godwin-Austen, 1907 (Eupulmonata: Ariophantidae) from south-eastern Myanmar, with descriptions of three new species. *Molluscan Research* 40(2): 183–204. <https://doi.org/10.1080/13235818.2020.1723041>
- Pholyotha A, Sutcharit C, Lin A, Panha S (2022a) Multigene phylogeny reveals the ribbed shell morphotypes in the land snail genus *Sarika* (Eupulmonata: Ariophantidae), with description of two new species from Thailand and Myanmar. *Contributions to Zoology (Amsterdam, Netherlands)* 91(2): 97–132. <https://doi.org/10.1163/18759866-bja10027>
- Pholyotha A, Sutcharit C, Lin A, Panha S (2022b) Uncovering local endemism from southeastern Myanmar: Description of the new karst-associated terrestrial snail genus *Burmochlamys* (Eupulmonata, Helicarionidae). *ZooKeys* 1110: 1–37. <https://doi.org/10.3897/zookeys.1110.82461>
- Pilsbry HA (1906) [1906–1907] Manual of Conchology, Structural and Systematic, with Illustrations of the Species, Series 2, Pulmonata. Volume 18: Achatinidae: Stenogyrinae and Coeliaxinae. Conchological Section, Academy of Natural Sciences of Philadelphia, Philadelphia, 1–357. [pls 1–51] [pp 1–272, pls 1–34 (1906); pp 273–357, pls 35–51 (1907)] [https://doi.org/10.1175/1520-0493\(1907\)35%3C357a:LP%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1907)35%3C357a:LP%3E2.0.CO;2)
- Pilsbry HA (1908) [1908–1910] Manual of Conchology, Structural and Systematic, with Illustrations of the Species, Series 2, Pulmonata. Volume 20: Caecilioides, *Glessula* and Partulidae. Conchological Section, Academy of Natural Sciences of Philadelphia, Philadelphia, 1–336. [pls 1–43] [pp 1–64, pls 1–10 (1908); pp 65–314, pls 22–36 (1909); pp 315–336, pls 37–43 (1910)] <https://www.biodiversitylibrary.org/item/16293>
- Preece RC, White TS, Raheem DC, Ketchum H, Ablett J, Taylor H, Naggs F (2022) William Benson and the golden age of malacology in British India: Biography, illustrated catalogue and evaluation of his molluscan types. *Tropical Natural History (Supplement 6)*: 1–434.
- Preston HB (1910) Description of new shells in the collection of the Indian Museum from Burma, Siam, and the Bay of Bengal. *Records of the Indian Museum* 5(1): 33–36. <https://doi.org/10.26515/rzsi/v5/i1/1910/163757>
- Raheem DC, Taylor H, Ablett J, Preece RC, Aravind NA, Naggs F (2014) A systematic revision of the land snails of the Western Ghats of India. *Tropical Natural History (Supplement 4)*: 1–294.
- Ramakrishna, Mitra SC, Dey A (2010) Annotated checklist of Indian land molluscs. Occasional paper no. 306. Zoological Survey of India, 359 pp.

- Rang S (1831) Description des coquilles terrestres recueillies pendant un voyage à la côte occidentale d'Afrique, et au Brésil. *Annales des Sciences Naturelles* 24(93): 5–63.
- Reeve LA (1849a, 1850) [1849–1850] *Conchologia Iconica: Illustrations of the shells of molluscous animals*. Volume 5, Monograph of the genus *Achatina*, plates 1–23. Lovell Reeve and Co., London. [pls 1–19 (1849); pls 20–23 (1850)]. <https://www.biodiversitylibrary.org/page/8967091>
- Reeve LA (1849b) [1848–1850] *Conchologia Iconica: Illustrations of the shells of molluscous animals*. Volume 5, Monograph of the genus *Bulimus*, plates 1–89. Lovell Reeve and Co., London. [pls 1–57 (1848); pls 58–84 (1849); pls 85–89 (1850)] <https://www.biodiversitylibrary.org/item/35904>
- Schileyko AA (1999) Treatise on recent terrestrial pulmonate molluscs Part 4. Draparnaudiidae, Caryodidae, Macrocyclidae, Acavidae, Clavatoridae, Dorcasiidae, Sculptariidae, Corillidae, Plectopylidae, Megalobulimidae, Strophocheilidae, Cerionidae, Achatinidae, Subulinidae, Glessulidae, Micractaeonidae, Ferrussaciidae. *Ruthenica* (Supplement 2): 437–564.
- Schileyko AA (2011) Checklist of land pulmonate molluscs of Vietnam (Gastropoda: Stylommatophora). *Ruthenica* 21(1): 1–68.
- Schileyko AA, Kuznetsov AG (1996) A new genus of the Subulinidae (Pulmonata) from Nepal. *Ruthenica* 5(2): 158–160.
- Semper C (1874) [1870–1885] *Reisen im Archipel der Philippinen, Theil 2. Wissenschaftliche Resultate*. Band 3, Landmollusken. Wiesbaden: Kreidel, 327 pp. [pp 129–168 (1874)]
- Solem A (1966) Some non-marine mollusks from Thailand. *Spolia Zoologia Musei Havnensis* 24: 1–110.
- Sutcharit C, Pholyotha A (2023) Hidden lineages in the mountains: The genus *Glyptaulax* Gude, 1914 and *Maelamaodiscus* gen. nov. (Heterobranchia: Stylommatophora: Charopidae and Ariophantidae) with description of two new species from western Thailand. *Tropical Natural History* (Supplement 7): 123–138.
- Sutcharit C, Jeratthitikul E, Pholyotha A, Lin A, Panha S (2020a) Molecular phylogeny reveals high diversity and endemism in the limestone karst-restricted land snail genus *Sophina* Benson, 1859 from Myanmar (Eupulmonata: Helicarionidae), with description of four new species. *Journal of Zoological Systematics and Evolutionary Research* 58(4): 957–981. <https://doi.org/10.1111/jzs.12420>
- Sutcharit C, Thach P, Chhuoy S, Ngor PB, Jeratthitikul E, Siritwut W, Panha S (2020b) Annotated checklist of the land snail fauna from southern Cambodia (Mollusca, Gastropoda). *ZooKeys* 948: 1–46. <https://doi.org/10.3897/zookeys.948.51671>
- Swainson W (1840) *A treatise on malacology; or the natural classification of shells and shell-fish*. Longman, London, 419 pp. <https://doi.org/10.5962/bhl.title.8027>
- Tenison-Woods JE (1888) Malaysian land and freshwater Mollusca. *Proceedings of the Linnean Society of New South Wales* 3: 1003–1095. <https://doi.org/10.5962/bhl.title.13026>
- Theobald W (1870) Descriptions of some new land shells from the Shan States and Pegu. *Journal of the Asiatic Society of Bengal* 39(1): 395–402.
- Theobald W (1878) Notes on the land and freshwater shells of Kashmir, more particularly of the Jhilum valley below Srinagar and the hills north of Jammu. *Journal of the Asiatic Society of Bengal* 47(2): 141–149.
- Theobald W, Stoliczka F (1872) Notes on Barmese and Arakanese land shells, with descriptions of a few new species. *Journal of the Asiatic Society of Bengal* 41(2): 329–334.

- Thiele J (1931) [1929–1935] Handbuch der systematischen Weichtierkunde. Jena, Gustav Fischer, 1154 pp. [Volume 1, pp 1–376 (1929), pp 377–778 (1931), pp 779–1022 (1934); Volume 2, pp 779–1022 (1935)]
- Tongkerd P, Lwin N, Páll-Gergely B, Chanabun R, Pholyotha A, Prasankok P, Seesamut T, Siriwut W, Srisonchai R, Sutcharit C, Panha S (2024) Contributions of a small collection of terrestrial microsnails (Pupilloidea, Hypselostomatidae) from Myanmar with description of three new species. *ZooKeys* 1195: 157–197. <https://doi.org/10.3897/zookeys.1195.112112>
- van Benthem Jutting WSS (1952) Systematic studies on the non-marine Mollusca of the Ind-Australian archipelago. *Treubia* 21: 291–435.
- van Benthem Jutting WSS (1959) Catalogue of the non-marine Mollusca of Sumatra and of its satellite islands. *Beaufortia* 7: 41–191.
- Vaught KC (1989) A Classification of the Living Mollusca. American Malacologists, Inc., Melbourne, FL, 195 pp.
- Vermeulen JJ, Liew TS, Schilthuizen M (2015) Additions to the knowledge of the land snails of Sabah (Malaysia, Borneo), including 48 new species. *ZooKeys* 531: 1–139. <https://doi.org/10.3897/zookeys.531.6097>
- von Martens E (1860) On the Mollusca of Siam. *Proceedings of the Zoological Society of London* 28: 6–18. <https://www.biodiversitylibrary.org/page/12866504>
- von Martens E (1864) Diagnosen neuer Heliceen aus dem ostasiatischen Archipel. *Monatsberichte der Königlichen Preussischen Akademien der Wissenschaften zu Berlin* 1864: 523–529. <https://www.biodiversitylibrary.org/page/35180469#page/541/>
- von Martens E (1895) Die gattung *Cylindrus* Fitz. *Archiv für Naturgeschichte* 61(1): 103–108. <https://www.biodiversitylibrary.org/item/47791>
- von Möllendorff OF (1891) On the land and freshwater shells of Perak. *Proceedings of the Zoological Society of London* 59(3): 330–348. <https://doi.org/10.1111/j.1096-3642.1891.tb01757.x>
- von Möllendorff OF (1894) On a collection of land-shells from the Samui Islands, Gulf of Siam. *Proceedings of the Zoological Society of London* 62: 146–156. <https://biodiversitylibrary.org/page/35990426>
- von Möllendorff OF (1899) Neue arten aus Hinterindien. *Nachrichtenblatt der Deutschen Malakozoologischen Gesellschaft* 31: 165, 166. <https://biodiversitylibrary.org/page/28228787>
- Wade CM, Mordan PB, Clarke B (2001) A phylogeny of the land snails (Gastropoda: Pulmonata). *Proceedings. Biological Sciences* 268(1465): 413–422. <https://doi.org/10.1098/rspb.2000.1372>
- Wade CM, Mordan PB, Naggs F (2006) Evolutionary relationships among the pulmonate land snails and slugs (Pulmonata, Stylommatophora). *Biological Journal of the Linnean Society* 87(4): 593–610. <https://doi.org/10.1111/j.1095-8312.2006.00596.x>
- Westerlund CA (1887) Fauna der in der paläarktischen Region, lebenden Binnenconchylien. III. Gen. *Buliminus*, *Sesteria*, *Pupa*, *Stenogyra* & *Cionella*. Lund, Hakan Ohlsson, 183 pp. <https://doi.org/10.5962/bhl.title.10301>
- Yu T-T, Páll-Gergely B, Salvador RB (2023) Caenogastropoda and Stylommatophora (Gastropoda) from the mid-Miocene Zhangpu amber of East Asia. *Palaeoworld* (Amsterdam) 32(4): 721–728. <https://doi.org/10.1016/j.palwor.2022.07.004>
- Zilch A (1959) Gastropoda, Euthyneura. In: Schindewolf OH (Ed) *Handbuch der Paläozoologie*, Band 6, Gastropoda. Gebrüder Borntraeger, Berlin, 201–400.
- Zilch A (1973) Die Typen und Typoide des Natur-Museums Senckenberg, 51: Mollusca: Achatinacea (2): Ferrussaciidae, Subulinidae. *Archiv für Molluskenkunde* 103: 99–152.