



**Rediscovery of *Pagodina bourguignati* Coutagne, 1881 in
France: an old and forgotten name for
Truncatellina beckmanni Quintana Cardona, 2010
(Gastropoda: Truncatellinidae)**

Louis Aureglia, Marin Marmier, Cédric Audibert, Daniel Pavon

► **To cite this version:**

Louis Aureglia, Marin Marmier, Cédric Audibert, Daniel Pavon. Rediscovery of *Pagodina bourguignati* Coutagne, 1881 in France: an old and forgotten name for *Truncatellina beckmanni* Quintana Cardona, 2010 (Gastropoda: Truncatellinidae). *Folia Malacologica*, 2026, 34 (3), pp.188-195. <10.12657/folmal.034.016>. <hal-05710687>

HAL Id: hal-05710687

<https://hal.science/hal-05710687v1>

Submitted on 4 Aug 2026

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

REDISCOVERY OF *PAGODINA BOURGUIGNATI* COUTAGNE, 1881 IN FRANCE: AN OLD AND FORGOTTEN NAME FOR *TRUNCATELLINA BECKMANNI* QUINTANA CARDONA, 2010 (GASTROPODA: TRUNCATELLINIDAE)

LOUIS AUREGLIA^{1*}, MARIN MARMIER², CÉDRIC AUDIBERT³, DANIEL PAVON⁴

¹ Aix Marseille Univ, IRD, LPED, 3 Place Victor Hugo, 13331 Marseille Cedex 03, France
(e-mail: louis.aureglia@gmail.com);  <https://orcid.org/0000-0001-9129-5016>

² Pôle Biodiversité régionale, Conservatoire d'espaces naturels de Provence-Alpes-Côte d'Azur, France

³ Centre "Louis Lortet" de conservation et d'étude des collections, Musée des Confluences, France

⁴ IMBE, Aix-Marseille Université, France;  <https://orcid.org/0009-0005-8126-8280>

* corresponding author

ABSTRACT: The taxonomic identity of *Pagodina bourguignati* Coutagne, 1881, described from the Vallon de Rognac in southern France, has long remained uncertain. The name was historically misunderstood, variously treated as a juvenile *Pagodulina*, an uncertain species, or a synonym of other taxa, but it was never critically reassessed based on the original specimen and associated archival evidence. Re-examination of the original material preserved in the Coutagne collection at the Musée des Confluences, together with archival documents associated with Coutagne's work, shows that the nominal taxon was based on a single specimen. That specimen, MDC.45030705, is therefore the holotype by monotypy. Its shell morphology, dimensions, and diagnostic apertural barriers are consistent with *Truncatellina beckmanni* Quintana Cardona, 2010, described from Menorca and recently reported from southern France. We therefore transfer Coutagne's species to *Truncatellina* and re-establish it as a valid species under the combination *Truncatellina bourguignati* (Coutagne, 1881), comb. nov., stat. rev. We further place *Truncatellina beckmanni* Quintana Cardona, 2010 in synonymy as a junior subjective synonym, syn. nov. Litter sampling conducted in 2026 confirms that this species still occurs in the Vallon de Rognac, its type locality, now known as Vabre de Malaga. We also propose the French vernacular name Maillotin trapu, translated here as "stout maillotin", in reference to the compact shell shape of the species.

KEYWORDS: Gastropoda; junior synonymy; nomenclature; Provence; Rognac; type material

INTRODUCTION

The identity of *Pagodina bourguignati* Coutagne, 1881 has remained unresolved since its original description from the Vallon de Rognac in the Bouches-du-Rhône, south-eastern France. After its publication, the name was initially accepted as a distinct taxon and was repeated in the French literature by [LOCARD \(1882, 1894, 1896\)](#) and later by [MARGIER \(1910\)](#). However, its status soon became unstable. [WESTERLUND \(1887\)](#) treated *P. bourguignati* as mere-

ly a juvenile of *Pagodulina pagodula*, thereby denying it specific validity. This interpretation was subsequently followed by [GERMAIN \(1913\)](#), who explicitly placed the taxon in synonymy with *P. pagodula*. The question nevertheless remained open, because later authors did not accept this interpretation. [TRYON & PILSBRY \(1926\)](#) expressly rejected [WESTERLUND'S \(1887\)](#) view and stated that the shell could not be a young *P. pagodula*, suggesting instead that it might



represent a vertiginid species. GERMAIN (1930) likewise reconsidered the taxon: although he retained it only with doubt, he stated that it had never been re-discovered, that it was certainly not a juvenile *P. pagodula*, and that it probably did not belong to *Pagodulina*, perhaps corresponding instead to an undetermined form of *Vertigo*. PILSBRY (1934) further suggested that “*Pagodina*” *bourguignati* might in fact be related to *Truncatellina*, and KLEMM (1939), after direct examination of a specimen, stated unambiguously that it had nothing to do with *Pagodulina* and should be removed from that genus as a *Truncatellina*. Despite these successive reinterpretations, no stable conclusion followed, and the name was never restored to practical use as a valid species. In later French checklists, the taxon was no longer maintained as distinct but was instead absorbed into the synonymy of *Pagodulina austeniana* by PAVON & BERTRAND (2005) and in TAXREF (GARGOMINY 2022), whereas MOLLUSCABASE (2026) still retained it under an unresolved status (“uncertain > unassessed”). In this sense, the taxon was not simply overlooked; rather, it remained historically unsettled, repeatedly cited, reinterpreted, or synonymised, but never critically reassessed based on the original specimen and associated archival evidence.

The taxonomic problem became more explicit after *Truncatellina beckmanni* Quintana Cardona, 2010, originally described from Menorca, was reported from the Bouches-du-Rhône as new to the French fauna by DOMMERGUES & GARGOMINY (2024). This recent French record concerns the same regional

context as COUTAGNE’s (1881) forgotten species and raises the question of whether *P. bourguignati* and *T. beckmanni* represent distinct taxa or the same species under different names. The comparison of the shell illustrated by QUINTANA CARDONA (2010), the extant specimen preserved in the Coutagne collection, and the newly collected topotypic material from Rognac provides a direct way to test this hypothesis.

More broadly, taxonomic work on *Truncatellina* remains largely fragmented and regional. Important contributions have addressed particular insular or regional faunas, for example in Macaronesia (HUTTERER & GROH 1991), Croatia (ŠTAMOL 1995), Greece (TRIANTIS et al. 2004), the Iberian Peninsula and north-west Africa (HOLYOAK et al. 2012), and more recently through critical morphometric reassessments of problematic taxa (KNEUBÜHLER et al. 2020) or local molecular re-evaluations (CUCHERAT & GUILLER 2021). In that context, the clarification of overlooked historical names remains necessary for nomenclatural stability and for accurate faunistic interpretation.

Accordingly, this paper re-examines the original material and archival evidence relating to *Pagodina bourguignati* Coutagne, 1881, documents newly collected topotypic material from its type locality, provides a morphometric comparison with *T. beckmanni* Quintana Cardona, 2010 and *T. callicratis* (Scacchi, 1833), determines the status of the type specimen, and evaluates whether *P. bourguignati* and *T. beckmanni* refer to the same species.

MATERIAL AND METHODS

The original material preserved in the Coutagne collection at the Musée des Confluences (Lyon) was examined from photographs and documentation supplied by the collection. The relevant specimen is catalogued as MDC.45030705 (Coutagne no. 834), labelled “Rognac, alluvions”, and attributable to Coutagne’s original gathering.

Archival documents associated with G. Coutagne’s collection were consulted, especially the collection notebooks 6J001 and 6J002, the letter dated 18 January 1881 to Bourguignat (6J053), and Coutagne’s manuscript drawing (6J016). These documents were used to assess whether Coutagne had one or several shells available, to verify the correspondence between the extant specimen and the published description, and to compare the archival dimensions with those given in the original description.

On 9 February 2026, an opportunistic field survey was conducted to confirm the presence of the species

described by COUTAGNE (1881) as *Pagodina bourguignati* and litter was collected in the Vabre de Malaga, referring to the locality described by Coutagne as Vallon de Rognac, municipality of Rognac, near the Étang de Berre (Bouches-du-Rhône, Provence, south-eastern France). The sample was processed using a Winkler extractor in the field. Approximately 1 L of litter was collected from sheltered litter accumulations beneath vegetation. It was then dried at 37 °C for 24 h, sieved through a column of mesh sizes, and manually sorted under a stereomicroscope. Fourteen shells referable to the species treated here were recovered from this sample. No visibly living specimens were obtained, and therefore no molecular or anatomical analyses were attempted. A simple localisation map was prepared to show the position of the two French localities discussed in this study: Vabre de Malaga, Rognac, and the Parc national des Calanques locality reported by DOMMERGUES & GARGOMINY (2024) (Fig. 1).




Fig. 1. Localisation of the French records of *Truncatellina bourguignati* (Coutagne, 1881) discussed in this study. (A) Position of the Bouches-du-Rhône department in southern France. (B) Detail of the Bouches-du-Rhône showing Vabre de Malaga, Rognac, corresponding to the type locality and newly sampled topotypic material, open circle, and the Parc national des Calanques locality reported as *T. beckmanni* by DOMMERGUES & GARGOMINY (2024), filled circle.

Morphometric comparison was based on shell length and shell width (Fig. 2). Measurements of the




Fig. 2. Morphometric comparison of shell length and shell width in *Truncatellina beckmanni*, *T. bourguignati* and *T. callicratis*. Shell length and shell width are expressed in millimetres. Circles correspond to specimens assigned to *T. beckmanni*, squares to *T. bourguignati*, and triangles to *T. callicratis*. Open black symbols indicate French specimens measured for the present study, whereas filled black symbols indicate comparative specimens taken from the literature. The red circle indicates the holotype of *T. beckmanni* Quintana Cardona, 2010, and the blue square indicates the holotype of *T. bourguignati* (Coutagne, 1881), based on Coutagne's manuscript measurement

French material were taken from calibrated photographs using ImageJ 1.54p. Shell length was measured as the maximum apical–basal dimension of the shell, and shell width as the maximum transverse diameter. These measurements were combined with the dimensions of the holotype of *P. bourguignati* recorded in Coutagne's manuscript note, the holotype of *T. beckmanni* given by QUINTANA CARDONA (2010), and individual published measurements of *T. beckmanni* and *T. callicratis* from HOLYOAK et al. (2012). The two specimens illustrated from the Calanques by DOMMERGUES & GARGOMINY (2024) were also included in the comparison. Population means, ranges and standard deviations not assignable to individual specimens were excluded from Fig. 2.

RESULTS

Recent litter sampling in the Vabre de Malaga (13340 Rognac, Bouches-du-Rhône; 43.48235, 5.24511) yielded 14 shells referable to the species treated here (Fig. 1). These shells match the morphology of the historical specimen preserved in the Coutagne collection and of the taxon here identified as *Truncatellina bourguignati* (Figs 3–10). The species is therefore confirmed to be still extant in its type locality.

Among the measured French shells, shell length ranged from 1.42 to 1.65 mm (mean 1.54 mm, SD = 0.06, n = 14) and shell width ranged from 0.82 to 0.97 mm (mean 0.90 mm, SD = 0.04, n = 14). The shell of the holotype of *P. bourguignati* measures 1.43 × 0.95 mm according to Coutagne's manuscript note. The shell length/width comparison places the French material attributed here to *T. bourguignati* within, or close to, the morphometric space occupied by published specimens of *T. beckmanni* (Fig. 2). The



holotype of *T. bourguignati* plots among the compact shells assigned to *T. beckmanni*. By contrast, the comparative *T. callicratis* specimens tend to show a more elongate shell for a similar or smaller shell width.

SYSTEMATICS

***Truncatellina bourguignati* (Coutagne, 1881), comb. nov., stat. rev. = *Truncatellina beckmanni* Quintana Cardona, 2010, junior subjective synonym, syn. nov.**

Vernacular name. French name proposed: Maillotin trapu; English translation: “stout maillotin”. The adjective “trapu” refers to the short, broad and compact shell shape of the species compared with the more elongate shell of several other regional *Truncatellina*.

Protonym. *Pagodina bourguignati* COUTAGNE, 1881: 39–40.

Type locality. “Le vallon de Rognac”, Bouches-du-Rhône, southern France.

Type material. Holotype (by monotypy): France, Bouches-du-Rhône, Vallon de Rognac, “Rognac, alluvions”, collected between July and December 1880, G. Coutagne leg., one shell, Musée des Confluences, Coll. Coutagne, MDC.45030705 (Coutagne no. 834).

Additional material examined. France, Bouches-du-Rhône, Vabre de Malaga, 13340 Rognac, 43.48235, 5.24511, litter sample, 9 February 2026, L. AUREGLIA leg., collection no. LA1607, 14 shells; France, Bouches-du-Rhône, Parc national des Calanques, Vallon de la Gineste, 13009 Marseille, 43.2375, 5.48098, 26 September 2025, M. MARMIER leg., 10 shells, one illustrated, shell height 1.5 mm, shell width 0.9 mm.

Type material of the junior synonym (from the original description). Holotype of *Truncatellina beckmanni* Quintana Cardona, 2010: shell deposited in the malacological collection of the Museu Diocesà de Menorca, registration number MDM-2703, from Lloc de Monges (Ciutadella de Menorca, Illes Balears); shell height 1.29 mm, shell width 0.79 mm.

Diagnosis. A very small, compact, strongly ribbed dentate species of *Truncatellina*, characterised by a short and broad shell, 5–6 strongly convex whorls, a deep suture, regular axial ribbing, a very narrow umbilicus, and a well-developed apertural dentition, notably with conspicuous parietal and palatal lamellae visible in frontal view. The species differs from *T. callicratis* by its more compact shell, lower height-to-diameter ratio, more strongly developed lamellae, especially the parietal and palatal lamellae, and more pronounced, regular ribbing. It differs

This descriptive pattern supports the interpretation that the French topotypic material, the Coutagne holotype and the published concept of *T. beckmanni* refer to the same compact morphotype.

from *Pagodulina* species historically associated with the name *P. bourguignati* by its truncatellinid shell shape and apertural barrier pattern.

Synonymy and nomenclatural history.

Isthmia – COUTAGNE, in litteris, 1881.

Pagodina bourguignati COUTAGNE, 1881: 30, 39–40, 51; original description and mention of the type locality.

Pagodina bourguignati – LOCARD 1882: 172.

Pupa pagodula – WESTERLUND 1887: 91, misidentification as a juvenile *Pagodulina pagodula*.

Pagodina bourguignati – LOCARD 1894: 327.

Pagodina bourguignati – LOCARD 1896: 215.

Pagodina bourguignati – MARGIER 1910: 44.

Pagodina pagodula – GERMAIN 1913: 183, placed in synonymy.

Pagodulina (?) *bourguignati* – TRYON & PILSBRY 1926: 179, considered possibly a vertiginid and explicitly not a juvenile *P. pagodula*.

Pagodulina pagodula ? var. *bourguignati* – GERMAIN 1930: 421, retained with doubt and considered possibly closer to *Vertigo*.

“*Pagodina*” *bourguignati* – PILSBRY 1934: 108, considered possibly referable to *Truncatellina*.

Truncatellina – KLEMM 1939: 256, explicitly removed from *Pagodulina*.


Pagodulina austeniana austeniana – PAVON & BERTRAND 2005: 39, treated as a synonym.

Pagodulina austeniana austeniana – GARGOMINY 2022, treated as a synonym in TaxRef.

Truncatellina beckmanni QUINTANA CARDONA 2010: 149–158, described from Menorca.

Truncatellina beckmanni – DOMMERGUES & GARGOMINY 2024: 1–3, reported from the Bouches-du-Rhône as new for France.

Description. Shell light brown to yellowish, relatively short and broad, giving it a compact appearance (Figs 3–10). It consists of 5–6 fairly rounded whorls, with a marked, deep suture. The aperture is subquadrangular, with a rounded lower part and provided with three well-developed lamellae: palatal, columellar, and parietal. Only the characteristic palatal and parietal lamellae are visible in frontal view, the columellar lamella ending before the peristome. The white palatal lamella is well-developed inside the shell and also visible in dorsal view. The peristome is discontinuous in the parietal region and reflect-



Figs 3–10. Composite plate illustrating *Truncatellina bourguignati* (Coutagne, 1881): 3–4 – holotype by monotypy, MDC.45030705, Musée des Confluences, Coutagne collection, Vallon de Rognac; 5–6 – archival drawing attributable to G. Coutagne (MS 6J016), corresponding to the holotype; 7–8 – shell from the population of the Parc national des Calanques; 9 – in situ photograph of a shell from the same Calanques population; 10 – shell from Vabre de Malaga, Rognac. The scale bar equals 1 mm and applies to Figs 3, 4, 7, 8 and 10 only

ed. The umbilicus is very narrow and not covered by the columellar margin. The shell surface bears thick, regular growth lines forming a characteristic ribbing. **Dimensions.** Measured French material: shell height 1.54 mm, range 1.42–1.65 mm, $n = 14$; shell width 0.90 mm, range 0.82–0.97 mm, $n = 14$. Holotype of *P. bourguignati*: 1.43×0.95 mm according to Coutagne's manuscript note, consistent with the dimensions given in the original description, 1.4×0.95 mm.

Habitat and ecology. The species appears to be associated with Mediterranean calcareous habitats, especially rocky or karstic slopes with limestone outcrops, scree, and accumulations of litter in sheltered microhabitats. At Vabre de Malaga, it was recovered from litter beneath a vegetation mosaic dominated by *Viburnum tinus*, *Pinus halepensis*, *Smilax aspera*, *Ruscus aculeatus*, and a few *Quercus ilex*. In the Calanques, it was found in a thermo-xeric shrubland on limestone with abundant stones and rock outcrops, in vegetation including *Quercus coccifera*, *Salvia rosmarinus*, *Smilax aspera*, and *Asparagus acutifolius*. The type locality in Menorca is likewise situated on calcareous substrate and was described as a small woodland with oleaster, juniper, and mastic. French habitats are illustrated in Figs 11–12.

Distribution. Western Mediterranean, with a still insufficiently documented distribution. Under the

present taxonomic interpretation, the species is known from Menorca, the southern Iberian Peninsula, and southern France. Published records attributable to this taxon exist from the Spanish provinces of Córdoba and Málaga, and from Portugal in Algarve, Estremadura, Ribatejo, and Beira Litoral. In France, it is currently known only from the Bouches-du-Rhône, in the Parc national des Calanques and at Vabre de Malaga, Rognac.

Remarks. Although the original description does not explicitly state the number of specimens examined, the archival documentation associated with G. Coutagne suggests that only a single shell was available to him. His collection notebooks (6J001 and 6J002) refer to only one specimen. In a letter dated 18 January 1881 (6J053) addressed to Bourguignat, Coutagne transmitted the specimen under the name *Isthmia*, which already indicates that he had recognised its truncatellinid affinities before publishing it in *Pagodina*. A manuscript drawing by Coutagne (6J016) matches the extant shell and gives dimensions of 1.43×0.95 mm (Figs 5–6). The specimen must therefore be treated as the holotype by monotypy, not as a syntype or lectotype.

The extant type specimen agrees closely with the shell morphology of *Truncatellina beckmanni* Quintana Cardona, 2010 and later documented by HOLYOAK



Figs 11–12. Habitats of *Truncatellina bourguignati* (Coutagne, 1881) in southern France: 11 – habitat at Vabre de Malaga, Rognac, corresponding to the type locality; 12 – habitat in the Parc national des Calanques, at the site where the species was collected and reported by DOMMERGUES & GARGOMINY (2024)

et al. (2012), especially in its short and robust shell, its strong ribbing, its low height relative to diameter, and its marked apertural armature. The dimensions of the Coutagne shell (1.4×0.95 mm in the original description; 1.43×0.95 mm in the manuscript note) fall within the morphometric space occupied by specimens assigned to *T. beckmanni* in the present comparison (Fig. 2). Recent topotypic material collected in 2026 at Vabre de Malaga, corresponding to Coutagne's historical Vallon de Rognac, matches

the same taxon and confirms that the species is still extant at its type locality.

Because *Pagodina bourguignati* Coutagne, 1881 is the oldest available name applicable to this species, it has priority over *Truncatellina beckmanni* Quintana Cardona, 2010. The valid name is therefore *Truncatellina bourguignati* (Coutagne, 1881), and *T. beckmanni* is here treated as its junior subjective synonym, syn. nov.

DISCUSSION

This study resolves a long-standing taxonomic problem concerning a minute Mediterranean truncatellinid that had remained difficult to interpret for more than a century. Its main result is not the description of a new taxon, but the reassessment of an old available name through the combined use of historical material, archival documentation, newly collected topotypic shells and morphometric comparison with related material. The present results provide a direct link between the forgotten name *Pagodina bourguignati* Coutagne, 1881 and the species later described as *Truncatellina beckmanni* Quintana Cardona, 2010. The combined evidence from the extant historical specimen, Coutagne's archival documentation, the manuscript drawing, and the newly collected topotypic material from the Vabre de Malaga (Vallon de Rognac) shows that a single, morphologically coherent taxon is involved. In particular, the archival documents remove any ambiguity concerning the type material by showing that Coutagne had only one shell at his disposal. The specimen preserved in the Musée des Confluences must therefore be regarded as the holotype by monotypy. Once this historical material is compared with the original concept of *T. beckmanni*

as defined by QUINTANA CARDONA (2010), the additional Iberian material documented by HOLYOAK et al. (2012), and the recent French material reported by DOMMERGUES & GARGOMINY (2024), the treatment of *T. beckmanni* as a junior subjective synonym of *T. bourguignati* becomes the most parsimonious interpretation. The name introduced in 2010 therefore applies to the same species and must be treated as a junior subjective synonym of the senior name *Truncatellina bourguignati* (COUTAGNE 1881).

This conclusion is necessarily based on shell morphology. No molecular or anatomical analyses were conducted because the newly collected material consisted only of empty shells, with no visibly living specimens recovered from the litter sample. This is a limitation, especially in a genus where shell characters may show convergence or limited diagnostic resolution, but no shell character currently allows the French topotypic material, the Coutagne holotype and the published concept of *T. beckmanni* to be separated. This nomenclatural clarification has immediate consequences for the taxonomy and faunistics of the French malacofauna. First, it restores an overlooked nineteenth-century name to practical



use and stabilises the nomenclature of this minute Mediterranean truncatellinid. Second, it shows that the French record published as *Truncatellina beckmanni* by DOMMERGUES & GARGOMINY (2024) should now be referred to *T. bourguignati* (Fig. 1). The rediscovery of the species in 2026 in the Vabre de Malaga (Vallon de Rognac) is also noteworthy, as it confirms that the species is still extant in its type locality more than a century after its original description. The present paper is intentionally limited to this taxonomic and nomenclatural point. A broader revision of western Mediterranean *Truncatellina*, ideally combining detailed conchological comparison with molecular sampling, would nevertheless be valuable, especially given the recurrent problems of shell-based over-splitting and hidden synonymy documented in the genus (HOLYOAK et al. 2012, KNEUBÜHLER et al.

2020, CUCHERAT & GUILLER 2021). Such a broader framework, however, is not required to establish the identity and priority of Coutagne's name in the present case.

ACKNOWLEDGEMENTS

We thank MAËLAN ADAM, KASSANDRA OGNO, ALAIN BERTRAND, and FRÉDÉRIC MAGNIN for helpful discussions concerning these specimens and the taxonomy of *Truncatellina* and *Pagodulina* more generally. We are also grateful to LAURENT GRIMA for kindly responding to our requests for consultation of the malacological collections of the Muséum d'histoire naturelle de Marseille. Finally, we thank the Parc national des Calanques for their support and LAUREEN KELLER for her help during the field surveys.

REFERENCES

- COUTAGNE G. 1881. Notes sur la faune malacologique du bassin du Rhône. Annales de la Société Linnéenne de Lyon 28: 1–55.
- CUCHERAT X., GUILLER A. 2021. Étude taxonomique du Maillotin de Bourgogne *Truncatellina arcyensis* Klemm, 1943. Conservatoire d'espaces naturels de Bourgogne, Gondécourt, France.
- DOMMERGUES J.-L., GARGOMINY O. 2024. *Truncatellina beckmanni*, nouvelle espèce pour la France observée dans les Calanques. MalaCo 20: 1–3.
- GARGOMINY O. 2022. TAXREF Version 4.17. UAR PatriNat (OFB-MNHN-CNRS-IRD), Paris.
<https://doi.org/10.15468/vqueam>
- GERMAIN L. 1913. Mollusques de la France et des régions voisines. Tome 2, Gastéropodes pulmonés et prosobranches terrestres et fluviatiles. Encyclopédie scientifique. Bibliothèque de zoologie. Vol. 2. O. Doin et fils, Paris.
- GERMAIN L. 1930. Mollusques terrestres et fluviatiles (première partie). Faune de France. Paris.
- HOLYOAK D. T., HOLYOAK G. A., TORRES ALBA J. S. 2012. A reassessment of the species of *Truncatellina* (Gastropoda: Vertiginidae) in the Iberian Peninsula and North-west Africa. Iberus 30(2): 7–33.
- HUTTERER R., GROH K. 1991. A review of Macaronesian *Truncatellina* (Gastropoda: Vertiginidae) with descriptions of four new species. Bocagiana (151): 1–19.
- KLEMM W. 1939. Zur rassenmäßigen Gliederung des Genus *Pagodulina* Clessin. Archiv für Naturgeschichte 8(2): 198–262.
- KNEUBÜHLER J., BAUR H., NEUBERT E. 2020. Deception in taxonomy – the example of *Truncatellina rothi* (Hesse, 1916) (Pupilloidea: Truncatellinidae). Basteria 84(4–6): 146–154.
- LOCARD A. 1882. Catalogue général des mollusques vivants de France: mollusques terrestres, des eaux douces et des eaux saumâtres. Prodrome de Malacologie Française. Vol. 1. H. Georg, Lyon.
<https://doi.org/10.5962/bhl.title.10669>
- LOCARD A. 1894. Les coquilles terrestres de France: description des familles, genres et espèces. Librairie J.-B. Baillièrre et fils, Paris.
- LOCARD A. 1896. Les coquilles terrestres de France. Description des familles, genres et espèces. Annales de la Société d'agriculture, sciences et industrie de Lyon 3: 5–258.
- MARGIER E. 1910. Mollusques de la famille des Pupidae (ancien genre Pupa-Drap.) qui vivent dans le Gard et les départements limitrophes. Bulletin de la Société d'étude des sciences naturelles de Nîmes 38: 43–48.
- MOLLUSCABASE eds. 2026. MolluscaBase. *Truncatellina* R. T. Lowe, 1852. Accessed at: <https://www.molluscabase.org/aphia.php?p=taxdetails&id=426420> on 10 March, 2026.
- PAVON D., BERTRAND A. 2005. Liste commentée des mollusques continentaux du département des Bouches-du-Rhône. Bulletin de la Société Linnéenne de Provence (56): 35–47.
- PILSBRY H. A. 1934. Geographic distribution of Pupillidae; Strobilopsidae, Valloniidae and Pleurodiscidae. Manual of Conchology, Structural and Systematic: With Illustrations of the Species. Second Series, Pulmonata. Vol. 28. Conchological Department, Academy of Natural Sciences of Philadelphia, Philadelphia.
- QUINTANA CARDONA J. 2010. *Truncatellina beckmanni* sp. nov. (Gastropoda: Pulmonata: Vertiginidae: Truncatellininae), una nueva especie endémica de Menorca (Islas Baleares, Mediterráneo occidental). Spira 3(3–4): 149–158.
- SCACCHI A. 1833. Osservazioni Zoologiche. Napoli, Tipi della Società Tipografica 1: 1–12 [February]; 2: 13–27 [May].
<http://biodiversitylibrary.org/page/35752103>



- ŠTAMOL V. 1995. *Truncatellina velkovrhi* n. sp. and *Truncatellina lussinensis* n. sp., two new species from Croatia (Gastropoda: Pulmonata: Vertiginidae). Archiv für Molluskenkunde 124(1–2): 97–101.
<https://doi.org/10.1127/arch.moll/124/1995/97>
- TRIANIS K. A., POKRYSZKO B. M., MYLONAS M., VARDINOYANNIS K. 2004. A new species of *Truncatellina* (Gastropoda: Vertiginidae) from Mount Ossa (=Kissavos) (Greece). Journal of Conchology 38(4): 393–397.
<https://doi.org/10.5962/p.408155>
- TRYON G. W. JR., PILSBRY H. A. 1926. Manual of conchology, structural and systematic: with illustrations of the species. Second Series, Pulmonata. Vol. 27. Philadelphia.
- WESTERLUND C. A. 1887. Fauna der in der paläarktischen Region (Europa, Kaukasien, Sibirien, Turan, Persien, Kurdistan, Armenien, Mesopotamien, Kleinasien, Syrien, Arabien, Egypten, Tripolis, Tunesien, Algerien und Marocco) lebenden Binnenconchylien. III. Gen. *Buliminus*, *Sesteria*, *Pupa*, *Stenogyra* & *Cionella*. Håkan Ohlsson, Lund.

Received: March 16th, 2026

Revised: May 20th, 2026

Accepted: May 21st, 2026

Published on-line: August 4th, 2026

