

Annotated checklist of the umagillid turbellarians infesting echinoids (Echinodermata)

Gilles Doignon¹ and Tom Artois²

¹ Laboratoire de Biologie Marine (CP 160/15), Université Libre de Bruxelles, 50 avenue F. D. Roosevelt, B-1050 Bruxelles, Belgium

² Hasselt University, Centre for Environmental Sciences, Research Group Biodiversity, Phylogeny & Population Studies, Agoralaan Gebouw D, B-3590 Diepenbeek, Belgium

Corresponding author : G. Doignon, e-mail : gdoignon@gmail.com

ABSTRACT. A literature survey of the taxonomically complex taxon *Syndesmis* Silliman, 1881 is presented, resulting in the recognition of 22 valid species and three species mentioned in several works but not yet described. For each of these species the host(s) are mentioned and the distribution is given. These results are compared with earlier checklists and the differences are discussed.

KEY WORDS : Biogeography, *Marcusella*, symbiosis, *Syndesmis*, *Syndisyrix*, Turbellaria.

INTRODUCTION

The family Umagillidae Wahl, 1910 (Platyhelminthes, Rhabdocoela) is the largest taxon of symbiotic turbellarians. It includes 68 valid species (WESTERVELT, 1981; CANNON, 1982, 1987; SHINN, 1987; CANNON, 1990; HERTEL et al., 1990; JANGOUX, 1990; KOZLOFF & WESTERVELT, 1990; WESTERVELT & KOZLOFF, 1990, 1992; MOENS et al., 1994; JONDELIUS, 1996; KOZLOFF, 1997; VASS & NAPPI, 1998). Most umagillids are common endosymbionts of echinoderms, particularly echinoids and holothuroids, and a few occur in sipunculids (see CANNON, 1982; JANGOUX, 1990). Our knowledge of the biology of umagillids is, however, very scant. For instance, although they are frequently found in the digestive tract or coelomic cavity of their hosts, almost nothing is known about the nature of their symbiotic relationships. The life cycle itself is completely elucidated for only four species : *Anopodium hymanae* Shinn, 1983 and *Wahlia pulchella* Kozloff & Shinn, 1987 (both infesting holothuroids) (SHINN, 1983a, 1985a, b, 1986a), *Fallacohospes inchoatus* Kozloff, 1965 (infesting crinoids) (SHINN, 1986b) and *Syndesmis franciscana* (Lehman, 1946) (infesting echinoids) (SHINN, 1980, 1981, 1983b). Most umagillid species are host-specific, though a few apparently infest several, often phylogenetically unrelated, echinoderm species (see CANNON, 1982; SHINN, 1984; JANGOUX, 1990).

Research on the biology of these worms is greatly hampered by the complex taxonomy of the group, which often leads to wrong assumptions. For instance, for almost sixty years all umagillids found in echinoids were attributed to *Syndesmis echinorum* François, 1886 (see VON GRAFF, 1903). Although already very early on, it was suggested that more species existed (see BRAUN, 1889), it was not until recently that *S. echinorum* was split into six valid species (see KOZLOFF & WESTERVELT, 1987, 1990; WESTERVELT & KOZLOFF, 1990, 1992; KOZLOFF, 1997). After

the description of the second recognized species of *Syndesmis*, *S. franciscana* (Lehman, 1946), many more species were described and some redescribed. Yet, authors often did not agree on the taxonomic importance of many characters. For instance, the validity of the genera *Syndisyrix* Lehman, 1946 and *Marcusella* Westblad, 1953 has been the subject of many impassioned debates (LEHMAN, 1946; MARCUS, 1949; STUNKARD & CORLISS, 1951; WESTBLAD, 1953; HYMAN, 1960; CANNON, 1982, 1987; KOZLOFF & WESTERVELT, 1987, 1990; HERTEL et al., 1990; WESTERVELT & KOZLOFF, 1990, 1992). In the most recent literature, all umagillid species infesting echinoids were brought together into the genus *Syndesmis* Silliman, 1881 (see CANNON, 1982; MOENS et al., 1994; GEVAERTS et al., 1995; JONDELIUS, 1996).

Taxonomic literature regarding the species of *Syndesmis* has become extremely entangling, and is almost inaccessible for novice and even experienced researchers who are not particularly interested in taxonomy. As a result, several attempts were made to review the extensive literature on these animals : STUNKARD & CORLISS (1951), HICKMAN (1956), JENNINGS (1971), BAREL & KRAMERS (1977), CANNON (1982), HERTEL et al. (1990), JANGOUX (1990), VASS & NAPPI (1998). Yet, in recent years, several works (KOZLOFF & WESTERVELT, 1987, 1990; HERTEL et al., 1990; WESTERVELT & KOZLOFF, 1990, 1992; KOZLOFF, 1997) showed that many specimens previously attributed to a given species, in fact belong to several distinct species and, consequently, some species are less opportunistic than previously thought, thereby making most earlier checklists outdated. Moreover, because of confusing synonymies, discrepancies in the spelling of species names, and because of an incomplete literature survey or bad translations of non-English literature, these reviews contain mistakes and contradictory information.

The new checklist we present here is intended to give an overview of all umagillid species infesting echinoids, based on a complete literature survey. As the names of

many species changed through time, because these were synonymized, misidentified or even misspelled, it is quite difficult to understand the literature in a way other than chronologically. Only a detailed reappraisal of all descriptions allows to attribute the most recently accepted name to species misidentified in older works, for example the "*S. echinorum*" of BAREL & KRAMERS (1970) that was later described as *S. albida* Kozloff & Westervelt, 1990 and *S. rubida* Kozloff & Westervelt, 1990. In this work, we follow the most modern views, e.g. the one that considers *Syndisyrix* a synonym of *Syndesmis* (MOENS et al., 1994; GEVAERTS et al., 1995; JONDELIUS, 1996). Similarly, we use the most recently published taxonomy of the echinoids (SMITH, 2003).

RESULTS AND DISCUSSION

To date, the 22 valid and three undescribed species of *Syndesmis* are known to infest a total of 31 species of echinoids belonging to the orders Clypeasteroidea and Spatangoidea (irregular sea urchins), Diadematoidea, Echinoidea and Temnopleuroidea (regular sea urchins) (Table 1). The taxon *Syndesmis* occurs in European waters (from the Mediterranean Sea to the Barents Sea), the Caribbean, the entire Pacific Ocean and the Indian Ocean. Some echinoid species (e.g., *Echinometra oblonga* (de Blainville, 1825)) may host up to three umagillid species and some species of *Syndesmis* are found in various (up to five) echinoid hosts. Interestingly, some species are found in hosts belonging to distinct families or even orders. For instance, *S. echinorum* is found in *Paracentrotus lividus* (Lamarck, 1816) (Echinidae) as well as *Sphaerechinus granularis* (Lamarck, 1816) (Toxopneustidae).

The superscript numbers following the species names in our checklist (Table 1) refer to major discrepancies with previous checklists, the details of which are presented here :

(1) : Based on a literature survey, JANGOUX (1990) mentioned the occurrence of *S. echinorum* in *Psammechinus microtuberculatus* (de Blainville, 1825). However, only VON GRAFF (1903) has studied this species of sea urchin (from Trieste, Italy) and he reported that all 30 specimens he inspected were uninfested. Apart from the records presented in Table 1, MEIXNER (1926) mentioned the occurrence of *S. echinorum* in sea urchins from Lessin Island (Croatia), but without specifying the host.

(2) : VON GRAFF (1903) mentioned the occurrence of *Syndesmis echinorum* in *Strongylocentrotus droebachiensis*

(Müller, 1776). This is impossible to verify owing to the lack of material. However, the fact that this species of echinoids is known to host two other morphologically similar and well documented species of *Syndesmis*, *S. franciscana* and *S. inconspicua* Westervelt & Kozloff, 1992 (see SHINN, 1984; WESTERVELT & KOZLOFF, 1992), suggests that the species VON GRAFF (1903) observed was probably not *S. echinorum*. VON GRAFF (1903) also reported *S. echinorum* in *Sphaerechinus granularis* from Bergen (Norway), but as this species of sea urchins is limited from the Mediterranean to the south of the English Channel, this observation is clearly erroneous. He probably misidentified the echinoid species, which could be *Echinus esculentus* Linnaeus, 1758, which harbours two distinct umagillid species (*S. rubida* and *S. albida*, see remarks 7 and 8).

(3) : Originally, *Syndesmis franciscana* was described as *Syndisyrix franciscanus* (LEHMAN, 1946; SHINN, 1980), but was also mentioned as *Syndesmis franciscanus* (GIESE, 1958) and *Syndesmus franciscus* (BERGER & PROFANT, 1961). We regard *Syndesmis franciscana* as the correct name of this species, the genus name being feminine.

(4) : *Syndesmis antillarum* Stunkard & Corliss, 1951 was sometimes confused with *S. franciscana* (WESTBLAD, 1953; BARNES, 1969). Before its description, the species was also mentioned as *Syndesmus* sp. (POWERS, 1935). See also remark 9 about *S. collongistyla* Hertel et al., 1990.

(5) : *Syndesmis dendrastrorum* Stunkard & Corliss, 1951, *S. atriovillosa* Westblad, 1953 and *S. pallida* Hickman, 1956 were previously attributed to *Marcusella* (WESTBLAD, 1953; HICKMAN, 1956), a genus synonymized with *Syndesmis* by CANNON (1982). The first species is reported as *S. dendrastomum* in JANGOUX (1990).

(6) : *Syndesmis compacta* Komschlies & Vande Vusse, 1980, *S. mammilata* Komschlies & Vande Vusse, 1980, *S. philippinensis* Komschlies & Vande Vusse, 1980 and *S. alcalai* Komschlies & Vande Vusse, 1980 are treated here as valid despite of their incomplete species descriptions. The second species is reported as *S. mammillata* in JANGOUX (1990).

(7) : For a long time, *Syndesmis albida* and *S. rubida* were confused with *Syndesmis echinorum* (BAREL & KRAMERS, 1970, 1977), also reported as *Syndesmus echinorum* (SHIPLEY, 1901). It is impossible to know if WESTBLAD (1926), BRUCE et al. (1963) and LYONS (1973) observed *S. rubida* or *S. albida*. See also remark 8.

^a This list updates previous reviews and reattributes outdated taxonomic names used in original works to the valid species. Superscript numbers following species names refer to notes in Results and Discussion.

Umagillid species : Some species have been described as *Syndisyrix* or *Marcusella* by various authors but, without complete reappraisal of the group, validity of these genera is still matter of discussion. All species are thus attributed to the genus *Syndesmis*. Species are presented in the chronological order of their description.

Sea urchin host : Species in bold are irregular sea urchins. Synonyms : *Echinus esculentus* is also known in the literature on umagillids as *Echinus sphaera*; *Paracentrotus lividus* as *Toxopneustes lividus* or *Strongylocentrotus lividus*; *Strongylocentrotus pallidus* as *S. echinoides*. Locality : BS Barents Sea; C Caribbeans; EA East Atlantic; EP East Pacific; IO Indian Ocean; M : Mediterranean Sea; WP West Pacific.

References : "/" separates the authors having studied the species in the different localities mentioned. "?" is used when the species mentioned by the author is not clearly identified but probably belongs to the one named in this list. Authors in bold are the authority to which to refer for the original description of the species.

TABLE 1
List of the umagillids infesting sea urchins, with details about host species infested, localities and references.^a

Umagillid species	Sea urchin host	Locality	References
<i>Syndesmis echinorum</i> ⁽¹⁾	<i>Paracentrotus lividus</i>	M: Banyuls (France) / Marseille (France) / EA: Roscoff (France) / Galicia (Spain)	FRANCOIS, 1886; KOZLOFF & WESTERVELT, 1987; Pers. obs. / BRIOT, 1906 / CHERBONNIER, 1951; BAREL & KRAMERS, 1970?; Pers. obs. / LAMAS SECO & RODRIGUEZ BABIO, 1978; CHERIAN et al., 1992; Pers. obs. / RUSSO, 1895; WAHL, 1909 / VON GRAFF, 1903 / BRIOT, 1906 / KOZLOFF & WESTERVELT, 1987; Pers. obs.
	<i>Sphaerechinus granularis</i>	M: Naples (Italy) / Trieste (Italy), Umag (Croatia) / Marseille (France) / Banyuls (France)	VON GRAFF, 1903
	<i>Strongylocentrotus droebachiensis</i> ? ⁽²⁾	BS: Alexandrowsk (Russia)	LAMAS SECO & RODRIGUEZ BABIO, 1978
	<i>Psammechinus militaris</i>	EA: Galicia (Spain)	LEHMAN, 1946; GESE, 1958; JENNINGS & METTRICK, 1968; BARNES, 1969; METTRICK & JENNINGS, 1969; HERTEL et al., 1990
<i>Syndesmis franciscana</i> ⁽³⁾	<i>Strongylocentrotus franciscanus</i>	EP: California	LEHMAN, 1946; GESE, 1958; JENNINGS & METTRICK, 1968; BARNES, 1969; METTRICK & JENNINGS, 1969; HERTEL et al., 1990
	<i>Strongylocentrotus pallidus</i>	EP: Washington State	ORHIEL, 1952; SHINN, 1980?, 1981, 1983b, 1984; SHINN & CLONEY, 1986
	<i>Strongylocentrotus droebachiensis</i>	EP: Washington State	SHINN, 1980?, 1981, 1983b, 1984; SHINN & CLONEY, 1986; WESTERVELT & KOZLOFF, 1992; SHINN, 1993
	<i>Strongylocentrotus purpuratus</i>	EP: California	LEHMAN, 1946?; GESE, 1958; BARNES, 1969; METTRICK & JENNINGS, 1969; METTRICK & BODDINGTON, 1972; HOLT & METTRICK, 1975; HERTEL et al., 1990
	<i>Lytechinus anamesus</i>	EP: California	BARNES, 1969; HERTEL et al., 1990
	<i>Diadema antillarum</i>	C: Dry Tortugas, Florida / Bermuda / Unknown	POWERS, 1935; STUNKARD & CORLISS, 1951 / SNYDER, 1980 / WESTBLAD, 1953
<i>Syndesmis antillarum</i> ⁽⁴⁾		EP: California / Washington State	STUNKARD & CORLISS, 1950, 1951; SMITH, 1973 / ORHIEL, 1952; SHINN, 1981, 1984, 1988
<i>Syndesmis dendrostrorum</i> ⁽⁵⁾	<i>Dendroaster excentricus</i>	EA: Plymouth (UK)	WESTBLAD, 1953
<i>Syndesmis atrovillosa</i> ⁽⁵⁾	<i>Spatangus purpureus</i>	WP: Tasmania / Sydney	HICKMAN, 1956 / MCGRAE, 1959?; ROHDE & WATSON, 1988
<i>Syndesmis punicea</i>	<i>Helicodidaris erythrogramma</i>	WP: Tasmania	HICKMAN, 1956
	<i>Amblypneustes ovum</i>	WP: Tasmania	HICKMAN, 1956
<i>Syndesmis pallida</i> ⁽⁵⁾	<i>Echinocardium cordatum</i>	IO: Madagascar / WP: Philippines	HYMAN, 1960 / KOMSCHLIES & VANDE VUSSE, 1980a
<i>Syndesmis glambulosa</i>	<i>Diadema setosum</i>	WP: Philippines / IO: Seychelles	KOMSCHLIES & VANDE VUSSE, 1980a / MARTENS & DE CLERCK, 1994
	<i>Echinothrix calamaris</i>	C: Saint-Barthélemy	MARCUS, 1968; HERTEL et al., 1990; HERTEL & DUSZYNSKI, 1991
<i>Syndesmis evelinae</i>	Unknown, <i>Echinometra lucunter</i>	WP: Philippines	KOMSCHLIES & VANDE VUSSE, 1980a
<i>Syndesmis compacta</i> ⁽⁶⁾	<i>Echinometra oblonga</i>	WP: Philippines	KOMSCHLIES & VANDE VUSSE, 1980b
<i>Syndesmis mammilata</i> ⁽⁶⁾	<i>Echinometra oblonga</i>	WP: Philippines	KOMSCHLIES & VANDE VUSSE, 1980b
<i>Syndesmis philippinensis</i> ⁽⁶⁾	<i>Echinometra oblonga</i>	WP: Philippines	KOMSCHLIES & VANDE VUSSE, 1980b
<i>Syndesmis atcala</i> ⁽⁶⁾	<i>Heterocentrotus mammillatus</i>	EP: Banyuls (France)	WESTERVELT & KOZLOFF, 1990; Pers. obs.
<i>Syndesmis aethiopharynx</i>	<i>Paracentrotus lividus</i>	EA: Plymouth (UK), Roscoff (France)	BAREL & KRAMERS, 1970; KOZLOFF & WESTERVELT, 1990
<i>Syndesmis albida</i> ⁽⁷⁾	<i>Echinus esculentus</i>	EA: Plymouth (UK), Roscoff (France)	GEDDES, 1880?; SULLIMAN, 1881?; CUENOT, 1892?, 1900? (Roscoff); SHIPLEY, 1901? (Plymouth); BAREL & KRAMERS, 1970; KOZLOFF & WESTERVELT, 1990
<i>Syndesmis rubida</i> ^{(7), (8)}	<i>Echinus esculentus</i>	C: Saint-Barthélemy, Jamaica	HERTEL et al., 1990
	<i>Echinometra lucunter</i>	C: Jamaica	NAPPI & CRAWFORD, 1984; HERTEL et al., 1990
	<i>Echinometra viridis</i>	IO: Jamaica / Puerto Rico / Haiti	JENNINGS & METTRICK, 1968; METTRICK & JENNINGS, 1969; NAPPI & CRAWFORD, 1984; ALLISON et al., 1987; UBELAKER et al., 1988; HERTEL et al., 1990; VASS & NAPPI, 1998 / HYMAN, 1960; JONES & CANTON, 1970 / JONES & CANTON, 1970
	<i>Lytechinus variegatus</i>	C: Jamaica	HERTEL et al., 1990
	<i>Lytechinus williamsi</i>	C: Puerto Rico	JONES & CANTON, 1970
	<i>Triplaneustes ventricosus</i>	EP: Washington State	ORHIEL, 1952?; SHINN, 1981?, 1983b?, 1984?; WESTERVELT & KOZLOFF, 1992
<i>Syndesmis inconspicua</i> ⁽¹⁰⁾	<i>Strongylocentrotus droebachiensis</i>	EP: Washington State	SHINN, 1981?, 1983b?, 1984?
<i>Syndesmis neglecta</i> ⁽¹¹⁾	<i>Strongylocentrotus pallidus</i> ?	EP: Washington State / California	BERGER & PROFANT, 1961?; WESTERVELT & KOZLOFF, 1992 / GESE, 1958; BOOLOOTIAN et al., 1959?; HYMAN, 1960
<i>Syndesmis longicanalis</i>	<i>Alloccentrotus fragilis</i>	IO: Kenya	MOENS & MARTENS, 1992; GEVAERTS et al., 1993
	<i>Triplaneustes granilla</i>	IO: Kenya	MOENS & MARTENS, 1992; GEVAERTS et al., 1994; GEVAERTS et al., 1995
	<i>Toxopneustes pileolus</i>	IO: Kenya	MOENS & MARTENS, 1992; GEVAERTS et al., 1993; MOENS et al., 1994; GEVAERTS et al., 1995
<i>Syndesmis camoni</i>	<i>Ammotrochus arachnoides</i>	IO: Western Australia	JONDELIUS, 1996
<i>Syndesmis echinicauda</i> ⁽¹²⁾	<i>Echinus acutus</i>	M: Banyuls, Port-Vendres (France)	FRANCOIS, 1886; KOZLOFF, 1997
<i>Syndesmis</i> sp 1	<i>Echinus chloroticus</i>	WP: New Zealand	MCGRAE, 1959
<i>Syndesmis</i> sp 2 ⁽¹³⁾	<i>Strongylocentrotus franciscanus</i>	EP: California	LEHMAN, 1946; BARNES, 1969; METTRICK & JENNINGS, 1969
	<i>Strongylocentrotus purpuratus</i>	EP: California	BARNES, 1969; METTRICK & JENNINGS, 1969
	<i>Lytechinus anamesus</i>	EP: California	BARNES, 1969; METTRICK & JENNINGS, 1969
<i>Syndesmis</i> sp 3	<i>Diadema savigny</i>	IO: Kenya	MARTENS & DE CLERCK, 1994

(8) : *Syndesmis rubida* can be distinguished from *S. albidia* by the colour : the former is brown red while the latter is pale pink to white (KOZLOFF & WESTERVELT, 1990). According to their descriptions, GEDDES (1880), SILLIMAN (1881), CUÉNOT (1892, 1900) and SHIPLEY (1901) likely observed *S. rubida*.

(9) : Prior to its description, *Syndesmis collongistyla* was confused with *S. franciscana* and *S. antillarum* (JENNINGS & METTRICK, 1968; JONES & CANTON, 1970; NAPPI & CRAWFORD, 1984; ALLISON et al., 1987).

(10) : *Syndesmis inconspicua* was mentioned as *S. "echinorum"* in older literature (ORIHÉL, 1952; SHINN, 1981, 1983b, 1984). This species was reported in *Strongylocentrotus droebachiensis* and *S. pallidus* (Sars, 1871) but not described until WESTERVELT & KOZLOFF (1992). These authors investigated specimens from *S. droebachiensis* only.

(11) : *Syndesmis neglecta* Westervelt & Kozloff, 1992 was previously listed as *Syndesmis franciscanus* (see GIESE, 1958), *Syndesmus franciscanus* (BOOLOOTIAN et al., 1959), *Syndesmis franciscana* (HYMAN, 1960) and *Syndesmus franciscus* (BERGER & PROFANT, 1961).

(12) : Specimens of *Syndesmis echiniacuti* Kozloff, 1997 were perhaps also used by FRANÇOIS (1886) in the original description of *Syndesmis echinorum* (see KOZLOFF & WESTERVELT, 1987 and KOZLOFF, 1997).

(13) : *Syndesmis* sp. 2 could be *S. inconspicua*. BARNES (1969) stated its close similarity with *S. echinorum*.

We hope this survey will be a useful tool for further research on this ecologically important group. Recent investigations have demonstrated that literature accounts of intraspecific variation in *Syndesmis* species found in distinct hosts or from distant locations should be treated sceptically, as some of these cases may in fact be instances of interspecific variation.

Re-evaluation of the current species, and the description of new species should be based on a thorough morphological study of the taxon as a whole (including the re-examination of all material deposited in collections), combined with molecular data. Unfortunately, confusions and misidentifications have persisted for years because some works were based on material of poor quality and sometimes lacked the study of histological sections. In some cases, researchers described a species, but failed to deposit type material in a collection. As a consequence, comparison between species can be very difficult, if not impossible, unless new material is collected. Yet, with the updated checklist presented here, future researchers will have immediate access to literature and a complete overview of *Syndesmis* species and their corresponding host(s) and geographical distribution.

ACKNOWLEDGEMENTS

We thank Prof. Michel Jangoux (Université Libre de Bruxelles) for valuable comments on the manuscript and Mr. Herwig Rammer (ULB) for help with translation of German literature. Gilles Doignon's work was supported by grants of the Fonds pour la Recherche dans l'Industrie et l'Agriculture (F.R.I.A.), of the Fonds National de la Recherche Scientifique (F.N.R.S.) de Belgique and of the Van Buuren Foundation. This paper is a contribution of the "Centre Interuniversitaire de Biol-

ogie Marine" (CIBIM), Belgium and to project G.0235.02 financed by the FWO-VL (National Fund for Scientific Research-Flanders, Belgium).

REFERENCES

- ALLISON, V.F., J.E. UBELAKER, E.M. MORGAN & D.W. DUSZYNSKI (1987). Ultrastructure of the epidermis of a rhabdocoel turbellarian *Syndisyrix* sp. *Parasitol. Res.*, 74 : 133-141.
- BAREL, C.D.N. & P.G.N. KRAMERS (1970). Notes on associates of echinoderms from Plymouth and the coast of Brittany. *Proc. Kon. Ned. Akad. Wet.*, 73 : 159-170.
- BAREL, C.D.N. & P.G.N. KRAMERS (1977). A survey of the echinoderm associates of the north-east atlantic area. *Zool. Verh., Leiden*, 156 : 1-159.
- BARNES, A.T. (1969). Two endoparasitic turbellarians from California echinoids. Ph. D. Thesis, University of California, Santa Barbara, 188 pp.
- BERGER, J. & R.J. PROFANT (1961). The entocommensal ciliate fauna of the pink sea urchin, *Allocentrotus fragilis*. *J. Parasit.*, 47 : 417-418.
- BOOLOOTIAN, R.A., A.C. GIESE, J.S. TUCKER & A. FARMANFARMALIAN (1959). A contribution to the biology of a deep sea echinoid, *Allocentrotus fragilis* (Jackson). *Biol. Bull.*, 116 (3) : 362-372.
- BRAUN, M. (1889). Ueber parasitische Strudelwürmer. *Zentbl. Bakt. ParasitK.*, 5 : 41-44.
- BRIOT, A. (1906). Sur les turbellariés parasites des oursins (*Syndesmis echinorum* François). *C. R. Soc. Biol.*, 60 : 1158-1159.
- BRUCE, J.R., J.S. COLMAN & N.S. JONES (1963). *Marine fauna of the Isle of Man and its surrounding areas*. Liverpool University Press, Liverpool.
- CANNON, L.R.G. (1982). Endosymbiotic Umagillids (Turbellaria) from holothurians of the Great Barrier Reef. *Zool. Scr.*, 11 (3) : 173-188.
- CANNON, L.R.G. (1987). Two new rhabdocoel turbellarians, *Umagilla pacifica* sp. n. and *U. karlingi* sp. n. (Umagillidae), endosymbiotic with holothurians (Echinodermata) from the Great Barrier Reef; and a discussion of sclerotic structures in the female system of the Umagillidae. *Zool. Scr.*, 16 (4) : 297-303.
- CANNON, L.R.G. (1990). *Anoplodium* (Rhabdocoela, Platyhelminthes) endosymbiotes of sea cucumbers from Australia and New Zealand. *Zool. Scr.*, 19 (4) : 395-402.
- CHERBONNIER, G. (1951). Inventaire de la faune marine de Roscoff, Echinodermes. *Trav. Stat. Biol. Roscoff*, 4 (15) : 1-15.
- CIFRIAN, B., P. GARCIA-CORRALES & S. MARTINEZ-ALOS (1992). Intracytoplasmic ciliary elements in epidermal cells of *Syndesmis echinorum* and *Paravortex cardii* (Platyhelminthes, Dalyellioida). *J. Morphol.*, 213 : 147-157.
- CUÉNOT, L. (1892). Commensaux et parasites des échinodermes (deuxième note). *Rev. Biol. Nord France*, 5 : 1-23.
- CUÉNOT, L. (1900). La comatule, l'oursin. In : BOUTAN (ed), *Zoologie descriptive (Vol. 1)*, pp 265-299.
- FRANÇOIS, P. (1886). Sur le *Syndesmis*, nouveau type de Turbellariés décrit par M. W.-A. Silliman. *C. R. Acad. Sci.*, 103 : 752-754.
- GEDDES, P. (1880). Observations sur le fluide périsvécéral des oursins. *Arch. Zool. Exp. Gén.*, 8 : 483-496.
- GEVAERTS, H., J.B. MOENS, E.E. MARTENS & E.R. SCHOCKAERT (1993). First data on the ultrastructure of the bursal valve of *Syndesmis longicanalis* from sea urchins from the Kenyan coast. Paper presented at the VIIth International Symposium on Biology of Turbellaria, Turku (Finland).
- GEVAERTS, H., J.B. MOENS, E.E. MARTENS & E.R. SCHOCKAERT (1995). Hard parts in the female system of *Syndesmis longicanalis* (Platyhelminthes, Rhabdocoela, Umagillidae) are

- basement membrane derivatives. *Inv. Biol.*, 114 (4) : 279-284.
- GEISE, A.C. (1958). Incidence of *Syndesmis* in the gut of two species of sea urchins. *Anat. Rec.*, 132 : 441-442.
- GRAFF, L. VON (1903). Die Turbellarien als Parasiten und Wirte. Festschr. Universität, Graz.
- HERTEL, L.A. & D.W. DUSZYNSKI (1991). *Syndisyrix evelinae* (Marcus, 1968) n. comb., from the rock-boring urchin, *Echinometra lucunter*, from St. Barthélemy. *J. Parasit.*, 77 (4) : 638-639.
- HERTEL, L.A., D.W. DUSZYNSKI & J.E. UBELAKER (1990). Turbellarians (Umagillidae) from Caribbean urchins with a description of *Syndisyrix collongistyla* n. sp. *Trans. Am. Micr. Soc.*, 109 (3) : 273-281.
- HICKMAN, V.V. (1956). Parasitic Turbellaria from Tasmanian Echinoidea. *Pap. Proc. R. Soc. Tasmania*, 90 : 169-181.
- HOLT, P.A. & D.F. METTRICK (1975). Ultrastructural studies of the epidermis and gastrodermis of *Syndesmis franciscana* (Turbellaria : Rhabdocoela). *Can. J. Zool.*, 53 : 536-549.
- HYMAN, L.H. (1960). New and known umagillid rhabdocoels from echinoderms. *Am. Mus. Novit.*, 1984 : 1-14.
- JANGOUX, M. (1990). Diseases of Echinodermata. In: KINNE (ed), *Diseases of Marine Animals (Vol. 3)*, Biologischen Anstalt Helgoland, Hamburg : 439-567.
- JENNINGS, J.B. (1971). Parasitism and commensalism in the Turbellaria. *Adv. Parasitol.*, 9 : 1-32.
- JENNINGS, J.B. & D.F. METTRICK (1968). Observations on the ecology, morphology and nutrition of the rhabdocoel turbellarian *Syndesmis franciscana* (Lehman, 1946) in Jamaica. *Carib. J. Sci.*, 8 (1-2) : 57-69.
- JONDELIUS, U. (1996). Three echinoderm inhabiting flatworms (Platyhelminthes, Rhabdocoela) from Western Australia. *Belg. J. Zool.*, 126 (1) : 37-48.
- JONES, I. & C.E. CANTON (1970). Additional observations on the distribution of *Syndesmis franciscana* in the Caribbean. *Carib. J. Sci.*, 10 (1-2) : 71-72.
- KOMSCHLIES, K.L. & F.J. VANDE VUSSE (1980a). *Syndesmis compacta* sp. n. and redescription of *S. glandulosa* Hyman 1960 (Turbellaria : Umagillidae) from Philippine sea urchins. *J. Parasit.*, 66 (4) : 664-666.
- KOMSCHLIES, K.L. & F.J. VANDE VUSSE (1980b). Three new species of *Syndesmis* Silliman 1881 (Turbellaria : Umagillidae) from Philippine sea urchins. *J. Parasit.*, 66 (4) : 659-663.
- KOZLOFF, E.N. (1997). *Syndesmis echiniacuti* sp. nov. (Turbellaria : Neorhabdocoela : Umagillidae), from the Mediterranean sea urchin *Echinus acutus*. *Cah. Biol. Mar.*, 38 : 97-102.
- KOZLOFF, E.N. & C.A. JR WESTERVELT (1987). Redescription of *Syndesmis echinorum* François, 1886 (Turbellaria : Neorhabdocoela : Umagillidae), with comments on distinctions between *Syndesmis* and *Syndisyrix*. *J. Parasit.*, 73 (1) : 184-193.
- KOZLOFF, E.N. & C.A. JR WESTERVELT (1990). *Syndesmis rubida* sp. nov. and *S. albida* sp. nov. (Turbellaria : Neorhabdocoela : Umagillidae) from the sea urchin *Echinus esculentus*. *Cah. Biol. Mar.*, 31 : 323-332.
- LAMAS SECO, A. & C. RODRIGUEZ BABIO (1978). Estudio monografico de *Syndesmis echinorum* François 1886 (Turbellaria, Rhabdocoela) endoparasito de equinoideos del litoral gallego. *Rev. Iber. Parasitol.*, 38 (1-2) : 165-192.
- LEHMAN, H.E. (1946). A histological study of *Syndisyrix franciscanus*, gen. et sp. nov., an endoparasitic rhabdocoel of the sea urchin, *Strongylocentrotus franciscanus*. *Biol. Bull.*, 91 : 295-311.
- LYONS, K.M. (1973). The epidermis and sense organs of the Monogenea and some related groups. *Adv. Parasitol.*, 9 : 193-232.
- MARCUS, E. (1949). Turbellaria Brasileiros (7). *Bol. Fac. Filos. Ciênc. Let. Univ. Sao Paulo*, 14 : 7-156.
- MARCUS, E. (1968). A new *Syndesmis* from Saint-Barthélemy, Lesser Antilles (Neorhabdocoela). *Stud. Fauna Curaçao and Other Carib. Isls.*, 25 : 134-139.
- MARTENS, E.E. & G.G. DE CLERCK (1994). Interstitial and parasitic platyhelminthes from the coast of the Seychelles. *Oceanic reefs of the Seychelles, Netherlands Indian Ocean Programme*, 2 : 1-10.
- MCRAE, A. (1959). *Evechinus chloroticus* (Val.), an endemic New Zealand echinoid. *Trans. R. Soc. N. Z.*, 86 : 205-267.
- MEIXNER, J. (1926). Beitrag zur Morphologie und zum System der Turbellaria-Rhabdocoela : II. Über *Typhlorhynchus nanus* Laidlaw und die parasitischen Rhabdocölen nebst Nachträgen zu den Calyptorhynchia. *Z. Morphol. Ökol. Tiere*, 5 : 527-624.
- METTRICK, D.F. & M.J. BODDINGTON (1972). Amino acid pools of *Syndesmis franciscana* (Turbellaria : Platyhelminthes) and of host coelomic fluid. *Can. J. Zool.*, 50 (4) : 411-413.
- METTRICK, D.F. & J.B. JENNINGS (1969). Nutrition and chemical composition of the rhabdocoel turbellarian *Syndesmis franciscana*, with notes on the taxonomy of *S. antillarum*. *J. Fish. Res. Bd. Can.*, 26 : 2669-2679.
- MOENS, J.B. & E.E. MARTENS (1992). *Syndesmis longicanalis* sp. nov., an umagillid turbellarian from sea urchins from the Kenyan coast. Paper presented at Dodonea Symposium, Ghent (Belgium).
- MOENS, J.B., E.E. MARTENS & E.R. SCHOCKAERT (1994). *Syndesmis longicanalis* sp. nov., an umagillid turbellarian (Platyhelminthes) from echinoids from the Kenyan coast. *Belg. J. Zool.*, 124 (2) : 105-114.
- NAPPI, A.J. & J.A. CRAWFORD (1984). The occurrence and distribution of a syndesmid (Turbellaria : Umagillidae) in Jamaican sea urchins. *J. Parasit.*, 70 (4) : 595-597.
- ORIHIEL, T.C. (1952). Entocommensal rhabdocoels from echinoids of Puget Sound. Master of Science Thesis, University of Washington, Seattle.
- POWERS, P.B.A. (1935). Studies on the ciliates of sea-urchins. *Papers Tortugas Lab.*, 29 : 295-326.
- ROHDE, K. & N. WATSON (1988). Ultrastructure of epidermis and sperm of the turbellarian *Syndisyrix punicea* (Hickman, 1956) (Rhabdocoela : Umagillidae). *Aust. J. Zool.*, 36 : 131-139.
- RUSSO, A. (1895). Sulla morfologia del *Syndesmis echinorum* François. *Ric. Lab. Anat. Norm. Univ. Roma*, 4 : 43-68.
- SHINN, G.L. (1980). Reproduction of *Syndisyrix franciscanus*, a flatworm symbiont of sea urchins. *Am. Zool.*, 20 (4) : 892.
- SHINN, G.L. (1981). The diet of three species of umagillid neorhabdocoel turbellarians inhabiting the intestine of echinoids. *Hydrobiologia*, 84 : 155-162.
- SHINN, G.L. (1983a). *Anoplodium hymanae* sp. n., an umagillid turbellarian from the coelom of *Stichopus californicus*, a Northeast Pacific holothurian. *Can. J. Zool.*, 61 (4) : 750-760.
- SHINN, G.L. (1983b). The life history of *Syndisyrix franciscanus*, a symbiotic turbellarian from the intestine of echinoids, with observations on the mechanism of hatching. *Ophelia*, 22 (1) : 57-79.
- SHINN, G.L. (1984). Symbiotic flatworms : reproductive biology and life cycles of Umagillid turbellarians infesting echinoderms. Ph. D. Thesis, University of Washington, Seattle, 306 pp.
- SHINN, G.L. (1985a). Reproduction of *Anoplodium hymanae*, a turbellarian flatworm (Neorhabdocoela, Umagillidae) inhabiting the coelom of sea cucumbers; production of egg capsules, and escape of infective stages without evisceration of the host. *Biol. Bull.*, 169 : 182-198.
- SHINN, G.L. (1985b). Infection of new hosts by *Anoplodium hymanae*, a turbellarian flatworm (Neorhabdocoela, Umagillidae) inhabiting the coelom of the sea cucumber *Stichopus californicus*. *Biol. Bull.*, 169 : 199-214.

- SHINN, G.L. (1986a). Life history and function of the secondary uterus of *Wahlia pulchella*, an umagillid turbellarian from the intestine of a northeastern pacific sea cucumber (*Stichopus californicus*). *Ophelia*, 25 (2) : 59-74.
- SHINN, G.L. (1986b). Spontaneous hatching of *Fallacohospes inchoatus*, an umagillid flatworm from the northeastern Pacific crinoid *Florometra serratissima*. *Can. J. Zool.*, 64 : 2068-2071.
- SHINN, G.L. (1987). Two new species of umagillid flatworms from the 20-rayed antarctic crinoid *Promachocrinus kerguelensis*. *Can. J. Zool.*, 65 : 1001-1009.
- SHINN, G.L. (1988). Contrasting modes of reproduction in the polymorphic symbiotic flatworm *Syndesmis dendrastrorum* (Turbellaria : Neorhabdocoela : Umagillidae). *Ophelia*, 29 (1) : 59-70.
- SHINN, G.L. (1993). Formation of egg capsules by flatworms (phylum Platyhelminthes). *Trans. Am. Micr. Soc.*, 112 (1) : 18-34.
- SHINN, G.L. & R.A. CLONEY (1986). Egg capsules of a parasitic turbellarian flatworm : ultrastructure of hatching sutures. *J. Morphol.*, 188 : 15-28.
- SHIPLEY, A.E. (1901). On some parasites found in *Echinus esculentus*, L. *Q. J. Sci.*, 44 : 281-290.
- SILLIMAN, W.-A. (1881). Sur un nouveau type de Turbellariés. *C. R. Acad. Sci.*, 93 : 1087-1089.
- SMITH, A.B. (2003). *The Echinoid Directory*. World Wide Web electronic publication. <http://www.nhm.ac.uk/palaeontology/echinoids> [accessed 24 January 2005].
- SMITH, N.S. (1973). A new description of *Syndesmis dendrastrorum* (Platyhelminthes, Turbellaria), an intestinal rhabdocoel inhabiting the sand dollar *Dendraster excentricus*. *Biol. Bull.*, 145 : 598-606.
- SNYDER, R.D. (1980). Commensal turbellarians from Bermuda holothurians. *Can. J. Zool.*, 58 : 1741-1744.
- STUNKARD, H.W. & J.O. CORLISS (1950). Parasitic Turbellarians from Echinoderms. *J. Parasit.*, 36 (suppl.) : 31.
- STUNKARD, H.W. & J.O. CORLISS (1951). New species of *Syndesmis* and a revision of the family Umagillidae Wahl, 1910 (Turbellaria : Rhabdocoela). *Biol. Bull.*, 101 : 319-334.
- UBELAKER, J.E., V.F. ALLISON, J.E. RICKER & D.W. DUSZYNSKI (1988). The ultrastructure of cement glands in a rhabdocoel turbellarian *Syndisyrix* sp. *Parasitol. Res.*, 74 : 216-220.
- VASS, E. & A.J. NAPPI (1998). Occurrence and distribution of the rhabdocoel flatworm *Syndisyrix collongistyla* in the sea urchins *Lytechinus variegatus* and *Tripneustes ventricosus* in Discovery Bay, Jamaica. *Carib. J. Sci.*, 34 (1-2) : 137-141.
- WAHL, B. (1909). Untersuchungen über den Bau der parasitischen Turbellarien aus der Familie der Dalyelliiden (Vorticiden). II. Die genera *Umagilla* und *Syndesmis*. *Sber. Akad. Wiss. Wien*, 118 : 943-965.
- WESTBLAD, E. (1926). Parasitische Turbellarien von der Westküste Skandinaviens. *Zool. Anz.*, 68 : 212-216.
- WESTBLAD, E. (1953). New Turbellaria parasites in Echinoderms. *Ark. Zool.*, 2 (5) : 269-288.
- WESTERVELT, C.A. Jr (1981). *Collastoma kozloffi* sp. n., a neorhabdocoel turbellarian from the intestine of the sipunculan *Themiste dyscrita*. *J. Parasit.*, 67 (4) : 574-577.
- WESTERVELT, C.A. Jr & E.N. KOZLOFF (1990). *Syndesmis aethopharynx* sp. nov. (Turbellaria : Neorhabdocoela : Umagillidae), from the sea urchin *Paracentrotus lividus*, with notes on a probable third species from this host. *Cah. Biol. Mar.*, 31 : 431-437.
- WESTERVELT, C.A. Jr & E.N. KOZLOFF (1992). Two new species of *Syndesmis* (Turbellaria : Neorhabdocoela : Umagillidae) from the sea urchins *Strongylocentrotus droebachiensis* and *Allocentrotus fragilis*. *Cah. Biol. Mar.*, 33 : 115-124.

Received : February 1, 2005

Accepted : October 14, 2005