

BIOGEOCHEMICAL CYCLE IN A COCCOLITHOPHORID BLOOM

Kim Suykens, Bruno Delille and Alberto V. Borges

Unité d'océanographie Chimique, Université de Liège, Sart Tilman, Belgium

E-mail: kim.suykens@ulq.ac.be

The biogeochemical properties of an extensive bloom of the coccolithophore, *Emiliana huxleyi*, at the shelf break in the northern Gulf of Biscay was investigated in June 2006. Total Alkalinity (TA) values in the water column showed strong non-conservative behaviour indicative of the impact of calcification, with the highest TA anomalies (up to $26\mu\text{mol.kg}^{-1}$) in the high reflectance coccolith patch. Partial pressure of CO_2 (pCO_2) values ranged from 250 to $338\mu\text{atm}$ and the area was found to act as a sink for atmospheric CO_2 . Overall, $\text{pCO}_2@13^\circ\text{C}$ (pCO_2 normalized at a constant temperature of 13°C) in the water column was negatively related to TA anomalies in agreement with an overall production of CO_2 related to calcification. Hence, the calcifying phase of the *E. huxleyi* bloom decreased the sink of atmospheric pCO_2 , but did not reverse the direction of the flux. Rates of pelagic respiration up to $5.5\text{mmol O}_2.\text{m}^{-3}.\text{d}^{-1}$ suggested a close coupling between primary production and respiration and/or between organic carbon content and respiration. Benthic respiration rates were quite low and varied between 2 and $9\text{mmol O}_2.\text{m}^{-3}.\text{d}^{-1}$, in agreement with the fact that the study area consists of sandy sediments with low organic matter content. Benthic respiration was well correlated to the chlorophyll a content of the top 1cm of the sediment cores. Evidence was found for dissolution of CaCO_3 due to the acidification of superficial sediments in relation to the production of CO_2 and the oxidation of H_2S in the oxic layers.