



Promoting the Science of Ecology

Chronic Exposure of Blue Crabs, *Callinectes Sapidus*, To Sublethal Concentrations of DDT

Author(s): Jack I. Lowe

Source: *Ecology*, Vol. 46, No. 6 (Nov., 1965), pp. 899-900

Published by: Ecological Society of America

Stable URL: <http://www.jstor.org/stable/1934032>

Accessed: 28/05/2009 06:50

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=esa>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



Ecological Society of America is collaborating with JSTOR to digitize, preserve and extend access to *Ecology*.

<http://www.jstor.org>

and a reduction in the number of predators and competing species. This simplification of the food web, together with the low diversity index, results in marked population fluctuations. In contrast to the large numbers of *Collembola* per sample in Frenchman's Cave, Christiansen, Willson and Tecklin (1961) reported an average of 11 *Collembola* per 1,500 cc sampling unit from Hunter's Cave, Iowa, where the organic content of samples at no time exceeded 5%.

Finally, several species of invertebrates found in Frenchman's Cave are of particular interest since they are most often associated with caves. How such troglodites (facultative cave inhabitants) as the beetle *Quedius (Microsaurus)* sp., the antlike beetle *Brathinus nitidus*, and the cave orb weaver spider *Meta menardi* reached the gypsum caves of Hants County, Nova Scotia, emphasizes our ignorance of the means of dispersal utilized by organisms. Dispersal into Nova Scotia is complicated since the province is a peninsula connected to the continent by an extensive low-lying area presently covered by salt marshes and coniferous bogs. Bleakney (1958) noted that the special environments of this isthmus have limited the dispersal of many invertebrates and vertebrates, including flying forms of both. Obviously troglodites must have means of dispersal that transport them through unfavorable environments to the habitat to which they are physiologically adapted.

ACKNOWLEDGMENTS

The authors are indebted to C. J. Fox, soil zoologist at the Canada Agricultural Research Station, Kentville, N. S., for loaning the battery of Tullgren funnels employed in the study, and for advice and assistance in the techniques of soil sampling. The help of J. Basaraba, Professor of Microbiology, Acadia University, in determination of pH and organic content of cave detritus, is acknowledged. The following specialists kindly identified specimens and also checked our tentative identifications: E. E. Lindquist, Acarina; K. Christiansen and W. R. Richards, *Collembola*; E. C. Becker and T. C. Barr, Jr., Coleoptera; C. D. Dondale, Araneida. The study was supported by Canada National Research Council Grant A-2009

LITERATURE CITED

Babos, S. 1964. Die Zeckenfauna Mitteleuropas. Akademiai Kiado, Budapest.

- Barr, Thomas C., Jr. 1960. Speciation and raiation in cavernicoles. *Am. Midland Naturalist* 64(1): 1-160.
- Bleakney, J. Sherman. 1958. A zoogeographical study of the amphibians and reptiles of Eastern Canada. *Nat. Mus. Canada Bull.* 155.
- Bohnsack, Kurt K. 1954. A study of the forest floor arthropods of an oak hickory woods in Southern Michigan. Unpublished Doctoral Dissertation. Univ. Michigan Library.
- Calder, Dale R. 1964. A study of the soil mesofauna of several different environments. Unpublished report at Canada Agricultural Research Station, Kentville, Nova Scotia.
- Cameron, Austin W. 1958. Mammals of the islands of the Gulf of St. Lawrence. *Nat. Mus. Canada Bull.* 154.
- Christiansen, K. 1964. Bionomics of *Collembola*. *Ann. Rev. Entomol.* 9: 147-178.
- , Mary Willson and Jerry Tecklin. 1961. Preliminary study of the microarthropod ecology of Hunter's Cave. *Bull. Nat. Speleol. Soc.* 23(2): 63-70.
- Crossley, D. A., Jr. and Kurt K. Bohnsack. 1960. Long-term ecological study in the Oak Ridge area: III. The oribatid mite fauna in pine litter. *Ecology* 41: 628-638.
- Dogiel and Efremoff. 1925. Versuch einer quantitativen Untersuchung der Bodenbevölkerung im Fichtenwalde. *Travaux de la société des naturalistes de Leningrad* 55: 97-110.
- Kevan, D. Keith McE. 1962. Soil Animals. H. F. and G. Witherby Ltd., London.
- Lunn, E. T. 1939. The ecology of the forest floor with particular reference to microarthropods. Unpublished doctoral dissertation Northwestern University Library.
- Naglitsch, F. 1962. Untersuchungen über die Collembolenfauna unter Luzernebeständen auf verschiedenen Böden. *Wis. Z. Karl-Marx Univ., Leipzig, Math.—Naturw., Reihe*, 11.
- Prest, Walter Henry. 1912. Report on cave examination in Hants County, N. S. *Proc. and Trans. N. S. Inst. Sci.* 13: 87-94.
- Tullgren, A. H. 1918. Ein sehr einfacher Ausleseapparat für terricole Tierformen. *Z. Angew. Entomol.* 4: 149-150.
- Vandiel, A. 1960. Biospeleologie. La biologie des animaux cavernicoles.

CHRONIC EXPOSURE OF BLUE CRABS, *CALLINECTES SAPIDUS*, TO SUBLETHAL CONCENTRATIONS OF DDT

JACK I. LOWE

U. S. Bureau of Commercial Fisheries Biological Laboratory, Gulf Breeze, Florida

Abstract. Juvenile blue crabs, *Callinectes sapidus*, were reared in flowing seawater containing sublethal concentrations of DDT. Crabs fed, molted, and grew for 9 months in seawater containing 0.25 ppb (micrograms/liter) DDT but could survive only a few days in water containing DDT in excess of 0.5 ppb.

Commercially important marine crustaceans are extremely sensitive to many of the commonly used insecticides (United States Department of the Interior 1963, 1964), but preliminary studies show that blue crabs, *Callinectes sapidus*, will tolerate low concentrations of DDT for long periods.

This note describes the results of an experiment in

which a small number of blue crabs was exposed continuously for 9 months to a sublethal concentration of DDT in flowing seawater. Such chronic exposures are of interest because blue crabs, especially the juveniles, spend much of their lives in shallow estuarine waters which may occasionally be polluted with insecticides.

This experiment was conducted at the Bureau of Com-

mercial Fisheries Biological Laboratory, Gulf Breeze, Florida, between April and December 1964. Facilities are available for conducting bioassays in flowing seawater which is pumped directly from Santa Rosa Sound.

Six juvenile blue crabs averaging 27 mm in carapace width were placed in each of three plastic aquaria with a capacity of approximately 20 liters. Each aquarium was divided into six compartments, 150 × 150 mm, to prevent cannibalism and facilitate the keeping of records on individual crabs. Seawater flowed through each aquarium at a constant rate of 154 liters/hr. Acetone stock solutions of DDT were mixed in aspirator bottles and metered into two of the aquaria at a rate of 0.5 ml per minute to obtain the desired concentration. This constant-flow apparatus was illustrated in detail by Lowe (1964). The crabs in the third aquarium were maintained as controls with only seawater flowing through the aquarium. In preliminary bioassays blue crabs tolerated a significantly higher concentration of acetone in the test solution than was used in this experiment.

The crabs were fed frozen fish throughout the experiment. Pieces of fish flesh of approximately the same size were placed daily in each crab's compartment after the removal of any uneaten food. Additional food in the form of plankton was possibly obtained from the unfiltered seawater.

During preliminary screening tests juvenile blue crabs survived only a few days in flowing seawater containing DDT in excess of 0.5 ppb. Therefore, concentrations of 0.5 and 0.25 ppb were chosen as test solutions for the chronic exposures.

Although only 4 of 18 crabs (22%) survived the entire 9 months of captive life, most appeared to feed, molt, and grow normally. Records of molting and final size of individual crabs are itemized in Table I because of the paucity of this type of information in the literature. There was no difference in behavior between the control crabs and those in the 0.25-ppb DDT solution, but frequent observations of the crabs in the 0.5-ppb solution showed this concentration to be near the threshold of tolerance. Two of the crabs in the 0.5-ppb solution became paralyzed and died after 2 weeks of exposure. These and other crabs dying in the 0.5-ppb solution exhibited typical symptoms of insecticide poisoning (extreme irritability, increased sensitivity to external stimuli, and eventual paralysis) before death.

All crabs surviving the DDT exposure molted as frequently as the controls. One crab that survived the 0.5-ppb exposure and two from the 0.25-ppb concentration molted at least six times, the maximum number of molts for any of the control crabs. The average number of molts during the first 8 weeks of the experiment was 2.5 for the crabs in both the control and 0.25 ppb DDT; no deaths occurred during this period. The crabs surviving the entire 36 weeks of captive life occupied approximately 75% of their individual compartments and had barely enough room to move around. The size variation among

TABLE I. Survival and growth records of individual blue crabs held in captivity

Test solution	Initial carapace width (mm)	Weeks of survival	Total number of molts	Carapace width at death (mm)
Control	25	36	6	121 ^a
	25	22	5	87
	25	20	4	64
	25	16	3	61
	28	12	3	50
	25	11	3	49
0.25 ppb DDT	27	36	7	136 ^a
	27	36	6	135 ^a
	30	29	5	90
	23	26	6	114
	28	17	4	71
	25	12	3	51
0.50 ppb DDT	29	36 ^b	6	114 ^a
	24	20	4	53
	25	8	2	48
	30	7	2	47
	25	2	1	—
	30	2	0	—

^a Crabs alive at end of experiment.

^b 23 weeks in 0.5 ppb followed by 13 weeks in 0.25 ppb DDT.

individuals was as expected. Data on crabs from Chesapeake Bay show that linear measurements of an instar may vary by as much as 50% of the mean dimension (Tyler and Cargo 1963). Two of the crabs surviving 36 weeks of exposure to DDT were females; half-moon-shaped abdomens indicated the ultimate instar.

Any conclusions drawn from this experiment must be made with caution because of the small number of animals used. The data suggest that a natural crab population could exist in estuarine waters chronically polluted with low levels of DDT, but that a sudden slight increase in concentration could be disastrous.

LITERATURE CITED

- Lowe, Jack I. 1964. Chronic exposure of spot, *Leiostomus xanthurus*, to sublethal concentrations of toxaphene in seawater. Trans. Amer. Fish. Soc. 93(4): 396-399.
- Tyler, Albert V., and David G. Cargo. 1963. Size relations of two instars of the blue crab, *Callinectes sapidus*. Chesapeake Science, 4(1): 52-54.
- United States Department of the Interior. 1963. Pesticide-wildlife studies—A review of Fish and Wildlife Service investigations during 1961 and 1962. Circular 167. 109 p.
- United States Department of the Interior. 1964. Pesticide-wildlife studies, 1963—A review of Fish and Wildlife Service investigations during the calendar year. Circular 199. 130 p.